Oral selection: all rows

Oral sampling site OR oral disease. Red: increased; blue: decreased positive-z profile fraction. Numbers are increased minus decreased fractions; max n = largest per-trait study count in each row. Dark border: BH-adjusted q <= 0.05 (0 oral cells); 2 oral cells met nominal Fisher p < 0.01; original 3144-cell family retained. Displayed cells retain source support (>=3 positive-z profiles in either direction and >=2 studies) and require |effect|>0.01. Gray: unavailable, below source support, or at/below the absolute-effect cutoff; it does not indicate nonsignificance. Source tests are profile-level diagnostics.

Aggressive periodontitis | Oral | max n=1
Chronic periodontitis | Oral | max n=1
Dental caries | Oral | max n=2 -0.29
Dental fluorosis | Oral | max n=1
Gingivitis | Oral | max n=2 -0.38 -0.38 -0.25
Halitosis | Oral | max n=1
Oral cavity carcinoma | Oral | max n=1
Oral leukoplakia | Oral | max n=1
Oral mucositis | Oral | max n=3 -0.20 +0.30 -0.07 -0.12
Oral squamous cell carcinoma | Oral | max n=7 +0.22 -0.06 -0.11 +0.12 -0.05
Periapical Periodontitis | Oral | max n=1
Periodontitis | Oral | max n=7 +0.14 -0.06 -0.04 +0.23
Periodontitis,Type II diabetes mellitus | Oral | max n=1
Root caries | Oral | max n=1
Abnormality on pulmonary function testing | Oral | max n=1
Acne | Oral | max n=1
Acute myeloid leukemia | Oral | max n=2
Age | Oral | max n=2
Age,Human papilloma virus infection | Oral | max n=1
Age-related macular degeneration | Oral | max n=1
Aging | Oral | max n=1
Air pollution | Oral | max n=1
Alcohol drinking | Oral | max n=1
Antimicrobial agent | Oral | max n=2
Arthritis | Oral | max n=4 +0.33 +0.67
Atopic asthma | Oral | max n=1
Behcet's syndrome | Oral | max n=1
Breast cancer | Oral | max n=1
Breast ductal carcinoma in situ | Oral | max n=1
CD4-positive T-lymphocyte count | Oral | max n=1
COVID-19 | Oral | max n=2
Cannabis use | Oral | max n=1
Carotid artery intima media thickness | Oral | max n=1
Chronic hepatitis B virus infection | Oral | max n=2 +0.20 +0.20 +0.40 +0.20
Chronic kidney disease | Oral | max n=1
Chronic otitis media | Oral | max n=1
Coinfection,Age | Oral | max n=1
Colorectal adenoma | Oral | max n=1
Colorectal cancer | Oral | max n=5 +0.14 +0.16 +0.71 +0.29 -0.07 -0.16
Diet | Oral | max n=1
Digestive system cancer | Oral | max n=1
Down syndrome | Oral | max n=1
Endometriosis | Oral | max n=1
Esophageal cancer | Oral | max n=3 -0.20 -0.20 -0.20 -0.40
Esophageal squamous cell carcinoma | Oral | max n=3 +0.14 -0.06 +0.23 +0.23
Ethnic group | Oral | max n=1
Gastric cancer | Oral | max n=4 -0.26 +0.03 +0.03 -0.03
Gestational diabetes | Oral | max n=1
Glaucoma | Oral | max n=3 -0.33 +0.33 -0.33
HDL cholesterol | Oral | max n=1
HIV infection | Oral | max n=3 +0.42
HIV infection,Human papilloma virus infection | Oral | max n=1
HIV infection,Oral candidiasis | Oral | max n=2
Head and neck carcinoma | Oral | max n=2 -0.33
Head and neck squamous cell carcinoma | Oral | max n=3 -0.04 -0.16 -0.03 -0.46
Health study participation | Oral | max n=1
Health trait | Oral | max n=1
Household income | Oral | max n=1
Human papilloma virus infection | Oral | max n=3 +0.55 +0.40 -0.10 +0.35
Human papillomavirus | Oral | max n=1
Hyperglycemia | Oral | max n=1
Hypertension | Oral | max n=2 -0.20 -0.20
Hypertriglyceridemia | Oral | max n=1
Hypertrophy | Oral | max n=1
Hypothyroidism | Oral | max n=1
Illness severity status | Oral | max n=1
Iron deficiency anemia | Oral | max n=1
Irritable bowel syndrome | Oral | max n=1
Ischemic stroke | Oral | max n=1
Juvenile idiopathic arthritis | Oral | max n=1
Lifestyle measurement | Oral | max n=1
Lung cancer | Oral | max n=2
Menopause | Oral | max n=1
Mucositis | Oral | max n=1
Nasopharyngeal squamous cell carcinoma | Oral | max n=1
Nicotine dependence,Smoking behavior | Oral | max n=1
Obesity | Oral | max n=2
Open-angle glaucoma | Oral | max n=1
Oral ulcer | Oral | max n=1
Osteoarthritis,Rheumatoid arthritis | Oral | max n=1
Oxygen | Oral | max n=1
Pancreatic carcinoma | Oral | max n=2
Parkinson's disease | Oral | max n=2 +0.40 +0.20 +1.00
Peptic esophagitis | Oral | max n=1
Phenylketonuria | Oral | max n=1
Physical activity | Oral | max n=1
Pituitary Gland Adenoma | Oral | max n=1
Population | Oral | max n=1
Post-COVID-19 disorder | Oral | max n=1
Pregnancy | Oral | max n=1
Premature birth | Oral | max n=1
Primary bacterial infectious disease | Oral | max n=1
Psoriasis | Oral | max n=2 -0.58 -0.75
Respiratory tract infectious disease | Oral | max n=1
Response to allogeneic hematopoietic stem cell transplant | Oral | max n=2
Response to combination chemotherapy | Oral | max n=1
Response to surgery | Oral | max n=1
Rheumatoid arthritis | Oral | max n=9 +0.22 +0.17 -0.17
Risk factor | Oral | max n=1
SARS coronavirus | Oral | max n=1
Sensory perception of bitter taste | Oral | max n=1
Sjogren syndrome | Oral | max n=1
Sleep Disorder | Oral | max n=1
Smoking behavior | Oral | max n=8 +0.13 +0.22 +0.32 -0.12 +0.06
Smoking behaviour measurement | Oral | max n=1
Smoking status measurement | Oral | max n=3 -0.17 +0.67 +0.33 +0.67 -0.50 -0.17
Socioeconomic status | Oral | max n=1
Specimen | Oral | max n=1
Squamous cell carcinoma | Oral | max n=5 +0.10 -0.50 +0.12 -0.05
Systemic lupus erythematosus | Oral | max n=1
Temporomandibular joint disorder | Oral | max n=1
Time | Oral | max n=1
Tongue neoplasm | Oral | max n=1
Treatment | Oral | max n=1
Type II diabetes mellitus | Oral | max n=5 +0.43 +0.63 +0.43 +0.13
Viral load | Oral | max n=1
Waist circumference | Oral | max n=1
Condition | body-site pair
facultatively aerobic
cellulose catabolic process
coccus shaped
sphere shaped
arsenate reduction
moderately halophilic
Six shared traits
1.00
0.75
0.50
0.25
0.00
−0.25
−0.50
−0.75
−1.00
Increased - decreased positive-z profile fraction