

1□:1□ ▼■





Report Blood Glucose

Save






mg/dL























Recheck **Blood Glucose**

Your blood glucose is very low (<60).



Drink a cup of fruit juice



Drink a cup of (non-diet) soda



Take 8 glucose tablets




Eat 30g of other quick sugar

Next Step



















Eat Quick Sugar





Recheck **Blood Glucose**


15:00

Next Step

















mg/dL

Next Step












Next Step





Eat Quick Sugar 2 Wait 15 Minutes

Recheck **Blood Glucose**




mg/dL

Next Step














Your blood glucose (112 mg/dL) is back to normal.

If you plan to eat within 1 hour:

Eat 15g of slow carbs now (6 saltine crackers or 3 graham cracker squares)

If you plan to eat after more than 1 hour:

Eat 30g of slow carbs now (1 peanut butter sandwich)


Finish











Your blood glucose (112 mg/dL) is back to normal.

If you plan to eat within 1 hour:

Eat 15g of slow carbs now (6 saltine crackers or 3 graham cracker squares)

If you plan to eat after more than 1 hour:


Eat 30g of slow carbs now (1 peanut butter sandwich)


Finish











Your blood glucose (112 mg/dL) is back to normal.

If you plan to eat within 1 hour:

Eat 15g of slow carbs now (6 saltine crackers or 3 graham cracker squares)

If you plan to eat after more than 1 hour:

Eat 30g of slow carbs now (1 peanut butter sandwich)

Finish







