

The synthesis and biological evaluation of a library of
autoinducer-antibiotic conjugates

Lois Overvoorde



Sidney Sussex College

University of Cambridge

September 2018

Supervised by Prof. David Spring

This dissertation is submitted for the degree of Doctor of Philosophy

Contents

1	Acknowledgements	2
2	Declaration	4
3	Abstract	5
4	Nomenclature	6
5	Introduction	9
5.1	Antibiotic resistance	9
5.2	Sideromycins	10
5.3	Synthetic siderophore-antibiotic conjugates	13
5.4	Autoinducer-antibiotic conjugates	16
5.4.1	Quorum sensing	16
5.4.1.1	<i>Vibrio fischeri</i>	16
5.4.1.2	<i>Pseudomonas aeruginosa</i>	18
5.4.2	Autoinducers	21
5.4.3	Autoinducer efflux	21
5.4.4	Antibiotics	21
5.4.5	Antibiotic efflux	22
5.4.6	Conjugate efflux and antibiotic action	22
5.5	Autoinducer analogue-ciprofloxacin conjugates	22
6	Aims	25
7	Results and discussion: autoinducer-antibiotic conjugates	26
7.1	Overview	26
7.1.1	Azido autoinducer derivatives	26
7.1.2	Alkynyl antibiotic derivatives	26
7.1.3	Synthesis of the conjugates	27
8	Results and discussion: autoinducer analogue-ciprofloxacin conjugates	27
8.1	Overview	27
8.1.1	Head groups	27
8.1.2	Library construction	28
9	References	30

1 Acknowledgements

Firstly, I would like to thank David Spring for the opportunity to work on this very interesting topic. I would also like to thank Hannah Sore for guidance and support through the later stages of the project, and Eddy Sotelo and Bin Yu for collaboration and useful discussions. Thank you to Mark Eldridge and Suzie Forrest for help with learning biochemical techniques, to Martin Welch for guidance and advice on data interpretation and especially to Tom O'Brien for stepping in to do some of the testing. Thanks also to Matt Pond, Melvyn Orriss, Nic Davies and Naomi Hobbs for help with equipment and glassware and to Jill Vaughan and Tommy Osberger

for proof-reading. Most importantly, I would like to thank Yssy Baker for proof-reading, help, support, advice and encouragement.

2 Declaration

This dissertation describes work carried out in the Department of Chemistry, University of Cambridge under the supervision of Prof. David Spring, and in the Department of Biochemistry, University of Cambridge under the supervision of Dr Martin Welch. This dissertation is the result of my own work and includes nothing that is the outcome of work done in collaboration except as specified in the text. The dissertation does not exceed the word limit specified by the Physics and Chemistry Degree Committee.

Lois Overvoorde September 2018

3 Abstract

Bacterial resistance to antibiotics is becoming a serious global health threat, and the discovery of new, safe and effective antibiotics is required urgently.¹⁻³ A new class of antibiotic, namely siderophore-antibiotic conjugates, has shown promise in initial studies.^{4,5} Siderophores are used by bacteria for iron uptake, and so attaching antibiotics to them allows the antibiotic to be carried across cell membranes. This study investigates conjugates designed using a similar approach, but using bacterial autoinducers⁶ instead of siderophores. Autoinducers are required for coordination of bacterial behaviours and are involved in the control of swarming, virulence factor production and biofilm formation.

The library was synthesised in two halves which were then coupled together using a copper(I)-catalysed azide-alkyne cycloaddition.^{7,8} The autoinducers were functionalised with azide groups and the antibiotics were functionalised with alkynes. The quorum sensing molecules produced by *Pseudomonas aeruginosa* were investigated as it is a significant human pathogen⁹ which displays high resistance to many antibiotics¹⁰ and uses quorum sensing to coordinate its group behaviours.¹¹ Azido analogues of these autoinducers were coupled with alkyne analogues of ciprofloxacin, which was chosen as it is commonly used against *P. aeruginosa*¹² but resistance to it is developing,¹³ and trimethoprim. It was hoped that the autoinducers would aid retention of the antibiotic in the cell, thus potentially increasing its potency or even restoring its efficacy against resistant strains.

swarming
needs
to be
men-
tioned
else-
where

analogues

4 Nomenclature

J	Coupling constant in Hz
m/z	Mass to charge ratio in Daltons
R_f	Retention factor
Ac	Acetate
AIP	Autoinducing peptide
aq.	Aqueous
atm	Atmosphere(s)
BHL	Butyryl homoserine lactone = C ₄ -HSL 19
Boc	<i>tert</i> -Butyloxycarbonyl
Cip	Ciprofloxacin
conc.	Concentrated
COSY	Correlation spectroscopy
d	Day(s)
Da	Daltons
DBU	1,8-Diazabicyclo[5.4.0]undec-7-ene
DIPEA	<i>N,N</i> -Diisopropylethylamine
DMAP	4-Dimethylaminopyridine
DMF	Dimethylformamide
DMP	Dess-Martin periodinane
DMSO	Dimethylsulfoxide
EDC	1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide
eq.	Equivalents
ESI	Electrospray ionization
Et	Ethyl
FT	Fourier transform
h	Hour(s)
HCTL	Homocysteine thiolactone
HHQ	2-Heptylquinolin-4(1H)-one
HMBC	Heteronuclear multiple-bond correlation spectroscopy
HMQC	Heteronuclear multiple-quantum correlation spectroscopy

HOBt 1-Hydroxybenzotriazole
 HPLC High-performance liquid chromatography
 HRMS High resolution mass spectroscopy
 HSL Homoserine lactone
 Hz Hertz
 IR Infrared
 LCMS Liquid chromatography mass spectroscopy
 LCT Liquid chromatography time-of-flight
 lit. Literature value
 M Molar
 m.p. Melting point
 Me Methyl
 MIC Minimum inhibitory concentration
 min Minute(s)
 mol Mole(s)
 Ms Methanesulfonyl
 NMR Nuclear magnetic resonance
 OdDHL *N*-(3-Oxododecanoyl)-homoserine lactone = 3-oxo-C₁₂-HSL **20**
 P.E. Petroleum ether
 PAI-1 *Pseudomonas* autoinducer 1 = 3-oxo-C₁₂-HSL **20**
 PAI-2 *Pseudomonas* autoinducer 2 = C₄-HSL **19**
 Pd/C Palladium on carbon
 PQS *Pseudomonas* Quinolone Signal
 Q-TOF Quadrupole time-of-flight
 r.t. Room temperature
 s Second(s)
 SAM *S*-adenosyl-L-methionine
 SAR Structure activity relationship
 sat. Saturated
 sp. Species
 TBAF Tetrabutylammonium fluoride
 TBDMS *tert*-Butyldimethylsilyl

TEA	Triethylamine
Tf	Trifluoromethanesulfonyl
TFA	Trifluoroacetic acid
THF	Tetrahydrofuran
THPTA	Tris(3-hydroxypropyltriazolylmethyl)amine
TLC	Thin layer chromatography
TMS	Trimethylsilyl
Ts	<i>para</i> -Toluenesulfonyl
UV	Ultraviolet

5 Introduction

5.1 Antibiotic resistance

Antibiotics add, on average, twenty years to a person's life.² However, antibiotic resistance is increasing alarmingly and is now recognised as a major threat to global health.^{1,2} Antibiotic discovery had its heyday in the 1940s to 60s, which saw the discovery of many new classes of antibiotic. Since then, the rate of discovery of new classes has slowed and resistance to existing treatments has increased.

The story of how Alexander Fleming discovered penicillin by accidentally allowing a Petri dish containing *Staphylococcus aureus* to become contaminated with *Penicillium* mould whilst he was on holiday in Suffolk² is well known to many scientists. The initial serendipitous discovery of penicillin occurred in 1928 and was reported in 1929,¹⁴ but it was not until 1943 that the drug was mass produced thanks to the research of Ernst Chain and Howard Florey. Unfortunately, bacterial resistance to penicillin was being found in hospitals by the late 1940s.^{15,16} This alarmingly quick emergence of resistance is a common phenomenon for antibiotics (see Table 1) as bacteria have multiple resistance mechanisms against antibacterial agents. These mechanisms can be broken down into five main categories:^{2,3}

1. The bacterium may inactivate the drug before it can cause damage, for example the hydrolysis of β -lactam antibiotics such as penicillin by β -lactamase enzymes.
2. The bacterium may produce a membrane, cell wall or biofilm which does not allow the drug to pass through, for example biofilm formation may allow bacterial resistance to antibiotics to increase 1000-fold compared with bacteria in suspension culture.¹⁷
3. The bacterium may pump antibacterial molecules out of its cell membrane using efflux pumps, for example the mexAB and mexXY pumps used by *Pseudomonas aeruginosa*.¹⁰
4. Mutations may cause the target of the antibacterial molecule to alter such that the molecule no longer effectively binds the target, for example the alteration of penicillin binding proteins which are involved in the final stages of peptidoglycan biosynthesis in the cell walls of MRSA and other penicillin-resistant bacteria.¹⁸
5. The bacterium may switch to using a metabolic pathway which does not involve the target of the antibacterial molecule, for example sulfonamide resistance may be achieved by taking in folic acid from the environment rather than synthesising it using *para*-aminobenzoic acid - a process which is blocked by sulfonamides.¹⁹

Antibiotic	Introduction	Resistance
Sulfonamides	1930s	1940s
Penicillin	1943	1946
Streptomycin	1943	1959
Chloramphenicol	1947	1959
Tetracycline	1948	1953
Erythromycin	1952	1988
Vancomycin	1956	1988
Methicillin	1960	1961
Ampicillin	1961	1973
Trimethoprim	1962	1972
Cephalosporins	1960s	late 1960s
Ciprofloxacin	1987	1988
Linezolid	2000	1997
Daptomycin	2003	2005

Table 1: A timeline of when various antibiotics were first introduced and when resistance to them first appeared.^{20–25}

The current pipeline of new antibiotics is widely thought to be worryingly inadequate.^{?, 26} Significant changes in how we use the antibiotics we already have, as well as investments in the discovery of new ones, are required. Antibiotics currently in late-stage clinical trials nearly all rely on non-novel mechanisms of action,²⁶ and so it is almost inevitable that resistance to them will develop quickly, as it has done for their predecessors.

Strategies to avoid resistance Peptide antibiotics kill bacteria quickly by the physical disruption of cell membranes, peptide antibiotics may not face the rapid emergence of resistance.²⁷ Strategy to stop virulence rather than kill²⁰ Toxin function/delivery, E.g. this doesn't kill it but stops it killing mice²⁸ Quorum sensing: LuxR homologs²⁰ *S. aureus* AgrC Inhibitory autoinducing peptides²⁰ Conjugates: sideromycins, synthetics, AI-AB gauguly

5.2 Sideromycins

Siderophores are peptides or small molecules used by microorganisms to chelate iron for the purposes of 'iron mining'.²⁹ Soluble iron is often scarce but it is crucial for many cellular processes including respiration and DNA synthesis. Siderophores are synthesised by the microorganisms and secreted into the extracellular environment where they bind to Fe^{3+} , often with exceptionally high affinities. The iron-bound siderophores are then brought back into the cell by active transport and the iron is released, either by reduction of the Fe^{3+} to Fe^{2+} or by enzymatic degradation of the siderophore. Siderophores have a wide range of structures (see Figure 1 and Figure 2), possibly so one species can avoid its siderophores being taken up by another species.³⁰

strategies
to get
new
antibi-
otics,
inc
sideromyc
and
syn-
thetic
link in
better




Figure 1: Iron-siderophore complexes: Deferoxamine B **1**³¹ (*Streptomyces pilosus* and *Streptomyces coelicolor*), rhodotorulic acid **2**³² (*Rhodotorula pilimanae*), fusarinine C **3**³³ (*Fusarium roseum*), enterobactin **4**³¹ (*Escherichia coli* and enteric bacteria), ferrichrome **5**³⁴ (*Ustilago sphaerogena*, *U. maydis*, *Aspergillus niger*, *A. quadricinctus*, *A. duricaulis* and *Penicillium resticolosum*), yersiniabactin **6**³¹ (*Yersinia pestis*).




Figure 2: Iron-siderophore complexes: pyoverdine PaA **7**^{31,35} (*P. aeruginosa*, PAO1 strain) and pyochelin **8**^{36,37} (*P. aeruginosa*). Note that pyochelin **8** is a tetradentate ligand, hence the iron ion has two sites which can bind other ligands.

Siderophore-antibiotic conjugates are produced naturally by some bacteria and are known as sideromycins⁴ (see Figure 3). Bacteria produce these molecules to attack other bacteria by hijacking their siderophore uptake mechanisms to introduce toxic compounds.

For example, albomycin **9** (see Figure 3) is a sideromycin produced by *Actinomyces subtropicus* and *Streptomyces griseus*^{38,39} which has been used to treat infections caused by various bacteria including *Yersinia enterocolitica* and *Streptococcus pneumoniae* in mice and humans.^{40,41} Albomycin **9** contains a siderophore coupled to a nucleoside antibiotic via a peptide linker. The siderophore section is structurally similar to ferrichrome **5** (see Figure 1), a siderophore produced by various fungi, but also taken up by bacteria including *Escherichia coli*, *Salmonella typhimurium* and *P. aeruginosa*.^{34,42} It has been shown that because of the structural similarity to ferrichrome **5**, *E. coli* will also take up albomycin **9**.³⁸ The linker is hydrolysed in the cytoplasm of the *E. coli*, releasing the active nucleoside antibiotic. This leads to 500-fold concentration of the antibiotic within the *E. coli* cells, enough to have significant effect on growth.

The success of albomycin⁴⁰ and other sideromycins such as salmycin A^{29,43,44} and ferrimycin A1^{45,46} has served as encouragement to many researchers to explore synthetic siderophore-antibiotic conjugates, which will be discussed in the next section.




Figure 3: Iron-sideromycin complexes: Albomycin **9**^{29,47} (*Actinomyces subtropicus* and *Streptomyces griseus*), salmycin A^{29,43,44} (*Streptomyces violaceus*) and ferrimycin²⁹ (*Streptomyces griseoflavus*).

5.3 Synthetic siderophore-antibiotic conjugates

Sideromycins served as inspiration for the design, synthesis and biological evaluation of a wide range of synthetic siderophore-antibiotic conjugates.⁴ Antibiotics used include β -lactams,^{48–50} nucleosides,⁵¹ glycopeptides⁵² and macrolides.⁵³ Sideromycin-fluoroquinolone conjugates have also been studied by several groups,^{54–56} including conjugates with linkers which can be cleaved^{55,56} in a similar manner to albomycin.³⁸ Some of these showed comparable activity to the parent antibiotic, but it is not clear whether attachment of the siderophore improved uptake or whether the conjugates acted as classical prodrugs.

β -lactam-sideromycin conjugates have been more widely investigated and show good activity *in vitro*, however, resistance can evolve by loss of the TonB transporter or of the relevant siderophore receptor, e.g. Cir and Fiu for catecholate siderophores or FhuA for hydroxamate siderophores.⁴ Recently a conjugate (Ent-Amp **12**, see Figure 4) of enterobactin and ampicillin joined using a copper(I)-catalyzed azide-alkyne cycloaddition has been shown to have increased activity against pathogenic *E. coli* when compared to native ampicillin.⁵⁷ Other work has focused on monocyclic β -lactams, for example pirazmonam **13** and U-78608 **14**, which show high

potency against Gram-negative bacteria including *P. aeruginosa*,^{58,59} Monocyclic β -lactams are generally fairly stable to β -lactamase activity, which is an advantage compared with many bicyclic β -lactams.

Three siderophore-antibiotic conjugates are reported as being in clinical trials:⁵ MC-1 **15**,⁶⁰ BAL30072 **16**⁴ (see Figure 4) and cefiderocol **17**.^{61,62}

MC-1 **15** is reported as being "in clinical phases of development",⁵ but no reports of studies in humans could be found. However, experiments in mice have been promising.⁶⁰ BAL30072 **16** is a siderophore- β -lactam conjugate which showed initial promise as it is a poor substrate for β -lactamases, and resistance due to loss of transport proteins is infrequent.⁴ However, it is unclear whether it will progress further in trials as it causes liver toxicity.⁶³ Cefiderocol **17** is a cephalosporin-catechol conjugate in phase 1 trials. Recent results indicate that 'single and 35 multiple intravenous doses of cefiderocol at up to 2000 mg were well tolerated in healthy 36 subjects'.⁶²

These examples show that siderophore-antibiotic conjugates are a promising strategy to deliver antibiotics across bacterial membranes, but it is worth noting that conjugation to a siderophore may lead to loss of activity, or resistance may be acquired by loss of transport proteins. Encouragingly though, albomycin **9**-resistant mutants have been shown to be less virulent,⁴¹ indicating that bacteria may lose out either by susceptibility to the antibiotic or by loss of fitness due to decreased iron transport.

Building on these positive examples, it is hoped that the strategy of conjugating a molecule which is important for virulence⁶⁴ with an antibiotic can be extended to conjugates of autoinducers and antibiotics in a similar 'Trojan horse' approach.




Figure 4: Examples of siderophore-antibiotic conjugates: Ent-Amp **12**,⁵⁷ pirazmonam **13**,^{58,59} U-78608 **14**,^{58,59} MC-1 **15**,⁶⁰ BAL30072 **16**⁴ and cefiderocol **17**.^{61,62}

5.4 Autoinducer-antibiotic conjugates

This study has extended the conjugation strategy discussed above by creating autoinducer-antibiotic conjugates. It was hypothesised that attaching an autoinducer to a known antibiotic could lead to increased cellular retention of the antibiotic, and could potentially restore function against resistant strains. The work is divided into two main sections. The first section focuses on conjugates of three *P. aeruginosa* autoinducers (see Figure 8) with ciprofloxacin and trimethoprim (see Figure 10). The second section focuses on conjugates of homoserine lactone analogues with ciprofloxacin (see 5.5).

5.4.1 Quorum sensing

A quorum is defined as ‘A fixed minimum number of members of an assembly or society that must be present at any of its meetings to make the proceedings of that meeting valid.’⁶⁵ A similar concept is used in bacterial signalling, whereby group behaviour is only triggered when a certain minimum concentration of bacteria has been reached. Examples of group behaviour include bioluminescence, the production of virulence factors and biofilm formation.⁶ It is advantageous for bacteria to coordinate such behaviours as they would be ineffective, and therefore a waste of resources, when carried out by a single bacterium. The process by which bacteria determine the concentration of similar bacteria in their vicinity, and act on that information, is known as quorum sensing.

Quorum sensing has since been observed in many species of bacteria, including *Vibrio fischeri*, *P. aeruginosa*, *Agrobacterium tumefaciens*, *Erwinia carotovora*, *Streptococcus pneumoniae*, *Bacillus subtilis*, *Staphylococcus aureus*, *Vibrio harveyi*, *Escherichia coli*, *Myxococcus xanthus*, *Salmonella enterica*, *Yersinia enterocolitica*, *Aeromonas sp.* and *Acinetobacter sp.*^{6,66–74} Many of these bacteria are significant causes of disease and death in humans, for example, in a typical year in the U.S. *P. aeruginosa* causes 6,700 multidrug-resistant infections and 440 deaths, methicillin-resistant *S. aureus* causes 80,500 severe infections and 11,300 deaths and non-typhoidal *Salmonella* causes 1.2 million illnesses, 23,000 hospitalisations and 450 deaths.¹

5.4.1.1 *Vibrio fischeri*

The first example of quorum sensing was discovered in *V. fischeri*, a symbiotic bacterium that produces bioluminescence in the photophore of the Hawaiian bobtail squid, *Euprymna scolopes*^{66,73,74} (see Figure 5). This bacterium receives amino acids^{75,76} from its host in exchange for producing light which the squid uses for counterillumination, to camouflage itself.⁷⁷

If a low population of *V. fischeri* were present in the photophore, the light that the bacteria could produce would be insufficient to provide counterillumination. Therefore, the bacteria conserve resources by not producing light. However, if there is a high population of *V. fischeri* it is useful for them all to produce light, as this incentivises the squid to provide them with nutrients.



Figure 5: ‘Euprymna scolopes, South shore of Oahu, Hawaii’ by Jamie Foster. Licensed under CC BY-SA 3.0 via Commons.

V. fischeri uses the LuxR-LuxI system to sense cell density. This system is seen as a paradigm of quorum sensing, and a simplified explanation of it is presented to show typical features of such a system (see Figure 6).




Figure 6: The LuxR-LuxI quorum sensing system in *V. fischeri*.

V. fischeri senses cell concentration by the detection of 3-oxo-C₆-HSL **18**⁷⁸ (see Figure 7), a freely diffusible⁷⁹

molecule which is synthesised by LuxI^{80,81} and secreted by all *V. fischeri* cells⁸² at a low basal level.⁶⁶ When the bacterial population density, and hence the concentration of 3-oxo-C₆-HSL **18**, reaches a threshold, 3-oxo-C₆-HSL **18** binds to LuxR,^{83–85} a receptor which is also synthesised at a low basal level.




Figure 7: 3-oxo-C₆-HSL **18**.

The LuxR complex binds to the *lux* operator, upregulating production of LuxI and hence 3-oxo-C₆-HSL **18**, and luciferase enzymes and hence blue-green light.^{86–88} Production of more 3-oxo-C₆-HSL **18** enables a positive feedback loop, reinforcing the effect of high population density on 3-oxo-C₆-HSL **18** concentration and hence light production. This is the reason that 3-oxo-C₆-HSL **18** is known as an autoinducer.

The system also contains a negatively feedback loop to avoid excessive expression of proteins: at high concentrations of 3-oxo-C₆-HSL **18** production of LuxR is inhibited.⁸⁹ Such balancing effects, as well as interactions with other quorum sensing and metabolic systems, are very common.

5.4.1.2 *Pseudomonas aeruginosa*

Another well-studied example of quorum sensing is in *P. aeruginosa*.^{11,90,91} *P. aeruginosa* is a Gram-negative opportunistic pathogen which typically infects immunocompromised individuals such as those with cystic fibrosis, neutropenia and AIDS. It can infect the pulmonary and urinary tracts as well being the most frequent cause of burn wound infections and the most frequent coloniser of medical devices such as catheters.⁹ Multidrug-resistant *P. aeruginosa* is classified as a ‘serious threat’ by the United States Centers for Disease Control and Prevention¹ and carbapenem-resistant *P. aeruginosa* is classified as ‘priority 1: critical’ by the World Health Organisation.⁹²

P. aeruginosa has a low susceptibility to many antibiotics and readily acquires antibiotic resistance by mutation or horizontal gene transfer.⁹³ It is difficult for antibiotics to cross into cells due to low cell membrane permeability⁹⁴ and biofilm formation,⁹⁵ and they are pumped out again by its multiple chromosomally encoded multidrug efflux pumps.¹⁰ *P. aeruginosa* biofilms are more resistant to many drugs including ciprofloxacin **24** and trimethoprim **25** compared with planktonic cells.^{95,96} This high level of antibiotic resistance makes *P. aeruginosa* an important target for drug discovery.

Quorum sensing in *P. aeruginosa* involves a complex interplay of five signalling molecules (see Figure 8) and various proteins (see Figure 9).^{11,90,91} These can be broken down into three main, interacting systems: Las, Rhl and Pqs.




Figure 8: *P. aeruginosa* autoinducers.

In the Las system, LasI⁹⁷ synthesises the 3-oxo-C₁₂-HSL **20**⁹⁸ autoinducer. 3-oxo-C₁₂-HSL **20** binds LasR,⁹⁹ and this complex upregulates the production of LasI¹⁰⁰ (thus causing autoinduction) as well as alkaline protease,¹⁰¹ elastase,⁹⁹ exotoxin A,¹⁰¹ HCN¹⁰² and LasA protease.¹⁰³ The LasR complex is also important in late-stage biofilm formation,⁷⁰ and upregulates the Rhl¹⁰⁴ and Pqs systems.^{105, 106}

In the Rhl system, RhlI¹⁰⁷ synthesises the C₄-HSL **19**¹⁰⁸ autoinducer. C₄-HSL **19** binds RhlR,¹⁰⁹ and this complex upregulates the production of RhlI¹⁰⁰ (again causing autoinduction), alkaline protease,¹¹⁰ elastase,¹⁰⁷ haemolysin,¹¹⁰ HCN,^{102, 110} LasA protease,¹⁰⁷ LecA,¹¹¹ pyocyanin^{107, 110} and rhamnolipids.¹⁰⁷ The RhlR complex also downregulates the Pqs system.^{106, 112} The Rhl system is controlled by both the Las and Pqs systems, as production of both RhlR and RhlI is upregulated by the LasR complex¹⁰⁴ and production of both RhlR is upregulated by the PqsR complex.¹¹³

In the Pqs system, the main autoinducer, PQS **22**,¹¹⁴ is synthesised by multiple enzymes. PhnAB,¹¹⁵ PqsA, PqsBC, PqsD^{116, 117} and PqsE^{118, 119} produce the precursor HHQ **21**, and PqsH converts HHQ **21** to PQS **22**. PQS **22**¹⁰⁶ or HHQ **21** binds PqsR,¹²⁰ and either complex can upregulate the synthesis of HHQ **21** causing autoinduction. The PqsR-PQS complex upregulates the production of chitinase,¹²¹ elastase,¹¹⁴ HCN,¹²¹ LecA,¹²² pyocyanin^{105, 123} and pyoverdine,¹²³ as well as increasing biofilm production¹²² and vesicle formation.¹²⁴ The PqsR-PQS complex also upregulates production of RhlR, so the Pqs system has control over the Rhl system.¹¹³ The Pqs system is controlled by both the Las and Rhl systems, as production of PqsR¹⁰⁶ and PqsH¹⁰⁵ is upregulated by the LasR complex and production of PqsA, PqsBC, PqsD, PqsE¹¹² and PqsR¹⁰⁶ is downregulated by the RhlR complex.

add
num-
bers
man-
ually
when
sorted




Figure 9: Quorum sensing in *P. aeruginosa*.^{11,90,91}

In addition to the above systems, AI-2 (see Figure 8), an interspecies signalling molecule,¹²⁵ is known to increase biofilm production and virulence in *P. aeruginosa*.^{126,127} This is thought to be achieved by interaction with the Las and Rhl systems, but the exact mechanism is not known.

In summary, *P. aeruginosa* uses the autoinducers shown in Figure 8 as part of three interacting quorum sensing systems to coordinate virulence and biofilm production, and this makes these autoinducers interesting therapeutic targets. Quorum sensing has been successfully targeted using many different modulators,^{90,128} but this study takes a slightly different approach. Inspired by the success of various siderophore-antibiotic conjugates (see 5.3), a library of autoinducer-antibiotic conjugates was synthesised, in the hope that the importance of autoinducers in harmful cellular behaviours would lead to increased activity of the conjugates (see 5.4).

5.4.2 Autoinducers

The *P. aeruginosa* autoinducers (see Figure 8) were chosen as *P. aeruginosa* is a significant human pathogen which shows high antibiotic resistance and utilises quorum sensing to coordinate pathogenic behaviours (see 5.4.1.2). Specifically, C₄-HSL **19**, HHQ **21** and PQS **22** derivatives were chosen as they were considered to be the most synthetically tractable.

5.4.3 Autoinducer efflux

Autoinducers must be exported from the cell in order to be used for intercellular communication, and the five known *P. aeruginosa* autoinducers are exported by various different transport mechanisms. The mechanism is not well known for HHQ **21** or AI-2 **23**, but it is known that PQS **22** is exported in vesicles,¹²⁹ C₄-HSL **19** passively diffuses in and out of cells,¹³⁰ and 3-oxo-C₁₂-HSL **20** is taken up passively, accumulates in the cell membrane and is actively pumped out by efflux pumps. The difference in transport mechanism for C₄-HSL **19** and 3-oxo-C₁₂-HSL **20** is thought to be largely due to chain length rather than the 3-oxo modification, as a shorter-chain version, 3-oxo-C₆-HSL **18** has been shown to be freely diffusible through *V. fischeri* membranes.⁷⁹

3-oxo-C₁₂-HSL **20** is exported primarily via the MexAB-OprM efflux system.^{10,131} The increased removal of 3-oxo-C₁₂-HSL **20** from the cell by upregulation of the MexAB-OprM system leads to decreased production of additional 3-oxo-C₁₂-HSL **20** (as the positive feedback loop is disrupted, see 5.4.1.2), and hence decreased production of pyocyanin, elastase and casein protease. It is expected that MexAB-OprM upregulation would also disrupt biofilm formation as a decrease in 3-oxo-C₁₂-HSL **20** levels would disrupt Las-mediated quorum sensing,¹³² but no direct studies of this could be found.

5.4.4 Antibiotics

Ciprofloxacin **24** and trimethoprim **25** (see Figure 10) were chosen as the antibiotic sides of the conjugates.

Ciprofloxacin **24** is second-generation fluoroquinolone antibiotic used to treat both Gram-positive and Gram-negative bacterial infections including *P. aeruginosa*.^{12,133} Ciprofloxacin **24** inhibits DNA replication by binding to DNA gyrase and topoisomerase IV.¹³⁴

Trimethoprim (see Figure 10) is a dihydrofolate reductase inhibitor used primarily to treat bladder infections.¹³⁵ It is active against several significant human pathogens including *Streptococcus pneumoniae* and *Haemophilus influenzae*, but not against *P. aeruginosa*. It was primarily chosen in this study as it was considered easy to functionalise, but also to test the feasibility of creating antibiotic activity against *P. aeruginosa*.




Figure 10: The antibiotics used in this section.

5.4.5 Antibiotic efflux

Ciprofloxacin **24** enters *P. aeruginosa* by diffusion,¹³⁶ but is pumped out by efflux pumps.¹³⁷ In the planktonic state several efflux pumps are known to pump out ciprofloxacin **24**, including MexAB–OprM, MexCD–OprJ, MexEF–OprN, MexXY–OprM, MexJK–OprM and MexVW–OprM.¹⁰ However, in biofilms only MexEF–OprN has an effect.¹³⁸

Trimethoprim **25** is mainly exported by the MexAB–OprM,¹³⁹ MexCD–OprJ¹⁴⁰ and MexEF–OprN¹⁴¹ multidrug efflux systems^{10,142} in the planktonic state. It is not known which pumps are used to export trimethoprim **25** from biofilms, but biofilms do show increased resistance to it.⁹⁶

5.4.6 Conjugate efflux and antibiotic action

There are two ways in which the conjugates could disrupt *P. aeruginosa* growth:

1. *P. aeruginosa* could develop resistance to an autoinducer-antibiotic conjugate by upregulation of its export mechanism, but this would also lead to increased export of the native autoinducer, thus disrupting the quorum sensing system and hence biofilm formation and virulence.^{11,131,132} For HSL conjugates this would mean upregulation of the MexAB–OprM pump, as this is the pump used for export of 3-oxo-C₁₂-HSL **20**.^{10,131} For PQS conjugates this would mean upregulation of vesicle formation.¹²⁹
2. The autoinducer section could make the conjugate a poor substrate for the antibiotic section's usual efflux mechanism, leading to accumulation of the conjugate within cells and hence increased antibacterial activity. For autoinducer-ciprofloxacin conjugates acting on planktonic *P. aeruginosa* this would mean the conjugate being a poor substrate of the various efflux pumps listed in the previous section. For autoinducer-ciprofloxacin conjugates acting on biofilms this would mean the conjugate being a poor substrate of MexEF–OprN (the sole exporter of ciprofloxacin **24** in biofilms¹³⁸ and not an exporter of HSLs **19** or **20**, or PQS **22**¹⁰). This mechanism could in principal work for trimethoprim **25** as well, but it is not known which pumps are active against this antibiotic in biofilms.

5.5 Autoinducer analogue-ciprofloxacin conjugates

Following on from the library of compounds based on *P. aeruginosa* autoinducers, a series of conjugates based on *analogues* of HSL were planned. This strategy was inspired by a paper¹⁴³ and patent¹⁴⁴ by Ganguly *et al.*, who synthesised and characterised a conjugate **101** of methyl ciprofloxacin with homocysteine thiolactone (see Figure 11). Homocysteine thiolactone is an analogue of homoserine lactone with the ring oxygen replaced by sulfur, and has been used as the head group in several other known quorum sensing modulators.^{82,145–151}




Figure 11: The HCTL-CipMe conjugate **101** studied by Ganguly *et al.*^{143,144}

As part of their characterisation of the HCTL-CipMe conjugate **101**, Ganguly *et al.* found the minimum inhibitory concentration (MIC) of the conjugate in *P. aeruginosa* under standard planktonic conditions. The

MIC was found to be ten times higher for the conjugate vs. ciprofloxacin (50 vs. 5 μm), indicating that the conjugate was less effective than ciprofloxacin under planktonic conditions.

Ganguly *et al.* then investigated the effect of the conjugate on biofilms. The conjugate and ciprofloxacin were first added to dilute *P. aeruginosa* liquid culture at 25 μm . As expected, the culture failed to grow and form biofilm in the presence of ciprofloxacin, but did grow in the presence of the conjugate **101**. They then incubated cultures for 24 h, to allow biofilms to grow, before adding the compounds. In contrast, they found that the conjugate **101** disrupted the biofilm more effectively than ciprofloxacin. When the biofilm was grown for 48 or 72 hours the conjugate had similarly disruptive effects, whereas ciprofloxacin ‘did not show any significant antibacterial activity’.

These results are exciting as they hint that an autoinducer conjugate might be able to combat an established *P. aeruginosa* infection more effectively than the unmodified antibiotic. Ganguly *et al.* suggest that their conjugate is more effective than ciprofloxacin in penetrating biofilms, and/or better at avoiding being pumped out by multidrug efflux pumps. They posit that this could be due to the thiolactone head, as they also showed that unconjugated C₄-HCTL **26** (see Figure 12) has ‘either enhanced uptake or functional activity’ when compared with C₄-HSL **19**.

It is possible that the conjugate **101** has higher activity against biofilms when compared with ciprofloxacin **24** because conjugate **101** avoids being pumped out by multidrug efflux pumps, or selects for the survival of mutants with upregulated efflux pumps, and hence disrupted quorum sensing systems (see 5.4.6).

While one might expect the conjugate **101** to behave like C₄-HSL **19**, and hence passively diffuse in and out of cells, it is possible that its transport more closely resembles that of 3-oxo-C₁₂-HSL **20**. 3-oxo-C₁₂-HSL **20**’s accumulation in membranes and interaction with efflux pumps is thought to be based primarily on tail chain length (see 5.4.3), and the ciprofloxacin half of the conjugate **101** could be seen as a long tail, especially as the carboxylic acid is methylated and hence less polar.




Figure 12: C₄-HSL **19** and C₄-HCTL **26**. Note that Ganguly *et al.* tested the *S* enantiomer of C₄-HCTL **26**, but used a racemic mixture in their HCTL-CipMe conjugate.

While the results found by Ganguly *et al.* show promise, they only test one conjugate, and do not include controls to show that the HCTL group specifically is necessary for the enhanced effect. It was therefore decided to build on this work by synthesising a series of ciprofloxacin conjugates with head groups known as part of quorum sensing modulators.^{128, 152}

The activity of the chosen head groups against *P. aeruginosa* receptors when coupled with the native C₄ and 3-oxo-C₁₂ tails is summarised in Table 2. It is hoped that high activity of these molecules should correlate with high activity of their ciprofloxacin conjugates. This is not a comprehensive list of active head groups, and other possible choices are covered in ??.









Head group		
	Partial agonist and antagonist against LasR. ¹⁴⁹ Shown to increase biofilm formation in <i>P. aeruginosa</i> . ¹⁴³	Strong agonist against LasR, with comparable activity to the native ligand. ^{146, 147, 149, 153}
	Partial agonist against LasR. ¹⁵²	Strong antagonist against LasR. ¹⁵²
	Poor agonist and antagonist against RhlR. ^{154, 155}	Strong antagonist against LasR. ¹⁵⁴
	Strong agonist against RhlR. ¹⁵⁴ <i>SS</i> enantiomer is more potent. ¹⁵⁵	Partial agonist against LasR. ¹⁵⁴
	Strong agonist against RhlR. ¹⁵⁴ <i>SS</i> enantiomer is more potent, with comparable activity to the native ligand. ¹⁵⁵	Strong agonist against LasR. ^{147, 154} <i>SS</i> enantiomer is more potent, with comparable activity to the native ligand. ¹⁵⁵
	Strong agonist against RhlR. ¹⁵⁴ <i>SS</i> enantiomer is more potent. ¹⁵⁵	Partial antagonist against LasR. ¹⁵⁴ Shown to reduce biofilm formation in <i>P. aeruginosa</i> . ¹⁵⁴

Table 2: Activities of autoinducers containing the chosen head groups when coupled with C₄ or 3-oxo-C₁₂ tails.

6 Aims

aims
here

7 Results and discussion: autoinducer-antibiotic conjugates

7.1 Overview

The first part of this project was focused on producing a library of autoinducer-antibiotic conjugates. *P. aeruginosa* autoinducers were used, in particular C₄-HSL **19**, HHQ **21** and PQS **22** (see Figure 8). Azido derivatives of these compounds were coupled to alkynyl derivatives of antibiotics, specifically ciprofloxacin **24** and trimethoprim **25** (see Figure 10), using a copper(I)-catalysed azide-alkyne cycloaddition.^{7,8}

7.1.1 Azido autoinducer derivatives

The structure-activity relationships in HHQ **21** and PQS **22** have been previously studied,^{156–158} and it was shown various substitutions on the benzene ring could be made without significantly decreasing activity. The 6-azido derivatives (see Figure 13) were chosen for this study as routes to them have previously been found.¹⁵⁹




Figure 13: The azido derivatives of HHQ **21** and PQS **22**: **36** and **47**.

Alteration of the lactone group of HSL derivatives is known to significantly decrease activity, especially where the number of H-bond donors or acceptors is altered.¹²⁸ Hence, the azide group was included on the tail.¹⁶⁰ Acyl tail length is known to play an important role in affinity,¹²⁸ so three derivatives of C₄-HSL **19** were synthesised: N₃-C₂-HSL **53**, N₃-C₄-HSL **56** and N₃-C₆-HSL **59** (see Figure 14).




Figure 14: The azido derivatives of C₄-HSL **19**: **53**, **56** and **59**.

7.1.2 Alkynyl antibiotic derivatives

The structure-activity relationships for ciprofloxacin have been investigated¹⁶¹ and modifications at the cyclopropane and piperazine groups were found not to cause loss of activity. It was decided an alkyne tail would be added onto the free NH of the piperazine ring, as this position is more synthetically accessible. Alkynyl ciprofloxacin derivative **66** (see Figure 15) was synthesised in this study (see ??), and two cleavable alkynyl ciprofloxacin derivatives **88** and **89** were synthesised by Dr Eddy Sotelo and combined with some of the azido HSL derivatives made in this study (see ?? and ??).




Figure 15: The alkyne ciprofloxacin derivatives **66**, **88** and **89**.

The choice to of alkyne tail attachment point on trimethoprim **25** (see Figure 16) is based on the use of that same point in a fluorogenic trimethoprim tag synthesised by Jing *et al.*¹⁶²





Figure 16: The alkyne trimethoprim derivative **69**.

7.1.3 Synthesis of the conjugates

A copper(I)-catalysed azide-alkyne cycloaddition,^{7,8} commonly referred to as a click reaction (although this is a more general term), was used to join each combination of autoinducer and antibiotic together (see Scheme 1).



Scheme 1: The construction of the triazole-linked autoinducer-antibiotic conjugate library using a copper(I)-catalysed azide-alkyne cycloaddition.

8 Results and discussion: autoinducer analogue-ciprofloxacin conjugates

8.1 Overview

8.1.1 Head groups

The head groups used in this study are shown in Figure 17. The cyclohexanol derivatives were synthesised as a diastereomerically pure racemate, whereas the cyclopentanol derivatives were synthesised as separate enantiomers. Unfortunately, cyclopentanone derivatives were not synthesised, and would be an obvious future addition to the library. The 2-methoxybenzene derivatives do not have precedents as quorum sensing modulators in the literature, but they were included so as to be compared with the 3-methoxybenzene derivatives.

make
this
actually
an
overview





Figure 17: The head groups used in this section.

8.1.2 Library construction

As Ganguly *et al.*⁷ synthesised their conjugate from Br-C₄-HCTL, it was envisaged that a branching strategy could be used to produce two sets of conjugates (see Scheme 2). The first set would be formed by the S_N2 reaction of the relevant bromide with methyl ciprofloxacin. The second set would be made by displacing the bromide with azide, then performing a click reaction with the alkynyl ciprofloxacin derivative **66** made previously to form the triazole-linked product. Ketone conjugates would be formed by oxidation of the alcohols.



Scheme 2

9 References

- [1] *Antibiotic Resistance Threats in the United States*. 2013.
- [2] S. C. Davies, *The Drugs Don't Work: A Global Threat*, Penguin Books Limited. 2013.
- [3] K. M. G. O'Connell, J. T. Hodgkinson, H. F. Sore, M. Welch, P. George, C. Salmond, D. R. Spring and G. P. C. Salmond. Combating multidrug-resistant bacteria: current strategies for the discovery of novel antibacterials. *Angewandte Chemie International Edition*, 52(41):10706–10733. 2013.
- [4] M. G. P. Page. Siderophore conjugates. *Annals of the New York Academy of Sciences*, 1277:115–126. 2013.
- [5] I. J. Schalk and G. L. A. Mislin. Bacterial Iron Uptake Pathways: Gates for the Import of Bactericide Compounds. 2017.
- [6] C. M. Waters and B. L. Bassler. Quorum sensing: cell-to-cell communication in bacteria. *Annual Review of Cell and Developmental Biology*, 21:319–346. 2005.
- [7] C. W. Tornøe, C. Christensen and M. Meldal. Peptidotriazoles on solid phase: [1,2,3]-triazoles by regio-specific copper(I)-catalyzed 1,3-dipolar cycloadditions of terminal alkynes to azides. *The Journal of Organic Chemistry*, 67(9):3057–3064. 2002.
- [8] V. V. Rostovtsev, L. G. Green, V. V. Fokin and K. B. Sharpless. A stepwise Huisgen cycloaddition process: copper(I)-catalyzed regioselective “ligation” of azides and terminal alkynes. *Angewandte Chemie International Edition*, 41(14):2596–2599. 2002.
- [9] G. P. Bodey, R. Bolivar, V. Fainstein and L. Jadeja. Infections caused by *Pseudomonas aeruginosa*. *Reviews of Infectious Diseases*, 5(2):279–313. 1983.
- [10] K. Poole. Efflux-mediated multiresistance in Gram-negative bacteria. *Clinical Microbiology and Infection*, 10(1):12–26. 2004.
- [11] J.-F. Dubern and S. P. Diggle. Quorum sensing by 2-alkyl-4-quinolones in *Pseudomonas aeruginosa* and other bacterial species. *Molecular BioSystems*, 4(9):882–888. 2008.
- [12] A. P. Macgowan, M. Wootton and H. A. Holt. The antibacterial efficacy of levofloxacin and ciprofloxacin against *Pseudomonas aeruginosa* assessed by combining antibiotic exposure and bacterial susceptibility. *Journal of Antimicrobial Chemotherapy*, 43:345–349. 1999.
- [13] H.-C. Su, K. Ramkissoon, J. Doolittle, M. Clark, J. Khatun, A. Secrest, M. C. Wolfgang and M. C. Giddings. The development of ciprofloxacin resistance in *Pseudomonas aeruginosa* involves multiple response stages and multiple proteins. *Antimicrobial Agents and Chemotherapy*, 54(11):4626–4635. 2010.
- [14] A. Fleming. On the antibacterial action of cultures of a penicillium, with special reference to their use in the isolation of *B. influenzae*. *The British Journal of Experimental Pathology*, 10(3):226–236. 1929.
- [15] M. Barber. Staphylococcal infection due to penicillin-resistant strains. *British Medical Journal*, 2(4534):863–865. 1947.
- [16] P. M. Rountree and E. F. Thomson. Incidence of penicillin-resistant and streptomycin-resistant staphylococci in a hospital. *The Lancet*, 254(6577):501–504. 1949.
- [17] P. S. Stewart and J. W. Costerton. Antibiotic resistance of bacteria in biofilms. *The Lancet*, 358(9276):135–138. 2001.

- [18] C. Fuda, M. Suvorov, S. B. Vakulenko and S. Mobashery. The basis for resistance to β -lactam antibiotics by penicillin-binding protein 2a of methicillin-resistant *Staphylococcus aureus*. *The Journal of Biological Chemistry*, 279(39):40802–40806. 2004.
- [19] O. Sköld. Sulfonamide resistance: mechanisms and trends. *Drug Resistance Updates*, 3(3):155–160. 2000.
- [20] A. E. Clatworthy, E. Pierson and D. T. Hung. Targeting virulence: a new paradigm for antimicrobial therapy. *Nature Chemical Biology*, 3(9):541–548. 2007.
- [21] S. R. Palumbi. Humans as the World’s Greatest Evolutionary Force. *Science*, 293(5536):1786–1790. 2001.
- [22] J. W. Ogle, L. B. Reller and M. L. Vasil. Development of resistance in *Pseudomonas aeruginosa* to imipenem, norfloxacin, and ciprofloxacin during therapy: proof provided by typing with a DNA probe. *The Journal of Infectious Diseases*, 157(4):743–748. 1988.
- [23] P. Huovinen. Resistance to Trimethoprim-Sulfamethoxazole. *Antimicrobial Resistance*, 32(11):1608–1614. 2001.
- [24] M. C. Birmingham, C. R. Rayner, A. K. Meagher, S. M. Flavin, D. H. Batts and J. J. Schentag. Linezolid for the treatment of multidrug-resistant , Gram-positive infections: experience from a compassionate-use program. *Clinical Infectious Diseases*, 36(2):159–168. 2003.
- [25] D. K. Lee, Y. Kim, K. S. Park, J. W. Yang, K. Kim and N. J. Ha. Antimicrobial activity of mupirocin, daptomycin, linezolid, quinupristin/dalfopristin and tigecycline against vancomycin-resistant enterococci (VRE) from clinical isolates in Korea (1998 and 2005). *Journal of Biochemistry and Molecular Biology*, 40(6):881–887. 2007.
- [26] H. W. Boucher, G. H. Talbot, J. S. Bradley, J. E. Edwards, D. Gilbert, L. B. Rice, M. Scheld, B. Spellberg and J. Bartlett, *Bad Bugs , No Drugs : No ESKAPE ! An Update from the Infectious Diseases Society of America*. 2009.
- [27] R. E. W. Hancock. Peptide antibiotics. 349:418–422. 1997.
- [28] L. Lin, B. Tan, P. Pantapalangkoor, T. Ho, B. Baquir, A. Tomaras, J. I. Montgomery, E. G. Barbacci, K. Hujer, R. A. Bonomo, L. Fernandez, R. E. W. Hancock, M. D. Adams, S. W. French, V. S. Buslon and B. Spellberg. Inhibition of LpxC Protects Mice from Resistant *Acinetobacter baumannii* by Modulating Inflammation and Enhancing Phagocytosis. *mBio*, 3(5):23–29. 2012.
- [29] R. C. Hider and X. L. Kong. Chemistry and biology of siderophores. *Natural Product Reports*, 27(5):637–657. 2010.
- [30] M. R. Seyedsayamdost, S. Cleto, G. Carr, H. Vlamakis, M. João Vieira, R. Kolter and J. Clardy. Mixing and matching siderophore clusters: structure and biosynthesis of serratiochelins from *Serratia sp.* V4. *Journal of the American Chemical Society*, 134(33):13550–135503. 2012.
- [31] T. Zheng and E. M. Nolan. Siderophore-based detection of Fe(III) and microbial pathogens. *Metallomics*, 4(9):866–880. 2012.
- [32] C. J. Carrano and K. N. Raymond. Synthesis and characterization of iron complexes of rhodotorulic acid: a novel dihydroxamate siderophore and potential chelating drug. *Journal of the Chemical Society, Chemical Communications*, (12):501–502. 1978.
- [33] M. B. Hossain, D. L. Eng-Wilmot, R. A. Loghry and D. van der Helm. Circular dichroism, crystal structure, and absolute configuration of the siderophore ferric *N,N',N''*-triacetylfusarinine, $\text{FeC}_3\text{9H}_5\text{7N}_6\text{O}_{15}$. *Journal of the American Chemical Society*, 102(18):5766–5773. 1980.

- [34] D. van der Helm, J. R. Baker, D. L. Eng-Wilmot, M. B. Hossain and R. A. Loghry. Crystal structure of ferrichrome and a comparison with the structure of ferrichrome A. *Journal of the American Chemical Society*, 102(12):4224–4231. 1980.
- [35] J.-m. Meyer. Pyoverdines: pigments, siderophores and potential taxonomic markers of fluorescent *Pseudomonas* species. *Archives of Microbiology*, 174:135–142. 2000.
- [36] K. Schlegel, J. Lex, K. Taraz and H. Budzikiewicz. The X-ray structure of the pyochelin Fe^{3+} complex. *Zeitschrift für Naturforschung*, 61c(3-4):263–266. 2006.
- [37] D. Cobessi, H. Celia and F. Pattus. Crystal Structure at High Resolution of Ferric-pyochelin and its Membrane Receptor FptA from *Pseudomonas aeruginosa*. pages 893–904. 2005.
- [38] A. Hartmann, H.-P. Fiedler and V. Braun. Uptake and conversion of the antibiotic albomycin by *Escherichia coli* K-12. *European Journal of Biochemistry*, 99(3):517–24. 1979.
- [39] H. Fiedler, F. Walz, A. Döhle and H. Zährner. Albomycin: studies on fermentation, isolation and quantitative determination. *Applied Microbiology and Biotechnology*, 21(6):341–347. 1985.
- [40] G. F. Gause. Recent studies on albomycin, a new antibiotic. *British Medical Journal*, 2(4949):1177–1179. 1955.
- [41] A. Pramanik, U. H. Stroehrer, J. Krejci, A. J. Standish, E. Bohn, J. C. Paton, I. B. Autenrieth and V. Braun. Albomycin is an effective antibiotic, as exemplified with *Yersinia enterocolitica* and *Streptococcus pneumoniae*. *International Journal of Medical Microbiology*, 297(6):459–469. 2007.
- [42] M. Hannauer, Y. Barda, G. L. A. Mislin, A. Shanzer and I. J. Schalk. The ferrichrome uptake pathway in *Pseudomonas aeruginosa* involves an iron release mechanism with acylation of the siderophore and recycling of the modified desferrichrome. *Journal of Bacteriology*, 192(5):1212–1220. 2010.
- [43] L. Vértessy, W. Aretz, H.-W. Fehlhaber and H. Kogler. Salmycin A–D, Antibiotika aus *Streptomyces violaceus*, DSM 8286, mit Siderophor-Aminoglycosid-Struktur. *Helvetica Chimica Acta*, 78(1):46–60. 1995.
- [44] V. Braun, A. Pramanik, T. Gwinner, M. Köberle and E. Bohn. Sideromycins: tools and antibiotics. *Biometals*, 22:3–13. 2009.
- [45] W. Sackmann, P. Reusser, L. Neipp, F. Kradolfer and F. Gross. Ferrimycin A, a new iron-containing antibiotic. *Antibiotics & Chemotherapy*, 12:34–45. 1962.
- [46] D. Gottlieb and P. D. Shaw, *Mechanism of Action*, Springer. 2012.
- [47] G. Benz, T. Schröder, J. Kurz, C. Wünsche, W. Karl, G. Steffens, J. Pfitzner and D. Schmidt. Constitution of the deferriform of the albomycins $\delta 1$, $\delta 2$ and ϵ . *Angewandte Chemie International Edition in English*, 21(7):527–528. 1982.
- [48] U. Möllmann, L. Heinisch, A. Bauernfeind, T. Köhler and D. Ankel-Fuchs. Siderophores as drug delivery agents: application of the "Trojan Horse" strategy. *Biometals*, 22(4):615–624. 2009.
- [49] C. Dini and J. Aszodi. Synthesis of a dihydroxythiophene analogue of catechospirines. *Bioorganic & Medicinal Chemistry Letters*, 10(4):349–352. 2000.
- [50] T. Kline, M. Fromhold, T. E. McKennon, S. Cai, J. Treiberg, N. Ihle, D. Sherman, W. Schwan, M. J. Hickey, P. Warrenner, P. R. Witte, L. L. Brody, L. Goltry, L. M. Barker, S. U. Anderson, S. K. Tanaka, R. M. Shawar, L. Y. Nguyen, M. Langhorne, A. Bigelow, L. Embuscado and E. Naeemi. Antimicrobial effects of novel siderophores linked to β -lactam antibiotics. *Bioorganic & Medicinal Chemistry*, 8(1):73–93. 2000.

- [51] Y. Lu and M. J. Miller. Syntheses and Studies of Multiwarhead Siderophore-5-fluorouridine Conjugates. *Bioorganic & Medicinal Chemistry*, 7(1999):3025–3038. 1999.
- [52] M. Ghosh and M. J. Miller. Synthesis and in vitro antibacterial activity of spermidine-based mixed catechol- and hydroxamate-containing siderophore–vancomycin conjugates. *Bioorganic & Medicinal Chemistry*, 4(1):43–48. 1996.
- [53] M. Ghosh and M. J. Miller. Design, synthesis, and biological evaluation of isocyanurate-based antifungal and macrolide antibiotic conjugates: iron transport-mediated drug delivery. *Bioorganic & Medicinal Chemistry*, 3(11):1519–1525. 1995.
- [54] S. R. Md-Saleh, E. C. Chilvers, K. G. Kerr, S. J. Milner, A. M. Snelling, J. P. Weber, G. H. Thomas, A.-K. Duhme-Klair and A. Routledge. Synthesis of citrate-ciprofloxacin conjugates. *Bioorganic & Medicinal Chemistry Letters*, 19(5):1496–1498. 2009.
- [55] F. Rivault, C. Liébert, A. Burger, F. Hoegy, M. A. Abdallah, I. J. Schalk and G. L. A. Mislin. Synthesis of pyochelin-norfloxacin conjugates. *Bioorganic & Medicinal Chemistry Letters*, 17(3):640–644. 2007.
- [56] C. Ji and M. J. Miller. Chemical syntheses and in vitro antibacterial activity of two desferrioxamine B-ciprofloxacin conjugates with potential esterase and phosphatase triggered drug release linkers. *Bioorganic & Medicinal Chemistry*, 20(12):3828–3836. 2012.
- [57] T. Zheng and E. M. Nolan. Enterobactin-Mediated Delivery of β -Lactam Antibiotics Enhances Antibacterial Activity Against Pathogenic Escherichia coli. *Journal of the American Chemical Society*. 2014.
- [58] G. E. Zurenko, S. E. Truesdell, B. H. Yagi, R. J. Mourey and A. L. Laborde. In vitro antibacterial activity and interactions with beta-lactamases and penicillin-binding proteins of the new monocarbam antibiotic U-78608. *Antimicrobial Agents and Chemotherapy*, 34(5):884–8. 1990.
- [59] J. M. Harrington, T. Gootz, M. Flanagan, M. Lall, J. O'Donnell, J. Winton, J. Mueller and A. L. Crumbliss. Characterization of the aqueous iron(III) chelation chemistry of a potential Trojan Horse antimicrobial agent: Chelate structure, stability and pH dependent speciation. *BioMetals*, 25(5):1023–1036. 2012.
- [60] C. J. McPherson, L. M. Aschenbrenner, B. M. Lacey, K. C. Fahnoe, M. M. Lemmon, S. M. Finegan, B. Tadakamalla, J. P. O. Donnell, J. P. Mueller and A. P. Tomaras. Clinically Relevant Gram-Negative Resistance Mechanisms Have No Effect on the Efficacy of MC-1 , a Novel Siderophore-Conjugated. 56(12):6334–6342. 2012.
- [61] A. Ito, T. Sato, M. Ota, M. Takemura, T. Nishikawa, S. Toba, N. Kohira, S. Miyagawa, N. Ishibashi, S. Matsumoto, R. Nakamura, M. Tsuji and Y. Yamanoa. In Vitro Antibacterial Properties of Cefiderocol, a Novel Siderophore Cephalosporin, against Gram-Negative Bacteria. *Antimicrobial Agents and Chemotherapy*, 62(1):1–11. 2018.
- [62] J. S. Yutaka Saisho, Takayuki Katsube, Scott White, Hiroyuki Fukase. Pharmacokinetics, Safety, and Tolerability of Cefiderocol, a Novel Siderophore Cephalosporin for Gram-Negative Bacteria, in Healthy Subjects. *Antimicrobial Agents and Chemotherapy*, (January). 2018.
- [63] F. Paech, S. Messner, J. Spickermann, M. Wind, A. Hortense, S. Hoffmann, A. Therese, W. Brett, A. H. Rachel, J. C. Jeff, W. Marc, S. Krähenbühl and M. Maurer. Mechanisms of hepatotoxicity associated with the monocyclic β - lactam antibiotic BAL30072. *Archives of Toxicology*, 91(11):3647–3662. 2017.
- [64] M. L. Vasil and U. A. Ochsner. The response of *Pseudomonas aeruginosa* to iron: genetics, biochemistry and virulence. *Molecular Microbiology*, 34(3):399–413. 1999.

- [65] *Oxford English Dictionary*, Oxford University Press. 2014.
- [66] M. B. Miller and B. L. Bassler. Quorum sensing in bacteria. *Annual Review of Microbiology*, 55:165–199. 2001.
- [67] W. C. Fuqua, S. C. Winans and E. P. Greenberg. MINIREVIEW Quorum Sensing in Bacteria : the LuxR-LuxI Family of Cell Density-Responsive Transcriptional Regulators. *Journal of Bacteriology*, 176(2):269–275. 1994.
- [68] S. Atkinson, C.-Y. Chang, R. E. Sockett, M. Cámara and P. Williams. Quorum Sensing in *Yersinia enterocolitica* Controls Swimming and Swarming Motility. *Journal of Bacteriology*, 188(4):1451–1461. 2006.
- [69] K.-G. Chan, S. D. Puthucheary, X.-Y. Chan, W.-F. Yin, C.-S. Wong, W.-S. S. Too and K.-H. Chua. Quorum sensing in *Aeromonas* species isolated from patients in Malaysia. *Current Microbiology*, 62(1):167–72. 2011.
- [70] K. Sauer, A. K. Camper, G. D. Ehrlich, J. W. Costerton and D. G. Davies. *Pseudomonas aeruginosa* Displays Multiple Phenotypes during Development as a Biofilm. *Journal of Bacteriology*, 184(4):1140–1154. 2002.
- [71] B. Michael, J. N. Smith, S. Swift and F. Heffron. SdiA of *Salmonella enterica* Is a LuxR Homolog That Detects Mixed Microbial Communities. *Journal of Bacteriology*, 183(19):5733–5742. 2001.
- [72] B. M. M. Ahmer. Cell-to-cell signalling in *Escherichia coli* and *Salmonella enterica*. *Molecular Microbiology*, 52(4):933–945. 2004.
- [73] K. H. Nealson, T. Platt and J. W. Hastings. Cellular Control of the Synthesis and Activity of the Bacterial Luminescent System. *Journal of Bacteriology*, 104(1):313–322. 1970.
- [74] K. L. Visick and E. G. Ruby. *Vibrio fischeri* and its host: it takes two to tango. *Current Opinion in Microbiology*, 9(6):632–638. 2006.
- [75] J. Graf and E. G. Ruby. Host-derived amino acids support the proliferation of symbiotic bacteria. *Proceedings of the National Academy of Sciences*, 95(4):1818–1822. 1998.
- [76] J. D. Lemus and M. J. McFall-Ngai. Alterations in the proteome of the *Euprymna scolopes* light organ in response to symbiotic *Vibrio fischeri*. *Applied and Environmental Microbiology*, 66(9):4091–4097. 2000.
- [77] B. W. Jones and M. K. Nishiguchi. Counterillumination in the Hawaiian bobtail squid, *Euprymna scolopes* Berry (Mollusca: Cephalopoda). *Marine Biology*, 144(6):1151–1155. 2004.
- [78] A. Eberhard, A. L. Burlingame, C. Eberhard, G. L. Kenyon, K. H. Nealson and N. J. Oppenheimer. Structural identification of autoinducer of *Photobacterium fischeri* luciferase. *Biochemistry*, 20(9):2444–2449. 1981.
- [79] H. B. Kaplan and E. P. Greenberg. Diffusion of autoinducer is involved in regulation of the *Vibrio fischeri* luminescence system. *Journal of Bacteriology*, 163(3):1210–1214. 1985.
- [80] M. R. Parsek, D. L. Val, B. L. Hanzelka, J. E. Cronan and E. P. Greenberg. Acyl homoserine-lactone quorum-sensing signal generation. *Proceedings of the National Academy of Sciences*, 96(8):4360–4365. 1999.
- [81] W. T. Watson, T. D. Minogue, D. L. Val, S. B. von Bodman and M. E. A. Churchill. Structural basis and specificity of acyl-homoserine lactone signal production in bacterial quorum sensing. *Molecular Cell*, 9(3):685–694. 2002.

- [82] A. L. Schaefer, B. L. Hanzelka, A. Eberhard and E. P. Greenberg. Quorum sensing in *Vibrio fischeri*: probing autoinducer-LuxR interactions with autoinducer analogs. *Journal of Bacteriology*, 178(10):2897–2901. 1996.
- [83] B. L. Hanzelka and E. P. Greenberg. Evidence that the N-terminal region of the *Vibrio Fischeri* LuxR protein constitutes an autoinducer binding domain. *Journal of Bacteriology*, 177(3):815–817. 1995.
- [84] S. H. Choi and E. P. Greenberg. The C-terminal region of the *Vibrio fischeri* LuxR protein contains an inducer-independent *lux* gene activating domain. *Proceedings of the National Academy of Sciences of the United States of America*, 88(24):11115–11119. 1991.
- [85] S. H. Choi and E. P. Greenberg. Genetic dissection of DNA binding and luminescence gene activation by the *Vibrio fischeri* LuxR protein. *Journal of Bacteriology*, 174(12):4064–4069. 1992.
- [86] J. H. Devine, G. S. Shadel and T. O. Baldwin. Identification of the operator of the lux regulon from the *Vibrio fischeri* strain ATCC7744. *Proceedings of the National Academy of Sciences*, 86(15):5688–5692. 1989.
- [87] J. Engebrecht, K. Nealson and M. Silverman. Bacterial bioluminescence: isolation and genetic analysis of functions from *Vibrio fischeri*. *Cell*, 32(3):773–781. 1983.
- [88] K. L. Visick, J. Foster, J. Doino, M. McFall-Ngai and E. G. Ruby. *Vibrio fischeri* lux genes play an important role in colonization and development of the host light organ. *Journal of Bacteriology*, 182(16):4578–4586. 2000.
- [89] P. V. Dunlap and J. M. Ray. Requirement for autoinducer in transcriptional negative autoregulation of the *Vibrio fischeri* *luxR* gene in *Escherichia coli*. *Journal of Bacteriology*, 171(6):3549–3552. 1989.
- [90] J. T. Hodgkinson. The synthesis of Pseudomonas Quinolone Signal analogues and their effects on quinolone signalling in *Pseudomonas aeruginosa*. PhD thesis, University of Cambridge. 2011.
- [91] P. N. Jimenez, G. Koch, J. A. Thompson, K. B. Xavier, R. H. Cool and W. J. Quax. The multiple signaling systems regulating virulence in *Pseudomonas aeruginosa*. *Microbiology and molecular biology reviews : MMBR*, 76(1):46–65. 2012.
- [92] S. Harbarth, G. Kahlmeter, J. Kluytmans, M. Mendelson, G. S. Hospital, C. Town, S. Africa, C. Pulcini, N. Singh, U. Theuretzbacher, C. U. S. Food, S. Spring, L. Grayson, C. Houchens, D. L. Monnet, M. Ouellette, J. B. Patel, N. Zealand, E. Carrara, A. Savoldi, D. Kattula and F. Burkert. GLOBAL PRIORITY LIST OF ANTIBIOTIC-RESISTANT BACTERIA TO GUIDE RESEARCH , DISCOVERY , AND DEVELOPMENT OF.
- [93] P. Cornelis, *Pseudomonas: Genomics and Molecular Biology*, Caister Academic Press. 2008.
- [94] H. Nikaido. Outer Membrane Barrier as a Mechanism of Antimicrobial Resistance. 33(11):1831–1836. 1989.
- [95] D. J. Evans, D. G. Allison, M. R. W. Brown and P. Gilbert. Susceptibility of *Pseudomonas aeruginosa* and *Escherichia coli* biofilms towards ciprofloxacin : effect of specific growth rate. (August):177–184. 1991.
- [96] M. E. Olson, H. Ceri, D. W. Morek, A. G. Buret and R. R. Read. Biofilm bacteria: formation and comparative susceptibility to antibiotics. *The Canadian Journal of Veterinary Research*, 66:86–92. 2002.
- [97] M. J. Wargo and D. A. Hogan. Examination of *Pseudomonas aeruginosa* *lasI* regulation and 3-oxo-C12-homoserine lactone production using a heterologous *Escherichia coli* system. *FEMS Microbiology Letters*, 273(1):38–44. 2007.

- [98] J. P. Pearson, K. M. Gray, L. Passador, K. D. Tucker, A. Eberhard, B. H. Iglewski and E. P. Greenberg. Structure of the autoinducer required for expression of *Pseudomonas aeruginosa* virulence genes. *Proceedings of the National Academy of Sciences of the United States of America*, 91(1):197–201. 1994.
- [99] M. J. Gambello and B. H. Iglewski. Cloning and characterization of the *Pseudomonas aeruginosa lasR* gene, a transcriptional activator of elastase expression. *Journal of Bacteriology*, 173(9):3000–3009. 1991.
- [100] E. C. Pesci, J. P. Pearson, P. C. Seed, E. C. Pesci, J. P. Pearson, P. C. Seed and B. H. Iglewski. Regulation of las and rhl quorum sensing in *Pseudomonas aeruginosa* Regulation of las and rhl Quorum Sensing in *Pseudomonas aeruginosa*. *Strain*, 179(10):3127–3132. 1997.
- [101] M. J. Gambello, S. Kaye and B. H. Iglewski. LasR of *Pseudomonas aeruginosa* Is a Transcriptional Activator of the Alkaline Protease Gene (apr) and an Enhancer of Exotoxin A Expression. 61(4):1180–1184. 1993.
- [102] G. Pessi and D. Haas. Transcriptional Control of the Hydrogen Cyanide Biosynthetic Genes hcnABC by the Anaerobic Regulator ANR and the Quorum-Sensing Regulators LasR and RhlR in *Pseudomonas aeruginosa*. 182(24):6940–6949. 2000.
- [103] D. S. Toder, M. J. Gambello and B. H. Iglewski. *Pseudomonas aeruginosa* LasA: a second elastase under the transcriptional control of lasR. *Molecular Microbiology*, 5(8):2003–2010. 1991.
- [104] A. Latifi, M. Foglino, K. Tanaka, P. Williams and A. Lazdunski. A hierarchical quorum-sensing cascade in *Pseudomonas aeruginosa* links the transcriptional activators LasR and RhlR (VsmR) to expression of the stationary-phase sigma factor RpoS. *Molecular Microbiology*, 21(6):1137–1146. 1996.
- [105] L. A. Gallagher, S. L. McKnight, M. S. Kuznetsova, E. C. Pesci and C. Manoil. Functions required for extracellular quinolone signaling by *Pseudomonas aeruginosa*. *Journal of Bacteriology*, 184(23):6472–6480. 2002.
- [106] D. S. Wade, M. W. Calfee, E. R. Rocha, E. A. Ling, E. Engstrom, J. P. Coleman and E. C. Pesci. Regulation of *Pseudomonas* Quinolone Signal Synthesis in *Pseudomonas aeruginosa*. 187(13):4372–4380. 2005.
- [107] J. M. Brint and D. E. Ohman. Synthesis of multiple exoproducts in *Pseudomonas aeruginosa* is under the control of RhlR-RhlI, another set of regulators in strain PAO1 with homology to the autoinducer-responsive LuxR-LuxI family. *Journal of Bacteriology*, 177(24):7155–7163. 1995.
- [108] J. P. Pearson, L. Passador, B. H. Iglewski and E. P. Greenberg. A second N-acylhomoserine lactone signal produced by *Pseudomonas aeruginosa*. *Proceedings of the National Academy of Sciences*, 92(5):1490–1494. 1995.
- [109] M. K. Winson, M. Camara, A. Latifi, M. Foglino, S. R. Chhabra, M. Daykin, M. Bally, V. Chapon, G. P. Salmond and B. W. Bycroft. Multiple N-acyl-L-homoserine lactone signal molecules regulate production of virulence determinants and secondary metabolites in *Pseudomonas aeruginosa*. *Proceedings of the National Academy of Sciences*, 92(20):9427–9431. 1995.
- [110] A. Latifi, M. K. Winson, M. Foglino, B. W. Bycroft, G. S. A. B. Stewart, A. Lazdunski and P. Williams. Multiple homologues of LuxR and LuxI control expression of virulence determinants and secondary metabolites through quorum sensing in *Pseudomonas aeruginosa* PAO1. *Molecular Microbiology*, 17(2):333–343. 1995.

- [111] K. Winzer, C. Falconer, N. C. Garber, S. P. Diggle, M. Camara and P. Williams. The *Pseudomonas aeruginosa* Lectins PA-IL and PA-IIL Are Controlled by Quorum Sensing and by RpoS. 182(22):6401–6411. 2000.
- [112] S. McGrath, D. S. Wade and E. C. Pesci. Dueling quorum sensing systems in *Pseudomonas aeruginosa* control the production of the *Pseudomonas* quinolone signal (PQS). 230:0–7. 2004.
- [113] S. L. McKnight, B. H. Iglewski and E. C. Pesci. The *Pseudomonas* Quinolone Signal Regulates *rhl* Quorum Sensing in *Pseudomonas aeruginosa*. 182(10):2702–2708. 2000.
- [114] E. C. Pesci, J. B. J. Milbank, J. P. Pearson, S. McKnight, A. S. Kende, E. P. Greenberg and B. H. Iglewski. Quinolone signaling in the cell-to-cell communication system of *Pseudomonas aeruginosa*. *Proceedings of the National Academy of Sciences*, 96(20):11229–11234. 1999.
- [115] J. M. Farrow and E. C. Pesci. Two distinct pathways supply anthranilate as a precursor of the *Pseudomonas* quinolone signal. *Journal of Bacteriology*, 189(9):3425–3433. 2007.
- [116] F. Lépine, E. Déziel, S. Milot and L. Rahme. A stable isotope dilution assay for the quantification of the *Pseudomonas* quinolone signal in *Pseudomonas aeruginosa* cultures. *Biochimica et Biophysica Acta (BBA) - General Subjects*, 1622(1):36–41. 2003.
- [117] F. Lépine, S. Milot, E. Déziel, J. He and L. G. Rahme. Electrospray/mass spectrometric identification and analysis of 4-hydroxy-2-alkylquinolines (HAQs) produced by *Pseudomonas aeruginosa*. *Journal of the American Society for Mass Spectrometry*, 15(6):862–869. 2004.
- [118] S. L. Drees and S. Fetzner. PqsE of *Pseudomonas aeruginosa* acts as pathway-specific thioesterase in the biosynthesis of alkylquinolone signaling molecules. *Chemistry & Biology*, 22(5):611–618. 2015.
- [119] J. Lin, J. Cheng, Y. Wang and X. Shen. The *Pseudomonas* Quinolone Signal (PQS): Not Just for Quorum Sensing Anymore. 8(July):1–9. 2018.
- [120] G. Xiao, E. Déziel, J. He, F. Lépine, B. Lesic, M.-H. Castonguay, S. Milot, A. P. Tampakaki, S. E. Stachel and L. G. Rahme. MvfR, a key *Pseudomonas aeruginosa* pathogenicity LTTR-class regulatory protein, has dual ligands. *Molecular Microbiology*, 62(6):1689–99. 2006.
- [121] E. Déziel, S. Gopalan, A. P. Tampakaki, F. Lépine, K. E. Padfield, M. Saucier, G. Xiao and L. G. Rahme. The contribution of MvfR to *Pseudomonas aeruginosa* pathogenesis and quorum sensing circuitry regulation : multiple quorum sensing-regulated genes are modulated without affecting lasRI , rhlRI or the production of N -acyl- L -homoserine lactones. 55:998–1014. 2005.
- [122] S. P. Diggle, K. Winzer, S. R. Chhabra, K. E. Worrall, M. Cámara and P. Williams. The *Pseudomonas aeruginosa* quinolone signal molecule overcomes the cell density-dependency of the quorum sensing hierarchy, regulates *rhl*-dependent genes at the onset of stationary phase and can be produced in the absence of LasR. *Molecular Microbiology*, 50(1):29–43. 2003.
- [123] S. P. Diggle, S. Matthijs, V. J. Wright, M. P. Fletcher, S. R. Chhabra, I. L. Lamont, X. Kong, R. C. Hider, P. Cornelis, M. Cámara and P. Williams. The *Pseudomonas aeruginosa* 4-Quinolone Signal Molecules HHQ and PQS Play Multifunctional Roles in Quorum Sensing and Iron Entrapment. *Chemistry and Biology*, 14(January):87–96. 2007.
- [124] L. Mashburn-Warren, J. Howe, K. Brandenburg and M. Whiteley. Structural requirements of the *Pseudomonas* quinolone signal for membrane vesicle stimulation. *Journal of Bacteriology*, 191(10):3411–3414. 2009.

- [125] C. S. Pereira, J. A. Thompson and K. B. Xavier. AI-2-mediated signalling in bacteria. *FEMS Microbiology Reviews*, 37(2):156–181. 2013.
- [126] H. Li, X. Li, Z. Wang, Y. Fu, Q. Ai, Y. Dong and J. Yu. Autoinducer-2 regulates *Pseudomonas aeruginosa* PAO1 biofilm formation and virulence production in a dose-dependent manner. *BMC Microbiology*, pages 1–8. 2015.
- [127] H. Li, X. Li, C. Song, Y. Zhang and Z. Wang. Autoinducer-2 Facilitates *Pseudomonas aeruginosa* PAO1 Pathogenicity in Vitro and in Vivo. 8(October):1–9. 2017.
- [128] W. R. J. D. Galloway, J. T. Hodgkinson, S. D. Bowden, M. Welch and D. R. Spring. Quorum sensing in Gram-negative bacteria: small-molecule modulation of AHL and AI-2 quorum sensing pathways. *Chemical Reviews*, 111(1):28–67. 2011.
- [129] C. Florez, J. E. Raab, A. C. Cooke and J. W. Schertzer. Membrane Distribution of the *Pseudomonas* Quinolone Signal Modulates Outer Membrane Vesicle Production in *Pseudomonas aeruginosa*. pages 1–13. 2017.
- [130] J. P. Pearson, C. Van Delden and B. H. Iglewski. Active Efflux and Diffusion Are Involved in Transport of *Pseudomonas aeruginosa* Cell-to-Cell Signals. *J Bacteriol*, 181(4):1203–1210. 1999.
- [131] K. Evans, L. Passador, R. Srikumar, E. Tsang, J. Nezezon and K. Poole. Influence of the MexAB-OprM multidrug efflux system on quorum sensing in *Pseudomonas aeruginosa*. *Journal of Bacteriology*, 180(20):5443–5447. 1998.
- [132] D. G. Davies, M. R. Parsek, J. P. Pearson, B. H. Iglewski, J. W. Costerton and E. P. Greenberg. The involvement of cell-to-cell signals in the development of a bacterial biofilm. *Science*, 280(5361):295–298. 1998.
- [133] C. M. Oliphant and G. M. Green. Quinolones: a comprehensive review. *American Family Physician*, 65(3):455–464. 2002.
- [134] K. Drlica and X. Zhao. DNA Gyrase , Topoisomerase IV , and the 4-Quinolones. 61(3):377–392. 1997.
- [135] R. N. Brogden, A. A. Carmine, R. C. Heel, T. M. Speight and G. S. Avery. Trimethoprim: A Review of its Antibacterial Activity, Pharmacokinetics and Therapeutic Use in Urinary Tract Infections. *Drugs*, 23(6):405–430. 1982.
- [136] R. A. Celesk and N. J. Robillard. Factors Influencing the Accumulation of Ciprofloxacin in *Pseudomonas aeruginosa*. 33(11):1921–1926. 1989.
- [137] K. Poole. MINIREVIEW Efflux-Mediated Resistance to Fluoroquinolones in Gram-Negative Bacteria. 44(9):2233–2241. 2000.
- [138] T. R. De Kievit, M. D. Parkins, R. J. Gillis, R. Srikumar, H. Ceri, K. Poole, B. H. Iglewski and D. G. Storey. Multidrug Efflux Pumps: Expression Patterns and Contribution to Antibiotic Resistance in *Pseudomonas aeruginosa* Biofilms. 45(6):1761–1770. 2001.
- [139] T. Köhler, M. Kok, M. Michea-hamzehpour, P. Plesiat, N. Gotoh, T. Nishino, L. K. Curty and J.-c. Pechere. Multidrug Efflux in Intrinsic Resistance to Trimethoprim and Sulfamethoxazole in *Pseudomonas aeruginosa*. 40(10):2288–2290. 1996.
- [140] K. Poole, N. Gotoh, H. Tsujimoto, Q. Zhao, A. Wada, T. Yamasaki, S. Neshat, J.-i. Yamagishi, X.-Z. Li and T. Nishino. Overexpression of the *mexC-mexD-oprJ* efflux operon in *nfxB*-type multidrug-resistant strains of *Pseudomonas aeruginosa*. *Molecular Microbiology*, 21(4):713–725. 1996.

- [141] T. Köhler, M. Michéa-Hamzehpour, U. Henze, N. Gotoh, L. K. Curty and J. C. Pechère. Characterization of MexE-MexF-OprN, a positively regulated multidrug efflux system of *Pseudomonas aeruginosa*. *Molecular Microbiology*, 23(2):345–354. 1997.
- [142] K. Poole. Multidrug Efflux Pumps and Antimicrobial Resistance in *Pseudomonas aeruginosa* and Related Organisms JMMB Symposium. *J. Mol. Microbiol. Biotechnol*, 3(2):255–264. 2001.
- [143] K. Ganguly, R. Wu, M. Ollivault-Shiflett, P. M. Goodwin, L. A. Silks and R. Iyer. Design, synthesis, and a novel application of quorum-sensing agonists as potential drug-delivery vehicles. *Journal of Drug Targeting*, 19(7):528–539. 2011.
- [144] R. Iyer, K. Ganguly and L. A. Silks. Synthetic analogs of bacterial quorum sensors. Los Alamos National Laboratory. 2012.
- [145] A. Eberhard, C. A. Widrig, P. Mcbath and J. B. Schineller. Analogs of the autoinducer of bioluminescence in *Vibrio fischeri*. *Archives of Microbiology*, 146(1):35–40. 1986.
- [146] L. Passador, K. D. Tucker, K. R. Guertin, M. P. Journet, A. S. Kende and B. H. Iglewski. Functional analysis of the *Pseudomonas aeruginosa* autoinducer PAI. *Journal of Bacteriology*, 178(20):5995–6000. 1996.
- [147] K. M. Smith, Y. Bu and H. Suga. Library Screening for Synthetic Agonists and Antagonists of a *Pseudomonas aeruginosa* Autoinducer. *Chemistry & Biology*, 10(6):563–571. 2003.
- [148] S. R. Chhabra, P. Stead, N. J. Bainton, G. P. Salmond, G. S. Stewart, P. Williams and B. W. Bycroft. Autoregulation of carbapenem biosynthesis in *Erwinia carotovora* by analogues of N-(3-oxohexanoyl)-L-homoserine lactone. *The Journal of Antibiotics*, 46(3):441–454. 1993.
- [149] C. E. McInnis and H. E. Blackwell. Thiolactone modulators of quorum sensing revealed through library design and screening. *Bioorganic & Medicinal Chemistry*, 19(16):4820–4828. 2011.
- [150] G. D. Geske, J. C. O. Neill, D. M. Miller, M. E. Mattmann and H. E. Blackwell. Modulation of Bacterial Quorum Sensing with Synthetic Ligands : Systematic Evaluation of N-Acylated Homoserine Lactones in Multiple Species and New Insights into Their Mechanisms of Action. *Journal of the American Chemical Society*, 129(44):13613–13625. 2007.
- [151] J. C. A. Janssens, K. Metzger, R. Daniels, D. Ptacek, T. Verhoeven, L. W. Habel, J. Vanderleyden, D. E. De Vos and S. C. J. De Keersmaecker. Synthesis of N -Acyl Homoserine Lactone Analogues Reveals Strong Activators of SdiA , the *Salmonella enterica* Serovar. *Applied and Environmental Microbiology*, 73(2):535–544. 2007.
- [152] J. T. Hodgkinson, W. R. J. D. Galloway, M. Wright, I. K. Mati, R. L. Nicholson, M. Welch and D. R. Spring. Design, synthesis and biological evaluation of non-natural modulators of quorum sensing in *Pseudomonas aeruginosa*. *Organic & Biomolecular Chemistry*, 10(30):6032. 2012.
- [153] M. E. Boursier, D. E. Manson, J. B. Combs, E. Helen and H. E. Blackwell. A comparative study of non-native N-acyl L-homoserine lactone analogs in two *Pseudomonas aeruginosa* quorum sensing receptors that share a common native ligand yet inversely regulate virulence. *Bioorganic & Medicinal Chemistry*, pages 1–17. 2018.
- [154] K. M. Smith, Y. Bu and H. Suga. Induction and Inhibition of *Pseudomonas aeruginosa* Quorum Sensing by Synthetic Autoinducer Analogs. *Chemistry & Biology*, 10(1):81–89. 2003.

- [155] G. J. Jog, J. Igarashi and H. Suga. Stereoisomers of *P. aeruginosa* Autoinducer Analog to Probe the Regulator Binding Site. *Chemistry and Biology*. 2006.
- [156] C. Lu, B. Kirsch, C. Zimmer, J. C. De Jong, C. Henn, C. K. Maurer, M. Müsken, S. Häussler, A. Steinbach and R. W. Hartmann. Discovery of antagonists of PqsR, a key player in 2-alkyl-4-quinolone- dependent quorum sensing in *Pseudomonas aeruginosa*. *Chemistry and Biology*, 19(3):381–390. 2012.
- [157] C. Lu, C. K. Maurer, B. Kirsch, A. Steinbach and R. W. Hartmann. Overcoming the unexpected functional inversion of a PqsR antagonist in *Pseudomonas aeruginosa*: An in vivo potent antivirulence agent targeting pqs quorum sensing. *Angewandte Chemie - International Edition*, 53(4):1109–1112. 2014.
- [158] J. Hodgkinson, S. D. Bowden, W. R. J. D. Galloway, D. R. Spring and M. Welch. Structure-activity analysis of the *Pseudomonas* quinolone signal molecule. *Journal of Bacteriology*, 192(14):3833–3837. 2010.
- [159] Y. R. Baker. Investigating quinolone based quorum sensing in *Pseudomonas aeruginosa* using a chemical proteomics approach. PhD thesis, University of Cambridge. 2015.
- [160] D. M. Stacy, S. T. Le Quement, C. L. Hansen, J. W. Clausen, T. Tolker-Nielsen, J. W. Brummond, M. Givskov, T. E. Nielsen and H. E. Blackwell. Synthesis and biological evaluation of triazole-containing N-acyl homoserine lactones as quorum sensing modulators. *Organic & Biomolecular Chemistry*, 11(6):938–954. 2013.
- [161] T. E. Renau, J. P. Sanchez, J. W. Gage, J. A. Dever, M. A. Shapiro, S. J. Gracheck and J. M. Domagala. Structure-activity relationships of the quinolone antibacterials against mycobacteria: effect of structural changes at N-1 and C-7. *Journal of Medicinal Chemistry*, 39(3):729–735. 1996.
- [162] C. Jing and V. W. Cornish. A fluorogenic TMP-tag for high signal-to-background intracellular live cell imaging. *ACS Chemical Biology*, 8(8):1704–12. 2013.

Todo list

swarming needs to be mentioned elsewhere	5
analogues,cleavables,bio	5
strategies to get new antibiotics, inc sideromycins and synthetic	10
link in better	10
pre-release paper	14
intro QS/AIs	16
better quality diagrams	17
add numbers manually when sorted	19
link	20
Boursier2018 pre-release, check for updates	24
aims here	25
make this actually an overview	27