



0.1 Cyclopentyl alcohol derivatives

0.1.1 Head-group synthesis




Scheme 1: a) AlMe_3 , CH_2Cl_2 , $0\text{ }^\circ\text{C}$. **113** : 32.1 %, **112** : 35.2 %. b) $\text{Pd}(\text{OH})_2$, MeOH, H_2 , 5 atm, r.t., 1 d, **114** : 100 %, **115** : 100 %.

0.1.2 Initial branching strategy




Scheme 2

Failed. Lots of side-products.




Scheme 3

0.1.3 TBS protection strategy



Scheme 4

Want to protect alcohol to stop side reactions




Scheme 5

Protection optimisation


Still get side-reactions when adding tail

0.1.4 Triazoles by two-step reaction

Talk about moving to two-step reaction.



Scheme 6




Scheme 7

Did click, then failed to deprotect.


0.1.5 Attaching the linker to ciprofloxacin first

Given the side-reactions and low yields associated with the literature synthesis of the S_N2 conjugates proposed by Ganguly et. al,² we investigated a second synthesis, building up the linker on the ciprofloxacin side before coupling with the head group (see ??).




Scheme 8

0.1.5.1 Synthesis of methyl-protected ciprofloxacin with linker with terminal carboxylate




Scheme 9: Synthesis of **130**. a) .


0.1.6 Triazoles from the chloride



Scheme 10




Scheme 11



Scheme 12

This worked.

0.2 Methyl 1-cyclopropyl-6-fluoro-4-oxo-7-(piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylate **92**



Ciprofloxacin **28** (10.0 g, 30 mmol, 1 eq.) and *p*-toluenesulfonic acid (8.60 mg, 44.5 mmol, 1.5 eq.) were refluxed in methanol (500 ml) for 72 h. The mixture was cooled to room temperature and NaHCO₃ (sat., aq., 100 ml) and water (300 ml) were added. The product was extracted with CH₂Cl₂ (2×400 ml). The combined organic fractions were dried over MgSO₄ and evaporated under reduced pressure. **92** was obtained as a white amorphous solid (9.16 g, 26.5 mmol, 83.3 %).

TLC R_f = 0.13 (5 % MeOH/CH₂Cl₂)

IR (neat) ν_{max} / cm⁻¹ = 2947.9 (C-H), 2834.9 (C-H), 1720.9 (ester C=O), 1616.8 (quinolone C=O)

¹H NMR (400 MHz, MeOD) δ / ppm = 8.55 (s, 1 H, *ortho* to C(=O)OCH₃), 7.71 (d, J = 13.5 Hz, 1 H, *ortho* to F), 7.41 (d, J = 7.2 Hz, 1 H, *meta* to F), 3.83 (s, 3 H, CH₃), 3.62 (tt, J = 7.4, 3.5 Hz, 1 H, NCH(CH₂)₂), 3.24 - 3.29 (m, 4 H, HN(CH₂CH₂)CH₂CH₂), 3.02 - 3.10 (m, 4 H, HN(CH₂)CH₂), 1.31 - 1.38 (m, 2 H, NCH(CH₂)₂), 1.12 - 1.20 (m, 2 H, NCH(CH₂)₂)


¹³C NMR (101 MHz, MeOD) δ / ppm = 175.2 (C(=O)CC(=O)OCH₃), 166.8 (C(=O)OCH₃), 154.9 (d, J = 248.0 Hz, *ipso* to F), 150.1 (C=CC(=O)OCH₃), 146.6 (d, J = 10.4 Hz, *ipso* to piperazine), 139.9 (*para* to F), 123.3 (d, J = 6.9 Hz, *para* to piperazine), 113.0 (d, J = 23.4 Hz, *ortho* to C=O and *ortho* to F), 110.1 (CC(=O)OCH₃), 107.1 (d, J = 3.5 Hz, *meta* to C=O and *meta* to F), 52.3 (CH₃), 51.7 (HN(CH₂CH₂)CH₂CH₂), 51.6 (HN(CH₂CH₂)CH₂CH₂), 46.5 (HN(CH₂)CH₂), 36.4 (NCH(CH₂)₂), 8.7 (NCH(CH₂)₂)

¹⁹F NMR (376.45 MHz, MeOD) δ / ppm = -124.8 (s, ciprofloxacin F)

HRMS (ESI⁺) m/z / Da = 346.1569, [M+H]⁺ found, [C₁₈H₂₁FN₃O₃]⁺ requires 346.1567

The data are consistent with the literature.¹

0.3 4-Bromo-*N*-(2-oxotetrahydrothiophen-3-yl)butanamide **94**



3-Aminodihydrothiophen-2(3*H*)-one hydrochloride **93** (15.0 g, 97.6 mmol, 1 eq.) and NaHCO₃ (16.4 g, 195 mmol, 2 eq.) were added to CH₂Cl₂ (150 ml) and water (150 ml). 4-Bromobutyryl chloride **58** (11.3 ml, 107 mmol, 1.1 eq.) was added dropwise over 45 min at 0 °C and the mixture was stirred for a further 1 h. The

organic layer was separated and the aqueous layer was extracted with a second portion of CH₂Cl₂ (150 ml). The combined organic layers were dried over MgSO₄ and evaporated under reduced pressure. **94** was obtained as a white, amorphous solid (22.7 g, 85.8 mmol, 87.9 %).

TLC R_f = 0.19 (50 % EtOAc/PE)

IR (neat) ν_{max} / cm⁻¹ = 3265.9 (amide N-H), 3063.2 (amide N-H), 1694.3 (thiolactone C=O), 1650.5 (amide C=O)


¹H NMR (400 MHz, CDCl₃) δ / ppm = 6.08 (d, J = 6.1 Hz, 1 H, NH), 4.54 (dt, J = 12.9, 6.5 Hz, 1 H, CHNH), 3.49 (t, J = 6.4 Hz, 2 H, CH₂Br), 3.37 (ddd, J = 12.2, 11.5, 5.3 Hz, 1 H, SCHH), 3.26 (ddd, J = 11.5, 6.9, 1.3 Hz, 1 H, SCHH), 2.91 (dddd, J = 12.5, 6.7, 5.3, 1.3 Hz, 1 H, SCH₂CHH), 2.45 (t, J = 7.4 Hz, 1 H, C(=O)CHH), 2.45 (t, J = 6.8 Hz, 1 H, C(=O)CHH), 2.20 (quin, J = 6.7 Hz, 1 H, C(=O)CH₂CH₂), 1.96 (dddd, J = 12.7, 12.5, 12.2, 7.0 Hz, 1 H, SCH₂CHH)

¹³C NMR (101 MHz, CDCl₃) δ / ppm = 205.4 (SC(=O)), 172.1 (NHC(=O)), 59.4 (CHNH), 34.1 (C(=O)CH₂), 33.1 (CH₂Br), 31.8 (SCH₂CH₂), 28.0 (C(=O)CH₂CH₂), 27.5 (SCH₂)

HRMS (ESI⁺) The compound does not ionise.

The compound has been synthesised previously^{2,3} but characterisation was not published.

0.4 Methyl 1-cyclopropyl-6-fluoro-4-oxo-7-(4-(4-oxo-4-((2-oxotetrahydrothiophen-3-yl)amino)butyl)piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylate **95**



Methyl 1-cyclopropyl-6-fluoro-4-oxo-7-(piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylate **92** (50 mg, 0.145 mmol, 1 eq.), 4-bromo-*N*-(2-oxotetrahydrothiophen-3-yl)butanamide **94** (34.5 mg, 0.145 mmol, 1 eq.) and K₂CO₃ (20 mg, 0.145 mmol, 1 eq.) were stirred in acetonitrile (2 ml) at 50 °C under argon. After 24 h a further portion of **94** (34.5 mg, 0.145 mmol, 1 eq.) was added. After another 24 h a further portion was added (69.0 mg, 0.290 mmol, 2 eq.). After another 24 h the temperature was raised so the mixture was at reflux. After a final 24 h the precipitate was filtered off and the filtrate was purified by column chromatography (SiO₂, 5-10 % MeOH/CH₂Cl₂) followed by preparatory HPLC (5-95 % acetonitrile/water over 20 min). **95** was obtained as a cream-coloured amorphous solid (9.4 mg, 0.018 mmol, 12.2 %).

TLC R_f = 0.47 (10 % MeOH/CH₂Cl₂)

IR (neat) ν_{max} / cm⁻¹ = 2944.2 (C-H), 2832.4 (C-H), 1722.4 (ester C=O), 1700.4 (thiolactone C=O), 1669.6 (amide C=O), 1617.3 (quinolone C=O)

¹H NMR (500 MHz, MeOD) δ / ppm = 8.53 (s, 1 H, *ortho* to C(=O)OCH₃), 7.68 (d, J=13.4 Hz, 1 H, *ortho* to F), 7.41 (d, J=7.3 Hz, 1 H, *meta* to F), 4.67 (dd, J=12.9, 6.9 Hz, 1 H, CHNH), 3.83 (s, 3 H, OCH₃), 3.61 (tt, J=6.9, 4.1 Hz, 1 H, NCH(CH₂)₂), 3.39 - 3.49 (m, 5 H, SCHH), 3.26 - 3.33 (m, 1 H, SCHH and CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 2.93 - 3.03 (m, 4 H, CH₂CH₂CH₂N(CH₂)CH₂), 2.79 (br. t, J=7.2, 7.2 Hz, 2 H, C(=O)CH₂CH₂CH₂), 2.59 (dddd, J=12.4, 6.9, 5.4, 1.4 Hz, 1 H, SCH₂CHH), 2.39 (t, J=7.20 Hz, 1 H, C(=O)CHH), 2.38 (t, J=6.94 Hz, 1 H, C(=O)CHH), 2.18 (qd, J=12.4, 7.0 Hz, 1 H, SCH₂CHH), 1.97 (quin, J=7.2 Hz, 2 H, C(=O)CH₂CH₂), 1.32 - 1.37 (m, 2 H, NCH(CHH)₂), 1.13 - 1.19 (m, 2 H, NCH(CHH)₂)


¹³C NMR (126 MHz, MeOD) δ / ppm = 207.0 (SC(=O)), 175.7 (NHC(=O)), 175.1 (C(=O)CC(=O)OCH₃), 166.6 (C(=O)OCH₃), 154.7 (d, J=249.0 Hz, *ipso* to F), 150.2 (s, CH=CC(=O)OCH₃), 145.6 (d, J=10.6 Hz, *ipso* to piperazine), 139.8 (*para* to F), 123.5 (d, J=6.9 Hz, *para* to piperazine), 113.1 (d, J=23.6 Hz, *ortho* to C=O and *ortho* to F), 110.0 (CC(=O)OCH₃), 107.4 (*meta* to C=O and *meta* to F), 60.2 (CHNH), 58.5 (C(=O)CH₂CH₂CH₂), 53.8 (CH₂CH₂CH₂N(CH₂)CH₂), 52.3 (OCH₃), 50.1 (CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 50.0 (CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 36.5 (NCH(CH₂)₂), 34.5 (C(=O)CH₂), 31.7 (SCH₂CH₂), 28.1 (SCH₂), 22.9 (C(=O)CH₂CH₂CH₂), 8.7 (NCH(CH₂)₂)

¹⁹F NMR (376.45 MHz, MeOD) δ / ppm = -125.4 (s, ciprofloxacin F)

HRMS (ESI⁺) m/z / Da = 531.2083, [M+H]⁺ found, [C₂₆H₃₂FN₄O₅S]⁺ requires 531.2077

The compound has been synthesised previously.^{2,3} Only HRMS characterisation was published, and this agrees with the result above.

0.5 4-Azido-*N*-(2-oxotetrahydrothiophen-3-yl)butanamide **96**



4-Bromo-*N*-(2-oxotetrahydrothiophen-3-yl)butanamide **94** (6.00 g, 27.0 mmol, 1 eq.) and NaN₃ (3.51 g, 54.1 mmol, 2 eq.) were refluxed in acetonitrile (120 ml) for 1.5 h. The solvent was evaporated under reduced pressure and the residue was partitioned between water (150 ml) and CH₂Cl₂ (150 ml). The aqueous layer was extracted twice more with CH₂Cl₂ (2×150 ml) and the combined organic fractions were dried with MgSO₄ and evaporated under reduced pressure. **96** was obtained as a yellow, sticky solid (4.60 g, 20.1 mmol, 89.3 %).

TLC R_f = 0.19 (50 % EtOAc/PE)

IR (neat) ν_{max} / cm⁻¹ = 3285.6 (N-H), 2963.9 (C-H), 2100.2 (azide), 1697.4 (thiolactone C=O), 1647.4 (amide C=O)


¹H NMR (400 MHz, CDCl₃) δ / ppm = 6.71 (d, J = 7.3 Hz, 1 H, NH), 4.54 (dt, J = 13.0, 7.0 Hz, 1 H, CHNH), 3.30 (t, J = 6.7 Hz, 2 H, CH₂N₃), 3.31 (td, J = 11.7, 5.3 Hz, 1 H, 1 H, SCHH), 3.19 (ddd, J = 11.3, 7.0, 1.2 Hz, 1 H, SCHH), 2.70 (dddd, J = 12.4, 6.8, 5.3, 1.2 Hz, 1 H, SCH₂CHH), 2.29 (t, J = 7.5 Hz, 1 H, C(=O)CHH), 2.28 (t, J = 7.1 Hz, 1 H, C(=O)CHH), 1.97 (qd, J = 12.4, 7.0 Hz, 1 H, SCH₂CHH), 1.85 (quin, J = 6.9 Hz, 2 H, C(=O)CH₂CH₂)

^{13}C NMR (101 MHz, CDCl_3) δ / ppm = 205.4 ($\text{SC}(=\text{O})$), 172.3 ($\text{NHC}(=\text{O})$), 59.4 (CHNH), 50.6 (CH_2N_3), 32.8 ($\text{C}(=\text{O})\text{CH}_2$), 31.8 (SCH_2CH_2), 27.5 (SCH_2), 24.6 ($\text{C}(=\text{O})\text{CH}_2\text{CH}_2$)

HRMS (ESI^+) m/z / Da = 251.0565, $[\text{M}+\text{Na}]^+$ found, $[\text{C}_8\text{H}_{12}\text{N}_4\text{NaO}_2\text{S}]^+$ requires 251.0573

The compound has not been reported previously.

0.6 1-Cyclopropyl-6-fluoro-4-oxo-7-(4-(4-(1-(4-oxo-4-((2-oxotetrahydrothiophen-3-yl)amino)butyl)-1*H*-1,2,3-triazol-4-yl)butyl)piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylic acid **97**



1-Cyclopropyl-6-fluoro-7-(4-(hex-5-yn-1-yl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylic acid **70** (15 mg, 36.7 μmol , 1 eq.) and 4-azido-*N*-(2-oxotetrahydrothiophen-3-yl)butanamide **96** (12.5 mg, 55.1 μmol , 1.5 eq.) were dissolved in 1:9:10 water/*t*-BuOH/DMSO (3 ml), and the mixture was degassed by bubbling N_2 through it. A solution of CuSO_4 and THPTA (182 μl , 18.2 μmol , 0.5 eq. 100 mM, aq.) was added, followed by a solution of sodium ascorbate (367 μl , 36.7 μmol , 1 eq., 100 mM, aq.). The mixture was stirred at r.t. under argon for 4 d. Water (10 ml) and 10 % *i*-PrOH/ CHCl_3 (10 ml) were added, the organic layer was separated and the aqueous layer was extracted again with 10 % *i*-PrOH/ CHCl_3 (2×10 ml). The combined organic layers were dried with MgSO_4 and evaporated under reduced pressure. The residue was purified by preparatory HPLC (5-95 % acetonitrile/water over 20 min). The combined pure fractions were evaporated under reduced pressure and then partitioned between NaHCO_3 (aq., sat., 50 ml) and 10 % *i*-PrOH/ CHCl_3 (50 ml). The organic layer was dried with MgSO_4 and evaporated under reduced pressure. **97** was obtained as a white amorphous solid (16.5 mg, 25.9 μmol , 70.6 %).

IR (neat) ν_{max} / cm^{-1} = 2918.8 (C-H), 1712.7 (carboxylic acid C=O and thiolactone C=O), 1657.6 (amide C=O), 1626.8 (quinolone C=O), 1616.2 (triazole)

^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ / ppm = 15.23 (br s, 1 H, $\text{C}(=\text{O})\text{OH}$), 8.66 (s, 1 H, *ortho* to $\text{C}(=\text{O})\text{OH}$), 8.23 (d, $J=8.5$ Hz, 1 H, NH), 7.90 (d, $J=13.4$ Hz, 1 H, *ortho* to F), 7.84 (s, 1 H, $\text{CH}=\text{CCH}_2$), 7.56 (d, $J=7.5$ Hz, 1 H, *meta* to F), 4.59 (ddd, $J=12.7, 8.4, 6.8$ Hz, 1 H, CHNH), 4.31 (t, $J=7.0$ Hz, 2 H, $\text{CH}_2\text{NCH}=\text{C}$), 3.80 - 3.86 (6.9, 4.0 Hz, 1 H, $\text{NCH}(\text{CH}_2)_2$), 3.34 - 3.37 (m, 1 H, SCHH), 3.32 (br t, $J=4.1$ Hz, 4 H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2\text{CH}_2$), 3.27 (ddd, $J=11.1, 6.9, 1.4$ Hz, 1 H, SCHH), 2.64 (t, $J=7.6$ Hz, 2 H, $\text{CH}=\text{CCH}_2$), 2.57 (br t, $J=4.7$ Hz, 4 H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2)\text{CH}_2$), 2.34 - 2.44 (m, 3 H, SCH_2CHH and $\text{CH}=\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_2$), 2.12 (t, $J=7.9$ Hz, 1 H, $\text{C}(=\text{O})\text{CHH}$), 2.12 (t, $J=7.0$ Hz, 1 H, $\text{C}(=\text{O})\text{CHH}$), 2.04 (m, 3 H, SCH_2CHH and $\text{C}(=\text{O})\text{CH}_2\text{CH}_2$), 1.64 (quin, $J=7.5$ Hz, 2 H, $\text{CH}=\text{CCH}_2\text{CH}_2$), 1.51 (quin, $J=7.5$ Hz, 2 H, $\text{CH}=\text{CCH}_2\text{CH}_2\text{CH}_2$), 1.28 - 1.34 (m, 2 H, $\text{NCH}(\text{CHH})_2$), 1.15 - 1.20 (m, 2 H, $\text{NCH}(\text{CHH})_2$)


^{13}C NMR (126 MHz, $\text{DMSO}-d_6$) δ / ppm = 205.6 ($\text{SC}(=\text{O})$), 176.4 ($\text{C}(=\text{O})\text{CC}(=\text{O})\text{OH}$), 171.4 ($\text{NHC}(=\text{O})$), 166.0 ($\text{C}(=\text{O})\text{OH}$), 153.1 (d, $J=249.3$ Hz, *ortho* to F), 148.0 ($\text{CH}=\text{CC}(=\text{O})\text{OH}$), 146.9 ($\text{CH}=\text{CCH}_2$), 145.3

(d, $J=10.1$ Hz, *ipso* to piperazine), 139.2 (*para* to F), 121.8 ($\text{CH}=\text{CCH}_2$), 118.6 (d, $J=7.7$ Hz, *para* to piperazine), 111.0 (d, $J=23.3$ Hz, *ortho* to C=O and *ortho* to F), 106.7 ($\text{CC}(=\text{O})\text{OH}$), 106.4 (d, $J=2.9$ Hz, *meta* to C=O and *meta* to F), 58.2 ($\text{SC}(=\text{O})\text{CHNH}$), 57.4 ($\text{CH}=\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 52.4 ($\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2$), 49.5 ($\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2\text{CH}_2$), 49.5 ($\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2\text{CH}_2$), 48.6 ($\text{CH}_2\text{NCH}=\text{C}$), 35.9 ($\text{NCH}(\text{CH}_2)_2$), 31.9 ($\text{NHC}(=\text{O})\text{CH}_2$), 30.1 (CH_2CHNH), 26.9 ($\text{CH}=\text{CCH}_2\text{CH}_2$), 26.8 (SCH_2), 25.9 ($\text{NHC}(=\text{O})\text{CH}_2\text{CH}_2$), 25.8 ($\text{CH}=\text{CCH}_2\text{CH}_2\text{CH}_2$), 25.0 ($\text{CH}=\text{CCH}_2$), 7.6 ($\text{NCH}(\text{CH}_2)_2$)

^{19}F NMR (376.45 MHz, MeOD) δ / ppm = -124.9 (s, ciprofloxacin F)

HRMS (ESI^+) m/z / Da = 640.2739, $[\text{M}+\text{H}]^+$ found, $[\text{C}_{31}\text{H}_{39}\text{FN}_7\text{O}_5\text{S}]^+$ requires 640. 2712

0.7 1-Cyclopropyl-6-fluoro-4-oxo-7-(4-(((4-(1-(4-oxo-4-((2-oxotetrahydrothiophen-3-yl)amino)butyl)-1*H*-1,2,3-triazol-4-yl)butanoyl)oxy)methoxy)carbonyl)piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylic acid **99**



1-Cyclopropyl-6-fluoro-7-(4-(((hex-5-ynoyloxy)methoxy)carbonyl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylic acid **98** (203 mg, 0.407 mmol, 1 eq.), 4-azido-*N*-(2-oxotetrahydrothiophen-3-yl)butanamide **96** (92.8 mg, 0.407 mmol, 1 eq.), CuI (40 mg, 0.190 mmol, 0.5 eq.) and DIPEA (0.356 ml, 0.264 mg, 2.04 mmol, 5 eq.) were stirred in CH_2Cl_2 (18.6 ml) at r.t. under Ar for 3 h. The mixture was filtered and the filtrate was dry-loaded onto SiO_2 and purified by column chromatography (SiO_2 , 5-10 % MeOH/ CH_2Cl_2). **99** was obtained as pale brown/yellow amorphous solid (14.7 mg, 20.2 μmol , 5.0 %).

TLC R_f = 0.40 (5 % CH_2Cl_2 /MeOH)

IR (neat) ν_{max} / cm^{-1} = 3054.9 (C-H), 1715.8 (carboxylic acid C=O and ester C=O), 1696.2 (carbamate C=O and thiolactone C=O), 1651.2 (amide C=O), 1629.2 (quinolone C=O)

^1H NMR (400 MHz, DMSO d_6) δ / ppm = 15.16 (br s, 1 H, $\text{C}(=\text{O})\text{OH}$), 8.65 (s, 1 H, *ortho* to $\text{C}(=\text{O})\text{OH}$), 8.21 (d, $J = 8.5$ Hz, 1 H, NH), 7.89 (d, $J = 13.1$ Hz, 1 H, *ortho* to F), 7.85 (s, 1 H, $\text{CH}=\text{CCH}_2$), 7.57 (d, $J = 7.4$ Hz, 1 H, *meta* to F), 5.74 (s, 1 H, OCH_2O), 4.58 (ddd, $J = 12.6, 8.1, 7.2$ Hz, 1 H, CHNH), 4.30 (t, $J = 6.9$ Hz, 2 H, $\text{C}(=\text{O})\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 3.80 (tt, $J = 6.9, 3.6$ Hz, 1 H, $\text{NCH}(\text{CH}_2)_2$), 3.62 (br t, $J = 5.2, 5.2$ Hz, 4 H, $\text{C}(=\text{O})\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2$), 3.38 (td, $J = 11.4, 5.5$ Hz, 1 H, SCHH), 3.34 (br. s, 4 H, $\text{C}(=\text{O})\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2\text{CH}_2$), 3.27 (ddd, $J = 11.0, 6.9, 1.6$ Hz, 1 H, SCHH), 2.64 (t, $J = 7.6$ Hz, 2 H, $\text{CH}=\text{CCH}_2$), 2.44 (t, $J = 7.5$ Hz, 2 H, $\text{CH}_2\text{C}(=\text{O})\text{O}$), 2.40 (dddd, $J = 12.3, 6.8, 5.4, 1.4$ Hz, 1 H, SCH_2CHH), 2.12 (t, $J = 7.8$ Hz, 1 H, $\text{NHC}(=\text{O})\text{CHH}$), 2.12 (t, $J = 6.8$ Hz, 1 H, $\text{NHC}(=\text{O})\text{CHH}$), 1.98 - 2.07 (m, 3 H, SCH_2CHH and $\text{NHC}(=\text{O})\text{CH}_2\text{CH}_2$), 1.86 (quin, $J = 7.5$ Hz, 2 H, $\text{CH}=\text{CCH}_2\text{CH}_2$), 1.29 - 1.36 (m, 2 H, $\text{NCH}(\text{CHH})_2$), 1.14 - 1.21 (m, 2 H, $\text{NCH}(\text{CHH})_2$)


^{13}C NMR (101 MHz, DMSO d_6) δ / ppm = 205.5 ($\text{SC}(=\text{O})$), 176.4 ($\text{C}(=\text{O})\text{CC}(=\text{O})\text{OH}$), 171.8 ($\text{C}(=\text{O})\text{OCH}_2\text{O}$), 171.3 ($\text{NHC}(=\text{O})$), 165.9 ($\text{C}(=\text{O})\text{OH}$), 152.8 (d, $J = 249.7$ Hz, *ipso* to F), 152.9 ($\text{OC}(=\text{O})\text{N}$), 148.1 ($\text{CH}=\text{CC}(=\text{O})$)

OH), 146.0 (CH=CCH₂), 144.9 (d, *J* = 9.6 Hz, *ipso* to piperazine), 139.1 (*para* to F), 122.0 (CH=CCH₂), 118.9 (d, *J* = 7.5 Hz, *para* to piperazine), 111.0 (d, *J* = 23.5 Hz, *ortho* to C=O and *ortho* to F), 106.8 (C(=O)OH, and *meta* to C=O and *meta* to F), 80.3 (OCH₂O), 58.2 (CHNH), 49.1 (C(=O)N(CH₂CH₂)CH₂CH₂), 49.1 (C(=O)N(CH₂CH₂)CH₂CH₂), 48.6 (C(=O)CH₂CH₂CH₂N), 43.4 (N(CH₂)CH₂), 43.0 (N(CH₂)CH₂), 35.9 (NCH(CH₂)₂), 32.7 (CH=CCH₂CH₂CH₂C(=O)), 31.8 (NHC(=O)CH₂), 30.1 (SCH₂CH₂), 26.8 (SCH₂), 25.8 (C(=O)CH₂CH₂CH₂N), 24.2 (CH=CCH₂CH₂CH₂C(=O)), 24.0 (CH=CCH₂CH₂CH₂C(=O)), 7.6 (NCH(CH₂)₂)

HRMS (ESI⁺) *m/z* / Da = 728.2502, [M+H]⁺ found, [C₃₃H₃₉FN₇O₉S]⁺ requires 728.2503

The compound has not been reported previously.

0.8 4-Bromo-*N*-(2-methoxyphenyl)butanamide **101**



2-Methoxyaniline **100** (9.12 ml, 10.0 g, 81.2 mmol, 1 eq.) and NaHCO₃ (8.19 g, 97.4 mmol, 1.2 eq.) were dissolved in water (100 ml) and CH₂Cl₂ (100 ml). The mixture was cooled to 0 °C and 4-bromobutyryl chloride **58** (9.40 ml, 15.1 g, 81.2 mmol, 1 eq.) was added dropwise over 15 min. The mixture was stirred at 0 °C for 1.5 h, then the aqueous layer was removed. The organic layer was dried with MgSO₄ and purified by column chromatography (SiO₂, 5-25 % EtOAc/P.E.). The combined pure fractions were dried with MgSO₄ and evaporated under reduced pressure. **101** was obtained as an initially colourless liquid which slowly turned blue then black if left out on the bench (11.0 g, 40.6 mmol, 50.0 %).

TLC *R_f* = 0.16 (10 % EtOAc/P.E.)

IR (neat) ν_{max} / cm⁻¹ = 3410.2 (N-H), 3313.4 (N-H), 2961.6 (C-H), 2939.5 (C-H), 2902.5 (C-H), 1676.4 (amide C=O)


¹H NMR (400 MHz, CDCl₃ d₁) δ / ppm = 8.32 (dd, *J* = 8.0, 1.7 Hz, 1 H, *ortho* to NH), 7.85 (br s, 1 H, NH), 7.02 (td, *J* = 7.9, 1.7 Hz, 1 H, *para* to NH), 6.93 (td, *J* = 7.7, 1.4 Hz, 1 H, *para* to OCH₃), 6.85 (dd, *J* = 8.1, 1.5 Hz, 1 H, *ortho* to OCH₃), 3.85 (s, 3 H, CH₃), 3.50 (t, *J* = 6.4 Hz, 2 H, CH₂Br), 2.56 (t, *J* = 7.1 Hz, 2 H, C(=O)CH₂), 2.25 (quin, *J* = 6.7 Hz, 2 H, C(=O)CH₂CH₂)

¹³C NMR (101 MHz, CDCl₃ d₁) δ / ppm = 169.4 (C(=O)), 147.6 (*ipso* to OCH₃), 127.2 (*ipso* to NH), 123.5 (*para* to NH), 120.7 (*para* to OCH₃), 119.6 (*ortho* to NH and *meta* to OCH₃), 109.8 (*ortho* to OCH₃ and *meta* to NH), 55.5 (CH₃), 35.4 (C(=O)CH₂), 33.1 (CH₂Br), 27.9 (C(=O)CH₂CH₂)

HRMS (ESI⁺) *m/z* / Da = 272.0287, [M+H]⁺ found, [C₁₁H₁₅BrNO₂]⁺ requires 272.0286

The compound has not been reported previously.

0.9 Methyl 1-cyclopropyl-6-fluoro-7-(4-(4-((2-methoxyphenyl)amino)-4-oxobutyl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylate **102**



Methyl 1-cyclopropyl-6-fluoro-4-oxo-7-(piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylate **92** (500 mg, 1.45 mmol, 1 eq.), 4-bromo-*N*-(2-methoxyphenyl)butanamide **101** (788 mg, 2.90 mmol, 2 eq.), DIPEA (1.28 ml, 950 mg, 7.35 mmol, 5 eq.), NaI (275 mg, 1.83 mmol, 1.3 eq.) and acetonitrile (10 ml) were stirred in a microwave reactor at 100 °C for 4 h. The mixture was dry-loaded onto SiO₂ and purified by column chromatography (SiO₂, 4 % MeOH/CH₂Cl₂). The combined pure fractions were dried with MgSO₄ and evaporated under reduced pressure. **102** was obtained as a bright pink glass (79.7 mg, 0.149 mmol, 10.2 %).

TLC R_f = 0.40 (10 % MeOH/CH₂Cl₂)

IR (neat) ν_{max} / cm⁻¹ = 2947.1 (C-H), 2833.7 (C-H), 1718.9 (ester C=O), 1685.3 (amide C=O), 1617.3 (quinolone C=O)


¹H NMR (400 MHz, CDCl₃ d₁) δ / ppm = 8.48 (s, 1 H, *ortho* to C(=O)OCH₃), 8.36 (d, J = 7.9 Hz, 1 H, *ortho* to NH), 7.87 - 7.99 (m, 2 H, *ortho* to F and NH), 7.19 (d, J = 6.5 Hz, 1 H, *meta* to F), 7.01 (t, J = 7.5 Hz, 1 H, *para* to NH), 6.93 (t, J = 7.7 Hz, 1 H, *para* to OCH₃), 6.85 (d, J = 7.9 Hz, 1 H, *ortho* to OCH₃), 3.88 (s, 3 H, C(=O)OCH₃), 3.85 (s, 3 H, aromatic OCH₃), 3.41 (tt, J = 6.9, 4.0 Hz, 1 H, NCH(CH₂)₂), 3.25 (br t, J = 5.0, 5.0 Hz, 4 H, C(=O)CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 2.67 (br t, J = 5.0, 5.0 Hz, 4 H, C(=O)CH₂CH₂CH₂N(CH₂)CH₂), 2.53 (t, J = 7.0 Hz, 2 H, C(=O)CH₂CH₂CH₂N), 2.47 (t, J = 7.1 Hz, 2 H, C(=O)CH₂CH₂CH₂N), 1.97 (quin, J = 6.8 Hz, 2 H, C(=O)CH₂CH₂CH₂N), 1.25 - 1.33 (m, 2 H, NCH(CH₂)₂), 1.07 - 1.14 (m, 2 H, NCH(CH₂)₂)

¹³C NMR (101 MHz, CDCl₃ d₁) δ / ppm = 172.9 (C(=O)CC(=O)OCH₃), 170.8 (NHC(=O)), 166.2 (C(=O)OCH₃), 153.3 (d, J = 248.0 Hz, *ipso* to F), 148.2 (C=CC(=O)OCH₃), 147.6 (*ipso* to OCH₃), 144.4 (d, J = 10.4 Hz, *ipso* to piperazine), 137.9 (*para* to F), 127.6 (*ipso* to NH), 123.4 (*para* to NH), 122.7 (d, J = 7.8 Hz, *para* to piperazine), 121.0 (*para* to OCH₃), 119.7 (*ortho* to NH and *meta* to OCH₃), 113.0 (d, J = 22.5 Hz, *ortho* to C=O and *ortho* to F), 109.8 (*ortho* to OCH₃ and *meta* to NH, and CC(=O)OCH₃), 104.7 (*meta* to C=O and *meta* to F), 57.2 (CH₂CH₂CH₂N), 55.6 (aromatic OCH₃), 52.7 (CH₂CH₂CH₂N(CH₂)CH₂), 51.9 (C(=O)OCH₃), 49.8 (CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 49.8 (CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 35.5 (CH₂CH₂CH₂N), 34.5 (NCH(CH₂)₂), 22.3 (CH₂CH₂CH₂N), 8.0 (NCH(CH₂)₂)

HRMS (ESI⁺) m/z / Da = 537.2523, [M+H]⁺ found, [C₂₉H₃₄FN₄O₅]⁺ requires 537.2513

The compound has not been reported previously.

0.10 4-Azido-*N*-(2-methoxyphenyl)butanamide **103**



4-Bromo-*N*-(2-methoxyphenyl)butanamide **101** (2.05 g, 7.51 mmol, 1 eq.) and NaN₃ (1.17 g, 18.0 mmol, 2.4 eq.) were refluxed in acetonitrile (100 ml) for 2 h. The mixture was cooled and filtered, and the filtrate was dry-loaded onto SiO₂ and purified by column chromatography using a Combiflash (SiO₂, 8-14 % then hold at 14 % EtOAc/P.E.). **103** was obtained as an initially colourless liquid which slowly turned blue then black if left out on the bench (0.469 g, 2.00 mmol, 26.7 %).

TLC R_f = 0.20 (25 % EtOAc/P.E.)

IR (neat) ν_{max} / cm⁻¹ = 3419.7 (N-H), 3329.6 (N-H), 2094.8 (azide), 1672.3 (amide C=O)


¹H NMR (400 MHz, CDCl₃ d₁) δ / ppm = 8.32 (dd, J = 7.9, 1.0 Hz, 1 H, *ortho* to NH), 7.86 (br s, 1 H, NH), 7.00 (td, J = 7.5, 1.5 Hz, 1 H, *para* to NH), 6.90 (td, J = 7.7, 1.1 Hz, 1 H, *para* to OCH₃), 6.83 (dd, J = 8.1, 1.4 Hz, 1 H, *ortho* to OCH₃), 3.81 (s, 3 H, CH₃), 3.33 (t, J = 6.7 Hz, 2 H, CH₂Br), 2.42 (t, J = 7.2 Hz, 2 H, C(=O)CH₂), 1.94 (quin, J = 6.9 Hz, 2 H, C(=O)CH₂CH₂)

¹³C NMR (101 MHz, CDCl₃ d₁) δ / ppm = 169.5 (C(=O)), 147.6 (*ipso* to OCH₃), 127.1 (*ipso* to NH), 123.4 (*para* to NH), 120.5 (*para* to OCH₃), 119.5 (*ortho* to NH and *meta* to OCH₃), 109.6 (*ortho* to OCH₃ and *meta* to NH), 55.2 (CH₃), 50.3 (CH₂N₃), 33.9 (C(=O)CH₂), 24.3 (C(=O)CH₂CH₂)

HRMS (ESI⁺) m/z / Da = 257.1010, [M+H]⁺ found, [C₁₁H₁₄N₄NaO₂]⁺ requires 257.1014

The data are consistent with the literature.⁴

0.11 1-Cyclopropyl-6-fluoro-7-(4-(4-(1-(4-((2-methoxyphenyl)amino)-4-oxobutyl)-1*H*-1,2,3-triazol-4-yl)butyl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylic acid **104**



1-Cyclopropyl-6-fluoro-7-(4-(hex-5-yn-1-yl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylic acid **70** (24.1 mg, 58.6 μ mol, 1 eq.) and 4-azido-*N*-(2-methoxyphenyl)butanamide **103** (13.7 mg, 58.5 μ mol, 1 eq.) were dissolved in water (3 ml), *t*-BuOH (9 ml) and CH₂Cl₂ (9 ml), and the mixture was degassed by bubbling through N₂. A solution of CuSO₄ and THPTA (117 μ l, 5.85 μ mol, 0.1 eq., 50 mM, aq.) was added, followed by a solution

of sodium ascorbate (234 μ l, 11.7 μ mol, 0.2 eq., 50 mM, aq.). The mixture was stirred at room temperature under argon for 16 h. Water (25 ml), CH_2Cl_2 (25 ml) and MeOH (5 ml) were added and the organic layer was separated off, dry-loaded onto SiO_2 and purified by column chromatography using a Combiflash (SiO_2 , 3-23 % MeOH/ CH_2Cl_2). The combined pure fractions were dried with MgSO_4 and evaporated under reduced pressure. **104** was obtained as a clear glass (14.7 mg, 22.8 μ mol, 39.0 %).

TLC R_f = 0.28 (10 % MeOH/ CH_2Cl_2)

IR (neat) ν_{max} / cm^{-1} = 2926.5 (C-H), 2846.6 (C-H), 1723.4 (carboxylic acid C=O), 1682.0 (amide C=O), 1625.8 (quinolone C=O), 1612.8 (triazole)

^1H NMR (400 MHz, CDCl_3) δ / ppm = 15.05 (br s, 1 H, C(=O)OH), 8.76 (s, 1 H, *ortho* to C(=O)OH), 8.31 (dd, J = 8.0, 1.7 Hz, 1 H, *ortho* to NH), 8.00 (d, J = 13.0 Hz, 1 H, *ortho* to F), 7.83 (br s, 1 H, NH), 7.37 (s, 1 H, CH=CCH₂), 7.35 (d, J = 7.2 Hz, 1 H, *meta* to F), 7.04 (td, J = 7.7, 1.7 Hz, 1 H, *para* to NH), 6.95 (td, J = 7.8, 1.5 Hz, 1 H, *para* to OCH₃), 6.88 (dd, J = 8.1, 1.4 Hz, 1 H, *ortho* to OCH₃), 4.47 (t, J = 6.7 Hz, 2 H, C(=O)CH₂CH₂CH₂N), 3.88 (s, 3 H, CH₃), 3.54 (tt, J = 6.9, 4.0 Hz, 1 H, NCH(CH₂)₂), 3.35 (br t, J = 4.7 Hz, 4 H, CH=CCH₂CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 2.76 (t, J = 7.5 Hz, 2 H, CH=CCH₂), 2.66 (t, J = 4.7 Hz, 4 H, CH=CCH₂CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 2.47 (t, J = 7.3 Hz, 2 H, CH=CCH₂CH₂CH₂CH₂N), 2.44 (t, J = 6.8 Hz, 2 H, C(=O)CH₂CH₂CH₂N), 2.32 (quin, J = 6.7 Hz, 2 H, C(=O)CH₂CH₂CH₂N), 1.75 (quin, J = 7.6 Hz, 2 H, CH=CCH₂CH₂CH₂CH₂N), 1.61 (quin, J = 7.5 Hz, 2 H, CH=CCH₂CH₂CH₂CH₂N), 1.35 - 1.42 (m, 2 H, NCH(CH₂)₂), 1.17 - 1.22 (m, 2 H, NCH(CH₂)₂)


^{13}C NMR (101 MHz, CDCl_3) δ / ppm = 177.1 (C(=O)CC(=O)OH), 169.5 (NHC(=O)), 167.0 (C(=O)OH), 153.7 (d, J = 251.4 Hz, *ipso* to F), 148.1 (CH=CCH₂), 147.8 (*ipso* to OCH₃), 147.3 (C=CC(=O)OH), 145.9 (d, J = 10.4 Hz, *ipso* to piperazine), 139.1 (*para* to F), 127.3 (*ipso* to NH), 123.9 (*para* to NH), 121.0 (*para* to OCH₃), 120.9 (CH=CCH₂), 119.7 (*para* to piperazine, and *ortho* to NH and *meta* to OCH₃), 112.4 (d, J = 23.4 Hz, *ortho* to C=O and *ortho* to F), 109.9 (*ortho* to OCH₃ and *meta* to NH), 108.1 (C(=O)OH), 104.7 (*meta* to C=O and *meta* to F), 58.1 (CH=CCH₂CH₂CH₂CH₂N), 55.6 (CH₃), 52.8 (CH=CCH₂CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 49.8 (CH=CCH₂CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 49.8 (CH=CCH₂CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 49.1 (C(=O)CH₂CH₂CH₂N), 35.2 (NCH(CH₂)₂), 33.8 (C(=O)CH₂CH₂CH₂N), 27.3 (CH=CCH₂CH₂CH₂CH₂N), 26.4 (CH=CCH₂CH₂CH₂CH₂N), 26.0 (C(=O)CH₂CH₂CH₂N), 25.5 (CH=CCH₂CH₂CH₂CH₂N), 8.2 (NCH(CH₂)₂)

^{19}F NMR (376.45 MHz, CDCl_3) δ / ppm = -120.7 (s, ciprofloxacin F)

HRMS (ESI^+) m/z / Da = 646.3132, $[\text{M}+\text{H}]^+$ found, $[\text{C}_{34}\text{H}_{41}\text{FN}_7\text{O}_5]^+$ requires 646.3153

The compound has not been reported previously.

0.12 4-Bromo-*N*-(3-methoxyphenyl)butanamide **106**



3-Methoxyaniline **105** (3.04 ml, 3.33 g, 27.1 mmol, 1 eq.) and NaHCO₃ (2.73 g, 32.5 mmol, 1.2 eq.) were dissolved in water (30 ml) and CH₂Cl₂ (30 ml). The mixture was cooled to 0 °C and 4-bromobutyryl chloride **58** (3.13 ml, 5.03 g, 27.1 mmol, 1 eq.) was added dropwise over 5 min. The mixture was stirred at 0 °C for 1 h, then the aqueous layer was removed. The organic layer was dry-loaded onto SiO₂ and purified by column chromatography using a Combiflash (SiO₂, 0-100 % EtOAc/P.E.). The combined pure fractions were dried with MgSO₄ and evaporated under reduced pressure. **106** was obtained as a pale pink amorphous solid (3.66 g, 13.5 mmol, 49.6 %).

TLC R_f = 0.18 (25 % EtOAc/P.E.)

IR (neat) ν_{max} / cm⁻¹ = 1670.9 (amide C=O)


¹H NMR (400 MHz, CDCl₃ d₁) δ / ppm = 8.45 (s, 1 H, NH), 7.27 (t, J = 2.2 Hz, 1 H, *ortho* to OCH₃ and *ortho* to NH), 7.14 (t, J = 8.1 Hz, 1 H, *meta* to OCH₃ and *meta* to NH), 7.02 (d, J = 8.3 Hz, 1 H, *para* to OCH₃), 6.62 (dd, J = 8.2, 2.1 Hz, 1 H, *para* to NH), 3.71 (s, 3 H, CH₃), 3.42 (t, J = 6.5 Hz, 2 H, CH₂Br), 2.51 (t, J = 6.9 Hz, 2 H, C(=O)CH₂), 2.19 (quin, J = 6.8 Hz, 2 H, C(=O)CH₂CH₂)

¹³C NMR (101 MHz, CDCl₃ d₁) δ / ppm = 170.3 (C(=O)), 159.9 (*ipso* to OCH₃), 139.0 (*ipso* to NH), 129.5 (*meta* to OCH₃ and *meta* to NH), 112.1 (*para* to OCH₃), 109.9 (*para* to NH), 105.7 (*ortho* to OCH₃ and *ortho* to NH), 55.2 (CH₃), 35.3 (C(=O)CH₂), 33.2 (CH₂Br), 28.0 (C(=O)CH₂CH₂)

HRMS (ESI⁺) The compound does not ionise.

The compound has not been reported previously.

0.13 Methyl 1-cyclopropyl-6-fluoro-7-(4-(4-((3-methoxyphenyl)amino)-4-oxobutyl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylate **107**



Methyl 1-cyclopropyl-6-fluoro-4-oxo-7-(piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylate **92** (500 mg, 1.45 mmol, 1 eq.), 4-bromo-*N*-(3-methoxyphenyl)butanamide **106** (788 mg, 2.90 mmol, 2 eq.), DIPEA (1.28 ml, 950 mg, 7.35 mmol, 5 eq.), NaI (275 mg, 1.83 mmol, 1.3 eq.) and acetonitrile (10 ml) were stirred in a microwave reactor at 100 °C for 4 h. The mixture was evaporated under reduced pressure and partitioned between CH₂Cl₂ (50 ml) and water (50 ml). The organic layer was separated off and the aqueous layer was extracted again with CH₂Cl₂ (50 ml). The combined organic layers were dried with MgSO₄ and purified by column chromatography (SiO₂, 0-4 % MeOH/CH₂Cl₂). The combined pure fractions were dried with MgSO₄ and evaporated under reduced pressure. **107** was obtained as an off-white amorphous solid (81.7 mg, 0.152 mmol, 10.5 %).

TLC R_f = 0.38 (10 % MeOH/CH₂Cl₂)

IR (neat) ν_{max} / cm^{-1} = 3270.8 (amide N-H) 2943.8 (C-H), 2817.0 (C-H), 1729.5 (ester C=O), 1682.0 (amide C=O), 1613.5 (quinolone C=O)

^1H NMR (400 MHz, CDCl_3) δ / ppm = 8.56 (s, 1 H, *ortho* to $\text{C}(=\text{O})\text{OCH}_3$), 8.06 (d, J = 13.3 Hz, 1 H, *ortho* to F), 8.02 (br s, 1 H, NH), 7.34 (t, J = 1.7 Hz, 1 H, *ortho* to OCH_3 and *ortho* to NH), 7.25 (d, J = 7.0 Hz, 1 H, *meta* to F), 7.20 (t, J = 8.2 Hz, 1 H, *meta* to OCH_3 and *meta* to NH), 6.98 (dd, J = 7.8, 1.7 Hz, 1 H, *para* to OCH_3), 6.65 (dd, J = 8.2, 2.1 Hz, 1 H, *para* to NH), 3.93 (s, 3 H, $\text{C}(=\text{O})\text{OCH}_3$), 3.80 (s, 3 H, aromatic OCH_3), 3.42 (tt, J = 6.8, 3.7 Hz, 1 H, $\text{NCH}(\text{CH}_2)_2$), 3.31 (br t, J = 4.3, 4.3 Hz, 4 H, $\text{C}(=\text{O})\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2\text{CH}_2$), 2.73 (br t, J = 4.5, 4.5 Hz, 4 H, $\text{C}(=\text{O})\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2\text{CH}_2$), 2.58 (t, J = 6.5 Hz, 2 H, $\text{C}(=\text{O})\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 2.48 (t, J = 6.8 Hz, 2 H, $\text{C}(=\text{O})\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 2.00 (quin, J = 6.8 Hz, 2 H, $\text{C}(=\text{O})\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 1.29 - 1.36 (m, 2 H, $\text{NCH}(\text{CHH})_2$), 1.11 - 1.17 (m, 2 H, $\text{NCH}(\text{CHH})_2$)


^{13}C NMR (101 MHz, CDCl_3) δ / ppm = 173.1 ($\text{C}(=\text{O})\text{CC}(=\text{O})\text{OCH}_3$), 170.9 ($\text{NHC}(=\text{O})$), 166.3 ($\text{C}(=\text{O})\text{OCH}_3$), 160.1 (*ipso* to OCH_3), 153.3 (d, J =250.1 Hz, *ipso* to F), 148.4 ($\text{C}=\text{CC}(=\text{O})\text{OCH}_3$), 144.1 (d, J =10.1 Hz, *ipso* to piperazine), 139.4 (*ipso* to NH), 138.0 (*para* to F), 129.6 (*meta* to NH and *meta* to OCH_3), 123.3 (d, J =6.4 Hz, *para* to piperazine), 113.4 (d, J =23.3 Hz, *ortho* to C=O and *ortho* to F), 111.8 (*para* to OCH_3), 110.0 ($\text{CC}(=\text{O})\text{OCH}_3$), 109.8 (*para* to NH), 105.5 (*ortho* to OCH_3 and *ortho* to NH), 105.0 (*meta* to C=O and *meta* to F), 57.0 ($\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 55.3 (aromatic OCH_3), 52.6 ($\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2\text{CH}_2$), 52.1 ($\text{C}(=\text{O})\text{OCH}_3$), 49.2 ($\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2\text{CH}_2$), 35.2 ($\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 34.6 ($\text{NCH}(\text{CH}_2)_2$), 21.7 ($\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 8.2 ($\text{NCH}(\text{CH}_2)_2$)

^{19}F NMR (376.45 MHz, MeOD) δ / ppm = -123.5 (s, ciprofloxacin F)

HRMS (ESI^+) m/z / Da = 537.2500, $[\text{M}+\text{H}]^+$ found, $[\text{C}_{29}\text{H}_{34}\text{FN}_4\text{O}_5]^+$ requires 537.2513

The compound has not been reported previously.

0.14 4-Azido-*N*-(3-methoxyphenyl)butanamide **108**



4-Bromo-*N*-(3-methoxyphenyl)butanamide **106** (2.05 g, 7.51 mmol, 1 eq.) and NaN_3 (1.17 g, 18.0 mmol, 2.4 eq.) were refluxed in acetonitrile (100 ml) for 7 h. The mixture was cooled and filtered, and the filtrate was dry-loaded onto SiO_2 and purified by column chromatography using a Combiflash (SiO_2 , 0-100 % EtOAc/P.E.). The combined pure fractions were dried with MgSO_4 and evaporated under reduced pressure. **108** was obtained as an straw-coloured liquid (0.294 g, 1.25 mmol, 16.7 %).

TLC R_f = 0.37 (50 % EtOAc/P.E.)

IR (neat) ν_{max} / cm^{-1} = 3298.3 (N-H), 2094.7 (azide), 1661.7 (amide C=O)

^1H NMR (400 MHz, MeOD) δ / ppm = 8.63 (br s, 1 H, NH), 7.26 (t, J = 2.3 Hz, 1 H, *ortho* to OCH_3 and *ortho* to NH), 7.15 (t, J = 8.1 Hz, 1 H, *meta* to OCH_3 and *meta* to NH), 7.01 (dd, J = 7.8, 1.6 Hz, 1 H, *para* to OCH_3), 6.63 (dd, J = 8.2, 1.9 Hz, 1 H, *para* to NH), 3.69 (s, 3 H, CH_3), 3.28 (t, J = 6.7 Hz, 2 H, CH_2N_3),


2.39 (t, $J = 7.4$ Hz, 2 H, C(=O)CH₂), 1.91 (quin, $J = 7.0$ Hz, 2 H, C(=O)CH₂CH₂)

¹³C NMR (101 MHz, MeOD) δ / ppm = 170.8 (C(=O)), 159.6 (*ipso* to OCH₃), 138.9 (*ipso* to NH), 129.2 (*meta* to OCH₃ and *meta* to NH), 112.3 (*para* to OCH₃), 109.5 (*para* to NH), 106.0 (*ortho* to OCH₃ and *ortho* to NH), 54.8 (CH₃), 50.4 (CH₂N₃), 33.6 (C(=O)CH₂), 24.4 (C(=O)CH₂CH₂)

HRMS (ESI⁺) The compound does not ionise.

The compound has not been reported previously.

0.15 1-Cyclopropyl-6-fluoro-7-(4-(4-(1-(4-((3-methoxyphenyl)amino)-4-oxobutyl)-1*H*-1,2,3-triazol-4-yl)butyl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylic acid **109**



1-Cyclopropyl-6-fluoro-7-(4-(hex-5-yn-1-yl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylic acid **70** (24.1 mg, 58.6 μ mol, 1 eq.) and 4-azido-*N*-(3-methoxyphenyl)butanamide **108** (13.7 mg, 58.5 μ mol, 1 eq.) were dissolved in water (1 ml), *t*-BuOH (9 ml) and CH₂Cl₂ (10 ml), and the mixture was degassed by bubbling through N₂. A solution of CuSO₄ and THPTA (58.5 μ l, 5.85 μ mol, 0.1 eq. 100 mM, aq.) was added, followed by a solution of sodium ascorbate (117 μ l, 11.7 μ mol, 0.2 eq., 100 mM, aq.). The mixture was stirred at room temperature under argon for 2 h, then the solvent was removed under reduced pressure. The residue was partitioned between water (15 ml) and CH₂Cl₂ (15 ml), and the aqueous layer was extracted a further four times with CH₂Cl₂ (4 \times 15 ml). The combined organic layers were dried with MgSO₄, dry-loaded onto SiO₂ and purified by column chromatography (SiO₂, 0-10 % MeOH/CH₂Cl₂). The combined pure fractions were dried with MgSO₄ and evaporated under reduced pressure. **109** was obtained as a clear glass (1.9 mg, 2.9 μ mol, 5.0 %).

TLC $R_f = 0.22$ (10 % MeOH/CH₂Cl₂)

IR (neat) ν_{max} / cm⁻¹ = 2922.8 (C-H), 2849.5 (C-H), 1725.8 (carboxylic acid C=O), 1684.7 (amide C=O), 1624.5 (quinolone C=O), 1612.2 (triazole)

¹H NMR (400 MHz, DMSO d₆) δ / ppm = 15.23 (br s, 1 H, C(=O)OH), 9.89 (s, 1 H, NH), 8.66 (s, 1 H, *ortho* to C(=O)OH), 7.90 (d, $J = 13.4$ Hz, 1 H, *ortho* to F), 7.88 (s, 1 H, CH=CCH₂), 7.55 (d, $J = 7.6$ Hz, 1 H, *meta* to F), 7.27 (t, $J = 2.1$ Hz, 1 H, *ortho* to C=O and *ortho* to F), 7.16 (t, $J = 8.1$ Hz, 1 H, *meta* to OCH₃ and *meta* to NH), 7.08 (d, $J = 7.8$ Hz, 1 H, *para* to OCH₃), 6.59 (ddd, $J = 8.1, 2.4, 0.7$ Hz, 1 H, *para* to NH), 4.36 (t, $J = 6.9$ Hz, 2 H, C(=O)CH₂CH₂CH₂N), 3.81 (tt, $J = 6.7, 4.0$ Hz, 1 H, NCH(CH₂)₂), 3.70 (s, 3 H, CH₃), 3.28 - 3.32 (m, 4 H, CH=CCH₂CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 2.64 (t, $J = 7.5$ Hz, 2 H, CH=CCH₂), 2.56 (m, $J = 4.2, 4.2$ Hz, 4 H, CH=CCH₂CH₂CH₂CH₂N(CH₂)CH₂), 2.38 (t, $J = 7.3$ Hz, 2 H, CH=CCH₂CH₂CH₂CH₂N), 2.30 (t, $J = 7.4$ Hz, 2 H, C(=O)CH₂CH₂CH₂N), 2.10 (quin, $J = 7.1$ Hz, 2 H, C(=O)CH₂CH₂CH₂N), 1.64 (quin, $J = 7.5$ Hz, 2 H, CH=CCH₂CH₂CH₂CH₂N), 1.51 (quin, $J = 7.2$ Hz, 2 H,

CH=CCH₂CH₂CH₂CH₂N), 1.27 - 1.33 (m, 2 H, NCH(CHH)₂), 1.15 - 1.20 (m, 2 H, NCH(CHH)₂)


¹³C NMR (101 MHz, DMSO d₆) δ / ppm = 176.3 (C(=O)CC(=O)OH), 170.1 (NHC(=O)), 165.9 (C(=O)OH), 159.4 (*ipso* to OCH₃), 153.0 (d, *J* = 248.6 Hz, *ipso* to F), 148.0 (CH=CCH₂), 146.9 (C=CC(=O)OH), 145.2 (d, *J* = 10.7 Hz, *ipso* to piperazine), 140.3 (*para* to F), 139.2 (*ipso* to NH), 129.4 (*meta* to OCH₃ and *meta* to NH), 121.7 (CH=CCH₂), 118.5 (d, *J* = 7.5 Hz, *para* to piperazine), 111.3 (*para* to OCH₃), 110.9 (d, *J* = 22.4 Hz, *ortho* to C=O and *ortho* to F), 108.4 (*para* to NH), 106.7 (CC(=O)OH), 106.3 (*meta* to C=O and *meta* to F), 104.8 (*ortho* to OCH₃ and *ortho* to NH), 57.3 (CH=CCH₂CH₂CH₂CH₂N), 54.9 (CH₃), 52.4 (CH=CCH₂CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 49.5 (CH=CCH₂CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 49.4 (CH=CCH₂CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 48.7 (C(=O)CH₂CH₂CH₂N), 35.8 (NCH(CH₂)₂), 32.9 (C(=O)CH₂CH₂CH₂N), 26.8 (CH=CCH₂CH₂CH₂CH₂N), 25.7 (CH=CCH₂CH₂CH₂CH₂N), 25.5 (C(=O)CH₂CH₂CH₂N), 24.9 (CH=CCH₂CH₂CH₂CH₂N), 7.6 (NCH(CH₂)₂)

¹⁹F NMR (376.45 MHz, DMSO d₆) δ / ppm = -121.5 (s, ciprofloxacin F)

HRMS (ESI⁺) *m/z* / Da = 646.3159, [M+H]⁺ found, [C₃₄H₄₁FN₇O₅]⁺ requires 646.3153

The compound has not been reported previously.

0.16 Methyl 7-(4-(4-(*tert*-butoxy)-4-oxobutyl)piperazin-1-yl)-1-cyclopropyl-6-fluoro-4-oxo-1,4-dihydroquinoline-3-carboxylate **122**



Methyl 1-cyclopropyl-6-fluoro-4-oxo-7-(piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylate **92** (200 mg, 0.579 mmol, 1 eq.), *tert*-butyl 4-bromobutanoate **121** (103 μl, 130 mg, 0.581 mmol, 1 eq.), NaI (86.9 mg, 0.580 mmol, 1 eq.), TEA (316 μl, 229 mg, 2.27 mmol, 4 eq.) and acetonitrile (10 ml) were stirred in a microwave reactor at 100 °C for 8 h. A second portion of *tert*-butyl 4-bromobutanoate **142** (103 μl, 130 mg, 0.581 mmol, 1 eq.) was added, and the mixture was stirred in the microwave reactor at 100 °C for a further 8 h. The mixture was then dry-loaded onto SiO₂ and purified by column chromatography (SiO₂, 0-4 % MeOH/CH₂Cl₂). **122** was obtained as a white amorphous solid (141 mg, 0.289 mmol, 49.9 %).

TLC *R_f* = 0.12 (4 % MeOH/CH₂Cl₂)

IR (neat) ν_{max} / cm⁻¹ = 2961.6 (C-H), 2830.5 (C-H), 1732.2 (*t*-Bu ester C=O) 1717.2 (ciprofloxacin ester C=O), 1620.6 (quinolone C=O)

¹H NMR (400 MHz, CDCl₃) δ / ppm = 8.39 (s, 1 H, *ortho* to C(=O)OCH₃), 7.82 (d, *J* = 13.3 Hz, 1 H, *ortho* to F), 7.17 (d, *J* = 7.2 Hz, 1 H, *meta* to F), 3.83 (s, 3 H, CH₃), 3.40 (tt, *J* = 7.2, 3.6 Hz, 1 H, NCH(CH₂)₂), 3.22 (t, *J* = 4.3 Hz, 4 H, CH₂N(CH₂CH₂)CH₂CH₂), 2.63 (t, *J* = 4.4 Hz, 4 H, CH₂N(CH₂)CH₂), 2.41 (t, *J* = 7.3 Hz, 2 H, CH₂N(CH₂)CH₂), 2.25 (t, *J* = 7.4 Hz, 2 H, CH₂CH₂CH₂N(CH₂)CH₂), 1.78 (quin, *J* = 7.3 Hz, 2 H, CH₂CH₂N(CH₂)CH₂), 1.41 (s, 9 H, C((CH₃)₃)), 1.24 (m, 2 H, NCH(CHH)₂), 1.09 (m, 2 H, NCH(CHH)₂)


^{13}C NMR (101 MHz, CDCl_3) δ / ppm = 172.7 ($\text{C}(=\text{O})\text{CC}(=\text{O})\text{OCH}_3$), 172.6 ($\text{C}(=\text{O})\text{OC}(\text{CH}_3)_3$), 165.9 ($\text{C}(=\text{O})\text{OCH}_3$), 153.1 (d, J = 249.7 Hz, *ipso* to F), 148.1 ($\text{C}=\text{CC}(=\text{O})\text{OCH}_3$), 144.3 (d, J = 10.4 Hz, *ipso* to piperazine), 137.7 (*para* to F), 122.5 (d, J = 6.9 Hz, *para* to piperazine) 112.6 (d, J = 22.5 Hz, *ortho* to C=O and *ortho* to F), 109.5 ($\text{CC}(=\text{O})\text{OCH}_3$) 104.7 (*meta* to C=O and *meta* to F), 80.0 ($\text{C}(\text{CH}_3)_3$), 57.4 ($\text{C}(=\text{O})\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 52.7 ($\text{C}(=\text{O})\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2)\text{CH}_2$), 51.7 (CH_3), 49.7 ($\text{C}(=\text{O})\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2\text{CH}_2$), 49.7 ($\text{C}(=\text{O})\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2\text{CH}_2$), 34.4 ($\text{NCH}(\text{CH}_2)_2$), 33.2 ($\text{C}(=\text{O})\text{CH}_2$), 28.0 ($\text{C}(\text{CH}_3)_3$), 22.0 ($\text{C}(=\text{O})\text{CH}_2\text{CH}_2$), 7.9 ($\text{NCH}(\text{CH}_2)_2$)

^{19}F NMR (376.45 MHz, CDCl_3) δ / ppm = -123.5 (s, ciprofloxacin F)

HRMS (ESI^+) m/z / Da = 488.2562, $[\text{M}+\text{H}]^+$ found, $[\text{C}_{26}\text{H}_{35}\text{FN}_3\text{O}_5]^+$ requires 488.2561

The compound has not been reported previously.

0.17 4-(4-(1-Cyclopropyl-6-fluoro-3-(methoxycarbonyl)-4-oxo-1,4-dihydroquinolin-7-yl)piperazin-1-yl)butanoic acid trifluoroacetate **123**



Methyl 7-(4-(4-(*tert*-butoxy)-4-oxobutyl)piperazin-1-yl)-1-cyclopropyl-6-fluoro-4-oxo-1,4-dihydroquinoline-3-carboxylate **122** (20 mg, 41.0 μmol) and TFA (0.2 ml) were stirred in CH_2Cl_2 (1.8 ml) at r.t. for 16 h then evaporated under reduced pressure. **123** was obtained as a white solid (21.4 mg, 39.2 μmol , 95.6 %).

mp T / $^\circ\text{C}$ = 225-231 (CH_2Cl_2 , decomposes)

IR (neat) ν_{max} / cm^{-1} = 1722.7 (ciprofloxacin ester C=O), 1699.0 (alkyl carboxylic acid C=O), 1673.3 (TFA C=O), 1614.6 (quinolone C=O)

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ / ppm = 8.47 (s, 1 H, *ortho* to $\text{C}(=\text{O})\text{OH}$), 7.80 (d, J = 13.2 Hz, 1 H, *ortho* to F), 7.47 (d, J = 7.4 Hz, 1 H, *meta* to F), 3.73 (s, 3 H, CH_3), 3.66 (tt, J = 7.2, 3.7 Hz, 1 H, $\text{NCH}(\text{CH}_2)_2$), 3.30 - 3.54 (br s, 8 H, $\text{CH}_2\text{N}(\text{CH}_2)\text{CH}_2$ and $\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2\text{CH}_2$) 3.13 - 3.22 (m, 2 H, $\text{CH}_2\text{N}(\text{CH}_2)\text{CH}_2$), 2.36 (t, J = 7.1 Hz, 2 H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2)\text{CH}_2$), 1.87 - 1.98 (m, 2 H, $\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2)\text{CH}_2$), 1.22 - 1.30 (m, 2 H, $\text{NCH}(\text{CH}_2)_2$), 1.06 - 1.15 (m, 2 H, $\text{NCH}(\text{CH}_2)_2$)


^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ / ppm = 173.5 ($\text{CH}_2\text{C}(=\text{O})\text{OH}$), 171.6 ($\text{C}(=\text{O})\text{CC}(=\text{O})\text{OCH}_3$), 164.9 ($\text{C}(=\text{O})\text{OCH}_3$), 158.2 (q, J = 31.5 Hz, $\text{CF}_3\text{C}(=\text{O})\text{OH}$), 152.5 (d, J = 247.6 Hz, *ipso* to F), 148.5 ($\text{C}=\text{CC}(=\text{O})\text{OH}$), 142.3 (d, J = 10.7 Hz, *ipso* to piperazine), 138.0 (*para* to F), 122.6 (d, J = 6.4 Hz, *para* to piperazine), 117.2 (q, J = 299.8 Hz, CF_3), 111.9 (d, J = 22.4 Hz, *ortho* to C=O and *ortho* to F), 109.1 ($\text{CC}(=\text{O})\text{OCH}_3$), 106.9 (*meta* to C=O and *meta* to F), 55.1 ($\text{C}(=\text{O})\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 51.4 (CH_3), 50.8 ($\text{C}(=\text{O})\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2)\text{CH}_2$), 46.7 ($\text{C}(=\text{O})\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2\text{CH}_2$), 46.7 ($\text{C}(=\text{O})\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2\text{CH}_2$), 34.9 ($\text{NCH}(\text{CH}_2)_2$), 30.6 ($\text{C}(=\text{O})\text{CH}_2$), 19.1 ($\text{C}(=\text{O})\text{CH}_2\text{CH}_2$), 7.6 ($\text{NCH}(\text{CH}_2)_2$)

^{19}F NMR (376.45 MHz, DMSO d_6) δ / ppm = -73.6 (s, CF_3), -124.6 (s, ciprofloxacin F)

HRMS (ESI^+) m/z / Da = 432.1921, $[\text{M}+\text{H}]^+$ found, $[\text{C}_{22}\text{H}_{27}\text{FN}_3\text{O}_5]^+$ requires 432.1935

The compound has not been reported previously.

0.18 (1*R*,2*R*)-2-(((*S*)-1-Phenylethyl)amino)cyclopentan-1-ol **112 and (1*S*,2*S*)-2-(((*S*)-1-phenylethyl)amino)cyclopentan-1-ol **113****



(*S*)-1-Phenylethan-1-amine **111** (7.85 ml, 7.38 g, 60.9 mmol, 1 eq.) was dissolved in CH_2Cl_2 (50 ml) and stirred rapidly at 0 °C. A solution of AlMe_3 (31 ml, 2.0 M in heptane, 60.9 mmol) was added dropwise and the mixture was stirred at 0 °C for 1 h. A solution of cyclohexene oxide **110** (5.71 ml, 5.50 g, 65.4 mmol, 1.1 eq.) in CH_2Cl_2 (50 ml) was then added dropwise, and the mixture was stirred at 0 °C for a further 3 h, followed by 48 h at r.t.. The mixture was cooled to 0 °C and NaF (11 g, 262 mmol, 4.3 eq.) was added portionwise, followed by water (7.00 ml, 7.00 g, 389 mmol, 6.4 eq.) and CH_2Cl_2 (50 ml). The suspension was allowed to warm to r.t. and stirred for 1 h, then filtered through Celite and washed with CH_2Cl_2 (500 ml). The filtrate was dried with K_2CO_3 , concentrated under reduced pressure and purified by column chromatography (SiO_2 , 20:5:1 hexane:EtOAc:TEA). **112** was obtained as a pale yellow oil (4.08 g, 19.9 mmol, 32.6 %). **113** was obtained as pale yellow crystals (4.48 g, 21.8 mmol, 35.8 %).

(1*R*,2*R*)-2-(((*S*)-1-Phenylethyl)amino)cyclopentan-1-ol **112**

TLC R_f = 0.25 (15:5:1 hexane:EtOAc:TEA)

IR (neat) ν_{max} / cm^{-1} = 3300.0 (br, O-H), 2959.7 (C-H), 2870.1 (C-H)

^1H NMR (400 MHz, CDCl_3) δ / ppm = 7.28 - 7.38 (m, 4 H, *ortho* and *meta* to CHCH_3), 7.21 - 7.28 (m, 1 H, *para* to CHCH_3), 3.83 (q, J = 6.6 Hz, 1 H, CHCH_3), 3.78 (q, J = 7.0 Hz, 1 H, CHOH), 2.62 (dt, J = 8.2, 7.2 Hz, 1 H, CHNH), 1.97 (quin, J = 6.7 Hz, 1 H, CH_2CHNH), 1.90 (quin, J = 6.9 Hz, 1 H, CH_2CHOH), 1.56 - 1.68 (m, $\text{CH}_2\text{CH}_2\text{CHOH}$), 1.43 (dq, J = 12.5, 8.0 Hz, 1 H, CH_2CHOH), 1.37 (d, J = 6.6 Hz, 3 H, CH_3), 1.25 - 1.36 (m, 1 H, CH_2CHNH)

^{13}C NMR (101 MHz, CDCl_3) δ / ppm = 144.75 (*ipso* to CHCH_3), 128.26 (*meta* to CHCH_3), 126.72 (*para* to CHCH_3), 126.30 (*ortho* to CHCH_3), 77.65 (CHOH), 63.38 (CHNH), 56.20 (CHCH_3), 31.74 (CH_2CHOH), 29.22 (CH_2CHNH), 24.58 (CH_3), 19.57 ($\text{CH}_2\text{CH}_2\text{CHOH}$)

HRMS (ESI^+) m/z / Da = 206.1554, $[\text{M}+\text{H}]^+$ found, $[\text{C}_{13}\text{H}_{20}\text{NO}]^+$ requires 206.1545

$[\alpha]_D^{20}$ / $^{\circ}10^{-1}\text{cm}^2\text{g}^{-1}$ = -92.8 (c / $\text{g}(100\text{ ml})^{-1}$ = 1.19, MeOH)

(1*S*,2*S*)-2-(((*S*)-1-Phenylethyl)amino)cyclopentan-1-ol 113

TLC R_f = 0.36 (15:5:1 hexane:EtOAc:TEA)

mp T / °C = 66-71.5 (hexane, EtOAc, TEA)

IR (neat) ν_{max} / cm^{-1} = 3150.0 (br, O-H), 2950.9 (C-H), 2868.2 (C-H)

^1H NMR (400 MHz, CDCl_3) δ / ppm = 7.28 - 7.34 (m, 4 H, *ortho* and *meta* to CHCH_3), 7.20 - 7.26 (m, 1 H, *para* to CHCH_3), 3.86 (q, J = 6.6 Hz, 1 H, CHCH_3), 3.85 (q, J = 6.6 Hz, 1 H, CHOH), 2.83 (td, J = 7.6, 5.7 Hz, 1 H, CHNH), 1.85 - 1.97 (m, 1 H, CHHCHOH), 1.77 (dtd, J = 12.9, 7.9, 7.9, 4.9 Hz, 1 H, CHHCHNH), 1.55 - 1.68 (m, 2 H, $\text{CH}_2\text{CH}_2\text{CHOH}$), 1.47 - 1.55 (m, 1 H, CHHCHOH), 1.36 (d, J = 6.6 Hz, 3 H, CH_3), 1.12 (dq, J = 12.7, 8.1 Hz, 1 H, CHHCHNH)


^{13}C NMR (101 MHz, CDCl_3) δ / ppm = 145.61 (*ipso* to CHCH_3), 128.08 (*meta* to CHCH_3), 126.61 (*para* to CHCH_3), 126.33 (*ortho* to CHCH_3), 77.43 (CHOH), 64.45 (CHNH), 56.62 (CHCH_3), 32.01 (CH_2CHOH), 30.56 (CH_2CHNH), 23.30 (CH_3), 20.06 ($\text{CH}_2\text{CH}_2\text{CHOH}$)

HRMS (ESI^+) m/z / Da = 206.1553, $[\text{M}+\text{H}]^+$ found, $[\text{C}_{13}\text{H}_{20}\text{NO}]^+$ requires 206.1545

$[\alpha]_D^{20}$ / $^{\circ}\text{10}^{-1}\text{cm}^2\text{g}^{-1}$ = -23.9 (c / g(100 ml)^{-1} = 0.96, MeOH)

The compounds have been synthesised previously,^{5,6} but NMR data were not published. The enantiomers of both compounds have also been synthesised previously, and the ^1H NMR data for these are consistent with the the above data.⁷

0.19 (1*R*,2*R*)-2-Aminocyclopentan-1-ol 114



(1*R*,2*R*)-2-(((*S*)-1-Phenylethyl)amino)cyclopentan-1-ol **112** (3.90 g, 19.0 mmol, 1 eq.), $\text{Pd}(\text{OH})_2$ (20 wt. % on C, moistened with 50 wt. % water, 1 g, 0.712 mmol, 0.04 eq.) and MeOH (50 ml) were stirred in a Paar hydrogenator at r.t. and 3 atm for 2 days. The mixture was then filtered through Celite and evaporated under reduced pressure. **114** was obtained as a yellow oil (1.92 g, 19.0 mmol, 100 %).

TLC R_f = 0.10 (10 % MeOH/ CH_2Cl_2)

IR (neat) ν_{max} / cm^{-1} = 3300.0 (br, O-H), 2958.3 (C-H), 2871.5 (C-H)

^1H NMR (400 MHz, MeOD) δ / ppm = 3.77 (ddd, J =6.6, 6.2, 5.6, 1 H, CHOH), 3.00 (td, J =7.3, 5.6 Hz, 1 H, CHNH_2), 2.00 (dtd, J =13.0, 7.7, 7.7, 5.6 Hz, 1 H, CHHCHNH_2), 1.97 (ddt, J =13.0, 8.7, 6.6, 6.6 Hz, 1 H, CHHCHOH), 1.63 - 1.77 (m, 2 H, $\text{CH}_2\text{CH}_2\text{CHOH}$), 1.53 (ddt, J =13.0, 9.5, 6.2, 6.2 Hz, 1 H, CHHCHOH), 1.37 (ddt, J =13.0, 8.3, 7.8, 7.8 Hz, 1 H, CHHCHNH_2)


^{13}C NMR (101 MHz, MeOD) δ / ppm = 80.7 ($\underline{\text{C}}\text{HOH}$), 60.8 ($\underline{\text{C}}\text{HNH}_2$), 33.2 ($\underline{\text{C}}\text{H}_2\text{CHOH}$), 32.1 ($\underline{\text{C}}\text{H}_2\text{CHNH}_2$), 21.2 ($\underline{\text{C}}\text{H}_2\text{CH}_2\text{CHOH}$)

HRMS (ESI⁺) m/z / Da = 102.0917, $[\text{M}+\text{H}]^+$ found, $[\text{C}_5\text{H}_{12}\text{NO}]^+$ requires 102.0913

$[\alpha]_D^{20}$ / $^{\circ}10^{-1}\text{cm}^2\text{g}^{-1}$ = -30.9 (c / $\text{g}(100\text{ ml})^{-1}$ = 1.5, EtOH)

The data are consistent with the literature.^{6,8}

0.20 (1*S*,2*S*)-2-Aminocyclopentan-1-ol **115**



(1*S*,2*S*)-2-(((*S*)-1-Phenylethyl)amino)cyclopentan-1-ol **113** (3.00 g, 14.6 mmol, 1 eq.), Pd(OH)₂ (20 wt. % on C, moistened with 50 wt. % water, 0.5 g, 0.356 mmol, 0.025 eq.) and MeOH (50 ml) were stirred in a Paar hydrogenator at r.t. and 2.5 atm for 2 days. The mixture was then filtered through Celite and evaporated under reduced pressure. **115** was obtained as a yellow oil (1.48 g, 14.6 mmol, 100 %).

TLC R_f = 0.10 (10 % MeOH/ CH_2Cl_2)

IR (neat) ν_{max} / cm^{-1} = 3300.0 (O-H), 2969.2 (C-H), 2872.7 (C-H)

^1H NMR (400 MHz, MeOD) δ / ppm = 3.77 (ddd, J =6.6, 6.2, 5.6, 1 H, $\underline{\text{C}}\text{HOH}$), 3.00 (td, J = 7.4, 5.6 Hz, 1 H, $\underline{\text{C}}\text{HNH}_2$), 2.00 (dtd, J = 13.0, 7.7, 7.7, 5.6 Hz, 1 H, $\underline{\text{C}}\text{HHCHNH}_2$), 1.97 (ddt, J = 13.0, 8.7, 6.4, 6.4 Hz, 1 H, $\underline{\text{C}}\text{HHCHOH}$), 1.64 - 1.77 (m, 2 H, $\underline{\text{C}}\text{H}_2\text{CH}_2\text{CHOH}$), 1.53 (ddt, J = 13.0, 9.5, 6.2, 6.2 Hz, 1 H, $\underline{\text{C}}\text{HHCHOH}$), 1.37 (ddt, J = 12.8, 8.5, 7.7, 7.7 Hz, 1 H, $\underline{\text{C}}\text{HHCHNH}_2$)


^{13}C NMR (101 MHz, MeOD) δ / ppm = 80.6 ($\underline{\text{C}}\text{HOH}$), 60.7 ($\underline{\text{C}}\text{HNH}_2$), 33.2 ($\underline{\text{C}}\text{H}_2\text{CHOH}$), 32.2 ($\underline{\text{C}}\text{H}_2\text{CHNH}_2$), 21.2 ($\underline{\text{C}}\text{H}_2\text{CH}_2\text{CHOH}$)

HRMS (ESI⁺) m/z / Da = 102.0915, $[\text{M}+\text{H}]^+$ found, $[\text{C}_5\text{H}_{12}\text{NO}]^+$ requires 102.0913

$[\alpha]_D^{20}$ / $^{\circ}10^{-1}\text{cm}^2\text{g}^{-1}$ = 33.4 (c / $\text{g}(100\text{ ml})^{-1}$ = 0.5, EtOH)

The data are consistent with the literature.^{6,8}

0.21 (1*S*,2*S*)-2-((*tert*-Butyldimethylsilyl)oxy)cyclopentan-1-amine **117**



(1*S*,2*S*)-2-Aminocyclopentan-1-ol **115** (0.480 g, 4.75 mmol) was stirred in dry CH₂Cl₂ (20 ml) under N₂ at 0 °C. TEA (3.14 ml, 2.28 g, 22.5 mmol, 5 eq.) was added dropwise, followed by TBSOTf (3 ml, 3.45 g, 13.1 mmol, 3 eq.) dropwise. The reaction was allowed to reach r.t. and stirred for 1 h. The reaction was quenched with NH₄Cl, diluted with CH₂Cl₂ (20 ml) and washed with water (20 ml). The organic phase was dried with Na₂SO₄, concentrated under reduced pressure and purified by column chromatography (SiO₂, 4 % MeOH/CH₂Cl₂). **117** was obtained as a yellow oil (1.00 g, 4.64 mmol, 97.7 %).

TLC R_f = 0.23 (10 % MeOH/CH₂Cl₂)


IR (neat) ν_{max} / cm⁻¹ = 2953.6 (C-H), 2931.1 (C-H), 2888.4 (C-H), 2858.8 (C-H), 1625.2 (N-H bend)

¹H NMR (400 MHz, CDCl₃) δ / ppm = 4.13 (q, J = 5.8 Hz, 1 H, CHOSi), 3.31 (td, J = 7.1, 5.2 Hz, 1 H, CHNH₂), 2.09 - 2.19 (m, 1 H, CHHCHNH₂), 1.97 (ddq, J = 8.8, 7.0, 6.0, 6.0, 6.0 Hz, 1 H, CHHCHOSi), 1.74 - 1.86 (m, 2 H, CH₂CH₂CHOSi), 1.64 - 1.74 (m, 1 H, CHHCHOSi), 1.58 (ddt, J = 13.2, 9.1, 6.0, 6.0 Hz, 1 H, CHHCHNH₂), 0.88 (s, 9 H, C(CH₃)₃), 0.09 (s, 3 H, SiCH₃), 0.07 (s, 3 H, SiCH₃)

¹³C NMR (101 MHz, CDCl₃) δ / ppm = 76.3 (CHOSi), 59.7 (CHNH), 32.2 (CH₂CHOSi), 26.8 (CH₂CHNH₂), 25.6 (C(CH₃)₃), 19.7 (CH₂CH₂CHOSi), 17.7 (C(CH₃)₃), -4.8 (SiCH₃), -5.2 (SiCH₃)

HRMS (ESI⁺) m/z / Da = 216.1785, [M+H]⁺ found, [C₁₁H₂₆NOSi]⁺ requires 216.1784 The compound has not been reported previously.

0.22 4-Chloro-*N*-((1*R*,2*R*)-2-hydroxycyclopentyl)butanamide **125**



(1*R*,2*R*)-2-Aminocyclopentan-1-ol **114** (500 mg, 4.94 mmol, 1 eq.), TEA (827 μ l, 600 mg, 5.93 mmol, 1.2 eq.) and CH₂Cl₂ (20 ml) were stirred at 0 °C and 4-chlorobutyl chloride **124** (608 μ l, 766 mg, 5.43 mmol, 1.1 eq.) was added dropwise over 5 min. The mixture was stirred at 0 °C for 30 min, then water (50 ml) was added. The organic layer was separated off, and the aqueous layer was extracted with CH₂Cl₂ (7×50 ml). The combined organic layers were dried with MgSO₄, concentrated under reduced pressure and purified by column chromatography (SiO₂, Et₂O). The combined pure fractions were dried with MgSO₄ and evaporated under reduced pressure. **125** was obtained as a white amorphous solid (651 mg, 3.16 mmol, 64.1 %).

TLC R_f = 0.35 (EtOAc)

IR (neat) ν_{max} / cm⁻¹ = 3277.6 (N-H and O-H), 2962.2 (C-H), 2876.0 (C-H), 1636.3 (amide C=O)

¹H NMR (400 MHz, CDCl₃) δ / ppm = 6.12 (br s, 1 H, NH), 4.42 (br s, 1 H, OH), 3.94 (q, J = 6.6 Hz, 1 H, CHOH), 3.82 (tt, J = 8.4, 5.3 Hz, 1 H, CHNH), 3.60 (t, J = 6.2 Hz, 2 H, CH₂Cl), 2.38 (t, J = 7.2 Hz, 2 H, CH₂C=O), 2.05 - 2.16 (m, 3 H, CHHCHNH and CH₂CH₂Cl), 1.96 - 2.04 (m, 1 H, CHHCHOH), 1.74 - 1.85 (m, 1 H, CHHCH₂CHOH), 1.58 - 1.73 (m, 2 H, CHHCH₂CHOH and CHHCHOH), 1.43 (dq, J = 12.7, 8.3 Hz, 1 H, CHHCHNH)


¹³C NMR (101 MHz, CDCl₃) δ / ppm = 173.8 (C=O), 79.4 (CHOH), 60.6 (CHNH), 44.4 (CH₂Cl), 32.8 (CH₂C=O), 32.4 (CH₂CHOH), 30.1 (CH₂CHNH), 28.0 (CH₂CH₂Cl), 21.1 (CH₂CH₂CHOH)

HRMS (ESI⁺) m/z / Da = 228.0787, [M+Na]⁺ found, [C₉H₁₆ClNNaO₂]⁺ requires 228.0762

$[\alpha]_D^{20}$ / °10⁻¹cm²g⁻¹ = -13.0 (c / g(100 ml)⁻¹ = 0.5, MeOH)

The compound has not been reported previously.

0.23 4-Chloro-*N*-((1*S*,2*S*)-2-hydroxycyclopentyl)butanamide **126**



(1*S*,2*S*)-2-Aminocyclopentanol **115** (72.3 mg, 716 μ mol, 1 eq.), TEA (500 μ l, 363 mg, 3.58 mmol, 5 eq.) and CH₂Cl₂ (5 ml) were stirred at 0 °C, and 4-chlorobutyryl chloride **124** (179 μ l, 226 mg, 1.60 mmol, 1.1 eq.) was added dropwise over 5 min. The mixture was stirred at 0 °C for 30 min, then water (10 ml) was added. The organic layer was separated off, and the aqueous layer was extracted with 10 % *i*-PrOH/CHCl₃ (2×10 ml). The combined organic layers were dried with MgSO₄, concentrated under reduced pressure and purified by column chromatography (SiO₂, Et₂O). The combined pure fractions were dried with MgSO₄ and evaporated under reduced pressure. **126** was obtained as a white amorphous solid (35.6 mg, 173 μ mol, 24.2 %).

TLC R_f = 0.35 (EtOAc)

¹H NMR (400 MHz, CDCl₃) δ / ppm = 6.05 (br s, 1 H, NH), 4.55 (br s, 1 H, OH), 3.95 (q, J=6.6 Hz, 1 H, CHOH), 3.82 (tt, J=8.4, 5.3 Hz, 1 H, CHNH), 3.60 (t, J=6.2 Hz, 2 H, CH₂Cl), 2.38 (t, J=7.0 Hz, 2 H, CH₂C=O), 2.05 - 2.17 (m, 3 H, CHHCHNH and CH₂CH₂Cl), 1.94 - 2.05 (m, 1 H, CHHCHOH), 1.74 - 1.86 (m, 1 H, CHHCH₂CHOH), 1.58 - 1.74 (m, 2 H, CHHCH₂CHOH and CHHCHCHOH), 1.42 (dq, J=12.5, 8.4 Hz, 1 H, CHHCHNH)


¹³C NMR (101 MHz, CDCl₃) δ / ppm = 173.8 (C=O), 79.4 (CHOH), 60.6 (CHNH), 44.4 (CH₂Cl), 32.8 (CH₂C=O), 32.4 (CH₂CHOH), 30.2 (CH₂CHNH), 28.0 (CH₂CH₂Cl), 21.2 (CH₂CH₂CHOH)

HRMS (ESI⁺) m/z / Da = 206.0939, [M+H]⁺ found, [C₉H₁₇ClNO₂]⁺ requires 206.0948

$[\alpha]_D^{20}$ / °10⁻¹cm²g⁻¹ = 10.0 (c / g(100 ml)⁻¹ = 0.05, MeOH)

The compound has not been reported previously.

0.24 4-Azido-*N*-((1*S*,2*S*)-2-((*tert*-butyldimethylsilyl)oxy)cyclopentyl)butanamide 119



(1*S*,2*S*)-2-((*tert*-Butyldimethylsilyl)oxy)cyclopentan-1-amine **117** (50 mg, 0.232 mmol, 1 eq.) and NaHCO₃ (22.0 mg, 0.262 mmol, 1.1 eq.) were added to CH₂Cl₂ (3 ml) and water (3 ml) at 0 °C, and 4-bromobutyryl chloride (25.3 ml, 40.5 mg, 0.219 mmol, 0.95 eq.) was added dropwise. The mixture was stirred for 3 h at 0 °C. The aqueous layer was removed and NaN₃ (100 mg, 1.54 mmol, 6.6 eq.) and DMF (3 ml) were added. The mixture was then stirred at 40 °C for 6 h. The solvents were then evaporated using a N₂ stream and the residue was purified by column chromatography (SiO₂, 0.5 % MeOH/CH₂Cl₂). The combined pure fractions were dried with MgSO₄ and evaporated under reduced pressure. **119** was obtained as a clear liquid (71 mg, 0.217 mmol, 99.2 %).

TLC R_f = 0.84 (1 % MeOH/CH₂Cl₂)

IR (neat) ν_{max} / cm⁻¹ = 3287.9 (N-H), 2953.4 (C-H), 2933.2 (C-H), 2882.7 (C-H), 2857.1 (C-H), 2094.9 (azide), 1639.4 (amide C=O)

¹H NMR (400 MHz, CDCl₃) δ / ppm = 5.35 (d, J = 5.1 Hz, 1 H, NH), 3.97 - 4.01 (m, 1 H, CHOSi), 3.93 - 3.98 (m, 1 H, CHNH), 3.35 (t, J = 6.6 Hz, 2 H, CH₂N₃), 2.24 (t, J = 7.0 Hz, 2 H, CH₂C=O), 2.09 - 2.19 (m, 1 H, CHHCHNH), 1.89 - 1.97 (quin, J = 6.8 Hz, 2 H, CH₂CH₂N₃), 1.74 - 1.84 (m, 2 H, CHHCHOSi and CHHCH₂CHOSi), 1.60 - 1.70 (m, 1 H, CHHCH₂CHOSi), 1.51 - 1.61 (m, 1 H, CHHCHOSi), 1.31 - 1.39 (m, 1 H, CHHCHNH), 0.87 (s, 9 H, C(CH₃)₃), 0.08 (s, 3 H, SiCH₃), 0.06 (s, 3 H, SiCH₃)


¹³C NMR (101 MHz, CDCl₃) δ / ppm = 171.17 (C=O), 77.80 (CHOSi), 58.36 (CHNH), 50.77 (CH₂N₃), 33.29 (CH₂C=O), 32.57 (CH₂CHOSi), 29.36 (CH₂CHNH), 25.72 (C(CH₃)₃), 24.77 (CH₂CH₂N₃), 20.40 (CH₂CH₂CHO Si), 17.95 (C(CH₃)₃), -4.75 (SiCH₃)

HRMS (ESI⁺) m/z / Da = 327.2221, [M+H]⁺ found, [C₁₅H₃₁N₄O₂Si]⁺ requires 327.2216

$[\alpha]_D^{20}$ / °10⁻¹cm²g⁻¹ = 12.4 (c / g(100 ml)⁻¹ = 0.5, MeOH)

The compound has not been reported previously.

0.25 4-Azido-*N*-((1*R*,2*R*)-2-hydroxycyclopentyl)butanamide 127



4-Chloro-*N*-((1*R*,2*R*)-2-hydroxycyclopentyl)butanamide **125** (200 mg, 0.972 mmol, 1 eq.) and NaN₃ (126 mg, 1.94 mmol, 2 eq.) were stirred in acetonitrile (4 ml) at 50 °C for 16 h. The solvent was then evaporated under reduced pressure and the residue was partitioned between water (20 ml) and 10 % *i*-PrOH/CHCl₃ (20 ml). The aqueous layer was extracted again with 10 % *i*-PrOH/CHCl₃ (3×20 ml) and the combined organic fractions were dried with MgSO₄ and evaporated under reduced pressure. **127** was obtained as white needles (181 mg, 0.852 mmol, 87.6 %).

TLC R_f = 0.35 (EtOAc)

mp T / °C = 56.0-59.5 (*i*-PrOH, CHCl₃)

IR (neat) ν_{max} / cm⁻¹ = 3279.9 (N-H and O-H), 2965.6 (C-H), 2875.4 (C-H), 2094.6 (azide), 1636.8 (amide C=O)

¹H NMR (400 MHz, CDCl₃) δ / ppm = 6.72 (d, J = 4.4 Hz, 1 H, NH), 4.82 (br. s., 1 H, OH), 3.88 (q, J = 6.6 Hz, 1 H, CHOH), 3.75 (tdd, J = 8.4, 8.4, 6.6, 4.4 Hz, 1 H, CHNH), 3.28 (t, J = 6.6 Hz, 2 H, CH₂N₃), 2.23 (t, J = 7.3 Hz, 2 H, CH₂C=O), 2.04 (dtd, J = 13.0, 8.0, 8.0, 4.9 Hz, 1 H, CHHCHNH), 1.92 (dtd, J = 13.0, 7.6, 7.6, 5.8 Hz, 1 H, CHHCHOH), 1.84 (quin, J = 7.0 Hz, 2 H, CH₂CH₂N₃), 1.59 - 1.77 (m, 2 H, CH₂CH₂CHOH), 1.54 (ddt, J = 12.7, 9.0, 6.7, 6.7 Hz, 1 H, CHHCHOH), 1.39 (dq, J = 12.9, 8.4 Hz, 1 H, CHHCHNH)


¹³C NMR (101 MHz, CDCl₃) δ / ppm = 173.8 (C=O), 78.8 (CHOH), 59.9 (CHNH), 50.5 (CH₂N₃), 32.5 (CH₂C=O), 32.0 (CH₂CHOH), 29.5 (CH₂CHNH), 24.6 (CH₂CH₂N₃), 20.7 (CH₂CH₂CHOH)

HRMS (ESI⁺) m/z / Da = 235.1174, [M+Na]⁺ found, [C₉H₁₆N₄NaO₂]⁺ requires 235.1171

$[\alpha]_D^{20}$ / °10⁻¹cm²g⁻¹ = -10.2 (c / g(100 ml)⁻¹ = 0.5, MeOH)

The compound has not been reported previously.

0.26 4-Azido-*N*-((1*S*,2*S*)-2-hydroxycyclopentyl)butanamide **128**



4-Chloro-*N*-((1*S*,2*S*)-2-hydroxycyclopentyl)butanamide **126** (35.0 mg, 0.170 mmol, 1 eq.) and NaN₃ (22.1 mg, 0.340 mmol, 2 eq.) were stirred in acetonitrile (2 ml) at 50 °C for 24 h. The reaction mixture was then partitioned between water (20 ml) and 10 % *i*-PrOH/CHCl₃ (5 ml). The aqueous layer was extracted again with 10 % *i*-PrOH/CHCl₃ (2×5 ml) and the combined organic fractions were dried with MgSO₄ and evaporated under reduced pressure. **128** was obtained as white needles (16.2 mg, 0.0764 mmol, 45.0 %).

TLC R_f = 0.35 (EtOAc)

IR (neat) ν_{max} / cm⁻¹ = 3286.7 (N-H and O-H), 2957.6 (C-H), 2930.6 (C-H), 2860.7 (C-H), 2094.7 (azide), 1642.2 (amide C=O)

¹H NMR (400 MHz, CDCl₃) δ / ppm = 5.82 (br s, 1 H, NH), 4.45 (br. s., 1 H, OH), 3.96 (q, J=6.6 Hz, 1 H, CHOH), 3.83 (tdd, J=8.5, 8.5, 6.0, 4.6 Hz, 1 H, CHNH), 3.37 (t, J=6.4 Hz, 2 H, CH₂N₃), 2.31 (t, J=7.2 Hz, 2 H, CH₂C=O), 2.09 - 2.19 (m, 1 H, CHHCHNH), 1.99 - 2.06 (m, 1 H, CHHCHOH), 1.90 - 1.97 (m, 2 H, CH₂CH₂N₃), 1.60 - 1.85 (m, 3 H, CH₂CHHCHOH), 1.42 (dq, J=12.8, 8.3 Hz, 1 H, CHHCHNH)


¹³C NMR (101 MHz, CDCl₃) δ / ppm = 173.8 (C=O), 79.7 (CHOH), 61.0 (CHNH), 50.7 (CH₂N₃), 32.8 (CH₂C=O), 32.6 (CH₂CHOH), 30.5 (CH₂CHNH), 24.7 (CH₂CH₂N₃), 21.3 (CH₂CH₂CHOH)

HRMS (ESI⁺) m/z / Da = 235.1178, [M+Na]⁺ found, [C₉H₁₆N₄NaO₂]⁺ requires 235.1171

$[\alpha]_D^{20}$ / °10⁻¹cm²g⁻¹ = 10.0 (c / g(100 ml)⁻¹ = 0.01, MeOH)

The compound has not been reported previously.

0.27 Methyl 1-cyclopropyl-6-fluoro-7-(4-(4-(((1*R*,2*R*)-2-hydroxycyclopentyl)amino)-4-oxobutyl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylate **129**



4-(4-(1-Cyclopropyl-6-fluoro-3-(methoxycarbonyl)-4-oxo-1,4-dihydroquinolin-7-yl)piperazin-1-yl)butanoic acid trifluoroacetate **123** (200 mg, 0.367 mmol, 1 eq.), (1*R*,2*R*)-2-aminocyclopentan-1-ol **114** (80 mg, 0.791 mmol, 2.1 eq.), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (112 mg, 0.584 mmol, 1.6 eq.), 1-hydroxybenzotriazole (96 mg, 0.710 mmol, 1.9 eq.) and DIPEA (192 μ l, 142 mg, 1.10 mmol, 3 eq.) were dissolved in DMF (5 ml) and stirred at r.t. for 16 h. The solvent was removed using a stream of N₂ and the residue was purified by preparatory HPLC (5-60 % acetonitrile/water over 12 min). The combined pure fractions were evaporated under reduced pressure and then partitioned between NaHCO₃ (aq., sat., 10 ml) and CH₂Cl₂ (10 ml). The organic layer was removed and the aqueous layer was extracted twice more with CH₂Cl₂ (2×10 ml). The combined organic fractions were dried with MgSO₄ and evaporated under reduced pressure. **129** was obtained as a white amorphous solid (73.0 mg, 0.142 mmol, 38.7 %).

TLC R_f = 0.43 (30 % MeOH/EtOAc)

IR (neat) ν_{max} / cm⁻¹ = 2972.9 (C-H), 2901.5 (C-H), 1728.4 (ester C=O), 1656.3 (amide C=O), 1612.9 (quinolone C=O)

¹H NMR (400 MHz, DMSO d₆) δ / ppm = 8.44 (s, 1 H, *ortho* to C(=O)OCH₃), 7.75 (d, J = 13.5 Hz, 1 H, *ortho* to F), 7.70 (d, J = 7.2 Hz, 1 H, CHNH), 7.43 (d, J = 7.5 Hz, 1 H, *meta* to F), 4.74 (d, J = 4.0 Hz, 1 H, CHOH), 3.78 - 3.82 (m, 1 H, CHOH), 3.74 - 3.78 (m, 1 H, CHNH), 3.74 (s, 3 H, CH₃), 3.65 (tt, J = 7.2, 3.9 Hz, 1 H, NCH(CH₂)₂), 3.25 (t, J = 4.8 Hz, 4 H, CH₂N(CH₂CH₂)CH₂CH₂), 2.57 (br s, 4 H, CH₂N(CH₂)CH₂), 2.34 (t, J = 7.4 Hz, 2 H, CH₂N(CH₂)CH₂), 2.11 (t, J = 7.4 Hz, 2 H, CH₂CH₂CH₂N(CH₂)CH₂), 1.92 (dddd, J = 13.0, 8.7, 7.3, 6.0 Hz, 1 H, CHHCHNH), 1.78 (dddd, J = 12.6, 8.9, 6.3, 6.3 Hz, 1 H, CHHCHOH), 1.69

(quin, $J = 7.3$ Hz, 2 H, $\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)_2$), 1.54 - 1.65 (m, 2 H, $\text{CH}_2\text{CH}_2\text{CHOH}$), 1.42 (ddt, $J = 13.1, 8.2, 5.3, 5.3$ Hz, 1 H, CHHCHOH), 1.32 (dddd, $J = 13.4, 8.5, 6.8, 5.8$ Hz, 1 H, CHHCHNH), 1.21 - 1.29 (m, 2 H, $\text{NCH}(\text{CHH})_2$), 1.07 - 1.13 (m, 2 H, $\text{NCH}(\text{CHH})_2$)

^{13}C NMR (101 MHz, DMSO d_6) δ / ppm = 171.9 ($\text{CH}_2\text{C}(=\text{O})\text{NH}$), 171.6 ($\text{C}(=\text{O})\text{CC}(=\text{O})\text{OCH}_3$), 165.0 ($\text{C}(=\text{O})\text{OCH}_3$), 152.6 (d, $J = 246.5$ Hz, *ipso* to F), 148.3 ($\text{C}=\text{CC}(=\text{O})\text{OCH}_3$), 143.9 (d, $J = 10.7$ Hz, *ipso* to piperazine), 138.1 (*para* to F), 121.8 (d, $J = 6.4$ Hz, *para* to piperazine), 111.5 (d, $J = 22.4$ Hz, *ortho* to C=O and *ortho* to F), 109.0 ($\text{CC}(=\text{O})\text{OCH}_3$), 106.2 (*meta* to C=O and *meta* to F), 76.3 (CHOH), 57.6 (CHNH), 57.2 ($\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 52.4 ($\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)_2$), 51.3 (CH_3), 49.6 ($\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)_2$), 34.8 ($\text{NCH}(\text{CH}_2)_2$), 33.3 ($\text{C}(=\text{O})\text{CH}_2$), 32.2 (CH_2CHOH), 29.5 (CH_2CHNH), 22.5 ($\text{C}(=\text{O})\text{CH}_2\text{CH}_2$), 20.6 ($\text{CH}_2\text{CH}_2\text{CHOH}$), 7.6 ($\text{NCH}(\text{CH}_2)_2$)


^{19}F NMR (376.45 MHz, DMSO d_6) δ / ppm = -124.3 (ciprofloxacin F)

HRMS (ESI⁺) m/z / Da = 515.2661, $[\text{M}+\text{H}]^+$ found, $[\text{C}_{27}\text{H}_{36}\text{FN}_4\text{O}_5]^+$ requires 515.2670

$[\alpha]_D^{20}$ / $^{\circ}\text{10}^{-1}\text{cm}^2\text{g}^{-1}$ = -6.0 (c / g(100 ml)^{-1} = 0.05, MeOH)

The compound has not been reported previously.

0.28 Methyl 1-cyclopropyl-6-fluoro-7-(4-(4-(((1*S*,2*S*)-2-hydroxycyclopentyl)amino)-4-oxobutyl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylate **130**



4-(4-(1-Cyclopropyl-6-fluoro-3-(methoxycarbonyl)-4-oxo-1,4-dihydroquinolin-7-yl)piperazin-1-yl)butanoic acid trifluoroacetate **123** (52.1 mg, 95.5 μmol , 1 eq.), (1*S*,2*S*)-2-aminocyclopentan-1-ol **115** (19.5 mg, 193 μmol , 2 eq.), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (29.7 mg, 155 μmol , 1.6 eq.), 1-hydroxybenzotriazole (25.8 mg, 191 μmol , 2 eq.) and DIPEA (33.3 μl , 24.7 mg, 191 μmol , 2 eq.) were dissolved in DMF (2 ml) and stirred at r.t. for 16 h. The solvent was removed using a stream of N_2 and the residue was purified by preparatory HPLC (5-50 % acetonitrile/water over 15 min). The combined pure fractions were evaporated under reduced pressure and then partitioned between NaHCO_3 (aq., sat., 5 ml) and CH_2Cl_2 (5 ml). The organic layer was removed and the aqueous layer was extracted twice more with CH_2Cl_2 (2 \times 5 ml). The combined organic fractions were dried with MgSO_4 and evaporated under reduced pressure. **130** was obtained as a white amorphous solid (4.9 mg, 9.5 μmol , 9.9 %).

TLC R_f = 0.38 (30 % MeOH/ CH_2Cl_2)

IR (neat) ν_{max} / cm^{-1} = 2937.7 (C-H), 1721.4 (ester C=O), 1620.5 (amide C=O and quinolone C=O)

^1H NMR (500 MHz, DMSO d_6) δ / ppm = 8.44 (s, 1 H, *ortho* to $\text{C}(=\text{O})\text{OCH}_3$), 7.75 (d, $J=13.5$ Hz, 1 H, *ortho*

to F), 7.69 (d, J=6.9 Hz, 1 H, CHNH), 7.43 (d, J=7.6 Hz, 1 H, *meta* to F), 4.73 (br s, 1 H, CHOH), 3.77 - 3.81 (m, 1 H, CHOH), 3.74 - 3.77 (m, 1 H, CHNH), 3.73 (s, 3 H, CH₃), 3.65 (tt, J=6.9, 4.0 Hz, 1 H, NCH(CH₂)₂), 3.24 (br. t, J=4.2, 4.2 Hz, 4 H, CH₂N(CH₂CH₂)CH₂CH₂), 2.55 (br t, J=5.0, 5.0 Hz, 4 H, CH₂N(CH₂)CH₂), 2.32 (t, J=7.2 Hz, 2 H, CH₂N(CH₂)CH₂), 2.10 (t, J=7.4 Hz, 2 H, CH₂CH₂CH₂N(CH₂)CH₂), 1.92 (dddd, J=13.0, 8.7, 7.3, 6.0 Hz, 1 H, CHHCHNH), 1.77 (ddt, J=12.6, 8.9, 6.3, 6.3 Hz, 1 H, CHHCHOH), 1.68 (quin, J=7.4 Hz, 2 H, CH₂CH₂N(CH₂)CH₂), 1.53 - 1.64 (m, 2 H, CH₂CH₂CHOH), 1.42 (ddt, J=12.9, 8.4, 5.2, 5.2 Hz, 1 H, CHHCHOH), 1.31 (ddt, J=13.0, 8.6, 6.4, 6.4 Hz, 1 H, CHHCHNH), 1.22 - 1.28 (m, 2 H, NCH(CHH)₂), 1.06 - 1.12 (m, 2 H, NCH(CHH)₂)

¹³C NMR (126 MHz, DMSO d₆) δ / ppm = 171.9 (NHC(=O)CH₂), 171.5 (C(=O)CC(=O)OCH₃), 165.0 (C(=O)OCH₃), 152.6 (d, J=247.4 Hz, *ipso* to F), 148.2 (C=CC(=O)OCH₃), 143.9 (d, J=10.3 Hz, *ipso* to piperazine), 138.1 (*para* to F), 121.7 (d, J=6.4 Hz, *para* to piperazine), 111.5 (d, J=23.0 Hz, *ortho* to C=O and *ortho* to F), 109.0 (CC(=O)OCH₃), 106.2 (*meta* to C=O and *meta* to F), 76.2 (CHOH), 57.6 (CHNH), 57.2 (CH₂CH₂CH₂N), 52.4 (CH₂CH₂CH₂N(CH₂)CH₂), 51.3 (CH₃), 49.6 (CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 49.6 (CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 34.7 (NCH(CH₂)₂), 33.2 (C(=O)CH₂), 32.2 (CH₂CHOH), 29.5 (CH₂CHNH), 22.5 (C(=O)CH₂CH₂), 20.6 (CH₂CH₂CHOH), 7.5 (NCH(CH₂)₂)


¹⁹F NMR (376.45 MHz, MeOD) δ / ppm = -125.5

HRMS (ESI⁺) *m/z* / Da = 515.2667, [M+H]⁺ found, [C₂₇H₃₆FN₄O₅]⁺ requires 515.2670

[α]_D²⁰ / °10⁻¹cm²g⁻¹ = 8.0 (*c* / g(100 ml)⁻¹ = 0.05, MeOH)

The compound has not been reported previously.

0.29 Methyl (*S*)-1-cyclopropyl-6-fluoro-4-oxo-7-(4-(4-oxo-4-((2-oxocyclopentyl)amino)butyl)piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylate **131**



Methyl 1-cyclopropyl-6-fluoro-7-(4-(4-(((1*S*,2*S*)-2-hydroxycyclopentyl)amino)-4-oxobutyl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylate **130** (20.0 mg, 38.9 μmol, 1 eq.) and Dess-Martin Periodane (32.8 mg, 77.4 μmol, 2 eq.) were stirred in CH₂Cl₂ (3 ml) for 6 h. The solvent was removed under reduced pressure and the residue was purified by preparatory HPLC (5-50 % acetonitrile/water over 10 min). The combined pure fractions were evaporated under reduced pressure, then NaHCO₃ (aq., sat., 30 ml) and 10 % *i*-PrOH/CHCl₃ (30 ml) were added. The organic layer was removed and dried with MgSO₄, then evaporated under reduced pressure. **131** was obtained as a white amorphous solid (11.3 mg, 22.0 μmol, 56.7 %).

¹H NMR (500 MHz, DMSO d₆) δ / ppm = 8.46 (s, 1 H, *ortho* to C(=O)OCH₃), 7.78 (d, J=13.5 Hz, 1 H, *ortho* to F), 7.45 (d, J=7.4 Hz, 1 H, *meta* to F), 4.02 (dt, J=11.1, 8.2 Hz, 1 H, CHNH), 3.73 (s, 3 H, CH₃), 3.65 (tt, J=6.9, 3.9 Hz, 1 H, NCH(CH₂)₂), 3.40 (s, 10 H, CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 2.05 - 2.29 (m, 5

H, NHC(=O)CH_2 , $\text{CH}_2\text{C(=O)CHNH}$ and CHHCHNH), 1.89 - 1.96 (m, 1 H, $\text{CHHCH}_2\text{CHNH}$), 1.69 - 1.80 (m, 3 H, $\text{CHHCH}_2\text{CHNH}$, CHHCHNH and $\text{NHC(=O)CH}_2\text{CH}_2$), 1.24 - 1.29 (m, 2 H, NCH(CHH)_2), 1.07 - 1.12 (m, 2 H, NCH(CHH)_2)

^{13}C NMR (126 MHz, DMSO d_6) δ / ppm = 215.2 (C(=O)CHNH), 171.7 (NHC(=O)CH_2), 171.7 (C(=O)CC(=O)OCH_3), 165.1 (C(=O)OCH_3), 152.6 (d, $J=246.6$ Hz, *ipso* to F), 148.4 (C=CC(=O)OCH_3), 138.1 (*para* to F), 109.1 (CC(=O)OCH_3), 56.3 (CHNH), 51.4 (CH_3), 35.6 ($\text{CH}_2\text{C(=O)CHNH}$), 34.8 ($\text{NCH(CH}_2)_2$), 28.8 (CH_2CHNH), 18.1 ($\text{CH}_2\text{CH}_2\text{CHNH}$), 7.6 ($\text{NCH(CH}_2)_2$)


^{19}F NMR (376.45 MHz, MeOD) δ / ppm = -124.3

HRMS (ESI $^+$) m/z / Da = 513.2495, $[\text{M}+\text{H}]^+$ found, $[\text{C}_{27}\text{H}_{34}\text{FN}_4\text{O}_5]^+$ requires 513.2513

$[\alpha]_D^{20}$ / $^{\circ}10^{-1}\text{cm}^2\text{g}^{-1}$ = 6.7 (c / $\text{g}(100\text{ ml})^{-1}$ = 0.075, MeOH)

The compound has not been reported previously.

0.30 7-(4-(4-(1-(4-(((1*S*,2*S*)-2-((*tert*-butyldimethylsilyl)oxy)cyclopentyl)amino)-4-oxobutyl)-1*H*-1,2,3-triazol-4-yl)butyl)piperazin-1-yl)-1-cyclopropyl-6-fluoro-4-oxo-1,4-dihydroquinoline-3-carboxylic acid **120**



1-Cyclopropyl-6-fluoro-7-(4-(hex-5-yn-1-yl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylic acid **70** (42.9 mg, 104 μmol , 1 eq.) and 4-azido-*N*-(((1*S*,2*S*)-2-((*tert*-butyldimethylsilyl)oxy)cyclopentyl)butanamide **119** (33.9 mg, 104 μmol , 1 eq.) were dissolved in 10 % water/*t*-BuOH (3 ml), and the mixture was degassed by bubbling N_2 through it. A solution of CuSO_4 and THPTA (104 μl , 10.4 μmol , 0.1 eq. 100 mM, aq.) was added, followed by a solution of sodium ascorbate (208 μl , 20.8 μmol , 0.2 eq., 100 mM, aq.). The mixture was stirred at room temperature under argon for 16 h, then solvent was removed under reduced pressure. The residue was partitioned between water (10 ml) and CH_2Cl_2 (10 ml), the organic layer was separated and the aqueous layer was extracted again with CH_2Cl_2 (10 ml). The combined organic layers were dried with MgSO_4 and evaporated under reduced pressure. **120** was obtained as a clear glass (67.1 mg, 90.9 μmol , 87.4 %).

IR (neat) ν_{max} / cm^{-1} = 2951.3 (C-H), 2929.2 (C-H), 2855.5 (C-H), 1741.0 (carboxylic acid C=O), 1640.3 (amide C=O), 1626.6 (quinolone C=O), 1612.3 (triazole)

^1H NMR (400 MHz, CDCl_3) δ / ppm = 8.67 (s, 1 H, *ortho* to C(=O)OH), 7.87 (d, J = 13.1 Hz, 1 H, *ortho* to F), 7.34 (s, 1 H, CH=CCH_2), 7.33 (d, J = 8.2 Hz, 1 H, *meta* to F), 5.92 (t, J = 6.6 Hz, 1 H, CHNH), 4.35 (t, J = 6.7 Hz, 2 H, $\text{CH}_2\text{NCH=C}$), 3.96 - 4.02 (m, 1 H, CHOSi), 3.90 - 3.96 (m, 1 H, CHNH), 3.55 (tt, J = 6.7, 4.0 Hz, 1 H, $\text{NCH(CH}_2)_2$), 3.34 (br t, J = 5.0 Hz, 4 H, $\text{CH}_2\text{N(CH}_2\text{CH}_2)_2$), 2.71 (t, J = 7.5 Hz, 2 H, CH=CCH_2), 2.66 (br s, 4 H, $\text{CH}_2\text{N(CH}_2)_2$), 2.46 (t, J = 7.3 Hz, 2 H, $\text{CH}_2\text{N(CH}_2)_2$), 2.03 - 2.22

1627.0 (quinolone C=O), 1613.0 (triazole)

¹H NMR (700 MHz, DMSO d₆) δ / ppm = 8.64 (s, 1 H, *ortho* to C(=O)OH), 7.87 (d, J = 13.3 Hz, 1 H, *ortho* to F), 7.84 (s, 1 H, CH=CCH₂), 7.75 (d, J = 7.1 Hz, 1 H, CHNH), 7.54 (d, J = 7.5 Hz, 1 H, *meta* to F), 4.73 (d, J = 3.8 Hz, 1 H, CHOH), 4.29 (t, J = 6.9 Hz, 2 H, CH₂NCH=C), 3.78 - 3.83 (m, 1 H, NCH(CH₂)₂), 3.75 - 3.78 (m, 1 H, CHOH), 3.71 - 3.75 (m, 1 H, CHNH), 3.31 (br t, J = 4.3 Hz, 4 H, CH₂N(CH₂CH₂)CH₂CH₂), 2.63 (t, J = 7.5 Hz, 2 H, CH=CCH₂), 2.56 (br t, J = 4.2 Hz, 4 H, CH₂N(CH₂)CH₂), 2.37 (t, J = 7.3 Hz, 2 H, CH₂N(CH₂)CH₂), 2.03 - 2.06 (m, 2 H, C(=O)CH₂), 1.97 - 2.02 (m, 2 H, C(=O)CH₂CH₂), 1.89 (dddd, J = 13.1, 8.9, 7.4, 5.7 Hz, 1 H, CHHCHNH), 1.75 (ddt, J = 13.0, 8.9, 6.4, 6.4 Hz, 1 H, CHHCHOH), 1.61 - 1.66 (m, 2 H, CH=CCH₂CH₂), 1.57 - 1.61 (m, 1 H, CHHCH₂CHOH), 1.54 - 1.57 (m, 1 H, CHHCH₂CHOH), 1.49 - 1.53 (m, 2 H, CH=CCH₂CH₂CH₂), 1.40 (ddt, J = 13.0, 8.4, 5.3, 5.3 Hz, 1 H, CHHCHOH), 1.29 - 1.32 (m, 2 H, NCH(CHH)₂), 1.25 - 1.29 (m, 1 H, CHHCHNH), 1.13 - 1.20 (m, 2 H, NCH(CHH)₂)

¹³C NMR (175 MHz, DMSO d₆) δ / ppm = 176.3 (C(=O)CC(=O)OH), 170.9 (NHC(=O)CH₂), 166.1 (C(=O)OH), 153.0 (d, J = 251.4 Hz, *ipso* to F), 147.9 (C=CC(=O)OH), 146.9 (CH=CCH₂), 145.2 (d, J = 8.7 Hz, *ipso* to piperazine), 139.2 (*para* to F), 121.7 (NCH=CCH₂), 118.7 (d, J = 5.8 Hz, *para* to piperazine), 111.0 (d, J = 23.3 Hz, *ortho* to C=O and *ortho* to F), 106.3 (*meta* to C=O and *meta* to F and CC(=O)OH), 76.2 (CHOH), 57.6 (CHNH), 57.4 (CH=CCH₂CH₂CH₂CH₂N), 52.5 (CH=CCH₂CH₂CH₂CH₂N(CH₂)CH₂), 49.5 (d, J = 4.4 Hz, CH=CCH₂CH₂CH₂CH₂N(CH₂)CH₂CH₂), 48.8 (CH₂NCH=CCH₂), 35.8 (NCH(CH₂)₂), 32.2 (CH₂CHOH), 32.0 (C(=O)CH₂), 29.5 (CH₂CHNH), 26.9 (CH=CCH₂CH₂), 26.0 (C(=O)CH₂CH₂), 25.8 (CH=CCH₂CH₂CH₂), 25.0 (CH=CCH₂), 20.5 (CH₂CH₂CHOH), 7.6 (NCH(CH₂)₂)


¹⁹F NMR (376.45 MHz, MeOD) δ / ppm = -122.1 (s, ciprofloxacin F)

HRMS (ESI⁺) m/z / Da = 624.3314, [M+H]⁺ found, [C₃₂H₄₃FN₇O₅]⁺ requires 624.3310

$[\alpha]_D^{20}$ / °10⁻¹cm²g⁻¹ = -3.6 (c / g(100 ml)⁻¹ = 0.0833, MeOH)

The compound has not been reported previously.

0.32 1-Cyclopropyl-6-fluoro-7-(4-(4-(1-(4-(((1*S*,2*S*)-2-hydroxycyclopentyl)amino)-4-oxobutyl)-1*H*-1,2,3-triazol-4-yl)butyl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylic acid **133**



1-Cyclopropyl-6-fluoro-7-(4-(hex-5-yn-1-yl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylic acid **70** (82.0 mg, 199 μ mol, 4 eq.) and 4-azido-*N*-(((1*S*,2*S*)-2-hydroxycyclopentyl)butanamide **128** (11.0 mg, 51.8 μ mol, 1 eq.) were dissolved in 10 % water/*t*-BuOH (3 ml), and the mixture was degassed by bubbling N₂ through it. A solution of CuSO₄ and THPTA (156 μ l, 15.6 μ mol, 0.3 eq. 100 mM, aq.) was added, followed by a solution of sodium ascorbate (312 μ l, 31.2 μ mol, 0.6 eq., 100 mM, aq.). The mixture was stirred at room temperature under

argon for 3 d. Water (10 ml) and 10 % *i*-PrOH/CHCl₃ (10 ml) were added, then the organic layer was separated and dried with MgSO₄ and evaporated under reduced pressure. The residue was purified by preparatory HPLC (5-95 % acetonitrile/water over 20 min). The combined pure fractions were evaporated under reduced pressure and then partitioned between NaHCO₃ (aq., sat., 10 ml) and 10 % *i*-PrOH/CHCl₃ (10 ml). The organic layer was dried with MgSO₄ and evaporated under reduced pressure. **133** was obtained as a white amorphous solid (7.2 mg, 11.5 μmol, 22.2 %).

IR (neat) ν_{max} / cm⁻¹ = 2954.9 (C-H), 2917.9 (C-H), 2850.2 (C-H), 1722.1 (carboxylic acid C=O), 1647.3 (amide C=O), 1626.7 (quinolone C=O) 1611.9 (triazole)

¹H NMR (400 MHz, DMSO d₆) δ / ppm = 15.22 (br s, 1 H, C(=O)OH), 8.67 (s, 1 H, *ortho* to C(=O)OH), 7.91 (d, J=13.3 Hz, 1 H, *ortho* to F), 7.84 (s, 1 H, CH=CCH₂), 7.74 (d, J=6.7 Hz, 1 H, CHNH), 7.56 (d, J=7.4 Hz, 1 H, *meta* to F), 4.71 (d, J=3.7 Hz, 1 H, CHOH), 4.29 (t, J=6.6 Hz, 2 H, CH₂NCH=C), 3.82 (tt, J=6.5, 4.3 Hz, 1 H, NCH(CH₂)₂), 3.69 - 3.79 (m, 2 H, CHOH and CHNH), 3.30 - 3.34 (m, 6 H, CH=CCH₂CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 2.64 (t, J=7.4 Hz, 2 H, CH=CCH₂), 1.95 - 2.08 (m, 4 H, C(=O)CH₂CH₂), 1.89 (dddd, J=12.8, 8.9, 7.4, 5.8 Hz, 1 H, CHHCHNH), 1.75 (ddt, J=12.7, 9.0, 6.2, 6.2 Hz, 1 H, CHHCHOH), 1.48 - 1.68 (m, 6 H, CH=CCH₂CH₂CH₂ and CH₂CH₂CHOH), 1.40 (ddt, J=13.0, 8.3, 5.3, 5.3 Hz, 1 H, CHHCHOH), 1.28 - 1.35 (m, 2 H, NCH(CHH)₂), 1.24 - 1.31 (m, 1 H, CHHCHNH), 1.15 - 1.21 (m, 2 H, NCH(CHH)₂)

¹³C NMR (101 MHz, DMSO d₆) δ / ppm = 176.4 (C(=O)CC(=O)OH), 170.9 (NHC(=O)CH₂), 166.0 (C(=O)OH), 153.0 (d, J=249.6 Hz, *ipso* to F), 148.1 (C=CC(=O)OH), 146.7 (CH=CCH₂), 145.2 (d, J=8.3 Hz, *ipso* to piperazine), 139.2 (*para* to F), 121.8 (NCH=CCH₂), 118.7 (*para* to piperazine), 111.0 (d, J=23.2 Hz, *ortho* to C=O and *ortho* to F), 106.7 (CC(=O)OH), 106.5 (*meta* to C=O and *meta* to F), 76.2 (CHOH), 57.5 (CHNH), 57.4 (br s, CH=CCH₂CH₂CH₂CH₂N), 52.3 (br s, CH=CCH₂CH₂CH₂CH₂N(CH₂)CH₂), 49.3 (br s, CH=CCH₂CH₂CH₂CH₂N(CH₂)CH₂), 48.8 (CH₂NCH=CCH₂), 35.9 (NCH(CH₂)₂), 32.2 (CH₂CHOH), 32.0 (C(=O)CH₂), 29.4 (CH₂CHNH), 26.7 (CH=CCH₂CH₂), 26.0 (C(=O)CH₂CH₂), 25.5 (CH=CCH₂CH₂CH₂), 24.9 (CH=CCH₂), 20.5 (CH₂CH₂CHOH), 7.6 (NCH(CH₂)₂)


¹⁹F NMR (376.45 MHz, MeOD) δ / ppm = -121.5

HRMS (ESI⁺) m/z / Da = 624.3298, [M+H]⁺ found, [C₃₂H₄₃FN₇O₅]⁺ requires 624.3310

$[\alpha]_D^{20}$ / °10⁻¹cm²g⁻¹ = -25.0 (c / g(100 ml)⁻¹ = 0.08, MeOH)

The compound has not been reported previously.

0.33 (*trans*)-2-Aminocyclohexan-1-ol **135**



Cyclohexene oxide **134** (10 ml, 9.70 g, 98.8 mmol, 1 eq.), NH₃ (90 ml, 35 % w/w aq., 27.7 g, 791 mmol, 8 eq.) and MeOH (100 ml) were stirred at r.t. for 72 h. The solvent was removed by blowing a stream of N₂ over it, followed by evaporation under high vacuum. **135** was obtained as a white amorphous solid (9.90 g, 85.2 mmol,

86.2 %)

TLC R_f = 0.04 (30 % MeOH/CH₂Cl₂)

IR (neat) ν_{max} / cm⁻¹ = 3350.4 (N-H), 3306.2 (br, O-H), 2926.9 (C-H), 2852.6 (C-H)


¹H NMR (400 MHz, CDCl₃) δ / ppm = 3.01 (td, J = 9.4, 4.8 Hz, 1 H, CHOH), 2.80 - 2.92 (m, 2 H, OH and NH₂), 2.35 (ddd, J = 11.1, 9.1, 4.1 Hz, 1 H, CHNH₂), 1.77 - 1.84 (m, 1 H, CHHCHOH), 1.69 - 1.76 (m, 1 H, CHHCHNH₂), 1.56 - 1.66 (m, 1 H, CHHCH₂CHOH), 1.45 - 1.56 (m, 1 H, CHHCH₂CHNH₂), 1.07 - 1.19 (m, 3 H, CHHCH₂CHOH, CHHCH₂CHNH₂ and CHHCHOH), 0.94 - 1.05 (m, 1 H, CHHCHNH₂)

¹³C NMR (101 MHz, CDCl₃) δ / ppm = 75.4 (CHOH), 56.6 (CHN₂), 33.8 (CH₂CHOH and CH₂CHN₂), 24.7 (CH₂CH₂CHN₂), 24.6 (CH₂CH₂CHOH)

HRMS (ESI⁺) m/z / Da = 116.1070, [M+H]⁺ found, [C₆H₁₄NO]⁺ requires 116.1070

The data are consistent with the literature.⁹

0.34 4-Chloro-*N*-((*trans*)-2-hydroxycyclohexyl)butanamide **136**



(*trans*)-2-Aminocyclohexan-1-ol **135** (1.04 g, 9.03 mmol, 1 eq.), TEA (1.65 ml, 1.20 g, 11.8 mmol, 1.3 eq.) and CH₂Cl₂ (50 ml) were stirred at 0 °C. 4-Chlorobutyryl chloride **124** (1.22 ml, 1.54 g, 10.9 mmol, 1.2 eq.) was added dropwise over 5 min. The mixture was stirred at 0 °C for 30 min, then water (50 ml) was added. The organic layer was separated off, and the aqueous layer was extracted with 10 % *i*-PrOH/CHCl₃ (2×50 ml). The combined organic layers were dried with MgSO₄, concentrated under reduced pressure and purified by column chromatography (SiO₂, 0-100 % EtOAc/Et₂O). The combined organic fractions were dried with MgSO₄ and evaporated under reduced pressure. **136** was obtained as white needles (1.51 g, 6.87 mmol, 76.1 %).

TLC R_f = 0.19 (Et₂O)

mp T / °C = 72.5-75.7 (*i*-PrOH, CHCl₃)

IR (neat) ν_{max} / cm⁻¹ = 3289.9 (N-H), 3250.0 (O-H), 2927.6 (C-H), 2857.1 (C-H), 1629.2 (amide C=O)

¹H NMR (400 MHz, MeOD) δ / ppm = 3.60 (t, J = 6.6 Hz, 2 H, CH₂Cl), 3.51 - 3.60 (m, 1 H, CHNH), 3.28 - 3.39 (m, 1 H, CHOH), 2.37 (td, J = 7.4, 2.3 Hz, 2 H, C(=O)CH₂), 2.06 (quin, J = 7.0 Hz, 2 H, C(=O)CH₂CH₂), 1.97 - 2.01 (m, 1 H, CHHCHOH), 1.85 - 1.93 (m, 1 H, CHHCHNH), 1.70 - 1.77 (m, 1 H, CHHCH₂CHOH), 1.64 - 1.70 (m, 1 H, CHHCH₂CHNH), 1.24 - 1.35 (m, 3 H, CHHCH₂CHOH, CHHCH₂CHNH and CHHCHOH), 1.13 - 1.25 (m, 1 H, CHHCHNH₂)


¹³C NMR (101 MHz, MeOD) δ / ppm = 175.0 (C(=O)), 74.1 (CHOH), 56.3 (CHNH), 45.3 (CH₂Cl), 35.6 (CH₂CHOH), 34.5 (C(=O)CH₂), 32.7 (CH₂CHNH), 30.1 (C(=O)CH₂CH₂), 25.8 (CH₂CH₂CHNH), 25.5

(CH₂CH₂CHOH)

HRMS (ESI⁺) m/z / Da = 242.0925, [M+Na]⁺ found, [C₁₀H₁₈ClNNaO₂]⁺ requires 242.0924

The compound has not been reported previously.

0.35 4-Azido-*N*-((*trans*)-2-hydroxycyclohexyl)butanamide **137**



4-Chloro-*N*-((*trans*)-2-hydroxycyclohexyl)butanamide **136** (345 mg, 1.57 mmol, 1 eq.) and NaN₃ (180 mg, 2.77 mmol, 1.75 eq.) were stirred in DMF (12 ml) at 50 °C for 16 h. Water (50 ml) and 10 % *i*-PrOH/CHCl₃ (50 ml) were added, and the organic layer was removed. The aqueous layer was extracted again with 10 % *i*-PrOH/CHCl₃ (50 ml) and the combined organic fractions were dried with MgSO₄. The solvent was evaporated under reduced pressure, and then by using a N₂ stream. **137** was obtained as large white prisms (347 mg, 1.53 mmol, 97.5 %).

TLC R_f = 0.23 (EtOAc)

mp T / °C = 74.5-75.7 (*i*-PrOH, CHCl₃)

IR (neat) ν_{max} / cm⁻¹ = 3299.0 (N-H), 3207.8 (O-H), 2944.3 (C-H), 2927.9 (C-H), 2859.2 (C-H), 2089.2 (azide), 1624.0 (amide C=O)


¹H NMR (400 MHz, MeOD) δ / ppm = 7.87 (d, J = 7.9 Hz, 1 H, NH), 5.27 (d, J = 4.3 Hz, 1 H, OH), 3.56 (td, J = 10.5, 4.4 Hz, 1 H, CHNH), 3.28 - 3.41 (m, 3 H, CHOH and CH₂N₃), 2.30 (td, J = 7.4, 2.7 Hz, 2 H, C(=O)CH₂), 1.95 - 2.03 (m, 1 H, CHHCHOH), 1.87 (m, 3 H, C(=O)CH₂CH₂ and CHHCHNH), 1.70 - 1.76 (m, 1 H, CHHCH₂CHOH), 1.63 - 1.70 (m, 1 H, CHHCH₂CHNH), 1.25 - 1.38 (m, 3 H, CHHCH₂CHOH, CHHCH₂CHNH and CHHCHOH), 1.14 - 1.24 (m, 1 H, CHHCHNH₂)

¹³C NMR (101 MHz, MeOD) δ / ppm = 175.1 (C(=O)), 74.0 (CHOH), 56.3 (CHNH), 52.0 (CH₂N₃), 35.5 (CH₂CHOH), 34.3 (C(=O)CH₂), 32.7 (CH₂CHNH), 26.3 (C(=O)CH₂CH₂), 25.8 (CH₂CH₂CHNH), 25.5 (CH₂CH₂CHOH)

HRMS (ESI⁺) m/z / Da = 249.1331, [M+Na]⁺ found, [C₁₀H₁₈N₄NaO₂]⁺ requires 249.1327

The compound has not been reported previously.

0.36 Methyl 1-cyclopropyl-6-fluoro-7-(4-(4-(((*trans*)-2-hydroxycyclohexyl)amino)-4-oxobutyl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylate **138**



4-(4-(1-Cyclopropyl-6-fluoro-3-(methoxycarbonyl)-4-oxo-1,4-dihydroquinolin-7-yl)piperazin-1-yl)butanoic acid trifluoroacetate **123** (200 mg, 0.367 mmol, 1 eq.), (*trans*)-2-aminocyclohexan-1-ol **135** (91.1 mg, 0.791 mmol, 2.1 eq.), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (112 mg, 0.584 mmol, 1.6 eq.), 1-hydroxybenzotriazole (96 mg, 0.710 mmol, 1.9 eq.) and DIPEA (192 μ l, 142 mg, 1.10 mmol, 3 eq.) were dissolved in DMF (5 ml) and stirred at r.t. for 16 h. The solvent was removed using a stream of N₂ and the residue was purified by preparatory HPLC (5-50 % acetonitrile/water over 10 min). The combined pure fractions were evaporated under reduced pressure and then partitioned between NaHCO₃ (aq., sat., 10 ml) and CH₂Cl₂ (10 ml). The organic layer was dried with MgSO₄ and evaporated under reduced pressure. **138** was obtained as a white amorphous solid (73.0 mg, 0.142 mmol, 38.7 %).

IR (neat) ν_{max} / cm⁻¹ = 3302.5 (N-H), 2929.8 (C-H), 2850.6 (C-H), 2832.9 (C-H), 1698.1 (ester C=O), 1646.4 (amide C=O), 1613.8 (quinolone C=O)

¹H NMR (400 MHz, MeOD) δ / ppm = 8.60 (s, 1 H, *ortho* to C(=O)OCH₃), 7.79 (d, J = 13.5 Hz, 1 H, *ortho* to F), 7.46 (d, J = 7.2 Hz, 1 H, *meta* to F), 3.84 (s, 3 H, CH₃), 3.62 - 3.68 (m, 1 H, NCH(CH₂)₂), 3.58 (td, J = 10.3, 4.2 Hz, 1 H, CHNH), 3.38 (br s, 4 H, CH₂N(CH₂CH₂)CH₂CH₂), 3.32 - 3.36 (m, 1 H, CHOH), 2.83 (br s, 4 H, CH₂N(CH₂)CH₂), 2.60 (t, J = 7.3 Hz, 2 H, C(=O)CH₂CH₂CH₂N), 2.32 (td, J = 7.1, 3.1 Hz, 2 H, C(=O)CH₂), 1.96 - 2.04 (m, 1 H, CHHCHOH), 1.87 - 1.96 (m, 3 H, CHHCHNH and C(=O)CH₂CH₂), 1.72 - 1.77 (m, 1 H, CHHCH₂CHOH), 1.66 - 1.72 (m, 1 H, CHHCH₂CHNH), 1.25 - 1.39 (m, 5 H, CHHCHOH, CHHCH₂CHOH, CHHCH₂CHNH and NCH(CHH)₂), 1.15 - 1.25 (m, 3 H, CHHCHOH and NCH(CHH)₂)


¹³C NMR (101 MHz, MeOD) δ / ppm = 175.8 (CH₂C(=O)NH), 175.3 (C(=O)CC(=O)OCH₃), 166.8 (C(=O)OCH₃), 154.9 (d, J = 248.8 Hz, *ipso* to F), 150.2 (C=CC(=O)OCH₃), 146.1 (d, J = 10.8 Hz, *ipso* to piperazine), 139.9 (*para* to F), 123.5 (d, J = 7.5 Hz, *para* to piperazine), 113.2 (d, J = 23.2 Hz, *ortho* to C=O and *ortho* to F), 110.2 (CC(=O)OCH₃), 107.2 (*meta* to C=O and *meta* to F), 74.1 (CHOH), 58.9 (C(=O)CH₂CH₂CH₂N), 56.4 (CHNH), 54.0 (C(=O)CH₂CH₂CH₂N(CH₂)CH₂), 52.3 (CH₃), 50.5 (d, J = 5.0 Hz, C(=O)CH₂CH₂CH₂N(CH₂)CH₂CH₂), 36.4 (NCH(CH₂)₂), 35.7 (CH₂CHOH), 35.1 (C(=O)CH₂), 32.8 (CH₂CHNH), 25.9 (CH₂CH₂CHNH), 25.5 (CH₂CH₂CHOH), 23.5 (C(=O)CH₂CH₂), 8.7 (NCH(CH₂)₂)

¹⁹F NMR (376.45 MHz, MeOD) δ / ppm = -124.7 (ciprofloxacin F)

HRMS (ESI⁺) m/z / Da = 529.2827, [M+H]⁺ found, [C₂₈H₃₈FN₄O₅]⁺ requires 529.2826

The compound has not been reported previously.

0.37 Methyl 1-cyclopropyl-6-fluoro-4-oxo-7-(4-(4-oxo-4-((2-oxocyclohexyl)amino)-butyl)piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylate **139**



Methyl 1-cyclopropyl-6-fluoro-7-(4-(4-(((*trans*)-2-hydroxycyclohexyl)amino)-4-oxobutyl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylate **138** (5.2 mg, 9.84 μ mol, 1 eq.) and Dess-Martin periodane (16.4 mg, 38.7 μ mol, 4 eq.) were stirred in CH_2Cl_2 (3 ml) for 6 h. The solvent was removed under reduced pressure and the residue was purified by preparatory HPLC (5-95 % acetonitrile/water over 20 min). The combined pure fractions were evaporated under reduced pressure to a volume of 20 ml, then NaHCO_3 (aq., sat., 30 ml) and 10 % *i*-PrOH/ CHCl_3 (30 ml) were added. The organic layer was dried with MgSO_4 and evaporated under reduced pressure. **139** was obtained as a white amorphous solid (3.6 mg, 6.8 μ mol, 69.1 %).

TLC R_f = 0.74 (30 % MeOH/ CH_2Cl_2)

IR (neat) ν_{max} / cm^{-1} = 2921.2 (C-H), 2851.6 (C-H), 1721.4 (ketone C=O), 1698.0 (ester C=O), 1639.3 (amide C=O), 1620.0 (quinolone C=O)

^1H NMR (400 MHz, DMSO d_6) δ / ppm = 8.45 (s, 1 H, *ortho* to C(=O)OCH₃), 7.87 (d, J = 6.2 Hz, 1 H, NH), 7.76 (d, J = 13.4 Hz, 1 H, *ortho* to F), 7.44 (d, J = 7.5 Hz, 1 H, *meta* to F), 4.42 (dddd, J = 13.0, 7.6, 6.0, 1.0 Hz, 1 H, CHNH), 3.73 (s, 3 H, CH₃), 3.65 (tt, J = 7.1, 3.9 Hz, 1 H, NCH(CH₂)₂), 3.25 (br s, 4 H, CH₂N(CH₂CH₂)CH₂CH₂), 2.58 (br s, 4 H, CH₂N(CH₂)CH₂), 2.45 - 2.53 (m, 1 H, CHHC(=O)CHNH), 2.36 (br s, 2 H, C(=O)CH₂CH₂CH₂N), 2.26 (dtt, J = 13.4, 2.6, 2.6, 1.6, 1.6 Hz, 1 H, CHHC(=O)CHNH), 2.16 - 2.22 (m, 2 H, C(=O)CH₂CH₂CH₂N), 2.12 (ddq, J = 12.7, 6.0, 2.8, 2.8, 2.8 Hz, 1 H, CHHCHNH), 2.00 (ddquin, J = 13.2, 6.0, 2.9, 2.9, 2.9, 2.9 Hz, 1 H, CHHCH₂C(=O)), 1.65 - 1.83 (m, 4 H, CH₂CH₂CHNH), 1.41 - 1.56 (m, 2 H, CHHCHNH and CHHCH₂C(=O)), 1.20 - 1.30 (m, 2 H, NCH(CH₂)₂), 1.05 - 1.13 (m, 2 H, NCH(CH₂)₂)


^{13}C NMR (101 MHz, DMSO d_6) δ / ppm = 207.5 (C(=O)CHNH), 171.7 (C(=O)CC(=O)OCH₃), 171.6 (CH₂C(=O)NH), 165.0 (C(=O)OCH₃), 152.6 (d, J = 247.6 Hz, *ipso* to F), 148.3 (C=CC(=O)OCH₃), 143.9 (br s, *ipso* to piperazine), 138.1 (*para* to F), 121.8 (d, J = 6.4 Hz, *para* to piperazine), 111.5 (d, J = 22.4 Hz, *ortho* to C=O and *ortho* to F), 109.0 (CC(=O)OCH₃), 106.3 (*meta* to C=O and *meta* to F), 57.0 (CHNH and C(=O)CH₂CH₂CH₂N), 52.3 (br s, C(=O)CH₂CH₂CH₂N(CH₂)CH₂), 51.3 (CH₃), 49.5 (br s, C(=O)CH₂CH₂CH₂N(CH₂)CH₂), 40.6 (CH₂C(=O)CHNH), 34.8 (NCH(CH₂)₂), 33.9 (CH₂CHNH), 32.9 (C(=O)CH₂CH₂CH₂N), 27.2 (CH₂CH₂C(=O)CHNH), 23.8 (CH₂CH₂CHNH), 22.4 (br s, C(=O)CH₂CH₂CH₂N), 7.6 (NCH(CH₂)₂)

^{19}F NMR (376.45 MHz, DMSO d_6) δ / ppm = -124.3 (ciprofloxacin F)

HRMS (ESI⁺) m/z / Da = 527.2654, $[\text{M}+\text{H}]^+$ found, $[\text{C}_{28}\text{H}_{36}\text{FN}_4\text{O}_5]^+$ requires 527.2670

The compound has not been reported previously.

0.38 1-Cyclopropyl-6-fluoro-7-(4-(4-(1-(4-(((*trans*)-2-hydroxycyclohexyl)amino)-4-oxobutyl)-1*H*-1,2,3-triazol-4-yl)butyl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylic acid **140**



1-Cyclopropyl-6-fluoro-7-(4-(hex-5-yn-1-yl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylic acid **70** (40 mg, 97.2 μmol , 1 eq.) and 4-azido-*N*-(((*trans*)-2-hydroxycyclohexyl)butanamide **137** (22.0 mg, 97.2 μmol , 1 eq.) were dissolved in 10 % water/*t*-BuOH (3 ml), and the mixture was degassed by bubbling N_2 through it. A solution of CuSO_4 and THPTA (97.2 μl , 9.72 μmol , 0.1 eq. 100 mM, aq.) was added, followed by a solution of sodium ascorbate (194 μl , 19.4 μmol , 0.2 eq., 100 mM, aq.). The mixture was stirred at r.t. under argon for 16 h. Water (50 ml) and 10 % *i*-PrOH/ CHCl_3 (50 ml) were added, then the organic layer was separated, dried with MgSO_4 and evaporated under reduced pressure. The residue was purified by preparatory HPLC (5-70 % acetonitrile/water over 15 min). The combined pure fractions were evaporated under reduced pressure and then partitioned between NaHCO_3 (aq., sat., 50 ml) and 10 % *i*-PrOH/ CHCl_3 (50 ml). The organic layer was dried with MgSO_4 and evaporated under reduced pressure. **140** was obtained as a white amorphous solid (30.3 mg, 47.5 μmol , 48.9 %).

IR (neat) ν_{max} / cm^{-1} = 3345.4 (N-H), 2927.6 (C-H), 2859.6 (C-H), 2814.7 (C-H), 1727.0 (carboxylic acid C=O), 1641.7 (amide C=O), 1625.8 (quinolone C=O), 1619.0 (triazole)

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ / ppm = 8.64 (s, 1 H, *ortho* to C(=O)OH), 7.86 (d, J = 13.9 Hz, 1 H, *ortho* to F), 7.84 (s, 1 H, $\text{CH}=\text{CCH}_2$), 7.64 (d, J = 8.1 Hz, 1 H, NH), 7.54 (d, J = 7.5 Hz, 1 H, *meta* to F), 4.54 (d, J = 4.7 Hz, 1 H, OH), 4.30 (t, J = 6.8 Hz, 2 H, C(=O) $\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 3.77 - 3.86 (m, 1 H, $\text{NCH}(\text{CH}_2)_2$), 3.33 - 3.40 (m, 1 H, CHNH), 3.31 (br t, J = 4.8, 4.8 Hz, 4 H, $\text{CH}=\text{CCH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2\text{CH}_2$), 3.14 - 3.24 (m, 1 H, CHOH), 2.63 (t, J = 7.4 Hz, 2 H, $\text{CH}=\text{CCH}_2$), 2.56 (br t, J = 4.6, 4.6 Hz, 4 H, $\text{CH}=\text{CCH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2)\text{CH}_2$), 2.38 (t, J = 6.9 Hz, 2 H, $\text{CH}=\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 2.04 - 2.08 (m, 2 H, C(=O) $\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 1.96 - 2.04 (m, 2 H, C(=O) $\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 1.78 - 1.87 (m, 1 H, CHHCHOH), 1.69 - 1.78 (m, 1 H, CHHCHNH), 1.63 (quin, J = 7.5 Hz, 2 H, $\text{CH}=\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 1.54 - 1.60 (m, 1 H, CHHCH_2OH), 1.51 (quin, J = 7.4 Hz, 2 H, $\text{CH}=\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 1.28 - 1.35 (m, 1 H, $\text{NCH}(\text{CHH})_2$), 1.11 - 1.22 (m, 5 H, $\text{NCH}(\text{CHH})_2$, CHHCHOH , $\text{CHHCH}_2\text{CHOH}$ and $\text{CH}_2\text{CH}_2\text{CHNH}$), 1.04 - 1.13 (m, 1 H, CHHCHNH)


^{13}C NMR (101 MHz, $\text{DMSO}-d_6$) δ / ppm = 176.4 (C(=O)CC(=O)OH), 170.9 ($\text{CH}_2\text{C}(\text{=O})\text{NH}$), 166.0 (C(=O)OH), 153.1 (d, J = 252.1 Hz, *ipso* to F), 148.0 (C=CC(=O)OH), 146.9 ($\text{CH}=\text{CCH}_2$), 145.3 (d, J = 10.0 Hz, *ipso* to piperazine), 139.2 (*para* to F), 121.8 ($\text{NCH}=\text{CCH}_2$), 118.5 (d, J = 8.3 Hz, *para* to piperazine), 110.9 (d, J = 23.2 Hz, *ortho* to C=O and *ortho* to F), 106.7 (CC(=O)OH), 106.3 (d, J = 3.3 Hz, *meta* to C=O and *meta* to F), 71.4 (CHOH), 57.4 ($\text{CH}=\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 54.2 (CHNH), 52.4 ($\text{CH}=\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2)\text{CH}_2$), 49.5 ($\text{CH}=\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2\text{CH}_2$), 49.5 ($\text{CH}=\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2)\text{CH}_2\text{CH}_2$), 48.8 (C(=O) $\text{CH}_2\text{CH}_2\text{CH}_2\text{NCH}=\text{C}$), 35.9 ($\text{NCH}(\text{CH}_2)_2$), 34.1 (CH_2CHOH), 32.3 (C(=O) $\text{CH}_2\text{CH}_2\text{CH}_2\text{NCH}=\text{C}$), 31.1 (CH_2CHNH), 26.9 ($\text{CH}=\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 26.1 (C(=O) $\text{CH}_2\text{CH}_2\text{CH}_2\text{NCH}=\text{C}$), 25.8 ($\text{CH}=\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 25.0 ($\text{CH}=\text{CCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{N}$), 24.2 ($\text{CH}_2\text{CH}_2\text{CHNH}$), 23.8 ($\text{CH}_2\text{CH}_2\text{CHOH}$), 7.6 ($\text{NCH}(\text{CH}_2)_2$)

¹⁹F NMR (376.45 MHz, DMSO d₆) δ / ppm = -121.4 (ciprofloxacin F)

HRMS (ESI⁺) *m/z* / Da = 638.3480, [M+H]⁺ found, [C₃₃H₄₅FN₇O₅]⁺ requires 638.3466

The compound has not been reported previously.

0.39 1-Cyclopropyl-6-fluoro-4-oxo-7-(4-(4-(1-(4-oxo-4-((2-oxocyclohexyl)amino)butyl)-1*H*-1,2,3-triazol-4-yl)butyl)piperazin-1-yl)-1,4-dihydroquinoline-3-carboxylic acid **141**



1-Cyclopropyl-6-fluoro-7-(4-(4-(1-(4-(((*trans*)-2-hydroxycyclohexyl)amino)-4-oxobutyl)-1*H*-1,2,3-triazol-4-yl)butyl)piperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylic acid **140** (15.0 mg, 23.6 mmol, 1 eq.) and Dess-Martin periodane (35.0 mg, 82.5 mmol, 3.5 eq.) were stirred in CH₂Cl₂ (3 ml) for 4 h. The solvent was removed under reduced pressure and the residue was purified by preparatory HPLC (5-70 % acetonitrile/water over 15 min). The combined pure fractions were evaporated under reduced pressure, then NaHCO₃ (aq., sat., 30 ml) and 10 % *i*-PrOH/CHCl₃ (30 ml) were added. The organic layer was dried with MgSO₄ and evaporated under reduced pressure. **141** was obtained as a clear gum (11.7 mg, 18.4 μmol, 78.0 %).

IR (neat) ν_{max} / cm⁻¹ = 2941.2 (C-H), 2859.8 (C-H), 1719.8 (carboxylic acid C=O and ketone C=O), 1656.8 (amide C=O), 1625.6 (quinolone C=O), 1613.5 (triazole)

¹H NMR (500 MHz, DMSO d₆) δ / ppm = 8.65 (s, 1 H, *ortho* to C(=O)OH), 7.94 (d, J=7.7 Hz, 1 H, NH), 7.88 (d, J=13.4 Hz, 1 H, *ortho* to F), 7.85 (s, 1 H, CH=CCH₂), 7.55 (d, J=7.3 Hz, 1 H, *meta* to F), 4.40 (dddd, J=12.8, 7.6, 6.1, 1.1 Hz, 1 H), 4.31 (t, J=7.0 Hz, 1 H, C(=O)CH₂CH₂CHHN), 4.31 (t, J=6.9 Hz, 1 H, C(=O)CH₂CH₂CH₂N), 3.74 - 3.84 (m, 1 H, NCH(CH₂)₂), 3.31 (br. s, 4 H, CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 2.64 (t, J=7.5 Hz, 2 H, CH=CCH₂), 2.56 (br t, J=5.0, 5.0 Hz, 4 H, CH₂CH₂CH₂N(CH₂)CH₂), 2.45 - 2.52 (m, 1 H, CHHC(=O)), 2.38 (t, J=7.1 Hz, 2 H, CH=CCH₂CH₂CH₂CH₂N), 2.25 (dt, J=13.4, 2.6, 2.6, 1.6, 1.6 Hz, 1 H, CHHC(=O)), 2.07 - 2.17 (m, 3 H, C(=O)CH₂CH₂CH₂N and CHHCNH), 1.96 - 2.05 (m, 3 H, C(=O)CH₂CH₂CH₂N and CHHCCH₂C(=O)), 1.68 - 1.81 (m, 2 H, CHHCCH₂CHNH), 1.64 (quin, J=7.5 Hz, 2 H, CH=CCH₂CH₂CH₂CH₂N), 1.40 - 1.56 (m, 5 H, CHHCCH₂C(=O), CHHCNH and CH=CCH₂CH₂CH₂CH₂N), 1.27 - 1.34 (m, 2 H, NCH(CH₂)₂), 1.13 - 1.20 (m, 2 H, NCH(CH₂)₂)

¹³C NMR (126 MHz, DMSO d₆) δ / ppm = 207.4 (C(=O)CHNH), 176.3 (C(=O)CC(=O)OH), 170.8 (CH₂C(=O)NH), 166.0 (C(=O)OH), 153.0 (d, J=246.4 Hz, *ipso* to F), 147.9 (C=CC(=O)OH), 146.8 (CH=CCH₂), 145.1 (d, J=10.1 Hz, *ipso* to piperazine), 139.1 (*para* to F), 121.7 (NCH=CCH₂), 118.7 (d, J=6.9 Hz, *para* to piperazine), 110.9 (d, J=23.0 Hz, *ortho* to C=O and *ortho* to F), 106.3 (CC(=O)OH, and *meta* to C=O and *meta* to F), 57.3 (CH=CCH₂CH₂CH₂CH₂N), 57.0 (CHNH), 52.4 (CH₂CH₂CH₂N(CH₂)CH₂), 49.5 (CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 49.5 (CH₂CH₂CH₂N(CH₂CH₂)CH₂CH₂), 48.7 (C(=O)CH₂CH₂CH₂NCH=C), 40.5 (CH₂C(=O)), 35.8 (NCH(CH₂)₂), 33.7 (CH₂CHNH), 31.8 (C(=O)CH₂CH₂CH₂NCH=C), 27.1 (CH₂CH₂C(=O)),

26.9 (CH=CCH₂CH₂CH₂CH₂N), 26.0 (C(=O)CH₂CH₂CH₂NCH=C), 25.7 (CH=CCH₂CH₂CH₂CH₂N), 24.9 (CH=CCH₂CH₂CH₂CH₂N), 23.8 (CH₂CH₂CHNH), 7.6 (NCH(CH₂)₂)

¹⁹F NMR (376 MHz, DMSO d₆) δ / ppm = -121.7 (s, ciprofloxacin F)

HRMS (ESI⁺) m/z / Da = 636.3303, [M+H]⁺ found, [C₃₃H₄₃FN₇O₅]⁺ requires 636.3310

The compound has not been reported previously.

1 References

- [1] K. Sachin, E.-M. Kim, S.-J. Cheong, H.-J. Jeong, S. T. Lim, M.-H. Sohn and D. W. Kim. Synthesis of N_4' - $[^{18}F]$ fluoroalkylated ciprofloxacin as a potential bacterial infection imaging agent for PET study. *Bioconjugate Chemistry*, 21(12):2282–2288. 2010.
- [2] K. Ganguly, R. Wu, M. Ollivault-Shiflett, P. M. Goodwin, L. A. Silks and R. Iyer. Design, synthesis, and a novel application of quorum-sensing agonists as potential drug-delivery vehicles. *Journal of Drug Targeting*, 19(7):528–539. 2011.
- [3] R. Iyer, K. Ganguly and L. A. Silks. Synthetic analogs of bacterial quorum sensors. Los Alamos National Laboratory. 2012.
- [4] R. Srinivasan, L. P. Tan, H. Wu, P.-Y. Yang, K. A. Kalesh and S. Q. Yao. High-throughput synthesis of azide libraries suitable for direct "click" chemistry and in situ screening. *Organic & Biomolecular Chemistry*, 7(9):1821. 2009.
- [5] J. Aubé, Michael S. Wolfe, R. K. Yantiss, S. M. Cook, F. Takusagawa, M. S. Wolfe, R. K. Yantiss, S. M. Cook and F. Takusagawa. Synthesis of Enantiopure N-tert-Butoxycarbonyl-2- aminocycloalkanones. *Synthetic Communications*, 22(20):3003–3012. 1992.
- [6] L. E. Overman, S. Sugai, L. E. Overman and S. Sugai. A Convenient Method for Obtaining trans -2-Aminocyclohexanol and trans -2-Aminocyclopentanol in Enantiomerically Pure Form. *The Journal of Organic Chemistry*, 50:4154–4155. 1985.
- [7] L. E. Overman and S. Sugai. Total Synthesis of (-)-Crinine. Use of Tandem Cationic Aza-Cope Rearrangement/Mannich Cyclizations for the Synthesis of Enantiomerically Pure Amaryllidaceae Alkaloids. *Helvetica Chimica Acta*, 68(3):745–749. 1985.
- [8] I. Schiffers, T. Rantanen, F. Schmidt, W. Bergmans, L. Zani and C. Bolm. Resolution of racemic 2-aminocyclohexanol derivatives and their application as ligands in asymmetric catalysis. *The Journal of Organic Chemistry*, 71(1):2320–2331. 2006.
- [9] F. Xue and C. T. Seto. Structure-activity studies of cyclic ketone inhibitors of the serine protease plasmin: Design, synthesis, and biological activity. *Bioorganic & Medicinal Chemistry*, 14:8467–8487. 2006.

Todo list