NEW WORLD SCREWWORM

Detection history across Guatemala, Mexico, and the United States

Carter Grout | Texas A&M University | cartergrout@tamu.edu

PROJECT OVERVIEW

NWS (*Cochliomyia hominivorax*) larvae feed on living tissue of animals and people. [1] The recent outbreak re-emerged in Mexico in November 2024, with the first U.S. animal case in June 2026, renewing the need to understand detection history. [2]

MY RESEARCH ROLE

Identify gaps in detection history and investigate possible routes and locations of entry into the United States before detection.


METHODS

Date-linked records were mapped in a timeline animation to examine changes in reported detections over time. Dates differ by source: outbreak start, publisher timestamp, or report date; they do not directly date transmission.

FINDINGS & NEXT STEPS

Mapped detection records
Guatemala: **152** points
Mexico: **26,480** points
U.S.: **54** reports (53 animal + 1 fly-trap)

The sparse detection history shown in northwestern Mexico is worth investigating alongside detections in New Mexico. Comparing dates and reporting coverage may help assess possible entry routes; this map alone does not establish a connection.



Reported NWS detections: Guatemala and Mexico points; U.S. county totals.

Data sources: Guatemala - WOAH WAHIS; Mexico - SENASICA; United States - USDA APHIS.

[1] USDA APHIS: NWS Prevention for Animals. [2] CDC: New World Screwworm Outbreak (cdc.gov/new-world-screwworm/situation-summary).