	Approach	*Community*	*Interactions*	*Techniques*	*Interaction types*	*Strengths*	*Limitations*
A. Realized interactions	**Area-based**	Field data	Field data	Diet analysis Metabarcoding Stable isotope	Food webs: predation herbivory scavenging propagule dispersal	Enables assessment of community network dynamics; captures local ecological context (e,g, abundance, phenology,…)	Labour- and cost-intensive; Scalability requires long-term monitoring networks
	Individual-based	Field data	Field data	Cameras Barcoding	Frugivory Pollination Herbivory Mycorrhizal Decomposition Host-parasite Host-pathogen	Enables multi-interaction assessment at the species level; repeatable for the same individuals; captures local ecological context	Labour- and cost-intensive. Primarily applicable to sessile or long-lived organisms
B. Potential interactions	**Co-occurrence**	Field data	Inferred	Species inventories (e.g. eDNA)	All interaction types	Scalable; allows reuse of existing diversity datasets	Risk of interaction overestimation; high uncertainty
Meta-network 	**Field meta-network inference**	Field data	Inferred	Combined field inventories with meta-network	Food web Frugivory Pollination Host-parasite	Reuses existing diversity data; enables Refinement of meta-network with field observations	Uncertainty depends on meta-network completeness; potential interaction-type bias
SDM 	**SDM meta-network inference**	Inferred	Inferred	Combined SDM with meta-network	Food web Frugivory Pollination Host-parasite	Spatially explicit; suitable for large-scale (e.g., national) assessments	High uncertainty from both community and interactions inference; interaction-type bias