

# Symmetric neural networks

YKY

May 16, 2019

## 1 General case for $y = Ax$

$$\boxed{\text{original}} \quad y_j = \sum_i a_{ij} x_i. \quad (1)$$

Equivariance implies:

$$\begin{aligned} \boxed{\text{swapped}} \quad y_j \cdot \sigma(x_j \ x_k) &= y_k \quad \boxed{\text{original}} \\ \sum_{i \neq j, k} a_{ij} x_i + a_{kj} x_j + a_{jj} x_k &= \sum_{i \neq j, k} a_{ik} x_i + a_{jk} x_j + a_{kk} x_k. \end{aligned} \quad (2)$$

Comparing coefficients yields:

$$\begin{aligned} a_{ij} &= a_{ik} & \forall j, k, (i \neq j, k) \\ a_{kj} &= a_{jk} & \forall j, k \\ a_{jj} &= a_{kk} & \forall j, k. \end{aligned} \quad (3)$$

In other words, the matrix  $A$  is of the form:

$$A = \alpha I + \beta 11^T. \quad (4)$$

## 2 Case for $y_k = A_k x \cdot x$

The general form of a “quadratic” vector function is:

$$y = (Ax) \cdot x + Bx + C. \quad (5)$$

We just focus on the quadratic term  $(Ax) \cdot x$ :

$$\boxed{\text{original}} \quad y_k = \sum_j \left[ \sum_i a_{ij}^k x_i \right] x_j. \quad (6)$$

Equivariance implies:

$$\boxed{\text{swapped}} \quad y_k \cdot \sigma(x_k \ x_h) = y_h \quad \boxed{\text{original}} \quad (7)$$

$$\sum_{j \neq h, k} \sum_{i \neq h, k} a_{ij}^k x_i x_j + a_{hh}^k x_k^2 + a_{kh}^k x_h x_k + a_{hk}^k x_k x_h + a_{kk}^k x_h^2 = \sum_{j \neq h, k} \sum_{i \neq h, k} a_{ij}^h x_i x_j + a_{hh}^h x_h^2 + a_{kh}^h x_k x_h + a_{hk}^h x_h x_k + a_{kk}^h x_k^2$$

which yields:

$$\begin{aligned} a_{ij}^h &= a_{ij}^k & \forall h, k, (i, j) \neq (h, k) \\ a_{hh}^h &= a_{kk}^k & \forall h, k \\ a_{hh}^k &= a_{kk}^h & \forall h, k \\ a_{kh}^k + a_{hk}^k &= a_{kh}^h + a_{hk}^h & \forall h, k. \end{aligned} \quad (8)$$