

A PIE Representation of coupled 2D PDEs and Stability Analysis using LPIs

Declan S. Jagt, Matthew M. Peet

Arizona State University
Tempe, AZ USA

American Control Conference
Atlanta, GA USA

June 8, 2022



Lyapunov Stability Analysis of PDEs Requires an Algebra of Operators on L_2

Consider a 2D PDE

$$\begin{aligned}\dot{u}(t, x, y) &= C[\partial_x + \partial_y]u(t, x, y), \\ (x, y) &\in [0, 1] \times [0, 1], \\ u(t, 0, y) &= u(t, x, 0) = 0.\end{aligned}$$

Letting $\mathbf{u}(t) = u(t) \in L_2^n [[0, 1] \times [0, 1]]$, stability can be certified with a quadratic Lyapunov Function (LF)

$$V(\mathbf{u}) = \langle \mathbf{u}, \mathcal{P}\mathbf{u} \rangle,$$

parameterized by some $\mathcal{P} : L_2^n \rightarrow L_2^n$.

Lyapunov Stability Analysis of PDEs Requires an Algebra of Operators on L_2

Consider a 2D PDE

$$\begin{aligned}\dot{u}(t, x, y) &= C[\partial_x + \partial_y]u(t, x, y), \\ (x, y) &\in [0, 1] \times [0, 1], \\ u(t, 0, y) &= u(t, x, 0) = 0.\end{aligned}$$

Letting $\mathbf{u}(t) = u(t) \in L_2^n[[0, 1] \times [0, 1]]$, stability can be certified with a quadratic Lyapunov Function (LF)

$$V(\mathbf{u}) = \langle \mathbf{u}, \mathcal{P}\mathbf{u} \rangle,$$

parameterized by some $\mathcal{P} : L_2^n \rightarrow L_2^n$.

Representing the PDE as

$$\begin{aligned}\dot{\mathbf{u}}(t) &= \mathcal{A}\mathbf{u}(t), \\ \mathbf{u}(t) &\in X \subset L_2^n[[0, 1] \times [0, 1]],\end{aligned}$$

the system is stable if and only if for some $\mathcal{P} > 0$,

$$\dot{V}(\mathbf{u}(t)) = \langle \mathbf{u}(t), [\mathcal{A}^*\mathcal{P} + \mathcal{P}\mathcal{A}]\mathbf{u}(t) \rangle \leq 0$$

along any solution $\mathbf{u}(t) \in X$.

Lyapunov Stability Analysis of PDEs Requires an Algebra of Operators on L_2

Consider a 2D PDE

$$\begin{aligned}\dot{u}(t, x, y) &= C[\partial_x + \partial_y]u(t, x, y), \\ (x, y) &\in [0, 1] \times [0, 1], \\ u(t, 0, y) &= u(t, x, 0) = 0.\end{aligned}$$

Letting $\mathbf{u}(t) = u(t) \in L_2^n[[0, 1] \times [0, 1]]$, stability can be certified with a quadratic Lyapunov Function (LF)

$$V(\mathbf{u}) = \langle \mathbf{u}, \mathcal{P}\mathbf{u} \rangle,$$

parameterized by some $\mathcal{P} : L_2^n \rightarrow L_2^n$.

Representing the PDE as

$$\begin{aligned}\dot{\mathbf{u}}(t) &= \mathcal{A}\mathbf{u}(t), \\ \mathbf{u}(t) &\in X \subset L_2^n[[0, 1] \times [0, 1]],\end{aligned}$$

the system is stable if and only if for some $\mathcal{P} > 0$,

$$\dot{V}(\mathbf{u}(t)) = \langle \mathbf{u}(t), [\mathcal{A}^*\mathcal{P} + \mathcal{P}\mathcal{A}]\mathbf{u}(t) \rangle \leq 0$$

along any solution $\mathbf{u}(t) \in X$.

How do we parameterize a set of operators \mathcal{P} that

- 1 act on infinite-dimensional states $\mathbf{u} \in L_2^n$,
- 2 is closed under addition and composition – i.e. is an **algebra**,
- 3 is suitably rich so as to avoid introducing significant conservatism?

An Algebra of Operators on **1D** States can be Parameterized by **3** Functions

For states $\mathbf{u} \in L_2^m[a, b]$ on a 1D domain $[a, b]$, we can parameterize an algebra of operators by 3 matrix-valued functions:



Definition 1 (1D-PI Operator)

For any $N = \{N_0, N_1, N_2\}$ with $N_i \in L_2^{n \times m}$, we define the associated 1D-PI operator

$$(\mathcal{P}[N]\mathbf{u})(x) = N_0(x)\mathbf{u}(x) + \int_a^x N_1(x, \theta)\mathbf{u}(\theta)d\theta + \int_x^b N_2(x, \theta)\mathbf{u}(\theta)d\theta \in L_2^n[a, b],$$

for arbitrary $\mathbf{u} \in L_2^m[a, b]$.

An Algebra of Operators on 1D States can be Parameterized by 3 Functions

For states $\mathbf{u} \in L_2^m[a, b]$ on a 1D domain $[a, b]$, we can parameterize an algebra of operators by 3 matrix-valued functions:



Definition 1 (1D-PI Operator)

For any $N = \{N_0, N_1, N_2\}$ with $N_i \in L_2^{n \times m}$, we define the associated 1D-PI operator

$$(\mathcal{P}[N]\mathbf{u})(x) = N_0(x)\mathbf{u}(x) + \int_a^x N_1(x, \theta)\mathbf{u}(\theta)d\theta + \int_x^b N_2(x, \theta)\mathbf{u}(\theta)d\theta \in L_2^n[a, b],$$

for arbitrary $\mathbf{u} \in L_2^m[a, b]$.


Proposition 2 (1D-PI Operators form an Algebra)

For any 1D-PI operators $\mathcal{B} : L_2^p \rightarrow L_2^n$ and $\mathcal{D} : L_2^m \rightarrow L_2^p$, there exists a 1D-PI operator $\mathcal{R} : L_2^{m \rightarrow n}$ such that for any $\mathbf{u} \in L_2^m$,

$$(\mathcal{B}(\mathcal{D}\mathbf{u}))(x) = (\mathcal{R}\mathbf{u})(x).$$

An Algebra of Operators on 2D States can be Parameterized by 9 Functions

Composing PI operators on $[a, b]$ and $[c, d]$,
we can define PI operators on the 2D
hypercube $\Omega := [a, b] \times [c, d]$ as:



Definition 3 (2D-PI Operator)


For any $N = \begin{bmatrix} N_{00} & N_{01} & N_{02} \\ N_{10} & N_{11} & N_{12} \\ N_{20} & N_{21} & N_{22} \end{bmatrix}$ with $N_{ij} \in L_2^{n \times m}$, we define the associated 2D-PI operator

$$\begin{aligned} (\mathcal{P}[N]\mathbf{u})(x, y) = & N_{00}(x, y)\mathbf{u}(x, y) + \int_a^x N_{01}(x, y, \theta)\mathbf{u}(\theta, y)d\theta + \int_x^b N_{02}(x, y, \theta)\mathbf{u}(\theta, y)d\theta \\ & + \int_c^y N_{10}(x, y, \nu)\mathbf{u}(x, \nu)d\nu + \int_a^x \int_c^y N_{11}(x, y, \theta, \nu)\mathbf{u}(\theta, \nu)d\theta d\nu + \int_x^b \int_c^y N_{12}(x, y, \theta, \nu)\mathbf{u}(\theta, \nu)d\theta d\nu \\ & + \int_y^d N_{20}(x, y, \nu)\mathbf{u}(x, \nu)d\nu + \int_a^x \int_y^d N_{21}(x, y, \theta, \nu)\mathbf{u}(\theta, \nu)d\theta d\nu + \int_x^b \int_y^d N_{22}(x, y, \theta, \nu)\mathbf{u}(\theta, \nu)d\theta d\nu \\ & \in L_2^n[\Omega] \end{aligned}$$

for arbitrary $\mathbf{u} \in L_2^m[\Omega]$.

An Algebra of Operators on 2D States can be Parameterized by 9 Functions

Composing PI operators on $[a, b]$ and $[c, d]$, we can define PI operators on the 2D hypercube $\Omega := [a, b] \times [c, d]$ as:



Definition 3 (2D-PI Operator)

For any $N = \begin{bmatrix} N_{00} & N_{01} & N_{02} \\ N_{10} & N_{11} & N_{12} \\ N_{20} & N_{21} & N_{22} \end{bmatrix}$ with $N_{ij} \in L_2^{n \times m}$, we define the associated 2D-PI operator

$$\begin{aligned} (\mathcal{P}[N]\mathbf{u})(x, y) = & N_{00}(x, y)\mathbf{u}(x, y) + \int_a^x N_{01}(x, y, \theta)\mathbf{u}(\theta, y)d\theta + \int_x^b N_{02}(x, y, \theta)\mathbf{u}(\theta, y)d\theta \\ & + \int_c^y N_{10}(x, y, \nu)\mathbf{u}(x, \nu)d\nu + \int_a^x \int_c^y N_{11}(x, y, \theta, \nu)\mathbf{u}(\theta, \nu)d\theta d\nu + \int_x^b \int_c^y N_{12}(x, y, \theta, \nu)\mathbf{u}(\theta, \nu)d\theta d\nu \\ & + \int_y^d N_{20}(x, y, \nu)\mathbf{u}(x, \nu)d\nu + \int_a^x \int_y^d N_{21}(x, y, \theta, \nu)\mathbf{u}(\theta, \nu)d\theta d\nu + \int_x^b \int_y^d N_{22}(x, y, \theta, \nu)\mathbf{u}(\theta, \nu)d\theta d\nu \\ & \in L_2^n[\Omega] \end{aligned}$$

for arbitrary $\mathbf{u} \in L_2^m[\Omega]$.

Proposition 4 (2D-PI Operators form an Algebra)

For any 2D-PI operators $\mathcal{B} : L_2^p \rightarrow L_2^n$ and $\mathcal{D} : L_2^m \rightarrow L_2^p$, there exists a 2D-PI operator $\mathcal{R} : L_2^{m \rightarrow n}$ such that for any $\mathbf{u} \in L_2^m$, $(\mathcal{B}(\mathcal{D}\mathbf{u}))(x) = (\mathcal{R}\mathbf{u})(x)$.

Consider a 2D PDE

$$\begin{aligned}\dot{u}(t, x, y) &= C[\partial_x + \partial_y]u(t, x, y), \\ (x, y) &\in [0, 1] \times [0, 1], \\ u(t, 0, y) &= u(t, x, 0) = 0.\end{aligned}$$

Consider a 2D PDE

$$\begin{aligned}\dot{u}(t, x, y) &= C[\partial_x + \partial_y]u(t, x, y), \\ (x, y) &\in [0, 1] \times [0, 1], \\ u(t, 0, y) &= u(t, x, 0) = 0.\end{aligned}$$

Then

- $\mathcal{A} = C[\partial_x + \partial_y]$,
- $X := \{u \in H_1^n \mid u(0, y) = u(x, 0) = 0\}$
where
 $H_1^n := \{\mathbf{u} \mid \partial_x \partial_y \mathbf{u} \in L_2^n\}$.

We can represent a linear PDE as

$$\begin{aligned}\dot{\mathbf{u}}(t) &= \mathcal{A}\mathbf{u}(t), \\ \mathbf{u}(t) &\in X \subset L_2^n[\Omega],\end{aligned}$$

where

- \mathcal{A} is a differential operator,
- X is a set defined by the boundary conditions (BCs) and continuity constraints.

Consider a 2D PDE

$$\begin{aligned}\dot{u}(t, x, y) &= C[\partial_x + \partial_y]u(t, x, y), \\ (x, y) &\in [0, 1] \times [0, 1], \\ u(t, 0, y) &= u(t, x, 0) = 0.\end{aligned}$$

Then

- $\mathcal{A} = C[\partial_x + \partial_y]$,
- $X := \{u \in H_1^n \mid u(0, y) = u(x, 0) = 0\}$
where
 $H_1^n := \{\mathbf{u} \mid \partial_x \partial_y \mathbf{u} \in L_2^n\}$.

We can represent a linear PDE as

$$\begin{aligned}\dot{\mathbf{u}}(t) &= \mathcal{A}\mathbf{u}(t), \\ \mathbf{u}(t) &\in X \subset L_2^n[\Omega],\end{aligned}$$

where

- \mathcal{A} is a differential operator,
- X is a set defined by the boundary conditions (BCs) and continuity constraints.

The system is stable if and only if for some $\mathcal{P} : L_2^n \rightarrow L_2^n$,

$$\begin{aligned}V(\mathbf{u}(t)) &= \langle \mathbf{u}(t), \mathcal{P}\mathbf{u}(t) \rangle > 0, \\ \dot{V}(\mathbf{u}(t)) &= \langle \mathbf{u}(t), [\mathcal{A}^* \mathcal{P} + \mathcal{P} \mathcal{A}]\mathbf{u}(t) \rangle \leq 0,\end{aligned}$$

along any solution $\mathbf{u}(t) \in X$.

Consider a 2D PDE

$$\begin{aligned}\dot{u}(t, x, y) &= C[\partial_x + \partial_y]u(t, x, y), \\ (x, y) &\in [0, 1] \times [0, 1], \\ u(t, 0, y) &= u(t, x, 0) = 0.\end{aligned}$$

Then

- $\mathcal{A} = C[\partial_x + \partial_y]$,
- $X := \{u \in H_1^n \mid u(0, y) = u(x, 0) = 0\}$
where
 $H_1^n := \{\mathbf{u} \mid \partial_x \partial_y \mathbf{u} \in L_2^n\}.$

We can represent a linear PDE as

$$\begin{aligned}\dot{\mathbf{u}}(t) &= \mathcal{A}\mathbf{u}(t), \\ \mathbf{u}(t) &\in X \subset L_2^n[\Omega],\end{aligned}$$

where

- \mathcal{A} is a differential operator,
- X is a set defined by the boundary conditions (BCs) and continuity constraints.

The system is stable if and only if for some $\mathcal{P} : L_2^n \rightarrow L_2^n$,

$$\begin{aligned}V(\mathbf{u}(t)) &= \langle \mathbf{u}(t), \mathcal{P}\mathbf{u}(t) \rangle > 0, \\ \dot{V}(\mathbf{u}(t)) &= \langle \mathbf{u}(t), [\mathcal{A}^* \mathcal{P} + \mathcal{P} \mathcal{A}]\mathbf{u}(t) \rangle \leq 0,\end{aligned}$$

along any solution $\mathbf{u}(t) \in X$.

Consider a 2D PDE

$$\begin{aligned}\dot{u}(t, x, y) &= C[\partial_x + \partial_y]u(t, x, y), \\ (x, y) &\in [0, 1] \times [0, 1], \\ u(t, 0, y) &= u(t, x, 0) = 0.\end{aligned}$$

Then

- $\mathcal{A} = C[\partial_x + \partial_y]$,
- $X := \{u \in H_1^n \mid u(0, y) = u(x, 0) = 0\}$
where
 $H_1^n := \{u \mid \partial_x \partial_y u \in L_2^n\}$.

We can represent a linear PDE as

$$\begin{aligned}\dot{\mathbf{u}}(t) &= \mathcal{A}\mathbf{u}(t), \\ \mathbf{u}(t) &\in X \subset L_2^n[\Omega],\end{aligned}$$

where

- \mathcal{A} is a differential operator,
- X is a set defined by the boundary conditions (BCs) and continuity constraints.

The system is stable if and only if for some $\mathcal{P} : L_2^n \rightarrow L_2^n$,

$$\begin{aligned}V(\mathbf{u}(t)) &= \langle \mathbf{u}(t), \mathcal{P}\mathbf{u}(t) \rangle > 0, \\ \dot{V}(\mathbf{u}(t)) &= \langle \mathbf{u}(t), [\mathcal{A}^*\mathcal{P} + \mathcal{P}\mathcal{A}]\mathbf{u}(t) \rangle \leq 0,\end{aligned}$$

along any solution $\mathbf{u}(t) \in X$.

Consider a 2D PDE

$$\begin{aligned}\dot{u}(t, x, y) &= C[\partial_x + \partial_y]u(t, x, y), \\ (x, y) &\in [0, 1] \times [0, 1], \\ u(t, 0, y) &= u(t, x, 0) = 0.\end{aligned}$$

Then

- $\mathcal{A} = C[\partial_x + \partial_y]$,
- $X := \{u \in H_1^n \mid u(0, y) = u(x, 0) = 0\}$
where
 $H_1^n := \{\mathbf{u} \mid \partial_x \partial_y \mathbf{u} \in L_2^n\}$.

We want to represent the system in a way that

- 1 is parameterized by PI operators;
- 2 incorporates the BCs and continuity conditions into the dynamics.

We can represent a linear PDE as

$$\begin{aligned}\dot{\mathbf{u}}(t) &= \mathcal{A}\mathbf{u}(t), \\ \mathbf{u}(t) &\in X \subset L_2^n[\Omega],\end{aligned}$$

where

- \mathcal{A} is a differential operator,
- X is a set defined by the boundary conditions (BCs) and continuity constraints.

The system is stable if and only if for some $\mathcal{P} : L_2^n \rightarrow L_2^n$,

$$\begin{aligned}V(\mathbf{u}(t)) &= \langle \mathbf{u}(t), \mathcal{P}\mathbf{u}(t) \rangle > 0, \\ \dot{V}(\mathbf{u}(t)) &= \langle \mathbf{u}(t), [\mathcal{A}^* \mathcal{P} + \mathcal{P} \mathcal{A}] \mathbf{u}(t) \rangle \leq 0,\end{aligned}$$

along any solution $\mathbf{u}(t) \in X$.

Consider a 2D PDE

$$\begin{aligned}\dot{u}(t, x, y) &= C[\partial_x + \partial_y]u(t, x, y), \\ (x, y) &\in [0, 1] \times [0, 1], \\ u(t, 0, y) &= u(t, x, 0) = 0.\end{aligned}$$

Consider a 2D PDE

$$\begin{aligned}\dot{u}(t, x, y) &= C[\partial_x + \partial_y]u(t, x, y), \\ (x, y) &\in [0, 1] \times [0, 1], \\ u(t, 0, y) &= u(t, x, 0) = 0.\end{aligned}$$

This system can be represented as

$$\begin{aligned}\dot{\mathbf{u}}(t) &= A_{10}\partial_x \mathbf{u}(t) + A_{01}\partial_y \mathbf{u}(t) \\ \mathbf{u}(t) &\in X,\end{aligned}$$

where $A_{10} = A_{01} = C$, and

$$X := \left\{ \mathbf{u} \in H_1^n \left[[0, 1] \times [0, 1] \right] \mid \begin{bmatrix} \mathbf{u}^{(0, y)} \\ \mathbf{u}_{(x, 0)} \end{bmatrix} = 0 \right\}$$

Linear, 2nd Order 2D PDEs are Parameterized by 9 Matrices

Consider a 2D PDE

$$\begin{aligned}\dot{\mathbf{u}}(t, x, y) &= C[\partial_x + \partial_y] \mathbf{u}(t, x, y), \\ (x, y) &\in [0, 1] \times [0, 1], \\ \mathbf{u}(t, 0, y) &= \mathbf{u}(t, x, 0) = 0.\end{aligned}$$

This system can be represented as

$$\begin{aligned}\dot{\mathbf{u}}(t) &= A_{10} \partial_x \mathbf{u}(t) + A_{01} \partial_y \mathbf{u}(t) \\ \mathbf{u}(t) &\in X,\end{aligned}$$

where $A_{10} = A_{01} = C$, and

$$X := \left\{ \mathbf{u} \in H_1^n([0, 1] \times [0, 1]) \mid \left[\begin{smallmatrix} \mathbf{u}^{(0,y)} \\ \mathbf{u}^{(x,0)} \end{smallmatrix} \right] = 0 \right\}$$

Any linear, 2nd order, 2D PDE can be represented as


$$\begin{aligned}\dot{\mathbf{u}}(t, x, y) &= A_{00} \begin{bmatrix} \mathbf{u}_0(t, x, y) \\ \mathbf{u}_1(t, x, y) \\ \mathbf{u}_2(t, x, y) \end{bmatrix} + A_{01} \partial_y \begin{bmatrix} \mathbf{u}_1(t, x, y) \\ \mathbf{u}_2(t, x, y) \end{bmatrix} + A_{02} \partial_y^2 \mathbf{u}_2(t, x, y) \\ &+ A_{10} \partial_x \begin{bmatrix} \mathbf{u}_1(t, x, y) \\ \mathbf{u}_2(t, x, y) \end{bmatrix} + A_{11} \partial_x \partial_y \begin{bmatrix} \mathbf{u}_1(t, x, y) \\ \mathbf{u}_2(t, x, y) \end{bmatrix} + A_{12} \partial_x \partial_y^2 \mathbf{u}_2(t, x, y) \\ &+ A_{20} \partial_x^2 \mathbf{u}_2(t, x, y) + A_{21} \partial_x^2 \partial_y \mathbf{u}_2(t, x, y) + A_{22} \partial_x^2 \partial_y^2 \mathbf{u}_2(t, x, y) \quad \forall (x, y) \in \Omega\end{aligned}$$

$$\mathbf{u}(t) \in X_{\mathcal{B}}[\Omega] := \left\{ \begin{bmatrix} \mathbf{u}_0 \\ \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix} \in \begin{bmatrix} L_2^{n_0}[\Omega] \\ H_1^{n_1}[\Omega] \\ H_2^{n_2}[\Omega] \end{bmatrix} \mid \mathcal{B} \Lambda_{\text{bf}} \mathbf{u} = 0 \right\}$$

where we define $H_k^n[\Omega] := \{ \mathbf{u} \mid \partial_x^{\alpha_1} \partial_y^{\alpha_2} \mathbf{u} \in L_2^n[\Omega], \forall \|\alpha\|_{\infty} \leq k \}$.

BCs in 2D can be Expressed in terms of a Standardized Boundary State

We represent the BCs as $\mathcal{B}\Lambda_{\text{bf}}\mathbf{u} = 0$,
where $\Lambda_{\text{bf}}\mathbf{u}$ is the *full boundary state*.



In 1D, the boundary values are all finite-dimensional.
Then, we can define

$$\Lambda_{\text{bf}}\mathbf{u} = \begin{bmatrix} \mathbf{u}_1(a) \\ \mathbf{u}_1(b) \\ \mathbf{u}_2(a) \\ \partial_x \mathbf{u}_2(a) \\ \mathbf{u}_2(b) \\ \partial_x \mathbf{u}_2(b) \end{bmatrix} \in \mathbb{R}^{2n_1+4n_2},$$


and impose boundary conditions as $B\Lambda_{\text{bf}}\mathbf{u} = 0$ for a matrix B .

BCs in 2D can be Expressed in terms of a Standardized Boundary State

We represent the BCs as $\mathcal{B}\Lambda_{\text{bf}}\mathbf{u} = 0$,
where $\Lambda_{\text{bf}}\mathbf{u}$ is the *full boundary state*.

In 2D, things are more complicated...

To define $\Lambda_{\text{bf}}\mathbf{u}$, we seek a “minimal”
representation of the boundary conditions.




BCs in 2D can be Expressed in terms of a Standardized Boundary State

We represent the BCs as $\mathcal{B}\Lambda_{\text{bf}}\mathbf{u} = 0$, where $\Lambda_{\text{bf}}\mathbf{u}$ is the *full boundary state*.

In 2D, things are more complicated...

To define $\Lambda_{\text{bf}}\mathbf{u}$, we seek a “minimal” representation of the boundary conditions.



Example 1

For $\mathbf{u}_1 \in H_1[\Omega]$, we can decompose

$$\mathbf{u}_1(x, c) = \mathbf{u}_1(a, c) + \int_a^x \partial_x \mathbf{u}_1(\theta, c) d\theta$$


So, we may enforce $\mathbf{u}_1(x, c) = 0$ as $\mathbf{u}_1(a, c) = 0$ and $\partial_x \mathbf{u}_1(x, c) = 0$.

BCs in 2D can be Expressed in terms of a Standardized Boundary State

We represent the BCs as $\mathcal{B}\Lambda_{\text{bf}}\mathbf{u} = 0$, where $\Lambda_{\text{bf}}\mathbf{u}$ is the *full boundary state*.

In 2D, things are more complicated...

To define $\Lambda_{\text{bf}}\mathbf{u}$, we seek a “minimal” representation of the boundary conditions.



Example 1

For $\mathbf{u}_1 \in H_1[\Omega]$, we can decompose

$$\mathbf{u}_1(x, c) = \mathbf{u}_1(a, c) + \int_a^x \partial_x \mathbf{u}_1(\theta, c) d\theta$$

So, we may enforce $\mathbf{u}_1(x, c) = 0$ as $\mathbf{u}_1(a, c) = 0$ and $\partial_x \mathbf{u}_1(x, c) = 0$.

Example 2

For $\mathbf{u}_2 \in H_2[\Omega]$, we can decompose

$$\mathbf{u}_2(b, y) = \mathbf{u}_2(b, d) + \partial_y \mathbf{u}_2(b, d)[y - d] + \int_d^y \partial_y^2 \mathbf{u}_2(b, \nu) d\nu$$


So, we may enforce $\mathbf{u}_2(b, y) = 0$ as $\mathbf{u}_2(b, d) = \partial_y \mathbf{u}_2(b, d) = 0$ and $\partial_y^2 \mathbf{u}_2(b, y) = 0$;

BCs in 2D can be Expressed in terms of a Standardized Boundary State

We represent the BCs as $\mathcal{B}\Lambda_{\text{bf}}\mathbf{u} = 0$, where $\Lambda_{\text{bf}}\mathbf{u}$ is the *full boundary state*.

In 2D, things are more complicated...

To define $\Lambda_{\text{bf}}\mathbf{u}$, we seek a “minimal” representation of the boundary conditions.



Example 1

For $\mathbf{u}_1 \in H_1[\Omega]$, we can decompose

$$\mathbf{u}_1(x, c) = \mathbf{u}_1(a, c) + \int_a^x \partial_x \mathbf{u}_1(\theta, c) d\theta$$

So, we may enforce $\mathbf{u}_1(x, c) = 0$ as $\mathbf{u}_1(a, c) = 0$ and $\partial_x \mathbf{u}_1(x, c) = 0$.

Example 2

For $\mathbf{u}_2 \in H_2[\Omega]$, we can decompose

$$\mathbf{u}_2(b, y) = \mathbf{u}_2(b, d) + \partial_y \mathbf{u}_2(b, d)[y - d] + \int_d^y \partial_y^2 \mathbf{u}_2(b, \nu) d\nu$$

So, we may enforce $\mathbf{u}_2(b, y) = 0$ as $\mathbf{u}_2(b, d) = \partial_y \mathbf{u}_2(b, d) = 0$ and $\partial_y^2 \mathbf{u}_2(b, y) = 0$;

For any PDE State, there Exists a Constraint-Free *Fundamental State*

We decompose the PDE state \mathbf{u} using the fundamental theorem of calculus:

$$\begin{aligned}\mathbf{u}_1(x, y) &= \mathbf{u}_1(a, c) + \int_a^x \partial_x \mathbf{u}_1(\theta, c) d\theta + \int_c^y \partial_y \mathbf{u}_1(a, \nu) d\nu \\ &\quad + \int_c^y \int_a^x \partial_x \partial_y \mathbf{u}_1(\theta, \nu) d\theta d\nu\end{aligned}$$

For any PDE State, there Exists a Constraint-Free *Fundamental State*

We decompose the PDE state \mathbf{u} using the fundamental theorem of calculus:

$$\begin{aligned}\mathbf{u}_1(x, y) &= \mathbf{u}_1(a, c) + \int_a^x \partial_x \mathbf{u}_1(\theta, c) d\theta + \int_c^y \partial_y \mathbf{u}_1(a, \nu) d\nu \\ &\quad + \int_c^y \int_a^x \partial_x \partial_y \mathbf{u}_1(\theta, \nu) d\theta d\nu\end{aligned}$$

For any PDE State, there Exists a Constraint-Free *Fundamental State*

We decompose the PDE state \mathbf{u} using the fundamental theorem of calculus:

$$\begin{aligned}\mathbf{u}_1(x, y) &= \mathbf{u}_1(a, c) + \int_a^x \partial_x \mathbf{u}_1(\theta, c) d\theta + \int_c^y \partial_y \mathbf{u}_1(a, \nu) d\nu \\ &\quad + \int_c^y \int_a^x \partial_x \partial_y \mathbf{u}_1(\theta, \nu) d\theta d\nu \\ \mathbf{u}_2(x, y) &= \mathbf{u}_2(a, c) + (x - a) \partial_x \mathbf{u}_2(a, c) \\ &\quad + (y - c) \partial_y \mathbf{u}_2(a, c) + (y - c)(x - a) \partial_x \partial_y \mathbf{u}_2(a, c) \\ &\quad + \int_a^x (x - \theta) \partial_x^2 \mathbf{u}_2(\theta, c) d\theta + \int_c^y (y - \nu) \partial_y^2 \mathbf{u}_2(a, \nu) d\nu \\ &\quad + (y - c) \int_a^x (x - \theta) \partial_x^2 \partial_y \mathbf{u}_2(\theta, c) d\theta + (x - a) \int_c^y (y - \nu) \partial_x \partial_y^2 \mathbf{u}_2(a, \nu) d\nu \\ &\quad + \int_c^y \int_a^x (y - \nu)(x - \theta) \partial_x^2 \partial_y^2 \mathbf{u}_2(\theta, \nu) d\theta d\nu\end{aligned}$$

For any PDE State, there Exists a Constraint-Free *Fundamental State*

We decompose the PDE state \mathbf{u} using the fundamental theorem of calculus:

$$\begin{aligned}\mathbf{u}_1(x, y) &= \mathbf{u}_1(a, c) + \int_a^x \partial_x \mathbf{u}_1(\theta, c) d\theta + \int_c^y \partial_y \mathbf{u}_1(a, \nu) d\nu \\ &\quad + \int_c^y \int_a^x \partial_x \partial_y \mathbf{u}_1(\theta, \nu) d\theta d\nu \\ \mathbf{u}_2(x, y) &= \mathbf{u}_2(a, c) + (x - a) \partial_x \mathbf{u}_2(a, c) \\ &\quad + (y - c) \partial_y \mathbf{u}_2(a, c) + (y - c)(x - a) \partial_x \partial_y \mathbf{u}_2(a, c) \\ &\quad + \int_a^x (x - \theta) \partial_x^2 \mathbf{u}_2(\theta, c) d\theta + \int_c^y (y - \nu) \partial_y^2 \mathbf{u}_2(a, \nu) d\nu \\ &\quad + (y - c) \int_a^x (x - \theta) \partial_x^2 \partial_y \mathbf{u}_2(\theta, c) d\theta + (x - a) \int_c^y (y - \nu) \partial_x \partial_y^2 \mathbf{u}_2(a, \nu) d\nu \\ &\quad + \int_c^y \int_a^x (y - \nu)(x - \theta) \partial_x^2 \partial_y^2 \mathbf{u}_2(\theta, \nu) d\theta d\nu\end{aligned}$$

Definition 5 (Fundamental State)

For an arbitrary PDE state $\mathbf{u} \in X_{\mathcal{B}}[\Omega]$, we define the corresponding fundamental state (PIE state) $\hat{\mathbf{u}} \in L_2^{n_0+n_1+n_2}[\Omega]$ as

$$\hat{\mathbf{u}} = \begin{bmatrix} \hat{\mathbf{u}}_0 \\ \hat{\mathbf{u}}_1 \\ \hat{\mathbf{u}}_2 \end{bmatrix} := \begin{bmatrix} \mathbf{u}_0 \\ \partial_x \partial_y \mathbf{u}_1 \\ \partial_x^2 \partial_y^2 \hat{\mathbf{u}}_2 \end{bmatrix} = \underbrace{\begin{bmatrix} I & & \\ & \partial_x \partial_y & \\ & & \partial_x^2 \partial_y^2 \end{bmatrix}}_{\mathcal{D}} \begin{bmatrix} \mathbf{u}_0 \\ \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix} = \mathcal{D}\mathbf{u}$$

For any PDE State, there Exists a Constraint-Free *Fundamental State*

We decompose the PDE state \mathbf{u} using the fundamental theorem of calculus:

$$\begin{aligned}
 \mathbf{u}_1(x, y) &= \mathbf{u}_1(a, c) + \int_a^x \partial_x \mathbf{u}_1(\theta, c) d\theta + \int_c^y \partial_y \mathbf{u}_1(a, \nu) d\nu \\
 &\quad + \int_c^y \int_a^x \partial_x \partial_y \mathbf{u}_1(\theta, \nu) d\theta d\nu \\
 \mathbf{u}_2(x, y) &= \mathbf{u}_2(a, c) + (x - a) \partial_x \mathbf{u}_2(a, c) \\
 &\quad + (y - c) \partial_y \mathbf{u}_2(a, c) + (y - c)(x - a) \partial_x \partial_y \mathbf{u}_2(a, c) \\
 &\quad + \int_a^x (x - \theta) \partial_x^2 \mathbf{u}_2(\theta, c) d\theta + \int_c^y (y - \nu) \partial_y^2 \mathbf{u}_2(a, \nu) d\nu \\
 &\quad + (y - c) \int_a^x (x - \theta) \partial_x^2 \partial_y \mathbf{u}_2(\theta, c) d\theta + (x - a) \int_c^y (y - \nu) \partial_x \partial_y^2 \mathbf{u}_2(a, \nu) d\nu \\
 &\quad + \int_c^y \int_a^x (y - \nu)(x - \theta) \partial_x^2 \partial_y^2 \mathbf{u}_2(\theta, \nu) d\theta d\nu
 \end{aligned}
 \quad \Lambda_{bc} \mathbf{u} = \begin{bmatrix} \mathbf{u}_1(a, c) \\ \mathbf{u}_2(a, c) \\ \partial_x \mathbf{u}_2(a, c) \\ \partial_y \mathbf{u}_2(a, c) \\ \partial_x \partial_y \mathbf{u}_2(a, c) \\ \hline \partial_x \mathbf{u}_1(x, c) \\ \partial_x^2 \mathbf{u}_2(x, c) \\ \partial_x^2 \partial_y \mathbf{u}_2(x, c) \\ \hline \partial_y \mathbf{u}_1(a, y) \\ \partial_y^2 \mathbf{u}_2(a, y) \\ \partial_x \partial_y^2 \mathbf{u}_2(a, y) \end{bmatrix}$$

Definition 5 (Fundamental State)

For an arbitrary PDE state $\mathbf{u} \in X_{\mathcal{B}}[\Omega]$, we define the corresponding fundamental state (PIE state) $\hat{\mathbf{u}} \in L_2^{n_0+n_1+n_2}[\Omega]$ as

$$\hat{\mathbf{u}} = \begin{bmatrix} \hat{\mathbf{u}}_0 \\ \hat{\mathbf{u}}_1 \\ \hat{\mathbf{u}}_2 \end{bmatrix} := \begin{bmatrix} \mathbf{u}_0 \\ \partial_x \partial_y \mathbf{u}_1 \\ \partial_x^2 \partial_y^2 \hat{\mathbf{u}}_2 \end{bmatrix} = \underbrace{\begin{bmatrix} I & & \\ & \partial_x \partial_y & \\ & & \partial_x^2 \partial_y^2 \end{bmatrix}}_{\mathcal{D}} \begin{bmatrix} \mathbf{u}_0 \\ \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix} = \mathcal{D} \mathbf{u}$$

The PDE State can be Expressed in terms of the Fundamental State through PI Operators

$$\begin{aligned}
 \mathbf{u} &= \begin{bmatrix} \mathbf{u}_0 \\ \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix} \\
 &= \overbrace{\begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ I & 0 & 0 & 0 & 0 & f_a^x[I] & 0 & 0 & f_c^y[I] & 0 & 0 \\ 0 & I & [x-a] & [y-c] & [y-c][x-a] & 0 & f_a^x[x-\theta] & [y-c] & f_a^x[x-\theta] & 0 & f_c^y[y-\nu] & [x-a] & f_c^y[y-\nu] \end{bmatrix}}^{\mathcal{K}_1} \overbrace{\begin{bmatrix} \mathbf{u}_1(a, c) \\ \mathbf{u}_2(a, c) \\ \partial_x \mathbf{u}_2(a, c) \\ \partial_y \mathbf{u}_2(a, c) \\ \partial_x \partial_y \mathbf{u}_2(a, c) \\ \partial_x \mathbf{u}_1(\theta, c) \\ \partial_x^2 \mathbf{u}_2(\theta, c) \\ \partial_x^2 \partial_y \mathbf{u}_2(\theta, c) \\ \partial_y \mathbf{u}_1(a, \nu) \\ \partial_y^2 \mathbf{u}_2(a, \nu) \\ \partial_x \partial_y^2 \mathbf{u}_2(a, \nu) \end{bmatrix}}^{\Lambda_{bc} \mathbf{u}} \\
 &\quad + \overbrace{\begin{bmatrix} I & f_a^x[I] & f_c^y[I] \\ f_a^x[x-\theta] & f_c^y[y-\nu] \end{bmatrix}}^{\mathcal{K}_2} \overbrace{\begin{bmatrix} \mathbf{u}_0 \\ \partial_x \partial_y \mathbf{u}_1 \\ \partial_x^2 \partial_y^2 \mathbf{u}_2 \end{bmatrix}}^{\hat{\mathbf{u}}} = \mathcal{K}_1 \Lambda_{bc} \mathbf{u} + \mathcal{K}_2 \hat{\mathbf{u}}
 \end{aligned}$$

The PDE State can be Expressed in terms of the Fundamental State through PI Operators

$$\begin{aligned}
 \mathbf{u} &= \begin{bmatrix} \mathbf{u}_0 \\ \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix} \\
 &= \overbrace{\begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ I & 0 & 0 & 0 & 0 & f_a^x[I] & 0 & 0 & f_c^y[I] & 0 & 0 \\ 0 & I & [x-a] & [y-c] & [y-c][x-a] & 0 & f_a^x[x-\theta] & [y-c] & f_a^x[x-\theta] & 0 & f_c^y[y-\nu] & [x-a] & f_c^y[y-\nu] \end{bmatrix}}^{\mathcal{K}_1} \Lambda_{bc} \mathbf{u} \\
 &\quad + \overbrace{\begin{bmatrix} I & f_a^x[I] & f_c^y[I] \\ f_a^x[x-\theta] & f_c^y[y-\nu] \end{bmatrix}}^{\mathcal{K}_2} \overbrace{\begin{bmatrix} \mathbf{u}_0 \\ \partial_x \partial_y \mathbf{u}_1 \\ \partial_x^2 \partial_y^2 \mathbf{u}_2 \end{bmatrix}}^{\hat{\mathbf{u}}} = \mathcal{K}_1 \Lambda_{bc} \mathbf{u} + \mathcal{K}_2 \hat{\mathbf{u}}
 \end{aligned}$$

$\Lambda_{bc} \mathbf{u}$

- $\mathbf{u}_1(a, c)$
- $\mathbf{u}_2(a, c)$
- $\partial_x \mathbf{u}_2(a, c)$
- $\partial_y \mathbf{u}_2(a, c)$
- $\partial_x \partial_y \mathbf{u}_2(a, c)$
- $\partial_x \mathbf{u}_1(\theta, c)$
- $\partial_x^2 \mathbf{u}_2(\theta, c)$
- $\partial_x^2 \mathbf{u}_2(\theta, c)$
- $\partial_y \mathbf{u}_1(a, \nu)$
- $\partial_y^2 \mathbf{u}_2(a, \nu)$
- $\partial_x \partial_y^2 \mathbf{u}_2(a, \nu)$

Lemma 6

Let $\mathbf{u} \in \begin{bmatrix} L_2[\Omega] \\ H_1[\Omega] \\ H_2[\Omega] \end{bmatrix}$, and define $\hat{\mathbf{u}} = \mathcal{D}\mathbf{u}$. Then, there exist PI operators \mathcal{K}_1 and \mathcal{K}_2 such that

$$\mathbf{u} = \mathcal{K}_1 \Lambda_{bc} \mathbf{u} + \mathcal{K}_2 \hat{\mathbf{u}}.$$

The PDE State can be Expressed in terms of the Fundamental State through PI Operators

$$\begin{aligned}
 \mathbf{u} &= \begin{bmatrix} \mathbf{u}_0 \\ \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix} \\
 &= \overbrace{\begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ I & 0 & 0 & 0 & 0 & f_a^x[I] & 0 & 0 & f_c^y[I] & 0 & 0 \\ 0 & I & [x-a] & [y-c] & [y-c][x-a] & 0 & f_a^x[x-\theta] & [y-c] & f_a^x[x-\theta] & 0 & f_c^y[y-\nu] & [x-a] & f_c^y[y-\nu] \end{bmatrix}}^{\mathcal{K}_1} \Lambda_{bc} \mathbf{u} \\
 &\quad + \overbrace{\begin{bmatrix} I & f_a^x[I] & f_c^y[I] \\ f_a^x[x-\theta] & f_c^y[y-\nu] \end{bmatrix}}^{\mathcal{K}_2} \underbrace{\begin{bmatrix} \mathbf{u}_0 \\ \partial_x \partial_y \mathbf{u}_1 \\ \partial_x^2 \partial_y^2 \mathbf{u}_2 \end{bmatrix}}_{\hat{\mathbf{u}}} = \mathcal{K}_1 \Lambda_{bc} \mathbf{u} + \mathcal{K}_2 \hat{\mathbf{u}}
 \end{aligned}$$

$\Lambda_{bc} \mathbf{u}$
 $\mathbf{u}_1(a, c)$
 $\mathbf{u}_2(a, c)$
 $\partial_x \mathbf{u}_2(a, c)$
 $\partial_y \mathbf{u}_2(a, c)$
 $\partial_x \partial_y \mathbf{u}_2(a, c)$
 $\partial_x \mathbf{u}_1(\theta, c)$
 $\partial_x^2 \mathbf{u}_2(\theta, c)$
 $\partial_x^2 \partial_y \mathbf{u}_2(\theta, c)$
 $\partial_y \mathbf{u}_1(a, \nu)$
 $\partial_y^2 \mathbf{u}_2(a, \nu)$
 $\partial_x \partial_y^2 \mathbf{u}_2(a, \nu)$

Lemma 6

Let $\mathbf{u} \in \begin{bmatrix} L_2[\Omega] \\ H_1[\Omega] \\ H_2[\Omega] \end{bmatrix}$, and define $\hat{\mathbf{u}} = \mathcal{D}\mathbf{u}$. Then, there exist PI operators \mathcal{K}_1 and \mathcal{K}_2 such that

$$\mathbf{u} = \mathcal{K}_1 \Lambda_{bc} \mathbf{u} + \mathcal{K}_2 \hat{\mathbf{u}}.$$

Corollary 7

Let $\mathbf{u} \in \begin{bmatrix} L_2[\Omega] \\ H_1[\Omega] \\ H_2[\Omega] \end{bmatrix}$, and define $\hat{\mathbf{u}} = \mathcal{D}\mathbf{u}$. Then there exist PI operators \mathcal{H}_1 and \mathcal{H}_2 such that

$$\Lambda_{bf} \mathbf{u} = \mathcal{H}_1 \Lambda_{bc} \mathbf{u} + \mathcal{H}_2 \hat{\mathbf{u}}.$$

There Exists a Direct Map from Fundamental State to PDE State

For appropriate PI operators $\mathcal{H}_1, \mathcal{H}_2$, we can represent

$$\Lambda_{\text{bf}}\mathbf{u} = \mathcal{H}_1\Lambda_{\text{bc}}\mathbf{u} + \mathcal{H}_2\hat{\mathbf{u}},$$

where $\hat{\mathbf{u}} = \mathcal{D}\mathbf{u}$.

Given BCs $\mathcal{B}\Lambda_{\text{bf}}\mathbf{u} = 0$, it follows that

$$0 = \mathcal{B}\Lambda_{\text{bf}}\mathbf{u} = \mathcal{B}\mathcal{H}_1\Lambda_{\text{bc}}\mathbf{u} + \mathcal{B}\mathcal{H}_2\hat{\mathbf{u}},$$

and therefore

$$\Lambda_{\text{bc}}\mathbf{u} = -(\mathcal{B}\mathcal{H}_1)^{-1}\mathcal{B}\mathcal{H}_2\hat{\mathbf{u}}.$$

There Exists a Direct Map from Fundamental State to PDE State

For appropriate PI operators $\mathcal{H}_1, \mathcal{H}_2$, we can represent

$$\Lambda_{\text{bf}}\mathbf{u} = \mathcal{H}_1\Lambda_{\text{bc}}\mathbf{u} + \mathcal{H}_2\hat{\mathbf{u}},$$

where $\hat{\mathbf{u}} = \mathcal{D}\mathbf{u}$.

Given BCs $\mathcal{B}\Lambda_{\text{bf}}\mathbf{u} = 0$, it follows that

$$0 = \mathcal{B}\Lambda_{\text{bf}}\mathbf{u} = \mathcal{B}\mathcal{H}_1\Lambda_{\text{bc}}\mathbf{u} + \mathcal{B}\mathcal{H}_2\hat{\mathbf{u}},$$

and therefore

$$\Lambda_{\text{bc}}\mathbf{u} = -(\mathcal{B}\mathcal{H}_1)^{-1}\mathcal{B}\mathcal{H}_2\hat{\mathbf{u}}.$$

For appropriate PI operators $\mathcal{K}_1, \mathcal{K}_2$, we can also represent

$$\mathbf{u} = \mathcal{K}_1\Lambda_{\text{bc}}\mathbf{u} + \mathcal{K}_2\hat{\mathbf{u}},$$

and thus

$$\begin{aligned}\mathbf{u} &= \mathcal{K}_1\Lambda_{\text{bc}}\mathbf{u} + \mathcal{K}_2\hat{\mathbf{u}} \\ &= -\mathcal{K}_1(\mathcal{B}\mathcal{H}_1)^{-1}\mathcal{B}\mathcal{H}_2\hat{\mathbf{u}} + \mathcal{K}_2\hat{\mathbf{u}} \\ &= \underbrace{\left[\mathcal{K}_2 - \mathcal{K}_1(\mathcal{B}\mathcal{H}_1)^{-1}\mathcal{B}\mathcal{H}_2\right]}_{\mathcal{T}}\hat{\mathbf{u}} = \mathcal{T}\hat{\mathbf{u}}\end{aligned}$$

There Exists a Direct Map from Fundamental State to PDE State

For appropriate PI operators $\mathcal{H}_1, \mathcal{H}_2$, we can represent

$$\Lambda_{\text{bf}} \mathbf{u} = \mathcal{H}_1 \Lambda_{\text{bc}} \mathbf{u} + \mathcal{H}_2 \hat{\mathbf{u}},$$

where $\hat{\mathbf{u}} = \mathcal{D} \mathbf{u}$.

Given BCs $\mathcal{B} \Lambda_{\text{bf}} \mathbf{u} = 0$, it follows that

$$0 = \mathcal{B} \Lambda_{\text{bf}} \mathbf{u} = \mathcal{B} \mathcal{H}_1 \Lambda_{\text{bc}} \mathbf{u} + \mathcal{B} \mathcal{H}_2 \hat{\mathbf{u}},$$

and therefore

$$\Lambda_{\text{bc}} \mathbf{u} = -(\mathcal{B} \mathcal{H}_1)^{-1} \mathcal{B} \mathcal{H}_2 \hat{\mathbf{u}}.$$

For appropriate PI operators $\mathcal{K}_1, \mathcal{K}_2$, we can also represent

$$\mathbf{u} = \mathcal{K}_1 \Lambda_{\text{bc}} \mathbf{u} + \mathcal{K}_2 \hat{\mathbf{u}},$$

and thus

$$\begin{aligned} \mathbf{u} &= \mathcal{K}_1 \Lambda_{\text{bc}} \mathbf{u} + \mathcal{K}_2 \hat{\mathbf{u}} \\ &= -\mathcal{K}_1 (\mathcal{B} \mathcal{H}_1)^{-1} \mathcal{B} \mathcal{H}_2 \hat{\mathbf{u}} + \mathcal{K}_2 \hat{\mathbf{u}} \\ &= \underbrace{\left[\mathcal{K}_2 - \mathcal{K}_1 (\mathcal{B} \mathcal{H}_1)^{-1} \mathcal{B} \mathcal{H}_2 \right]}_{\mathcal{T}} \hat{\mathbf{u}} = \mathcal{T} \hat{\mathbf{u}} \end{aligned}$$

Theorem 8

Let \mathcal{B} be a given PI operator, and such that the operator $\mathcal{B} \mathcal{H}_1$ is invertible. Then, there exists a 2D-PI operator \mathcal{T} such that for any $\mathbf{u} \in X_{\mathcal{B}}[\Omega]$ and $\hat{\mathbf{u}} \in L_2[\Omega]$, we have

$$\mathbf{u} = \mathcal{T} \mathcal{D} \mathbf{u} \quad \text{and} \quad \hat{\mathbf{u}} = \mathcal{D} \mathcal{T} \hat{\mathbf{u}},$$

$$\text{where } \mathcal{D} := \begin{bmatrix} I & & \\ & \partial_x \partial_y & \\ & & \partial_x^2 \partial_y^2 \end{bmatrix}.$$

Any Well-Posed, Linear, 2nd Order, 2D PDE can be Equivalently Represented as a PIE

Using the relation $\mathbf{u} = \mathcal{T}\hat{\mathbf{u}}$, the PDE defined by $\{A_{ij}, \mathcal{B}\}$,

$$\dot{\mathbf{u}}(t) = \mathcal{A}\mathbf{u}(t) = A_{00}\mathbf{u}(t) + \dots + A_{22} \partial_x^2 \partial_y^2 \mathbf{u}_2(t),$$

$$\mathbf{u}(t) \in X_{\mathcal{B}}[\Omega] := \left\{ \begin{bmatrix} \mathbf{u}_0 \\ \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix} \in \begin{bmatrix} L_2^{n_0}[\Omega] \\ H_1^{n_1}[\Omega] \\ H_2^{n_2}[\Omega] \end{bmatrix} \mid \mathcal{B}\Lambda_{bf}\mathbf{u} = 0 \right\},$$

can be equivalently represented as a Partial Integral Equation (PIE)

$$\mathcal{T}\dot{\hat{\mathbf{u}}}(t) = \mathcal{A}\hat{\mathbf{u}}(t),$$

$$\hat{\mathbf{u}}(t) \in L_2[\Omega]$$

defined by $\{\mathcal{T}, \mathcal{A}\}$:

Lemma 9

For given $\{A_{ij}, \mathcal{B}\}$, let

$$\mathcal{T} := [\mathcal{K}_2 - \mathcal{K}_1(\mathcal{B}\mathcal{H}_1)^{-1}\mathcal{B}\mathcal{H}_2], \quad \text{and} \quad \mathcal{A} := \mathcal{A}\mathcal{T},$$

where \mathcal{A} is defined by $\{A_{ij}\}$, and where \mathcal{H}_1 , \mathcal{H}_2 , \mathcal{K}_1 and \mathcal{K}_2 are such that

$$\begin{aligned} \Lambda_{bf}\mathbf{u} &= \mathcal{H}_1\Lambda_{bc}\mathbf{u} + \mathcal{H}_2\hat{\mathbf{u}}, \\ \mathbf{u} &= \mathcal{K}_1\Lambda_{bc}\mathbf{u} + \mathcal{K}_2\hat{\mathbf{u}}. \end{aligned}$$

Then, for any $\hat{\mathbf{u}}_I \in L_2^n[\Omega]$, $\hat{\mathbf{u}}(t)$ solves the PIE defined by $\{\mathcal{T}, \mathcal{A}\}$ with the initial condition $\hat{\mathbf{u}}_I$ if and only if $\mathbf{u}(t) = \mathcal{T}\hat{\mathbf{u}}(t)$ solves the PDE defined by $\{A_{ij}, \mathcal{B}\}$ with the initial condition $\mathbf{u}_I = \mathcal{T}\hat{\mathbf{u}}_I$.

Stability of a PIE can be Tested as an LMI

For a PIE defined by $\{\mathcal{T}, \mathcal{A}\}$,

$$\begin{aligned}\mathcal{T}\dot{\mathbf{u}}(t) &= \mathcal{A}\mathbf{u}(t), \\ \mathbf{u}(t) &\in L_2^n[\Omega],\end{aligned}$$

a LF can be parameterized by a 2D-PI operator \mathcal{P} as

$$V(\hat{\mathbf{u}}(t)) = \langle \mathcal{T}\hat{\mathbf{u}}(t), \mathcal{P}\mathcal{T}\hat{\mathbf{u}}(t) \rangle_{L_2},$$

for which

$$\begin{aligned}\dot{V}(\mathbf{u}(t)) &= \langle \mathcal{T}\dot{\mathbf{u}}, \mathcal{P}\mathcal{T}\mathbf{u} \rangle_{L_2} + \langle \mathcal{T}\mathbf{u}, \mathcal{P}\mathcal{T}\dot{\mathbf{u}} \rangle_{L_2} \\ &= \langle \mathbf{u}, [\mathcal{A}^*\mathcal{P}\mathcal{T} + \mathcal{T}^*\mathcal{P}\mathcal{A}]\mathbf{u} \rangle_{L_2}.\end{aligned}$$

Then, stability of the PIE can be verified by solving the Linear PI Inequality (LPI)

$$\begin{aligned}\mathcal{P} &> 0, \\ \mathcal{A}^*\mathcal{P}\mathcal{T} + \mathcal{T}^*\mathcal{P}\mathcal{A} &\leq 0.\end{aligned}$$

Stability of a PIE can be Tested as an LMI

For a PIE defined by $\{\mathcal{T}, \mathcal{A}\}$,

$$\begin{aligned}\mathcal{T}\dot{\mathbf{u}}(t) &= \mathcal{A}\mathbf{u}(t), \\ \mathbf{u}(t) &\in L_2^n[\Omega],\end{aligned}$$

a LF can be parameterized by a 2D-PI operator \mathcal{P} as

$$V(\hat{\mathbf{u}}(t)) = \langle \mathcal{T}\hat{\mathbf{u}}(t), \mathcal{P}\mathcal{T}\hat{\mathbf{u}}(t) \rangle_{L_2},$$

for which

$$\begin{aligned}\dot{V}(\mathbf{u}(t)) &= \langle \mathcal{T}\dot{\mathbf{u}}, \mathcal{P}\mathcal{T}\mathbf{u} \rangle_{L_2} + \langle \mathcal{T}\mathbf{u}, \mathcal{P}\mathcal{T}\dot{\mathbf{u}} \rangle_{L_2} \\ &= \langle \mathbf{u}, [\mathcal{A}^*\mathcal{P}\mathcal{T} + \mathcal{T}^*\mathcal{P}\mathcal{A}]\mathbf{u} \rangle_{L_2}.\end{aligned}$$

Then, stability of the PIE can be verified by solving the Linear PI Inequality (LPI)

$$\begin{aligned}\mathcal{P} &> 0, \\ \mathcal{A}^*\mathcal{P}\mathcal{T} + \mathcal{T}^*\mathcal{P}\mathcal{A} &\leq 0.\end{aligned}$$

To enforce positivity of PI operators, let

$$\mathcal{P} := \mathcal{Z}^* P \mathcal{Z},$$

for some fixed 2D-PI operator \mathcal{Z} . Then if $P > 0$,

$$\begin{aligned}\langle \mathbf{u}, \mathcal{P}\mathbf{u} \rangle_{L_2} &= \langle \mathcal{Z}\mathbf{u}, P\mathcal{Z}\mathbf{u} \rangle_{L_2} \\ &= \left\langle P^{\frac{1}{2}}\mathcal{Z}\mathbf{u}, P^{\frac{1}{2}}\mathcal{Z}\mathbf{u} \right\rangle_{L_2} > 0,\end{aligned}$$

for any $\mathbf{u} \in L_2^n[\Omega]$.

Stability of a PIE can be Tested as an LMI

For a PIE defined by $\{\mathcal{T}, \mathcal{A}\}$,

$$\begin{aligned}\mathcal{T}\dot{\mathbf{u}}(t) &= \mathcal{A}\mathbf{u}(t), \\ \mathbf{u}(t) &\in L_2^n[\Omega],\end{aligned}$$

a LF can be parameterized by a 2D-PI operator \mathcal{P} as

$$V(\hat{\mathbf{u}}(t)) = \langle \mathcal{T}\hat{\mathbf{u}}(t), \mathcal{P}\mathcal{T}\hat{\mathbf{u}}(t) \rangle_{L_2},$$

for which

$$\begin{aligned}\dot{V}(\mathbf{u}(t)) &= \langle \mathcal{T}\dot{\mathbf{u}}, \mathcal{P}\mathcal{T}\mathbf{u} \rangle_{L_2} + \langle \mathcal{T}\mathbf{u}, \mathcal{P}\mathcal{T}\dot{\mathbf{u}} \rangle_{L_2} \\ &= \langle \mathbf{u}, [\mathcal{A}^*\mathcal{P}\mathcal{T} + \mathcal{T}^*\mathcal{P}\mathcal{A}]\mathbf{u} \rangle_{L_2}.\end{aligned}$$

Then, stability of the PIE can be verified by solving the Linear PI Inequality (LPI)

$$\begin{aligned}\mathcal{P} &> 0, \\ \mathcal{A}^*\mathcal{P}\mathcal{T} + \mathcal{T}^*\mathcal{P}\mathcal{A} &\leq 0.\end{aligned}$$

To enforce positivity of PI operators, let

$$\mathcal{P} := \mathcal{Z}^* P \mathcal{Z},$$

for some fixed 2D-PI operator \mathcal{Z} . Then if $P > 0$,

$$\begin{aligned}\langle \mathbf{u}, \mathcal{P}\mathbf{u} \rangle_{L_2} &= \langle \mathcal{Z}\mathbf{u}, P\mathcal{Z}\mathbf{u} \rangle_{L_2} \\ &= \left\langle P^{\frac{1}{2}}\mathcal{Z}\mathbf{u}, P^{\frac{1}{2}}\mathcal{Z}\mathbf{u} \right\rangle_{L_2} > 0,\end{aligned}$$

for any $\mathbf{u} \in L_2^n[\Omega]$.

As such, if there exist matrices $P > 0$ and $Q \leq 0$ such that, for given PI operators \mathcal{Z}_1 and \mathcal{Z}_2 ,

$$\begin{aligned}\mathcal{P} &= \mathcal{Z}_1^* P \mathcal{Z}_1, \\ \mathcal{A}^*\mathcal{P}\mathcal{T} + \mathcal{T}^*\mathcal{P}\mathcal{A} &= \mathcal{Z}_2^* Q \mathcal{Z}_2,\end{aligned}$$

then the PIE defined by $\{\mathcal{T}, \mathcal{A}\}$ is stable.

Stability of 2D PDEs can be Numerically Tested with PIETOOLS

Combining all steps, we can perform stability analysis of 2D PDEs using PIETOOLS (<https://control.asu.edu/pietools/>):

- 1 Represent the PDE in the standardized format by $\{A_{ij}, \mathcal{B}\}$:

```
PDE.n.n_pde = [n0,n1,n2];    PDE.dom = [a,b;c,d];    PDE.vars = [x,tt;y,nu];  
PDE.PDE.A = ...;            PDE.BC.Ebb = ...;
```

Stability of 2D PDEs can be Numerically Tested with PIETOOLS

Combining all steps, we can perform stability analysis of 2D PDEs using PIETOOLS (<https://control.asu.edu/pietools/>):

- 1 Represent the PDE in the standardized format by $\{A_{ij}, \mathcal{B}\}$:

```
PDE.n.n_pde = [n0,n1,n2];    PDE.dom = [a,b;c,d];    PDE.vars = [x,tt;y,nu];  
PDE.PDE.A = ...;            PDE.BC.Ebb = ...;
```

- 2 Convert the PDE to a PIE defined by $\{\mathcal{T}, \mathcal{A}\}$:

```
PIE = convert_PIETOOLS_PDE(PDE);  
T = PIE.T;          A = PIE.A;
```

Stability of 2D PDEs can be Numerically Tested with PIETOOLS

Combining all steps, we can perform stability analysis of 2D PDEs using PIETOOLS (<https://control.asu.edu/pietools/>):

- 1 Represent the PDE in the standardized format by $\{A_{ij}, \mathcal{B}\}$:

```
PDE.n.n_pde = [n0,n1,n2];    PDE.dom = [a,b;c,d];    PDE.vars = [x,tt;y,nu];  
PDE.PDE.A = ...;            PDE.BC.Ebb = ...;
```

- 2 Convert the PDE to a PIE defined by $\{\mathcal{T}, \mathcal{A}\}$:

```
PIE = convert_PIETOOLS_PDE(PDE);  
T = PIE.T;            A = PIE.A;
```

- 3 Test for existence of a PI operator $\mathcal{P} > 0$ such that $\mathcal{A}^* \mathcal{P} \mathcal{T} + \mathcal{T}^* \mathcal{P} \mathcal{A} \leq 0$:

```
prog = sosprogram([x y tt nu]);  
[prog, P] = poslpivar_2d(prog,n,dom,deg);  
P = P + eps;    Q = - A'*P*T - T'*P*A;  
prog = lpi_ineq_2d(prog,Q);  
prog = sossolve(prog);
```


Ex. 1: Advection Equation

For the simple advection equation

$$\dot{u}(t, x, y) = C\partial_x u(t, x, y) + C\partial_y u(t, x, y)$$

$$u(t, 0, y) = u(t, x, 0) = 0, \quad (x, y) \in [0, 1]^2$$

we have $\mathbf{u}(t) = \mathbf{u}_1(t) = u(t) \in H_1[[0, 1]^2]$.

The associated fundamental state is

$\hat{\mathbf{u}} = \partial_x \partial_y \mathbf{u}$, with corresponding PIE

$$\begin{aligned} & \int_0^x \int_0^y \dot{\hat{\mathbf{u}}}(t, \theta, \nu) d\nu d\theta \\ &= C \int_0^y \hat{\mathbf{u}}(t, x, \nu) d\nu + C \int_0^x \hat{\mathbf{u}}(t, \theta, y) d\theta. \end{aligned}$$

Ex. 1: Advection Equation

For the simple advection equation

$$\begin{aligned}\dot{u}(t, x, y) &= C \partial_x u(t, x, y) + C \partial_y u(t, x, y) \\ u(t, 0, y) &= u(t, x, 0) = 0, \quad (x, y) \in [0, 1]^2\end{aligned}$$

we have $\mathbf{u}(t) = \mathbf{u}_1(t) = u(t) \in H_1[[0, 1]^2]$.

The associated fundamental state is

$\hat{\mathbf{u}} = \partial_x \partial_y \mathbf{u}$, with corresponding PIE

$$\begin{aligned}\int_0^x \int_0^y \hat{\mathbf{u}}(t, \theta, \nu) d\nu d\theta \\ = C \int_0^y \hat{\mathbf{u}}(t, x, \nu) d\nu + C \int_0^x \hat{\mathbf{u}}(t, \theta, y) d\theta.\end{aligned}$$

Ex. 2: Wave Equation

The wave equation on $(x, y) \in [0, 1]^2$

$$\begin{aligned}\ddot{u}(x, y) &= \partial_x^2 u(x, y) + \partial_y^2 u(x, y) \\ u(0, y) &= \partial_x u(0, y) = u(x, 0) = \partial_y u(x, 0) = 0\end{aligned}$$

can be represented in the standardized format by defining $\mathbf{u}_1 = u$ and $\mathbf{u}_2 = \dot{u}$, as

$$\begin{aligned}\begin{bmatrix} \dot{\mathbf{u}}_1 \\ \dot{\mathbf{u}}_2 \end{bmatrix} &= \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \partial_x^2 \begin{bmatrix} \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \partial_y^2 \begin{bmatrix} \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix} \\ &\quad + \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix}.\end{aligned}$$

The associated fundamental state is $\hat{\mathbf{u}} = \partial_x^2 \partial_y^2 \mathbf{u}$, with corresponding PIE

$$\begin{aligned}\int_0^x \int_0^y \begin{bmatrix} (x-\theta)(y-\nu) & 0 \\ 0 & (x-\theta)(y-\nu) \end{bmatrix} \begin{bmatrix} \dot{\mathbf{u}}_1 \\ \dot{\mathbf{u}}_2 \end{bmatrix} d\nu d\theta \\ = \int_0^y \begin{bmatrix} 0 & 0 \\ (y-\nu) & 0 \end{bmatrix} \begin{bmatrix} \hat{\mathbf{u}}_1 \\ \hat{\mathbf{u}}_2 \end{bmatrix} d\nu + \int_0^x \begin{bmatrix} 0 & 0 \\ (x-\theta) & 0 \end{bmatrix} \begin{bmatrix} \hat{\mathbf{u}}_1 \\ \hat{\mathbf{u}}_2 \end{bmatrix} d\theta \\ + \int_0^x \int_0^y \begin{bmatrix} 0 & (x-\theta)(y-\nu) \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \hat{\mathbf{u}}_1 \\ \hat{\mathbf{u}}_2 \end{bmatrix} d\nu d\theta\end{aligned}$$

For BCs $u \equiv 0$ on $\partial\Omega$, stability can be verified with PIETOOLS.

Examples

Ex. 1: Advection Equation

For the simple advection equation

$$\begin{aligned}\dot{u}(t, x, y) &= C \partial_x u(t, x, y) + C \partial_y u(t, x, y) \\ u(t, 0, y) &= u(t, x, 0) = 0, \quad (x, y) \in [0, 1]^2\end{aligned}$$

we have $\mathbf{u}(t) = \mathbf{u}_1(t) = u(t) \in H_1[[0, 1]^2]$.

The associated fundamental state is

$\hat{\mathbf{u}} = \partial_x \partial_y \mathbf{u}$, with corresponding PIE

$$\begin{aligned}\int_0^x \int_0^y \hat{\mathbf{u}}(t, \theta, \nu) d\nu d\theta \\ = C \int_0^y \hat{\mathbf{u}}(t, x, \nu) d\nu + C \int_0^x \hat{\mathbf{u}}(t, \theta, y) d\theta.\end{aligned}$$

Ex. 3: Reaction-Diffusion Equation

For the 2D reaction-diffusion equation on $(x, y) \in [0, 1]^2$,

$$\begin{aligned}\dot{u}(x, y) &= u_{xx}(x, y) + u_{yy}(x, y) + \lambda u(x, y) \\ u(0, y) &= u(1, y) = u(x, 0) = u(x, 1) = 0,\end{aligned}$$

stability can be proven analytically for any $\lambda \leq 2\pi^2 = 19.739\dots$. Using PIETOOLS, stability was verified for any $\lambda \leq 19.736$.

Ex. 2: Wave Equation

The wave equation on $(x, y) \in [0, 1]^2$

$$\begin{aligned}\ddot{u}(x, y) &= \partial_x^2 u(x, y) + \partial_y^2 u(x, y) \\ u(0, y) &= \partial_x u(0, y) = u(x, 0) = \partial_y u(x, 0) = 0\end{aligned}$$

can be represented in the standardized format by defining $\mathbf{u}_1 = u$ and $\mathbf{u}_2 = \dot{u}$, as

$$\begin{aligned}\begin{bmatrix} \dot{\mathbf{u}}_1 \\ \dot{\mathbf{u}}_2 \end{bmatrix} &= \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \partial_x^2 \begin{bmatrix} \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \partial_y^2 \begin{bmatrix} \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix} \\ &\quad + \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \mathbf{u}_1 \\ \mathbf{u}_2 \end{bmatrix}.\end{aligned}$$

The associated fundamental state is $\hat{\mathbf{u}} = \partial_x^2 \partial_y^2 \mathbf{u}$, with corresponding PIE

$$\begin{aligned}\int_0^x \int_0^y \begin{bmatrix} (x-\theta)(y-\nu) & 0 \\ 0 & (x-\theta)(y-\nu) \end{bmatrix} \begin{bmatrix} \dot{\mathbf{u}}_1 \\ \dot{\mathbf{u}}_2 \end{bmatrix} d\nu d\theta \\ = \int_0^y \begin{bmatrix} 0 & 0 \\ (y-\nu) & 0 \end{bmatrix} \begin{bmatrix} \hat{\mathbf{u}}_1 \\ \hat{\mathbf{u}}_2 \end{bmatrix} d\nu + \int_0^x \begin{bmatrix} 0 & 0 \\ (x-\theta) & 0 \end{bmatrix} \begin{bmatrix} \hat{\mathbf{u}}_1 \\ \hat{\mathbf{u}}_2 \end{bmatrix} d\theta \\ + \int_0^x \int_0^y \begin{bmatrix} 0 & (x-\theta)(y-\nu) \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \dot{\mathbf{u}}_1 \\ \dot{\mathbf{u}}_2 \end{bmatrix} d\nu d\theta\end{aligned}$$

For BCs $u \equiv 0$ on $\partial\Omega$, stability can be verified with PIETOOLS.

