

ON THE CONVERGENCE OF DOUBLY STOCHASTIC PRIMAL-DUAL HYBRID GRADIENT METHOD

YIHENG XIAO* AND HUIKANG LIU†

Abstract. We study a block-structured class of convex-concave saddle-point problems in which both the primal and dual variables admit natural separable decompositions. Motivated by large-scale applications where a full update on either side can be computationally expensive, we propose a doubly stochastic primal-dual hybrid gradient method (DSPDHG) that performs randomized block updates on both primal and dual variables. The method extends classical PDHG and stochastic PDHG (SPDHG) schemes in a unified manner: it reduces to deterministic PDHG when all blocks are selected and to one-sided stochastic variants when only one side is randomized. For the general convex setting, we establish an $\mathcal{O}(1/K)$ ergodic convergence rate for the expected restricted primal-dual gap under suitable blockwise step-size conditions. We further analyze a restarted variant of DSPDHG under a quadratic growth condition in terms of the smoothed primal-dual gap. Under this regularity assumption, we prove linear convergence of the restarted outer iterates. Numerical evidence is provided to show that restarted DSPDHG with standard step sizes demonstrates competitive practical performance compared with PDHG, SPDHG, and their restarted variants.

Key words. convex optimization, primal-dual hybrid gradient, stochastic block-coordinate methods, saddle-point problems, restart, linear convergence

1. Introduction. Primal-dual first-order methods have become a standard tool for solving large-scale convex optimization problems with composite or constrained structure, particularly when accelerated with GPUs [4, 19, 9, 22, 16, 10, 14, 15, 11, 1, 17, 20, 26, 25]. Among them, the primal-dual hybrid gradient (PDHG) method [6, 7] is particularly attractive because each iteration only requires matrix-vector multiplications and proximal mappings, and thus scales favorably to problems for which projections or subproblem solves are expensive. In many applications, however, even a single full primal-dual update may still be computationally demanding when the underlying operator is complex and the number of variables is large.

When the primal or dual variables admit a natural block structure, a successful approach to reduce the per-iteration complexity of PDHG is exploiting randomization. In particular, stochastic PDHG (SPDHG) [5, 3] updates only a randomly selected subset of dual blocks at each iteration while retaining the simplicity of PDHG, and similar convergence guarantees under arbitrary samplings can still be established. Existing SPDHG-type methods, however, primarily exploit separability on the dual side. For many modern problems, the primal variable is also block structured, and a full primal update can be as costly as a full dual update. This naturally raises the following question: can one randomize *both* the primal and dual updates while preserving the favorable theoretical properties of PDHG and SPDHG?

In this paper, we answer this question affirmatively. We consider the block-structured convex optimization problem

$$(1.1) \quad \min_{x=(x_1, \dots, x_m)} \left\{ \sum_{i=1}^n f_i \left(\sum_{j=1}^m A_{ij} x_j \right) + \sum_{j=1}^m g_j(x_j) \right\},$$

where both the primal variable x and the dual variable induced by Fenchel duality are separable. We propose a *doubly stochastic primal-dual hybrid gradient* (DSPDHG) method, which at every iteration samples a subset of primal blocks and a subset of

*Shanghai Jiao Tong University (xyh051014@sjtu.edu.cn).

† Shanghai Jiao Tong University (hkl1u@sjtu.edu.cn).

dual blocks, and updates only the selected coordinates. The resulting method can be viewed as a genuine two-sided stochastic generalization of PDHG: it reduces to deterministic PDHG when all blocks are selected, and to singly stochastic variants when only one side is randomized.

The main technical challenge is that two independent random samplings interact through the bilinear coupling operator. In contrast to the classical SPDHG setting, one must control not only the stochastic dual correction but also the stochastic primal correction, as well as the cross terms generated by the simultaneous extrapolation of both variables. To address this difficulty, we introduce weighted Lyapunov functions adapted to the two sampling distributions and derive one-step inequalities that connect the virtual full updates with the actual random block updates. This allows us to establish both sublinear and linear convergence guarantees within a unified framework.

Our main contributions are summarized as follows.

General convex problems. For proper, lower semicontinuous, convex objectives and suitable blockwise step sizes, we establish an $\mathcal{O}(1/K)$ ergodic convergence rate for the expected primal–dual gap of DSPDHG. The result applies to general fixed-cardinality samplings on both the primal and dual sides and extends the usual singly stochastic PDHG analysis to the doubly stochastic regime.

Linear convergence under regularity. We further show that, under a smoothed quadratic growth condition [12, 22, 21, 18] on a smoothed primal–dual gap functional, a restarted version of DSPDHG converges linearly. This condition is weaker and more flexible than global strong convexity–strong concavity, and is tailored to the primal–dual geometry of the method.

A unified two-sided stochastic framework. Our analysis makes explicit how the primal sampling probabilities, the dual sampling probabilities, and the block interaction structure of the operator jointly determine the admissible step sizes and convergence behavior. In particular, the deterministic PDHG and one-sided stochastic PDHG schemes arise as special cases of our framework.

Related work. A closely related work is the doubly stochastic primal–dual coordinate method proposed in [24] for bilinear saddle-point problems, where linear convergence was established under strong convexity–concavity. However, the extrapolation parameter in that method depends explicitly on the strong convexity parameters. Another related method, proposed in [2] for problems with a sparse coupling operator A , is also formulated as a primal–dual coordinate descent scheme. Its randomness, however, is induced by a single sampling mechanism and is therefore closer in spirit to a singly stochastic method. In contrast, our approach is developed for general convex saddle-point problems, employs genuinely separate samplings on the primal and dual sides, and yields both an $\mathcal{O}(1/K)$ ergodic convergence guarantee and linear convergence of the restarted scheme under a smoothed quadratic growth condition.

The rest of the paper is organized as follows. In section 2, we introduce notation, the saddle-point formulation, and the general assumption used throughout the paper. Section 3 presents the DSPDHG algorithm and the basic one-step descent estimate. Section 4 proves the ergodic $\mathcal{O}(1/K)$ convergence rate in the general convex setting. Section 5 establishes linear convergence of the restarted scheme under the smoothed quadratic growth condition. Numerical experiments are reported in section 6.

2. Preliminaries.

2.1. Problem formulation and saddle-point representation. Let $\mathcal{X}_j$ ($j \in [m]$) and $\mathcal{Y}_i$ ($i \in [n]$) be finite-dimensional Euclidean spaces, and define

$$\mathcal{X} := \prod_{j=1}^m \mathcal{X}_j, \quad \mathcal{Y} := \prod_{i=1}^n \mathcal{Y}_i, \quad \mathcal{Z} := \mathcal{X} \times \mathcal{Y}.$$

For $x \in \mathcal{X}$ and $y \in \mathcal{Y}$, we write

$$x = (x_1, \dots, x_m), \quad y = (y_1, \dots, y_n),$$

with $x_j \in \mathcal{X}_j$ and $y_i \in \mathcal{Y}_i$. Throughout the paper, we equip all product spaces with the canonical inner products and their induced norms.

We consider the structured convex optimization problem

$$(2.1) \quad \min_{x \in \mathcal{X}} \left\{ \sum_{i=1}^n f_i \left(\sum_{j=1}^m A_{ij} x_j \right) + \sum_{j=1}^m g_j(x_j) \right\},$$

where each $f_i : \mathcal{Y}_i \rightarrow \mathbb{R} \cup \{+\infty\}$ and $g_j : \mathcal{X}_j \rightarrow \mathbb{R} \cup \{+\infty\}$ is proper, lower semi-continuous, and convex, and each $A_{ij} : \mathcal{X}_j \rightarrow \mathcal{Y}_i$ is linear. For each $i \in [n]$, let $f_i^* : \mathcal{Y}_i \rightarrow \mathbb{R} \cup \{+\infty\}$ denote the Fenchel conjugate of f_i . For convenience, define

$$g(x) := \sum_{j=1}^m g_j(x_j), \quad f^*(y) := \sum_{i=1}^n f_i^*(y_i),$$

and let $A : \mathcal{X} \rightarrow \mathcal{Y}$ denote the block operator

$$(Ax)_i = \sum_{j=1}^m A_{ij} x_j, \quad i \in [n].$$

We also use the row and column block operators

$$A_i := (A_{i1}, \dots, A_{im}) : \mathcal{X} \rightarrow \mathcal{Y}_i, \quad A^j := \begin{bmatrix} A_{1j} \\ \vdots \\ A_{nj} \end{bmatrix} : \mathcal{X}_j \rightarrow \mathcal{Y}.$$

By Fenchel duality, (2.1) admits the saddle-point reformulation

$$(2.2) \quad \min_{x \in \mathcal{X}} \max_{y \in \mathcal{Y}} \Phi(x, y) := g(x) + \sum_{i=1}^n \langle A_i x, y_i \rangle - f^*(y).$$

We impose the following standard assumptions throughout the paper.

Assumption 2.1. The functions $\{f_i\}_{i=1}^n$ and $\{g_j\}_{j=1}^m$ are proper, lower semicontinuous, and convex. The saddle-point set $\mathcal{Z}^*$ of (2.2) is nonempty.

A pair $z^* = (x^*, y^*) \in \mathcal{Z}$ is a saddle point of (2.2) if and only if it satisfies the KKT system

$$(2.3) \quad 0 \in \begin{bmatrix} \partial g(x^*) + A^\top y^* \\ \partial f^*(y^*) - Ax^* \end{bmatrix}.$$

Equivalently,

$$-(A^j)^\top y^* \in \partial g_j(x_j^*), \quad j \in [m] \quad A_i x^* \in \partial f_i^*(y_i^*), \quad i \in [n].$$

2.2. Proximal mappings and block samplings. For a proper, lower semicontinuous, convex function h on a Euclidean space $\mathcal{U}$ and a positive definite self-adjoint operator M , we define the proximal mapping

$$\text{prox}_{Mh}(u) := \arg \min_{v \in \mathcal{U}} \left\{ h(v) + \frac{1}{2} \|v - u\|_{M^{-1}}^2 \right\},$$

where $\|w\|_M^2 := \langle Mw, w \rangle$. Let $\tau_j > 0$ for $j \in [m]$ and $\sigma_i > 0$ for $i \in [n]$, and define the block-diagonal operators

$$\tau := \text{diag}(\tau_1 I, \dots, \tau_m I), \quad \sigma := \text{diag}(\sigma_1 I, \dots, \sigma_n I).$$

The doubly stochastic algorithm uses two random block samplings at iteration $k+1$: a primal sampling $S^{k+1} \subseteq [m]$ and a dual sampling $T^{k+1} \subseteq [n]$. We write

$$p_j := \mathbb{P}(j \in S^{k+1}), \quad q_i := \mathbb{P}(i \in T^{k+1}),$$

and assume that these probabilities are strictly positive. Associated with the sampling probabilities, we define

$$P := \text{diag}(p_1 I, \dots, p_m I), \quad Q := \text{diag}(q_1 I, \dots, q_n I).$$

Given the extrapolated iterates $\bar{y}^k$ and $\bar{x}^{k+1}$, it is convenient to introduce the corresponding *virtual full updates*

$$(2.4) \quad \hat{x}_j^{k+1} = \text{prox}_{\tau_j g_j}(x_j^k - \tau_j (A^j)^\top \bar{y}^k), \quad \hat{y}_i^{k+1} = \text{prox}_{\sigma_i f_i^*}(y_i^k + \sigma_i A_i \bar{x}^{k+1}),$$

for all $j \in [m]$ and $i \in [n]$. The actual algorithm updates only the coordinates in S^{k+1} and T^{k+1} ; the operators P^{-1} and Q^{-1} then appear naturally in the extrapolation formulas and in the unbiased representation of the stochastic iterates.

2.3. Weighted metrics and primal–dual gap functionals. The analysis is carried out in the weighted norm

$$(2.5) \quad \|z\|_V^2 := \|x\|_{\tau^{-1}P^{-1}}^2 + \|y\|_{\sigma^{-1}Q^{-1}}^2, \quad z = (x, y) \in \mathcal{Z}.$$

We also use the primal–dual gap kernel

$$(2.6) \quad H(\bar{x}, \bar{y}, x, y) = H(\bar{z}, z) := \Phi(\bar{x}, y) - \Phi(x, \bar{y}),$$

which is nonnegative when z is a saddle point. For a set $B \subseteq \mathcal{Z}$, the restricted primal–dual gap is defined by

$$(2.7) \quad G_B(\bar{x}, \bar{y}) := \sup_{z=(x,y) \in B} H(\bar{x}, \bar{y}, x, y).$$

This quantity will be used to state the ergodic $\mathcal{O}(1/K)$ convergence result in the general convex setting.

To capture the linear convergence regime, we further define the smoothed gap functional

$$(2.8) \quad G_\mu(\bar{z}, \dot{z}) := \sup_{z \in \mathcal{Z}} \left\{ H(\bar{x}, \bar{y}, x, y) - \frac{\mu}{2} \|z - \dot{z}\|_V^2 \right\}, \quad \bar{z} = (\bar{x}, \bar{y}), \quad \dot{z} \in \mathcal{Z},$$

where $\mu > 0$ is a smoothing parameter.

Algorithm 3.1 Doubly stochastic primal–dual hybrid gradient (DSPDHG)

```

1: Input: stepsizes  $\{\tau_j\}_{j=1}^m$ ,  $\{\sigma_i\}_{i=1}^n$ , initial points  $x^0 \in \mathcal{X}$ ,  $y^0 \in \mathcal{Y}$ , and set  $\bar{y}^0 = y^0$ 
2: for  $k = 0, 1, 2, \dots$  do
3:   Draw a primal block set  $S^{k+1} \subseteq [m]$ 
4:   Update  $x_j^{k+1} = \begin{cases} \text{prox}_{\tau_j g_j}(x_j^k - \tau_j (A^j)^\top \bar{y}^k), & j \in S^{k+1}, \\ x_j^k, & j \notin S^{k+1} \end{cases}$ 
5:   Set  $\bar{x}^{k+1} = x^{k+1} + (P^{-1} - I)(x^{k+1} - x^k)$ 
6:   Draw a dual block set  $T^{k+1} \subseteq [n]$ 
7:   Update  $y_i^{k+1} = \begin{cases} \text{prox}_{\sigma_i f_i^*}(y_i^k + \sigma_i A_i \bar{x}^{k+1}), & i \in T^{k+1}, \\ y_i^k, & i \notin T^{k+1} \end{cases}$ 
8:   Set  $\bar{y}^{k+1} = y^{k+1} + Q^{-1}(y^{k+1} - y^k)$ 
9:
10: end for

```

3. The DSPDHG Algorithm. We now present the doubly stochastic primal–dual hybrid gradient method. In contrast to classical SPDHG, which randomizes only the dual update, the proposed method randomizes *both* the primal and dual block updates. This is particularly useful when both sides of the saddle-point formulation admit a block decomposition and full updates on either side are expensive.

The method maintains both the current iterates (x^k, y^k) and the extrapolated variables $(\bar{x}^k, \bar{y}^k)$. Given $(x^k, \bar{y}^k)$, the primal update is first performed on the sampled primal coordinates, followed by an extrapolation in the primal variable. Then the dual update is carried out on the sampled dual coordinates, followed by an extrapolation in the dual variable.

The intuition behind these two extrapolations is as follows. The full updates $\hat{x}^{k+1}$ and $\hat{y}^{k+1}$ given by (2.4) are not computed in practice, but it is easy to see that

$$\begin{aligned} \hat{x}^{k+1} &= P^{-1}\mathbb{E}[x^{k+1} \mid \mathcal{F}_k] + (I - P^{-1})x^k, \\ \hat{y}^{k+1} &= Q^{-1}\mathbb{E}[y^{k+1} \mid \mathcal{F}_{k+\frac{1}{2}}] + (I - Q^{-1})y^k. \end{aligned}$$

where $\mathcal{F}_k$ and $\mathcal{F}_{k+\frac{1}{2}}$ are the sigma-fields generated by $\{S^1, \dots, S^k, T^1, \dots, T^k\}$ and $\{S^1, \dots, S^{k+1}, T^1, \dots, T^k\}$, respectively. Therefore, the actual stochastic updates can be represented as unbiased blockwise realizations of (2.4) after rescaling by P^{-1} and Q^{-1} , i.e.,

$$(3.1) \quad \mathbb{E}[\bar{x}^{k+1} \mid \mathcal{F}_k] = \hat{x}^{k+1}, \quad \text{and} \quad \mathbb{E}[\bar{y}^{k+1} \mid \mathcal{F}_{k+\frac{1}{2}}] = \hat{y}^{k+1} + \mathbb{E}[y^{k+1} - y^k \mid \mathcal{F}_{k+\frac{1}{2}}]$$

where the last term $\mathbb{E}[y^{k+1} - y^k \mid \mathcal{F}_{k+\frac{1}{2}}]$ works as the extrapolation in the dual variables of the standard PDHG method.

Our convergence analysis follows the standard PDHG philosophy of deriving a one-step estimate from the proximal optimality conditions, but the doubly stochastic setting requires additional correction terms to account for the mismatch between virtual full updates and realized random block updates.

3.1. A one-step estimate.

Recall the primal–dual gap kernel

$$H(\bar{x}, \bar{y}, x, y) := \Phi(\bar{x}, y) - \Phi(x, \bar{y}).$$

The next lemma is a one-step estimate for the virtual full updates. It is obtained by applying the proximal optimality condition separately to the primal and dual virtual updates.

LEMMA 3.1. *Suppose Assumption 2.1 holds. For any $k \geq 0$ and any $(x, y) \in \mathcal{Z}$, the virtual full updates satisfy*

$$\begin{aligned} H(\hat{x}^{k+1}, \hat{y}^{k+1}, x, y) &\leq \langle A\hat{x}^{k+1}, y \rangle - \langle Ax, \hat{y}^{k+1} \rangle \\ &\quad + \langle A(x - \hat{x}^{k+1}), \bar{y}^k \rangle + \langle A\bar{x}^{k+1}, \hat{y}^{k+1} - y \rangle \\ &\quad + \frac{1}{2} \|x^k - x\|_{\tau^{-1}}^2 - \frac{1}{2} \|\hat{x}^{k+1} - x^k\|_{\tau^{-1}}^2 - \frac{1}{2} \|\hat{x}^{k+1} - x\|_{\tau^{-1}}^2 \\ &\quad + \frac{1}{2} \|y^k - y\|_{\sigma^{-1}}^2 - \frac{1}{2} \|\hat{y}^{k+1} - y^k\|_{\sigma^{-1}}^2 - \frac{1}{2} \|\hat{y}^{k+1} - y\|_{\sigma^{-1}}^2. \end{aligned} \quad (3.2)$$

Proof. We use the standard proximal inequality: if $\hat{u} = \text{prox}_{\mu h}(u)$, then

$$h(v) - h(\hat{u}) \geq \langle \mu^{-1}(u - \hat{u}), v - \hat{u} \rangle, \quad \forall v.$$

Applying this inequality to

$$\hat{x}_j^{k+1} = \text{prox}_{\tau_j g_j}(x_j^k - \tau_j (A^j)^\top \bar{y}^k),$$

and using the identity

$$2\langle a - b, c - b \rangle = \|a - b\|^2 + \|c - b\|^2 - \|a - c\|^2,$$

we obtain, for each $j \in [m]$,

$$\begin{aligned} g_j(\hat{x}_j^{k+1}) - g_j(x_j) &\leq \langle A^j(x_j - \hat{x}_j^{k+1}), \bar{y}^k \rangle \\ &\quad + \frac{1}{2\tau_j} \|x_j^k - x_j\|^2 - \frac{1}{2\tau_j} \|\hat{x}_j^{k+1} - x_j^k\|^2 - \frac{1}{2\tau_j} \|\hat{x}_j^{k+1} - x_j\|^2. \end{aligned}$$

Summing over j gives

$$\begin{aligned} g(\hat{x}^{k+1}) - g(x) &\leq \langle A(x - \hat{x}^{k+1}), \bar{y}^k \rangle + \frac{1}{2} \|x^k - x\|_{\tau^{-1}}^2 \\ &\quad - \frac{1}{2} \|\hat{x}^{k+1} - x^k\|_{\tau^{-1}}^2 - \frac{1}{2} \|\hat{x}^{k+1} - x\|_{\tau^{-1}}^2. \end{aligned} \quad (3.3)$$

Similarly, applying the same proximal inequality to

$$\hat{y}_i^{k+1} = \text{prox}_{\sigma_i f_i^*}(y_i^k + \sigma_i A_i \bar{x}^{k+1}),$$

we obtain, after summing over $i \in [n]$,

$$\begin{aligned} f^*(\hat{y}^{k+1}) - f^*(y) &\leq \langle A\bar{x}^{k+1}, \hat{y}^{k+1} - y \rangle + \frac{1}{2} \|y^k - y\|_{\sigma^{-1}}^2 \\ &\quad - \frac{1}{2} \|\hat{y}^{k+1} - y^k\|_{\sigma^{-1}}^2 - \frac{1}{2} \|\hat{y}^{k+1} - y\|_{\sigma^{-1}}^2. \end{aligned} \quad (3.4)$$

Finally, by the definition of the gap kernel, we have

$$H(\hat{x}^{k+1}, \hat{y}^{k+1}, x, y) = \langle A\hat{x}^{k+1}, y \rangle - \langle Ax, \hat{y}^{k+1} \rangle + g(\hat{x}^{k+1}) - g(x) + f^*(\hat{y}^{k+1}) - f^*(y).$$

Substituting (3.3) and (3.4) into the above identity yields (3.2). $\square$

The main difficulty is to convert (3.2), which is written in terms of the virtual full updates, into an inequality involving the actual stochastic iterates. For fixed $(x, y) \in \mathcal{Z}$, define

$$\begin{aligned} \Gamma_k := & \langle A(\bar{x}^{k+1} - x), Q^{-1}(y^{k+1} + y^{k-1} - 2y^k) \rangle \\ & + \frac{1}{2} \|x^k - x\|_{\tau^{-1}P^{-1}}^2 - \frac{1}{2} \|x^{k+1} - x^k\|_{\tau^{-1}P^{-1}}^2 - \frac{1}{2} \|x^{k+1} - x\|_{\tau^{-1}P^{-1}}^2 \\ & + \frac{1}{2} \|y^k - y\|_{\sigma^{-1}Q^{-1}}^2 - \frac{1}{2} \|y^{k+1} - y^k\|_{\sigma^{-1}Q^{-1}}^2 - \frac{1}{2} \|y^{k+1} - y\|_{\sigma^{-1}Q^{-1}}^2. \end{aligned} \quad (3.5)$$

Here and below we use the convention $y^{-1} = y^0$.

LEMMA 3.2. Suppose Assumption 2.1 holds. For all $k \geq 0$ and all $(x, y) \in \mathcal{Z}$,

$$\mathbb{E}[H(\hat{x}^{k+1}, \hat{y}^{k+1}, x, y)] \leq \mathbb{E}[\Gamma_k]. \quad (3.6)$$

Proof. Recall that $\mathcal{F}_k$ and $\mathcal{F}_{k+\frac{1}{2}}$ are the sigma-fields generated by $S^1, \dots, S^k, T^1, \dots, T^k$ and $S^1, \dots, S^{k+1}, T^1, \dots, T^k$, respectively. By the definitions of the stochastic updates and extrapolations,

$$\bar{x}^{k+1} = x^k + P^{-1}(x^{k+1} - x^k), \quad \bar{y}^k = y^k + Q^{-1}(y^k - y^{k-1}),$$

and hence

$$\mathbb{E}[\bar{x}^{k+1} \mid \mathcal{F}_k] = \hat{x}^{k+1}. \quad (3.7)$$

Similarly, since

$$\mathbb{E}[y^{k+1} - y^k \mid \mathcal{F}_{k+\frac{1}{2}}] = Q(\hat{y}^{k+1} - y^k),$$

we have

$$\hat{y}^{k+1} = y^k + Q^{-1}\mathbb{E}[y^{k+1} - y^k \mid \mathcal{F}_{k+\frac{1}{2}}]. \quad (3.8)$$

We first compare the quadratic terms. Define $\hat{\mathcal{Q}}_k$ and $\mathcal{Q}_k$ to be the quadratic terms in (3.2) and (3.5) respectively.

$$\begin{aligned} \hat{\mathcal{Q}}_k := & \frac{1}{2} \|x^k - x\|_{\tau^{-1}}^2 - \frac{1}{2} \|\hat{x}^{k+1} - x^k\|_{\tau^{-1}}^2 - \frac{1}{2} \|\hat{x}^{k+1} - x\|_{\tau^{-1}}^2 \\ & + \frac{1}{2} \|y^k - y\|_{\sigma^{-1}}^2 - \frac{1}{2} \|\hat{y}^{k+1} - y^k\|_{\sigma^{-1}}^2 - \frac{1}{2} \|\hat{y}^{k+1} - y\|_{\sigma^{-1}}^2, \\ \mathcal{Q}_k := & \frac{1}{2} \|x^k - x\|_{\tau^{-1}P^{-1}}^2 - \frac{1}{2} \|x^{k+1} - x^k\|_{\tau^{-1}P^{-1}}^2 - \frac{1}{2} \|x^{k+1} - x\|_{\tau^{-1}P^{-1}}^2 \\ & + \frac{1}{2} \|y^k - y\|_{\sigma^{-1}Q^{-1}}^2 - \frac{1}{2} \|y^{k+1} - y^k\|_{\sigma^{-1}Q^{-1}}^2 - \frac{1}{2} \|y^{k+1} - y\|_{\sigma^{-1}Q^{-1}}^2. \end{aligned} \quad (3.9)$$

For the primal part, the blockwise update rule implies

$$\begin{aligned} & \mathbb{E} \left[\frac{1}{2} \|x^k - x\|_{\tau^{-1}P^{-1}}^2 - \frac{1}{2} \|x^{k+1} - x^k\|_{\tau^{-1}P^{-1}}^2 - \frac{1}{2} \|x^{k+1} - x\|_{\tau^{-1}P^{-1}}^2 \mid \mathcal{F}_k \right] \\ & = \frac{1}{2} \|x^k - x\|_{\tau^{-1}}^2 - \frac{1}{2} \|\hat{x}^{k+1} - x^k\|_{\tau^{-1}}^2 - \frac{1}{2} \|\hat{x}^{k+1} - x\|_{\tau^{-1}}^2. \end{aligned} \quad (3.10)$$

Indeed, this is because for any $\mathcal{F}_k$ measurable random vector x ,

$$\forall j \in [m], \quad \mathbb{E}[\|x_j^{k+1} - x_j\|^2 \mid \mathcal{F}_k] = p_j \|\hat{x}_j^{k+1} - x_j\|^2 + (1 - p_j) \|x_j^k - x_j\|^2.$$

229 The same argument applied conditionally on $\mathcal{F}_{k+\frac{1}{2}}$ gives

$$\begin{aligned}
 230 \quad & \mathbb{E} \left[\frac{1}{2} \|y^k - y\|_{\sigma^{-1}Q^{-1}}^2 - \frac{1}{2} \|y^{k+1} - y^k\|_{\sigma^{-1}Q^{-1}}^2 - \frac{1}{2} \|y^{k+1} - y\|_{\sigma^{-1}Q^{-1}}^2 \middle| \mathcal{F}_{k+\frac{1}{2}} \right] \\
 231 \quad (3.11) \quad & = \frac{1}{2} \|y^k - y\|_{\sigma^{-1}}^2 - \frac{1}{2} \|\hat{y}^{k+1} - y^k\|_{\sigma^{-1}}^2 - \frac{1}{2} \|\hat{y}^{k+1} - y\|_{\sigma^{-1}}^2.
 \end{aligned}$$

232 Combining (3.10) and (3.11), we have

$$233 \quad (3.12) \quad \mathbb{E}[\hat{Q}_k] = \mathbb{E}[Q_k]$$

234 It remains to rewrite the bilinear terms in (3.2). Denote

$$235 \quad B_k := \langle A\hat{x}^{k+1}, y \rangle - \langle Ax, \hat{y}^{k+1} \rangle + \langle A(x - \hat{x}^{k+1}), \bar{y}^k \rangle + \langle A\bar{x}^{k+1}, \hat{y}^{k+1} - y \rangle.$$

236 Using (3.7) and the fact that y and $\bar{y}^k$ are $\mathcal{F}_k$ -measurable, we obtain

$$\begin{aligned}
 237 \quad \mathbb{E}[B_k] &= \mathbb{E}[\langle A\bar{x}^{k+1}, y \rangle - \langle Ax, \hat{y}^{k+1} \rangle + \langle A(x - \bar{x}^{k+1}), \bar{y}^k \rangle + \langle A\bar{x}^{k+1}, \hat{y}^{k+1} - y \rangle] \\
 238 \quad &= \mathbb{E}[\langle A(\bar{x}^{k+1} - x), \hat{y}^{k+1} - \bar{y}^k \rangle].
 \end{aligned}$$

239 By (3.8) and the definition of $\bar{y}^k$,

$$240 \quad \hat{y}^{k+1} - \bar{y}^k = Q^{-1} \mathbb{E}[y^{k+1} - y^k \mid \mathcal{F}_{k+\frac{1}{2}}] - Q^{-1}(y^k - y^{k-1}).$$

241 Since $\bar{x}^{k+1}$ is $\mathcal{F}_{k+\frac{1}{2}}$ -measurable, it follows that

$$242 \quad (3.13) \quad \mathbb{E}[B_k] = \mathbb{E}[\langle A(\bar{x}^{k+1} - x), Q^{-1}(y^{k+1} + y^{k-1} - 2y^k) \rangle].$$

243 Taking expectations in (3.2) and substituting (3.12), (3.13), we obtain exactly

$$244 \quad \mathbb{E}[H(\hat{x}^{k+1}, \hat{y}^{k+1}, x, y)] \leq \mathbb{E}[\Gamma_k],$$

245 which proves the claim. $\square$

246 **3.2. Step-size conditions.** The doubly stochastic setting generates coupling
 247 terms involving both the sampled primal and dual coordinates. To control these
 248 terms, we introduce the block operator constants

$$249 \quad (3.14) \quad \Lambda_{r,s} := \max\{\|A_{IJ}\| : I \subseteq [n], J \subseteq [m], |I| = r, |J| = s\},$$

$$250 \quad (3.15) \quad \Lambda_r := \max\{\|A_I\| : I \subseteq [n], |I| = r\},$$

251 where A_{IJ} denotes the restriction of A to the row block set I and the column block
 252 set J , and A_I denotes the restriction of A to the row block set I . The norms above
 253 are the induced operator norms under the canonical Euclidean product norms.

254 We focus on fixed-cardinality samplings satisfying

$$255 \quad |S^{k+1}| = s, \quad |T^{k+1}| = r, \quad \forall k \geq 0.$$

256 For uniform fixed-cardinality samplings, the corresponding marginal probabilities are

$$257 \quad p_j = p = \frac{s}{m}, \quad q_i = q = \frac{r}{n}.$$

Assumption 3.3. We use uniform fixed-cardinality samplings and equal step sizes, namely

$$p_j = p, \quad q_i = q, \quad \tau_j = \tau, \quad \sigma_i = \sigma.$$

Besides, we assume that the sampling sets $S^k, T^k, k \geq 1$ are all independent. Moreover, the step sizes and sampling probabilities satisfy

$$(3.16) \quad \Lambda_{r,s} p^{-1/2} q^{-1/2} \tau^{1/2} \sigma^{1/2} = \gamma_1^2 < \frac{1}{2},$$

$$(3.17) \quad \Lambda_r p^{1/2} q^{-1/2} \tau^{1/2} \sigma^{1/2} = \gamma_2^2 \leq \frac{1}{2}.$$

Remark 3.4. The assumptions of uniform sampling and equal step sizes are imposed only to simplify the notation and the proof of Lemma 4.1. The analysis can be extended to nonuniform samplings and block-dependent step sizes by using an expected separable overapproximation (ESO) inequality, as in [5]. The remaining parts of the argument do not rely on uniform sampling or equal step sizes.

Remark 3.5. The constant $\frac{1}{2}$ in our step-size condition is more conservative than the constant 1 typically used in deterministic PDHG. This additional restriction arises from the stochastic updates on both the primal and dual sides. In our numerical experiments, however, we observed that the larger constant 1 also works well in practice.

Assumption 3.3 is a two-sided analogue of the standard SPDHG step-size restriction. The first condition controls the interaction between the stochastic primal and dual increments, while the second ensures positivity of the time-dependent Lyapunov functional used below.

4. General Convex Convergence. We now establish the basic convergence guarantees of DSPDHG in the general convex setting. Our goal is to prove an $\mathcal{O}(1/K)$ ergodic convergence rate for the expected restricted primal-dual gap.

4.1. Lyapunov functions. To telescope the one-step estimate, we introduce two weighted quadratic functionals. For $z = (x, y)$, define

$$(4.1) \quad V(z) = \frac{1}{4} \|x\|_{\tau^{-1}P^{-1}}^2 + \frac{1}{4} \|y\|_{\sigma^{-1}Q^{-1}}^2 + \langle AP^{-1}x, Q^{-1}y \rangle,$$

$$(4.2) \quad V_k(z) = \frac{1}{2} \|x\|_{\tau^{-1}P^{-1}}^2 + \frac{1}{2} \|y\|_{\sigma^{-1}Q^{-1}}^2 + \frac{1}{4} \|y^k - y^{k-1}\|_{\sigma^{-1}Q^{-1}}^2 - \langle Ax, Q^{-1}(y^k - y^{k-1}) \rangle.$$

Here and below, we use the convention $y^{-1} = y^0$. Under Assumption 3.3, these functionals satisfy the coercivity and descent properties needed for the telescoping argument.

LEMMA 4.1. *Suppose Assumption 3.3 holds. Then for all $k \geq 0$:*

$$(4.3) \quad V(x^{k+1} - x^k, y^k - y^{k-1}) \geq \frac{1 - 2\gamma_1^2}{4} \left(\|x^{k+1} - x^k\|_{\tau^{-1}P^{-1}}^2 + \|y^k - y^{k-1}\|_{\sigma^{-1}Q^{-1}}^2 \right),$$

$$(4.4) \quad V_k(z) \geq \frac{1}{4} \|x\|_{\tau^{-1}P^{-1}}^2 + \frac{1}{2} \|y\|_{\sigma^{-1}Q^{-1}}^2,$$

$$(4.5) \quad V_k(x^k - x, y^k - y) - V_{k+1}(x^{k+1} - x, y^{k+1} - y) \geq V(x^{k+1} - x^k, y^k - y^{k-1}) + \Gamma_k.$$

Proof. First, we use (3.16) to prove (4.3). By definition, we have

$$(4.6) \quad V(x^{k+1} - x^k, y^k - y^{k-1}) = \frac{1}{4} \|x^{k+1} - x^k\|_{\tau^{-1}P^{-1}}^2 + \frac{1}{4} \|y^k - y^{k-1}\|_{\sigma^{-1}Q^{-1}}^2 \\ + \langle AP^{-1}(x^{k+1} - x^k), Q^{-1}(y^k - y^{k-1}) \rangle.$$

Let $J = S^{k+1}, I = T^k$. Only for $j \in J$, $x_j^{k+1} - x_j^k$ is nonzero and only for $i \in I$, $y_i^k - y_i^{k-1}$ is nonzero. Therefore,

$$(4.7) \quad |\langle AP^{-1}(x^{k+1} - x^k), Q^{-1}(y^k - y^{k-1}) \rangle| = \left| \sum_{i=1}^n \sum_{j=1}^m \frac{1}{pq} \langle A_{ij}(x_j^{k+1} - x_j^k), y_i^k - y_i^{k-1} \rangle \right| \\ = \left| \sum_{i \in I} \sum_{j \in J} \tau^{\frac{1}{2}} \sigma^{\frac{1}{2}} p^{-\frac{1}{2}} q^{-\frac{1}{2}} \langle A_{ij}(x_j^{k+1} - x_j^k) \tau^{-\frac{1}{2}} p^{-\frac{1}{2}}, (y_i^k - y_i^{k-1}) \sigma^{-\frac{1}{2}} q^{-\frac{1}{2}} \rangle \right| \\ = \left(p^{-\frac{1}{2}} q^{-\frac{1}{2}} \tau^{\frac{1}{2}} \sigma^{\frac{1}{2}} \right) \cdot \left| \langle A_{IJ} p^{-\frac{1}{2}} \tau^{-\frac{1}{2}} (x_J^{k+1} - x_J^k), \sigma^{-\frac{1}{2}} q^{-\frac{1}{2}} (y_I^k - y_I^{k-1}) \rangle \right| \\ \leq \left(p^{-\frac{1}{2}} q^{-\frac{1}{2}} \tau^{\frac{1}{2}} \sigma^{\frac{1}{2}} \right) \cdot \|A_{IJ}\| \cdot \|p^{-\frac{1}{2}} \tau^{-\frac{1}{2}} (x_J^{k+1} - x_J^k)\| \cdot \|\sigma^{-\frac{1}{2}} q^{-\frac{1}{2}} (y_I^k - y_I^{k-1})\| \\ \leq \frac{\gamma_1^2}{2} \left(\|p^{-\frac{1}{2}} \tau^{-\frac{1}{2}} (x_J^{k+1} - x_J^k)\|^2 + \|\sigma^{-\frac{1}{2}} q^{-\frac{1}{2}} (y_I^k - y_I^{k-1})\|^2 \right) \\ = \frac{\gamma_1^2}{2} \left(\|x^{k+1} - x^k\|_{\tau^{-1}P^{-1}}^2 + \|y^k - y^{k-1}\|_{\sigma^{-1}Q^{-1}}^2 \right).$$

Combining (4.6) and (4.7), we get (4.3).

Second, we use (3.17) to prove (4.4). It is obvious that we only need to show

$$(4.8) \quad \frac{1}{4} \|x\|_{\tau^{-1}P^{-1}}^2 + \frac{1}{4} \|y^k - y^{k-1}\|_{\sigma^{-1}Q^{-1}}^2 - \langle Ax, Q^{-1}(y^k - y^{k-1}) \rangle \geq 0$$

Similar to the previous computation, we have

$$(4.9) \quad |\langle Ax, Q^{-1}(y^k - y^{k-1}) \rangle| = \left| \sum_{i \in I} \sum_{j=1}^m \sigma^{\frac{1}{2}} \tau^{\frac{1}{2}} p^{\frac{1}{2}} q^{-\frac{1}{2}} \langle A_{ij} x_j \tau^{-\frac{1}{2}} p^{-\frac{1}{2}}, (y_i^k - y_i^{k-1}) \sigma^{-\frac{1}{2}} q^{-\frac{1}{2}} \rangle \right| \\ = \left(p^{\frac{1}{2}} q^{-\frac{1}{2}} \tau^{\frac{1}{2}} \sigma^{\frac{1}{2}} \right) \cdot \left| \langle p^{-\frac{1}{2}} \tau^{-\frac{1}{2}} A_I x, \sigma^{-\frac{1}{2}} q^{-\frac{1}{2}} (y_I^k - y_I^{k-1}) \rangle \right| \\ \leq \frac{\gamma_2^2}{2} \left(\|p^{-\frac{1}{2}} \tau^{-\frac{1}{2}} x\|^2 + \|\sigma^{-\frac{1}{2}} q^{-\frac{1}{2}} (y_I^k - y_I^{k-1})\|^2 \right) \\ \leq \frac{1}{4} \|x\|_{\tau^{-1}P^{-1}}^2 + \frac{1}{4} \|y^k - y^{k-1}\|_{\sigma^{-1}Q^{-1}}^2.$$

This inequality directly leads to (4.8).

Finally, we use (3.16) to prove (4.5). We can compute

$$(4.10) \quad V_k(x^k - x, y^k - y) - V_{k+1}(x^{k+1} - x, y^{k+1} - y) - V(x^{k+1} - x^k, y^k - y^{k-1}) - \Gamma_k \\ = \frac{1}{4} \|x^{k+1} - x^k\|_{\tau^{-1}P^{-1}}^2 + \frac{1}{4} \|y^{k+1} - y^k\|_{\sigma^{-1}Q^{-1}}^2 \\ + \langle A(I - P^{-1})(x^{k+1} - x^k), Q^{-1}(y^{k+1} - y^k) \rangle.$$

308 Following the same procedure as in (4.7), and letting $\gamma_3^2 = \Lambda_{r,s} \cdot (p^{-\frac{1}{2}} - p^{\frac{1}{2}})q^{-\frac{1}{2}}\tau^{\frac{1}{2}}\sigma^{\frac{1}{2}}$,
 309 we have

$$\begin{aligned} & |\langle A(I - P^{-1})(x^{k+1} - x^k), Q^{-1}(y^{k+1} - y^k) \rangle| \\ 310 \quad (4.11) \quad & \leq \frac{\gamma_3^2}{2} \left(\|x^{k+1} - x^k\|_{\tau^{-1}P^{-1}}^2 + \|y^{k+1} - y^k\|_{\sigma^{-1}Q^{-1}}^2 \right) \\ & \leq \frac{1}{4} \left(\|x^{k+1} - x^k\|_{\tau^{-1}P^{-1}}^2 + \|y^{k+1} - y^k\|_{\sigma^{-1}Q^{-1}}^2 \right), \end{aligned}$$

311 where the last inequality follows from $\gamma_3^2 \leq \gamma_1^2 \leq \frac{1}{2}$. This inequality directly leads to
 312 (4.5). $\square$

313 Combining Lemmas 3.2 and 4.1 yields the fundamental recursive estimate.

314 **LEMMA 4.2.** *Under Assumption 2.1 and Assumption 3.3, for every $k \geq 0$ and*
 315 *$(x, y) \in \mathcal{Z}$,*

$$\begin{aligned} 316 \quad \mathbb{E}[H(\hat{x}^{k+1}, \hat{y}^{k+1}, x, y)] & \leq \mathbb{E}[V_k(x^k - x, y^k - y)] - \mathbb{E}[V_{k+1}(x^{k+1} - x, y^{k+1} - y)] \\ 317 \quad (4.12) \quad & \quad - \mathbb{E}[V(x^{k+1} - x^k, y^k - y^{k-1})]. \end{aligned}$$

318 *In particular, if $z^* = (x^*, y^*) \in \mathcal{Z}^*$ is a saddle point, then*

$$\begin{aligned} & \mathbb{E}[V_k(x^k - x^*, y^k - y^*)] \leq d_0, \quad \forall k \geq 0, \\ 319 \quad (4.13) \quad & \sum_{k=0}^{\infty} \mathbb{E}[V(x^{k+1} - x^k, y^k - y^{k-1})] \leq d_0, \end{aligned}$$

320 where $d_0 = \mathbb{E}[V_0(x^0 - x^*, y^0 - y^*)]$. Consequently,

$$321 \quad (4.14) \quad \mathbb{E}\|z^k - z^*\|_V^2 \leq 2\mathbb{E}\|z^0 - z^*\|_V^2, \quad \forall k \geq 0,$$

322 and

$$323 \quad (4.15) \quad \mathbb{E}\left[\sum_{k=0}^{\infty} \|z^{k+1} - z^k\|_V^2\right] \leq \frac{2}{1 - 2\gamma_1^2} \mathbb{E}\|z^0 - z^*\|_V^2.$$

324 *Proof.* Using (3.6), (4.5), we derive (4.12). By the definition of saddle point, we
 325 have $H(\hat{x}^{k+1}, \hat{y}^{k+1}, x^*, y^*) \geq 0$. Therefore,

$$\begin{aligned} 326 \quad (4.16) \quad & 0 \leq \mathbb{E}[V_k(x^k - x^*, y^k - y^*)] - \mathbb{E}[V_{k+1}(x^{k+1} - x^*, y^{k+1} - y^*)] \\ & \quad - \mathbb{E}[V(x^{k+1} - x^k, y^k - y^{k-1})]. \end{aligned}$$

327 The remaining claims follow by combining (4.16) and (4.3), (4.4). $\square$

328 **4.2. A realized one-step bound.** To derive a rate for the actual stochastic
 329 iterates rather than the virtual full updates, we need one additional decomposition.
 330 Define

$$331 \quad g_{P^{-1}}(\bar{x}) := \sum_{j=1}^m \frac{1}{p_j} g_j(\bar{x}_j), \quad f_{Q^{-1}}^*(\bar{y}) := \sum_{i=1}^n \frac{1}{q_i} f_i^*(\bar{y}_i),$$

332 and

$$333 \quad g_{P^{-1}-I}(\bar{x}) := \sum_{j=1}^m \left(\frac{1}{p_j} - 1 \right) g_j(\bar{x}_j), \quad f_{Q^{-1}-I}^*(\bar{y}) := \sum_{i=1}^n \left(\frac{1}{q_i} - 1 \right) f_i^*(\bar{y}_i).$$

334 For $z = (x, y)$, define

$$335 \quad (4.17) \quad D_g^{P^{-1}-I}(\bar{x}; z) := g_{P^{-1}-I}(\bar{x}) - g_{P^{-1}-I}(x) + \langle A(P^{-1} - I)(\bar{x} - x), y \rangle,$$

$$336 \quad (4.18) \quad D_{f^*}^{Q^{-1}-I}(\bar{y}; z) := f_{Q^{-1}-I}^*(\bar{y}) - f_{Q^{-1}-I}^*(y) - \langle Ax, (Q^{-1} - I)(\bar{y} - y) \rangle.$$

337 We also define the stochastic residuals

$$338 \quad (4.19) \quad u^k := \hat{x}^{k+1} - x^k - P^{-1}(x^{k+1} - x^k), \quad v^k := \hat{y}^{k+1} - y^k - Q^{-1}(y^{k+1} - y^k).$$

339 **LEMMA 4.3.** *Suppose Assumption 2.1 and 3.3 hold. For all $k \geq 0$ and $(x, y) \in \mathcal{Z}$,*
 340 *we have*

$$\begin{aligned} 341 \quad H(x^{k+1}, y^{k+1}, x, y) &\leq D_g^{P^{-1}-I}(x^k; z) - D_g^{P^{-1}-I}(x^{k+1}; z) \\ 342 &\quad + D_{f^*}^{Q^{-1}-I}(y^k; z) - D_{f^*}^{Q^{-1}-I}(y^{k+1}; z) \\ 343 &\quad + \langle x, u^k \rangle_{\tau^{-1}} + \langle y, v^k \rangle_{\sigma^{-1}} + \varepsilon_k \\ 344 &\quad + V_k(x^k - x, y^k - y) - V_{k+1}(x^{k+1} - x, y^{k+1} - y) \\ 345 \quad (4.20) &\quad - V(x^{k+1} - x^k, y^k - y^{k+1}). \end{aligned}$$

346 Moreover, $\mathbb{E}[\varepsilon_k] = 0$.

347 *Proof.* We first compare the virtual and realized gap kernels. By definition,

$$\begin{aligned} 348 \quad &H(\hat{x}^{k+1}, \hat{y}^{k+1}, x, y) - H(x^{k+1}, y^{k+1}, x, y) \\ 349 &= g(\hat{x}^{k+1}) - g(x^{k+1}) + \langle A(\hat{x}^{k+1} - x^{k+1}), y \rangle \\ 350 \quad (4.21) &\quad + f^*(\hat{y}^{k+1}) - f^*(y^{k+1}) + \langle Ax, y^{k+1} - \hat{y}^{k+1} \rangle. \end{aligned}$$

351 Substituting (4.21) into the virtual one-step inequality (3.2), and rearranging the
 352 separable terms, gives

$$\begin{aligned} &H(x^{k+1}, y^{k+1}, x, y) \leq g(x^{k+1}) - g(\hat{x}^{k+1}) + \langle A(x^{k+1} - \hat{x}^{k+1}), y \rangle \\ &\quad + f^*(y^{k+1}) - f^*(\hat{y}^{k+1}) + \langle Ax, \hat{y}^{k+1} - y^{k+1} \rangle \\ &\quad + \langle A\hat{x}^{k+1}, y \rangle - \langle Ax, \hat{y}^{k+1} \rangle \\ 353 \quad (4.22) &\quad + \langle A(x - \hat{x}^{k+1}), \bar{y}^k \rangle + \langle A\bar{x}^{k+1}, \hat{y}^{k+1} - y \rangle \\ &\quad + \frac{1}{2} \|x^k - x\|_{\tau^{-1}}^2 - \frac{1}{2} \|\hat{x}^{k+1} - x^k\|_{\tau^{-1}}^2 - \frac{1}{2} \|\hat{x}^{k+1} - x\|_{\tau^{-1}}^2 \\ &\quad + \frac{1}{2} \|y^k - y\|_{\sigma^{-1}}^2 - \frac{1}{2} \|\hat{y}^{k+1} - y^k\|_{\sigma^{-1}}^2 - \frac{1}{2} \|\hat{y}^{k+1} - y\|_{\sigma^{-1}}^2. \end{aligned}$$

354 For the first two rows in (4.22), we have the following equalities from direct
 355 computation.

$$\begin{aligned} &(4.23) \quad g(x^{k+1}) - g(\hat{x}^{k+1}) + \langle A(x^{k+1} - \hat{x}^{k+1}), y \rangle \\ &= D_g^{P^{-1}-I}(x^k; z) - D_g^{P^{-1}-I}(x^{k+1}; z) + R_g^k + \langle x^k - \hat{x}^{k+1} - P^{-1}(x^k - x^{k+1}), A^\top y \rangle, \\ 356 \quad &\quad f^*(y^{k+1}) - f^*(\hat{y}^{k+1}) + \langle Ax, \hat{y}^{k+1} - y^{k+1} \rangle \\ &= D_{f^*}^{Q^{-1}-I}(y^k; z) - D_{f^*}^{Q^{-1}-I}(y^{k+1}; z) + R_{f^*}^k + \langle Ax, \hat{y}^{k+1} - y^k - Q^{-1}(y^{k+1} - y^k) \rangle, \end{aligned}$$

where

$$\begin{aligned} R_g^k &:= g(x^k) - g(\hat{x}^{k+1}) + g_{P^{-1}}(x^{k+1}) - g_{P^{-1}}(x^k), \\ R_{f^*}^k &:= f^*(y^k) - f^*(\hat{y}^{k+1}) + f_{Q^{-1}}^*(y^{k+1}) - f_{Q^{-1}}^*(y^k). \end{aligned}$$

For the quadratic terms in (4.22), we have

$$(4.24) \quad \hat{\mathcal{Q}}_k = \mathcal{Q}_k + \hat{\mathcal{Q}}_k - \mathcal{Q}_k$$

where $\hat{\mathcal{Q}}_k, \mathcal{Q}_k$ are defined in (3.9). Letting $\Delta\mathcal{Q}_k = \hat{\mathcal{Q}}_k - \mathcal{Q}_k$, and using the identity

$$\|a - c\|_M^2 - \|b - c\|_M^2 = \|a\|_M^2 - \|b\|_M^2 + 2\langle c, b - a \rangle_M,$$

we obtain

$$(4.25) \quad \Delta\mathcal{Q}_k = \langle x, u^k \rangle_{\tau^{-1}} + \langle y, v^k \rangle_{\sigma^{-1}} + \delta_k,$$

where the zero-mean quadratic residual is

$$\begin{aligned} \delta_k &:= \frac{1}{2} (\|x^{k+1} - x^k\|_{\tau^{-1}P^{-1}}^2 - \|\hat{x}^{k+1} - x^k\|_{\tau^{-1}}^2) \\ &\quad + \frac{1}{2} (\|y^{k+1} - y^k\|_{\sigma^{-1}Q^{-1}}^2 - \|\hat{y}^{k+1} - y^k\|_{\sigma^{-1}}^2) \\ &\quad + \frac{1}{2} (\|x^{k+1}\|_{\tau^{-1}P^{-1}}^2 - \|x^k\|_{\tau^{-1}P^{-1}}^2 + \|x^k\|_{\tau^{-1}}^2 - \|\hat{x}^{k+1}\|_{\tau^{-1}}^2) \\ &\quad + \frac{1}{2} (\|y^{k+1}\|_{\sigma^{-1}Q^{-1}}^2 - \|y^k\|_{\sigma^{-1}Q^{-1}}^2 + \|y^k\|_{\sigma^{-1}}^2 - \|\hat{y}^{k+1}\|_{\sigma^{-1}}^2). \end{aligned}$$

Combining (4.22) (4.23) (4.24) and (4.25), we have

$$\begin{aligned} H(x^{k+1}, y^{k+1}, x, y) &\leq D_g^{P^{-1}-I}(x^k; z) - D_g^{P^{-1}-I}(x^{k+1}; z) \\ &\quad + D_{f^*}^{Q^{-1}-I}(y^k; z) - D_{f^*}^{Q^{-1}-I}(y^{k+1}; z) \\ &\quad + R_g^k + R_{f^*}^k + \mathcal{B}_k \\ &\quad + \mathcal{Q}_k + \langle x, u^k \rangle_{\tau^{-1}} + \langle y, v^k \rangle_{\sigma^{-1}} + \delta_k, \end{aligned}$$

where $\mathcal{B}_k$ denotes the remaining bilinear part, which is

$$\begin{aligned} \mathcal{B}_k &= \langle A\hat{x}^{k+1}, y \rangle - \langle Ax, \hat{y}^{k+1} \rangle + \langle A(x - \hat{x}^{k+1}), \bar{y}^k \rangle + \langle A\bar{x}^{k+1}, \hat{y}^{k+1} - y \rangle \\ &\quad + \langle x^k - \hat{x}^{k+1} - P^{-1}(x^k - x^{k+1}), A^\top y \rangle + \langle Ax, \hat{y}^{k+1} - y^k - Q^{-1}(y^{k+1} - y^k) \rangle \end{aligned}$$

Using

$$\bar{x}^{k+1} = x^k + P^{-1}(x^{k+1} - x^k), \quad \bar{y}^k = y^k + Q^{-1}(y^k - y^{k-1}),$$

a direct rearrangement of the bilinear terms gives

$$\begin{aligned} \mathcal{B}_k &= \langle A(\bar{x}^{k+1} - x), Q^{-1}(y^{k+1} + y^{k-1} - 2y^k) \rangle \\ &\quad + \langle A\bar{x}^{k+1}, v^k \rangle - \langle Au^k, \bar{y}^k \rangle. \end{aligned}$$

Combining (4.27) and (4.29), and using the definition of Γ_k , we obtain

$$\begin{aligned} H(x^{k+1}, y^{k+1}, x, y) &\leq D_g^{P^{-1}-I}(x^k; z) - D_g^{P^{-1}-I}(x^{k+1}; z) \\ &\quad + D_{f^*}^{Q^{-1}-I}(y^k; z) - D_{f^*}^{Q^{-1}-I}(y^{k+1}; z) \\ &\quad + \langle x, u^k \rangle_{\tau^{-1}} + \langle y, v^k \rangle_{\sigma^{-1}} + \varepsilon_k + \Gamma_k, \end{aligned}$$

where

$$(4.31) \quad \varepsilon_k := R_g^k + R_{f^*}^k + \delta_k + \langle A\bar{x}^{k+1}, v^k \rangle - \langle Au^k, \bar{y}^k \rangle.$$

It remains to verify that ε_k has zero expectation. This follows from the block update rules. Indeed, for the primal part of ε_k , we have

$$\mathbb{E}[u^k \mid \mathcal{F}_k] = 0, \quad \mathbb{E}[R_g^k \mid \mathcal{F}_k] = 0, \quad \mathbb{E}[\delta_k^x \mid \mathcal{F}_k] = 0.$$

For the dual part of ε_k , we have

$$\mathbb{E}[v^k \mid \mathcal{F}_{k+\frac{1}{2}}] = 0, \quad \mathbb{E}[R_{f^*}^k \mid \mathcal{F}_{k+\frac{1}{2}}] = 0, \quad \mathbb{E}[\delta_k^y \mid \mathcal{F}_{k+\frac{1}{2}}] = 0.$$

Here δ_k^x and δ_k^y are the x - and y -parts of δ_k , respectively. Since $\bar{y}^k$ is $\mathcal{F}_k$ -measurable and $\bar{x}^{k+1}$ is $\mathcal{F}_{k+\frac{1}{2}}$ -measurable, we also have

$$\mathbb{E}\langle Au^k, \bar{y}^k \rangle = 0, \quad \mathbb{E}\langle A\bar{x}^{k+1}, v^k \rangle = 0.$$

Therefore $\mathbb{E}[\varepsilon_k] = 0$.

Finally, combining (4.30) with (4.5) gives (4.20). $\square$

To control the accumulated correction terms in Lemma 4.3, we need to estimate $D_g^{P^{-1}-I}$ and $D_{f^*}^{Q^{-1}-I}$. These terms do not admit a direct descent structure and therefore cannot be handled by the Lyapunov argument alone.

The idea is to relate them to the primal–dual gap and to distances to the solution set using saddle-point optimality and convexity. To decouple these two effects, we allow two (possibly different) saddle points, leading to the following estimate.

LEMMA 4.4. *Suppose Assumption 2.1 holds. Let $z_1^* = (x_1^*, y_1^*)$ and $z_2^* = (x_2^*, y_2^*)$ be two saddle points. Define*

$$\underline{p} := \min_{j \in [m]} p_j, \quad \underline{q} := \min_{i \in [n]} q_i,$$

and

$$\begin{aligned} \gamma_4^2 &:= \left\| \tau^{1/2} (P^{-1/2} - P^{1/2}) A^\top Q^{1/2} \sigma^{1/2} \right\| \\ \gamma_5^2 &:= \left\| \tau^{1/2} P^{1/2} A^\top (Q^{-1/2} - Q^{1/2}) \sigma^{1/2} \right\| \end{aligned}$$

Then, for any positive integer K and any $z = (x, y) \in \mathcal{Z}$,

$$\begin{aligned} D_g^{P^{-1}-I}(x^0; z) - D_g^{P^{-1}-I}(x^K; z) &\leq \left(\frac{1}{\underline{p}} - 1 \right) (\Phi(x^0, y_1^*) - \Phi(x_1^*, y_1^*)) \\ &+ \frac{\gamma_4^2}{2} \left(2\|x^K - x_2^*\|_{\tau^{-1}P^{-1}}^2 + 2\|x^0 - x_2^*\|_{\tau^{-1}P^{-1}}^2 + \|y - y_1^*\|_{\sigma^{-1}Q^{-1}}^2 \right), \end{aligned}$$

and

$$\begin{aligned} D_{f^*}^{Q^{-1}-I}(y^0; z) - D_{f^*}^{Q^{-1}-I}(y^K; z) &\leq \left(\frac{1}{\underline{q}} - 1 \right) (\Phi(x_1^*, y_1^*) - \Phi(x_1^*, y^0)) \\ &+ \frac{\gamma_5^2}{2} \left(\|x - x_1^*\|_{\tau^{-1}P^{-1}}^2 + 2\|y^K - y_2^*\|_{\sigma^{-1}Q^{-1}}^2 + 2\|y^0 - y_2^*\|_{\sigma^{-1}Q^{-1}}^2 \right). \end{aligned}$$

413 *Consequently,*

$$414 \quad D_g^{P^{-1}-I}(x^0; z) - D_g^{P^{-1}-I}(x^K; z) + D_{f^*}^{Q^{-1}-I}(y^0; z) - D_{f^*}^{Q^{-1}-I}(y^K; z) \\ 415 \quad (4.34) \quad \leq \alpha H(x^0, y^0, x_1^*, y_1^*) + \beta \|z^K - z_2^*\|_V^2 + \beta \|z^0 - z_2^*\|_V^2 + \frac{\beta}{2} \|z - z_1^*\|_V^2,$$

416 *where*

$$417 \quad \alpha := \max \left\{ \frac{1}{\underline{p}} - 1, \frac{1}{\underline{q}} - 1 \right\}, \quad \beta := \max\{\gamma_4^2, \gamma_5^2\}.$$

418 *Proof.* We first prove the bound for the primal correction term. By definition,

$$419 \quad (4.35) \quad D_g^{P^{-1}-I}(x^0; z) - D_g^{P^{-1}-I}(x^K; z) = g_{P^{-1}-I}(x^0) - g_{P^{-1}-I}(x^K) \\ + \langle A(P^{-1} - I)(x^0 - x^K), y \rangle.$$

420 Since z_1^* is a saddle point, we have

$$421 \quad -(A^j)^\top y_1^* \in \partial g_j((x_1^*)_j), \quad j \in [m].$$

422 Hence, by convexity of g_j ,

$$423 \quad g_j(x_j^K) \geq g_j((x_1^*)_j) - \langle (A^j)^\top y_1^*, x_j^K - (x_1^*)_j \rangle.$$

424 Multiplying by $p_j^{-1} - 1$ and summing over j gives

$$425 \quad g_{P^{-1}-I}(x^K) \geq g_{P^{-1}-I}(x_1^*) - \langle A(P^{-1} - I)(x^K - x_1^*), y_1^* \rangle.$$

426 Substituting this inequality into (4.35), we obtain

$$427 \quad D_g^{P^{-1}-I}(x^0; z) - D_g^{P^{-1}-I}(x^K; z) \\ 428 \quad \leq g_{P^{-1}-I}(x^0) - g_{P^{-1}-I}(x_1^*) + \langle A(P^{-1} - I)(x^0 - x_1^*), y_1^* \rangle \\ 429 \quad (4.36) \quad + \langle A(P^{-1} - I)(x^0 - x^K), y - y_1^* \rangle.$$

430 For the first three terms on the right-hand side, using again the saddle-point subgr-
431 dient condition gives

$$432 \quad g_{P^{-1}-I}(x^0) - g_{P^{-1}-I}(x_1^*) + \langle A(P^{-1} - I)(x^0 - x_1^*), y_1^* \rangle \\ 433 \quad = \sum_{j=1}^m \left(\frac{1}{p_j} - 1 \right) [g_j(x_j^0) - g_j((x_1^*)_j) + \langle A^j(x_j^0 - (x_1^*)_j), y_1^* \rangle] \\ 434 \quad \leq \left(\frac{1}{\underline{p}} - 1 \right) [g(x^0) - g(x_1^*) + \langle A(x^0 - x_1^*), y_1^* \rangle] \\ 435 \quad (4.37) \quad = \left(\frac{1}{\underline{p}} - 1 \right) (\Phi(x^0, y_1^*) - \Phi(x_1^*, y_1^*)).$$

436 For the remaining cross term, by the definition of γ_4^2 ,

$$437 \quad |\langle A(P^{-1} - I)(x^0 - x^K), y - y_1^* \rangle| \leq \frac{\gamma_4^2}{2} \left(\|x^0 - x^K\|_{\tau^{-1}P^{-1}}^2 + \|y - y_1^*\|_{\sigma^{-1}Q^{-1}}^2 \right) \\ 438 \quad (4.38) \quad \leq \frac{\gamma_4^2}{2} \left(2\|x^K - x_2^*\|_{\tau^{-1}P^{-1}}^2 + 2\|x^0 - x_2^*\|_{\tau^{-1}P^{-1}}^2 + \|y - y_1^*\|_{\sigma^{-1}Q^{-1}}^2 \right).$$

Combining (4.36), (4.37), and (4.38) proves (4.32).

The proof of (4.33) is analogous. Indeed, since z_1^* is a saddle point,

$$A_i x_1^* \in \partial f_i^*((y_1^*)_i), \quad i \in [n],$$

and repeating the same argument with f_i^* , Q , and the operator norm constant γ_5^2 yields (4.33).

It remains to combine the two estimates. By the definition of the gap kernel,

$$H(x^0, y^0, x_1^*, y_1^*) = \Phi(x^0, y_1^*) - \Phi(x_1^*, y^0),$$

and therefore

$$H(x^0, y^0, x_1^*, y_1^*) = [\Phi(x^0, y_1^*) - \Phi(x_1^*, y_1^*)] + [\Phi(x_1^*, y_1^*) - \Phi(x_1^*, y^0)].$$

Summing (4.32) and (4.33), and using the definitions of α and β , yields (4.34). $\square$

To control the random linear terms in Lemma 4.3, we introduce an auxiliary sequence driven by the stochastic errors u^k and v^k . This allows us to bound their accumulated contribution uniformly over a bounded test region B .

LEMMA 4.5. *Suppose Assumptions 2.1 and 3.3 hold. Let $z^* \in \mathcal{Z}^*$ be a saddle point. Given $\tilde{z}^0 = (\tilde{x}^0, \tilde{y}^0)$, define*

$$\tilde{x}^{k+1} = \tilde{x}^k + P u^k, \quad \tilde{y}^{k+1} = \tilde{y}^k + Q v^k, \quad k \geq 0.$$

Then, for any $K \geq 1$, any $z = (x, y) \in \mathcal{Z}$, and any set $B \subseteq \mathcal{Z}$, we have

$$(4.39) \quad \sum_{k=0}^{K-1} \langle x, u^k \rangle_{\tau^{-1}} \leq \frac{1}{2} \|\tilde{x}^0 - x\|_{\tau^{-1}P}^2 + \frac{1}{2} \sum_{k=0}^{K-1} \|u^k\|_{\tau^{-1}P}^2 + \sum_{k=0}^{K-1} \langle \tilde{x}^k, u^k \rangle_{\tau^{-1}},$$

$$(4.40) \quad \sum_{k=0}^{K-1} \langle y, v^k \rangle_{\sigma^{-1}} \leq \frac{1}{2} \|\tilde{y}^0 - y\|_{\sigma^{-1}Q}^2 + \frac{1}{2} \sum_{k=0}^{K-1} \|v^k\|_{\sigma^{-1}Q}^2 + \sum_{k=0}^{K-1} \langle \tilde{y}^k, v^k \rangle_{\sigma^{-1}}.$$

Moreover,

$$(4.41) \quad \mathbb{E} \left[\sum_{k=0}^{\infty} \left(\|u^k\|_{\tau^{-1}P}^2 + \|v^k\|_{\sigma^{-1}Q}^2 \right) \right] \leq \frac{2}{1 - 2\gamma_1^2} \mathbb{E} \|z^0 - z^*\|_V^2.$$

Consequently,

$$(4.42) \quad \mathbb{E} \left[\sup_{z \in B} \left\{ \sum_{k=0}^{K-1} \langle x, u^k \rangle_{\tau^{-1}} + \langle y, v^k \rangle_{\sigma^{-1}} \right\} \right] \leq \mathbb{E} \sup_{z \in B} \left\{ \frac{1}{2} \|\tilde{x}^0 - x\|_{\tau^{-1}P}^2 + \frac{1}{2} \|\tilde{y}^0 - y\|_{\sigma^{-1}Q}^2 \right\} + \frac{1}{1 - 2\gamma_1^2} \mathbb{E} \|z^0 - z^*\|_V^2.$$

Proof. We first prove (4.39). By the definition of $\tilde{x}^{k+1}$,

$$\frac{1}{2} \|\tilde{x}^{k+1} - x\|_{\tau^{-1}P}^2 = \frac{1}{2} \|\tilde{x}^k - x\|_{\tau^{-1}P}^2 + \langle u^k, \tilde{x}^k - x \rangle_{\tau^{-1}} + \frac{1}{2} \|u^k\|_{\tau^{-1}P}^2.$$

Rearranging and summing over $k = 0, \dots, K-1$ gives

$$\sum_{k=0}^{K-1} \langle x - \hat{x}^k, u^k \rangle_{\tau^{-1}} \leq \frac{1}{2} \|\hat{x}^0 - x\|_{\tau^{-1}P}^2 + \frac{1}{2} \sum_{k=0}^{K-1} \|u^k\|_{\tau^{-1}P}^2,$$

which implies (4.39). The proof of (4.40) is identical.

Next, by the definition of u^k ,

$$u^k = \hat{x}^{k+1} - x^k - P^{-1}(x^{k+1} - x^k).$$

we have

$$\begin{aligned} \|u^k\|_{\tau^{-1}P}^2 &= \|x^k - \hat{x}^{k+1}\|_{\tau^{-1}P}^2 + \|x^{k+1} - x^k\|_{\tau^{-1}P}^2 \\ &\quad - 2\langle x^k - \hat{x}^{k+1}, x^k - x^{k+1} \rangle_{\tau^{-1}}. \end{aligned} \quad (4.43)$$

Since $x^k, \hat{x}^{k+1}$ are $\mathcal{F}_k$ measurable and $\mathbb{E}[x^k - x^{k+1} \mid \mathcal{F}_k] = P(x^k - \hat{x}^{k+1})$, we have

$$\begin{aligned} \mathbb{E}[\langle x^k - \hat{x}^{k+1}, x^k - x^{k+1} \rangle_{\tau^{-1}} \mid \mathcal{F}_k] &= \langle x^k - \hat{x}^{k+1}, P(x^k - \hat{x}^{k+1}) \rangle_{\tau^{-1}} \\ &= \|x^k - \hat{x}^{k+1}\|_{\tau^{-1}P}^2. \end{aligned} \quad (4.44)$$

Therefore,

$$\begin{aligned} \mathbb{E}[\|u^k\|_{\tau^{-1}P}^2] &= \mathbb{E}[\|x^{k+1} - x^k\|_{\tau^{-1}P}^2] - \mathbb{E}[\|x^k - \hat{x}^{k+1}\|_{\tau^{-1}P}^2] \\ &\leq \mathbb{E}[\|x^{k+1} - x^k\|_{\tau^{-1}P}^2]. \end{aligned} \quad (4.45)$$

Similarly,

$$\begin{aligned} \mathbb{E}[\|v^k\|_{\sigma^{-1}Q}^2] &= \mathbb{E}[\|y^{k+1} - y^k\|_{\sigma^{-1}Q}^2] - \mathbb{E}[\|y^k - \hat{y}^{k+1}\|_{\sigma^{-1}Q}^2] \\ &\leq \mathbb{E}[\|y^{k+1} - y^k\|_{\sigma^{-1}Q}^2]. \end{aligned} \quad (4.46)$$

Combining (4.45) and (4.46), we have

$$\begin{aligned} \mathbb{E} \left[\sum_{k=0}^{\infty} (\|u^k\|_{\tau^{-1}P}^2 + \|v^k\|_{\sigma^{-1}Q}^2) \right] &\leq \mathbb{E} \left[\sum_{k=0}^{\infty} \|z^{k+1} - z^k\|_V^2 \right] \leq \frac{2}{1 - 2\gamma_1^2} \mathbb{E} \|z^0 - z^*\|_V^2, \end{aligned} \quad (4.47)$$

where the last inequality comes from (4.15).

Finally, summing (4.39) and (4.40) and taking the supremum over $z \in B$, we obtain

$$\begin{aligned} &\sup_{z \in B} \left\{ \sum_{k=0}^{K-1} (\langle x, u^k \rangle_{\tau^{-1}} + \langle y, v^k \rangle_{\sigma^{-1}}) \right\} \\ &\leq \sup_{z \in B} \left\{ \frac{1}{2} \|\hat{x}^0 - x\|_{\tau^{-1}P}^2 + \frac{1}{2} \|\hat{y}^0 - y\|_{\sigma^{-1}Q}^2 \right\} \\ &\quad + \frac{1}{2} \sum_{k=0}^{K-1} (\|u^k\|_{\tau^{-1}P}^2 + \|v^k\|_{\sigma^{-1}Q}^2) + M_K, \end{aligned} \quad (4.48)$$

where

$$M_K := \sum_{k=0}^{K-1} (\langle \hat{x}^k, u^k \rangle_{\tau^{-1}} + \langle \hat{y}^k, v^k \rangle_{\sigma^{-1}}).$$

Since $\hat{x}^k$ and $\hat{y}^k$ are measurable with respect to $\mathcal{F}_k$ and u^k, v^k are conditionally mean-zero with respect to $\mathcal{F}_k$ and $\mathcal{F}_{k+\frac{1}{2}}$ respectively, $\mathbb{E}[M_K] = 0$. Taking expectation on (4.48) and using (4.41) gives (4.42). $\square$

4.3. Ergodic $\mathcal{O}(1/K)$ rate. Define the ergodic averages

$$(4.49) \quad X^K := \frac{1}{K} \sum_{k=1}^K x^k, \quad Y^K := \frac{1}{K} \sum_{k=1}^K y^k.$$

For a bounded set $B \subseteq \mathcal{Z}$, define the restricted primal–dual gap

$$(4.50) \quad G_B(\bar{x}, \bar{y}) := \sup_{z=(x,y) \in B} H(\bar{x}, \bar{y}, x, y).$$

The next theorem is the main result of this section.

THEOREM 4.6. *Suppose Assumptions 2.1 and 3.3 hold. Let $B \subseteq \mathcal{Z}$ be bounded. Then there exists a constant $C_B > 0$, depending only on the initial point, the saddle-point set, the step sizes, and B , such that*

$$(4.51) \quad \mathbb{E}[G_B(X^K, Y^K)] \leq \frac{C_B}{K}, \quad \forall K \geq 1.$$

Hence the ergodic sequence generated by DSPDHG converges at rate $\mathcal{O}(1/K)$ in the expected restricted primal–dual gap.

Proof. Fix $z = (x, y) \in B$. Since $H(\cdot, \cdot, x, y)$ is convex in its first two arguments, Jensen’s inequality gives

$$(4.52) \quad H(X^K, Y^K, x, y) \leq \frac{1}{K} \sum_{k=0}^{K-1} H(x^{k+1}, y^{k+1}, x, y).$$

Therefore,

$$(4.52) \quad K G_B(X^K, Y^K) \leq \sup_{z \in B} \sum_{k=0}^{K-1} H(x^{k+1}, y^{k+1}, x, y).$$

Summing the realized one-step inequality (4.20) from $k = 0$ to $K - 1$, using (4.52), and dropping the nonnegative terms

$$(4.53) \quad V_K(x^K - x, y^K - y), \quad \sum_{k=0}^{K-1} V(x^{k+1} - x^k, y^k - y^{k-1}),$$

we obtain

$$(4.53) \quad \begin{aligned} K \mathbb{E}[G_B(X^K, Y^K)] &\leq \mathbb{E} \left[\sup_{z \in B} \left\{ D_g^{P^{-1}-I}(x^0; z) - D_g^{P^{-1}-I}(x^K; z) \right. \right. \\ &\quad \left. \left. + D_{f^*}^{Q^{-1}-I}(y^0; z) - D_{f^*}^{Q^{-1}-I}(y^K; z) \right\} \right] \\ &\quad + \mathbb{E} \left[\sup_{z \in B} \left\{ \sum_{k=0}^{K-1} (\langle x, u^k \rangle_{\tau^{-1}} + \langle y, v^k \rangle_{\sigma^{-1}}) \right\} \right] \\ &\quad + \mathbb{E} \left[\sup_{z \in B} V_0(x^0 - x, y^0 - y) \right]. \end{aligned}$$

Here the martingale residuals ε_k vanish after taking expectation.

Since $y^{-1} = y^0$, the definition of V_0 gives

$$V_0(x^0 - x, y^0 - y) = \frac{1}{2} \|z - z^0\|_V^2.$$

Next, applying the weighted correction bound (4.34) with $z_1^* = z_2^* = z^* \in \mathcal{Z}^*$ yields

$$\begin{aligned} & \mathbb{E} \left[\sup_{z \in B} \left\{ D_g^{P^{-1}-I}(x^0; z) - D_g^{P^{-1}-I}(x^K; z) + D_{f^*}^{Q^{-1}-I}(y^0; z) - D_{f^*}^{Q^{-1}-I}(y^K; z) \right\} \right] \\ & \leq \alpha \mathbb{E}[H(x^0, y^0, x^*, y^*)] + \beta \mathbb{E}\|z^K - z^*\|_V^2 + \beta \mathbb{E}\|z^0 - z^*\|_V^2 + \frac{\beta}{2} \mathbb{E} \left[\sup_{z \in B} \|z - z^*\|_V^2 \right]. \end{aligned}$$

Moreover, by (4.42) with $\tilde{x}^0 = x^0$ and $\tilde{y}^0 = y^0$,

$$\begin{aligned} & \mathbb{E} \left[\sup_{z \in B} \left\{ \sum_{k=0}^{K-1} (\langle x, u^k \rangle_{\tau^{-1}} + \langle y, v^k \rangle_{\sigma^{-1}}) \right\} \right] \\ & \leq \frac{1}{2} \mathbb{E} \left[\sup_{z \in B} \|z - z^0\|_V^2 \right] + \frac{1}{1 - 2\gamma_1^2} \mathbb{E}\|z^0 - z^*\|_V^2. \end{aligned}$$

Combining (4.53), (4.54), and (4.55), and using

$$\mathbb{E}\|z^K - z^*\|_V^2 \leq 2 \mathbb{E}\|z^0 - z^*\|_V^2,$$

we obtain

$$\begin{aligned} K \mathbb{E}[G_B(X^K, Y^K)] & \leq \alpha \mathbb{E}[H(x^0, y^0, x^*, y^*)] + \left(3\beta + \frac{1}{1 - 2\gamma_1^2} \right) \mathbb{E}\|z^0 - z^*\|_V^2 \\ & + \frac{\beta}{2} \mathbb{E} \left[\sup_{z \in B} \|z - z^*\|_V^2 \right] + \mathbb{E} \left[\sup_{z \in B} \|z - z^0\|_V^2 \right]. \end{aligned}$$

The right-hand side is finite because B is bounded. Defining this quantity as C_B and dividing both sides by K proves (4.51). $\square$

Remark 4.7. An explicit admissible constant in Theorem 4.6 is

$$\begin{aligned} C_B & := \alpha \mathbb{E}[H(x^0, y^0, x^*, y^*)] + \left(3\beta + \frac{1}{1 - 2\gamma_1^2} \right) \mathbb{E}\|z^0 - z^*\|_V^2 \\ & + \frac{\beta}{2} \mathbb{E} \left[\sup_{z \in B} \|z - z^*\|_V^2 \right] + \mathbb{E} \left[\sup_{z \in B} \|z - z^0\|_V^2 \right], \end{aligned}$$

where $z^* \in \mathcal{Z}^*$ is arbitrary.

Remark 4.8. Theorem 4.6 extends the classical $\mathcal{O}(1/K)$ expected restricted-gap guarantee for SPDHG to the doubly stochastic setting. The main additional difficulty is that stochasticity appears on both primal and dual sides, which requires controlling two correction sequences, u^k and v^k , instead of a single one-sided correction term.

5. Linear Convergence under Smoothed Quadratic Growth. We now show that a restarted variant of DSPDHG achieves linear convergence under a regularity condition formulated in terms of a smoothed primal–dual gap. Unlike strong convexity–concavity assumptions imposed on specific components of the objective, the condition below is expressed directly in terms of the saddle-point geometry, and is therefore more flexible.

5.1. Smoothed quadratic growth. Recall the smoothed gap functional

$$G_\mu(\bar{z}, \dot{z}) := \sup_{z \in \mathcal{Z}} \left\{ H(\bar{x}, \bar{y}, x, y) - \frac{\mu}{2} \|z - \dot{z}\|_V^2 \right\}, \quad \bar{z} = (\bar{x}, \bar{y}), \dot{z} \in \mathcal{Z},$$

introduced in (2.8). The notion of smoothed quadratic growth with respect to G_μ was introduced in [12] and has since been used in several related works.

DEFINITION 5.1 (Smoothed quadratic growth). *Suppose $\mu > 0$. We say that the smoothed duality gap G_μ satisfies quadratic growth on a set $\mathcal{R}$ with parameter ζ if, for every $\bar{z} \in \mathcal{R}$ and every $z^* \in \mathcal{Z}^*$,*

$$(5.1) \quad G_\mu(\bar{z}, z^*) \geq \zeta \operatorname{dist}_V(\bar{z}, \mathcal{Z}^*)^2.$$

This condition lower bounds the smoothed gap by the squared distance to the solution set in the weighted geometry induced by $\|\cdot\|_V$. It plays the role of an error bound condition and serves as the key ingredient for establishing linear convergence.

Remark 5.2. The smoothed quadratic growth condition holds in a variety of settings, including strongly convex–strongly concave problems [12], linear programs [12], quadratic programs (QPs) [22], and piecewise linear quadratic (PLQ) functions [12]. In particular, the two classes of convex problems considered in our numerical experiments, namely soft-margin SVM and model predictive control (MPC), fall into the PLQ and QP categories, respectively, and therefore satisfy this condition automatically.

Other regularity conditions have also been considered in the literature, such as metric subregularity (see [3, Assumption 2]) and the two-sided quadratic functional growth (QFG) condition (see [23, Definition 2.7]). For quadratic programming, [22] showed that smoothed quadratic growth can be implied by metric subregularity. However, whether such an implication holds more generally remains open. Moreover, to the best of our knowledge, there is currently no result clarifying whether smoothed quadratic growth and two-sided QFG imply each other.

5.2. Restarted scheme. We consider a restarted outer scheme built from the ergodic output of DSPDHG. Given an outer iterate $\mathbf{Z}^t \in \mathcal{Z}$, we run DSPDHG for K inner iterations initialized at $z^0 = \mathbf{Z}^t$, and define

$$(5.2) \quad \mathbf{Z}^{t+1} = \frac{1}{K} \sum_{k=1}^K z^k = \left(\frac{1}{K} \sum_{k=1}^K x^k, \frac{1}{K} \sum_{k=1}^K y^k \right).$$

Equivalently, if $\text{DSPDHG}(\mathbf{Z}^t, K)$ denotes the one-epoch ergodic output obtained by running K iterations of Algorithm 3.1 from the initial point $\mathbf{Z}^t$, then

$$\mathbf{Z}^{t+1} = \text{DSPDHG}(\mathbf{Z}^t, K).$$

Assumption 5.3. There exist constants $\mu > 0$ and $\zeta > 0$, together with a compact set $\mathcal{R} \subseteq \mathcal{Z}$, such that G_μ satisfies quadratic growth on $\mathcal{R}$ with parameter ζ . Moreover, the outer iterates satisfy $\mathbf{Z}^t \in \mathcal{R}$ for all $t \geq 0$.

Remark 5.4. The requirement $\mathbf{Z}^t \in \mathcal{R}$ for all $t \geq 0$ may appear somewhat restrictive, but it is needed for the linear convergence analysis. In the strongly convex case, where one may take $\mathcal{R} = \mathcal{Z}$, this condition is automatic. In other settings, however, such as quadratic programming [22], the set $\mathcal{R}$ must be chosen as a proper compact

Algorithm 5.1 Restarted DSPDHG

```

1: Input: initial point  $\mathbf{Z}^0 \in \mathcal{Z}$  and epoch length  $K$ 
2: for  $t = 0, 1, 2, \dots$  do
3:   Run Algorithm 3.1 for  $K$  iterations with  $z^0 = \mathbf{Z}^t$ 
4:   Set  $\mathbf{Z}^{t+1} := \frac{1}{K} \sum_{k=1}^K z^k$ 
5: end for

```

subset of $\mathcal{Z}$, and verifying that the iterates remain in $\mathcal{R}$ can be nontrivial. The difficulty comes from the fact that the trajectory is only controlled in expectation, so one cannot exclude low-probability excursions far away from $\mathcal{R}$. A similar assumption is also imposed in [3, Assumption 2] in the linear convergence analysis of SPDHG.

5.3. Linear convergence of the restarted method. We can now state the main theorem of this section.

THEOREM 5.5. *Suppose Assumptions 2.1, 3.3, and 5.3 hold. Let $\{\mathbf{Z}^t\}_{t \geq 0}$ be generated by Algorithm 5.1. Assume that the epoch length K satisfies*

$$(5.3) \quad K \geq \max \left\{ \frac{4 + 2\beta}{\mu}, \frac{100}{\zeta} \left(\alpha\zeta + 5\beta + 2 + \frac{1}{1 - 2\gamma_1^2} \right), \frac{400\beta}{\zeta} \right\}.$$

Then the restarted DSPDHG scheme converges linearly. More precisely, there exists a constant $M > 0$ such that

$$(5.4) \quad \mathbb{E}[G_\mu(\mathbf{Z}^t, \Pi^{t-1})] \leq M 5^{-t}, \quad \forall t \geq 1,$$

where $\Pi^{t-1} := \text{proj}_{\mathcal{Z}^*}(\mathbf{Z}^{t-1})$. Consequently,

$$(5.5) \quad \mathbb{E}[\text{dist}_V(\mathbf{Z}^t, \mathcal{Z}^*)^2] \leq \frac{M}{\zeta} 5^{-t}, \quad \forall t \geq 1.$$

Remark 5.6. Theorem 5.5 shows that DSPDHG achieves a geometric rate once the smoothed quadratic growth condition holds. The restart mechanism converts the ergodic $\mathcal{O}(1/K)$ behavior of the inner method into a linear decrease of the smoothed gap and, consequently, of the distance to the saddle-point set.

The constants α and β come from the one-epoch estimate and the correction bounds. They are not needed to run the inner DSPDHG iterations; they only enter the theoretical lower bound on the restart length K .

The rest of this section presents the proof of Theorem 5.5.

5.3.1. A one-epoch estimate. We next derive an estimate for one restart epoch. Throughout this subsection, we write

$$Z^K := (X^K, Y^K) := \frac{1}{K} \sum_{k=1}^K z^k, \quad z^k = (x^k, y^k),$$

where Z^K denotes the ergodic output of one epoch, while z^K denotes the last inner iterate.

Summing the realized one-step inequality (4.20) from $k = 0$ to $K-1$, and using the

convexity of $H(\cdot, \cdot, x, y)$ in its first two arguments, we have that for any $z = (x, y) \in \mathcal{Z}$,

$$\begin{aligned}
 & KH(X^K, Y^K, x, y) \\
 & \leq D_g^{P^{-1}-I}(x^0; z) - D_g^{P^{-1}-I}(x^K; z) + D_{f^*}^{Q^{-1}-I}(y^0; z) - D_{f^*}^{Q^{-1}-I}(y^K; z) \\
 & + \sum_{k=0}^{K-1} (\langle x, u^k \rangle_{\tau-1} + \langle y, v^k \rangle_{\sigma-1} + \varepsilon_k) + V_0(x^0 - x, y^0 - y).
 \end{aligned}
 \tag{5.6}$$

This inequality is the starting point for the linear convergence analysis.

The following lemma gives a one-epoch estimate for the smoothed gap.

LEMMA 5.7. *Suppose Assumptions 2.1 and 3.3 hold. Then, for any $z_1^*, z_2^* \in \mathcal{Z}^*$ and any K satisfying $\mu K \geq 2\beta + 4$,*

$$\begin{aligned}
 \mathbb{E}[G_\mu(Z^K, z_2^*)] & \leq \frac{\alpha}{K} \mathbb{E}[G_\mu(z^0, z_1^*)] + \frac{\beta}{K} \mathbb{E}[\|z_1^* - z_2^*\|_V^2] \\
 & + \frac{1}{K} \left(3\beta + 2 + \frac{1}{1 - 2\gamma_1^2} \right) \mathbb{E}[\|z^0 - z_2^*\|_V^2].
 \end{aligned}
 \tag{5.7}$$

Proof. Starting from (5.6), we use (4.34) with two arbitrary saddle points z_1^*, z_2^* and (4.39), (4.40) with the auxiliary initialization $\tilde{x}^0 = x^0$ and $\tilde{y}^0 = y^0$. This gives, for every $z = (x, y) \in \mathcal{Z}$,

$$\begin{aligned}
 & KH(X^K, Y^K, x, y) \\
 & \leq \alpha H(x^0, y^0, x_1^*, y_1^*) + \beta \|z^K - z_2^*\|_V^2 + \beta \|z^0 - z_2^*\|_V^2 + \frac{\beta}{2} \|z - z_1^*\|_V^2 + \|z - z^0\|_V^2 \\
 & + \frac{1}{2} \sum_{k=0}^{K-1} (\|u^k\|_{\tau-1P}^2 + \|v^k\|_{\sigma-1Q}^2) + \sum_{k=0}^{K-1} (\langle \tilde{x}^k, u^k \rangle_{\tau-1} + \langle \tilde{y}^k, v^k \rangle_{\sigma-1} + \varepsilon_k).
 \end{aligned}
 \tag{5.8}$$

The expectation of the last summation is zero by $\mathbb{E}[M_K] = 0$ and $\mathbb{E}[\varepsilon_k] = 0$. Dividing (5.8) by K and subtracting $\frac{\mu}{2} \|z - z_2^*\|_V^2$ from both sides yields

$$\begin{aligned}
 & H(X^K, Y^K, x, y) - \frac{\mu}{2} \|z - z_2^*\|_V^2 \\
 & \leq \frac{\alpha}{K} H(x^0, y^0, x_1^*, y_1^*) + \frac{\beta}{K} \|z^K - z_2^*\|_V^2 + \frac{\beta}{K} \|z^0 - z_2^*\|_V^2 + \frac{\beta}{2K} \|z - z_1^*\|_V^2 \\
 & + \frac{1}{K} \|z - z^0\|_V^2 - \frac{\mu}{2} \|z - z_2^*\|_V^2 + \frac{1}{2K} \sum_{k=0}^{K-1} (\|u^k\|_{\tau-1P}^2 + \|v^k\|_{\sigma-1Q}^2) \\
 & + \frac{1}{K} \sum_{k=0}^{K-1} (\langle \tilde{x}^k, u^k \rangle_{\tau-1} + \langle \tilde{y}^k, v^k \rangle_{\sigma-1} + \varepsilon_k).
 \end{aligned}
 \tag{5.9}$$

Since $G_\mu(z^0, z_1^*) = \sup_z \{H(x^0, y^0, x, y) - \frac{\mu}{2} \|z - z_1^*\|_V^2\}$,

$$H(x^0, y^0, x_1^*, y_1^*) \leq G_\mu(z^0, z_1^*).$$

It remains to bound the quadratic terms involving z . Using

$$\|z - z_1^*\|_V^2 \leq 2\|z - z_2^*\|_V^2 + 2\|z_1^* - z_2^*\|_V^2,$$

we have

$$\begin{aligned} & \sup_{z \in \mathcal{Z}} \left\{ \frac{\beta}{2K} \|z - z_1^*\|_V^2 + \frac{1}{K} \|z - z^0\|_V^2 - \frac{\mu}{2} \|z - z_2^*\|_V^2 \right\} \\ (5.10) \quad & \leq \frac{\beta}{K} \|z_1^* - z_2^*\|_V^2 + \sup_{z \in \mathcal{Z}} \left\{ - \left(\frac{\mu}{2} - \frac{\beta}{K} \right) \|z - z_2^*\|_V^2 + \frac{1}{K} \|z - z^0\|_V^2 \right\}. \end{aligned}$$

Since $\mu K > 2\beta + 2$, the quadratic function in the last supremum is concave. The standard quadratic maximization formula gives

$$\begin{aligned} & \sup_{z \in \mathcal{Z}} \left\{ - \left(\frac{\mu}{2} - \frac{\beta}{K} \right) \|z - z_2^*\|_V^2 + \frac{1}{K} \|z - z^0\|_V^2 \right\} \\ (5.11) \quad & \leq \frac{\mu K - 2\beta}{K(\mu K - 2\beta - 2)} \|z^0 - z_2^*\|_V^2 \leq \frac{2}{K} \|z^0 - z_2^*\|_V^2. \end{aligned}$$

where the last inequality holds because of the assumption $\mu K \geq 2\beta + 4$. Combining (5.10) and (5.11), taking the supremum over $z \in \mathcal{Z}$ in (5.9), and then using the fact $H(z^0, z_1^*) \leq G_\mu(z^0, z_1^*)$, we obtain

$$\begin{aligned} G_\mu(z^K, z_2^*) & \leq \frac{\alpha}{K} G_\mu(z^0, z_1^*) + \frac{\beta}{K} \|z^K - z_2^*\|_V^2 + \frac{\beta}{K} \|z^0 - z_2^*\|_V^2 \\ & \quad + \frac{\beta}{K} \|z_1^* - z_2^*\|_V^2 + \frac{2}{K} \|z^0 - z_2^*\|_V^2 + \frac{1}{2K} \sum_{k=0}^{K-1} \left(\|u^k\|_{\tau^{-1}P}^2 + \|v^k\|_{\sigma^{-1}Q}^2 \right) \\ (5.12) \quad & \quad + \frac{1}{K} \sum_{k=0}^{K-1} \left(\langle \tilde{x}^k, u^k \rangle_{\tau^{-1}} + \langle \tilde{y}^k, v^k \rangle_{\sigma^{-1}} + \varepsilon_k \right). \end{aligned}$$

Taking expectations in (5.12), the last summation vanishes. Moreover, by the uniform second-moment bound (4.14),

$$\mathbb{E} \|z^K - z_2^*\|_V^2 \leq 2 \mathbb{E} \|z^0 - z_2^*\|_V^2,$$

and by the square-summability estimation (4.41),

$$\mathbb{E} \left[\sum_{k=0}^{K-1} \left(\|u^k\|_{\tau^{-1}P}^2 + \|v^k\|_{\sigma^{-1}Q}^2 \right) \right] \leq \frac{2}{1 - 2\gamma_1^2} \mathbb{E} \|z^0 - z_2^*\|_V^2.$$

Therefore,

$$\begin{aligned} \mathbb{E}[G_\mu(z^K, z_2^*)] & \leq \frac{\alpha}{K} \mathbb{E}[G_\mu(z^0, z_1^*)] + \frac{\beta}{K} \mathbb{E}[\|z_1^* - z_2^*\|_V^2] \\ (5.13) \quad & \quad + \frac{1}{K} \left(3\beta + 2 + \frac{1}{1 - 2\gamma_1^2} \right) \mathbb{E}[\|z^0 - z_2^*\|_V^2], \end{aligned}$$

which proves (5.7). $\square$

5.3.2. Inter-epoch recursion. We now derive a recursion across consecutive restart epochs.

LEMMA 5.8. *Suppose Assumptions 2.1, 3.3, and 5.3 hold. For each outer iterate, define $\Pi^t := \text{proj}_{\mathcal{Z}^*}(\mathbf{Z}^t)$. Then, for all K satisfying $\mu K \geq 2\beta + 4$ and $t \geq 2$,*

$$\begin{aligned} \mathbb{E}[G_\mu(\mathbf{Z}^{t+1}, \Pi^t)] & \leq \frac{4\beta}{K\zeta} \mathbb{E}[G_\mu(\mathbf{Z}^{t-1}, \Pi^{t-2})] \\ (5.14) \quad & \quad + \frac{1}{K\zeta} \left(\alpha\zeta + 5\beta + 2 + \frac{1}{1 - 2\gamma_1^2} \right) \mathbb{E}[G_\mu(\mathbf{Z}^t, \Pi^{t-1})]. \end{aligned}$$

660 *Proof.* Applying Lemma 5.7 to the epoch initialized at $z^0 = \mathbf{Z}^t$, with ergodic
 661 output $Z^K = \mathbf{Z}^{t+1}$, and choosing $z_2^* = \Pi^t$ and $z_1^* = \Pi^{t-1}$, we obtain, for all $t \geq 1$,

$$662 \quad \mathbb{E}[G_\mu(\mathbf{Z}^{t+1}, \Pi^t)]$$

$$663 \quad (5.15) \quad \leq \frac{\alpha}{K} \mathbb{E}[G_\mu(\mathbf{Z}^t, \Pi^{t-1})] + \frac{\beta}{K} \mathbb{E}[\|\Pi^t - \Pi^{t-1}\|_V^2] + \frac{c_0(K)}{K} \mathbb{E}[\text{dist}_V(\mathbf{Z}^t, \mathcal{Z}^*)^2],$$

664 where

$$665 \quad c_0(K) := 3\beta + 2 + \frac{1}{1 - 2\gamma_1^2}.$$

666 It remains to control the distance between two consecutive projected saddle
 667 points. Consider the previous epoch, initialized at $\mathbf{Z}^{t-1}$ and producing the ergodic
 668 output $\mathbf{Z}^t$. By the triangle inequality,

$$669 \quad \|\Pi^t - \Pi^{t-1}\|_V^2 \leq 2\|\mathbf{Z}^t - \Pi^t\|_V^2 + 2\|\mathbf{Z}^t - \Pi^{t-1}\|_V^2.$$

670 Taking expectations gives

$$671 \quad (5.16) \quad \mathbb{E}[\|\Pi^t - \Pi^{t-1}\|_V^2] \leq 2\mathbb{E}[\text{dist}_V(\mathbf{Z}^t, \mathcal{Z}^*)^2] + 2\mathbb{E}[\|\mathbf{Z}^t - \Pi^{t-1}\|_V^2].$$

672 Since $\mathbf{Z}^t$ is the ergodic average of the inner iterates of the previous epoch, Jensen's
 673 inequality yields

$$674 \quad \|\mathbf{Z}^t - \Pi^{t-1}\|_V^2 \leq \frac{1}{K} \sum_{k=1}^K \|z^k - \Pi^{t-1}\|_V^2,$$

675 where z^k denotes the inner iterates generated during the epoch initialized at $\mathbf{Z}^{t-1}$.

676 Applying the uniform bound in (4.14) to the previous epoch with $z^* = \Pi^{t-1}$ gives

$$677 \quad \mathbb{E}[\|z^k - \Pi^{t-1}\|_V^2] \leq 2\mathbb{E}[\|\mathbf{Z}^{t-1} - \Pi^{t-1}\|_V^2] = 2\mathbb{E}[\text{dist}_V(\mathbf{Z}^{t-1}, \mathcal{Z}^*)^2].$$

678 Therefore,

$$679 \quad (5.17) \quad \mathbb{E}\|\mathbf{Z}^t - \Pi^{t-1}\|_V^2 \leq 2\mathbb{E}[\text{dist}_V(\mathbf{Z}^{t-1}, \mathcal{Z}^*)^2]$$

680 Substituting this bound into (5.16), we obtain

$$681 \quad (5.18) \quad \mathbb{E}[\|\Pi^t - \Pi^{t-1}\|_V^2] \leq 2\mathbb{E}[\text{dist}_V(\mathbf{Z}^t, \mathcal{Z}^*)^2] + 4\mathbb{E}[\text{dist}_V(\mathbf{Z}^{t-1}, \mathcal{Z}^*)^2].$$

682 Combining (5.15) and (5.18), we get

$$683 \quad \mathbb{E}[G_\mu(\mathbf{Z}^{t+1}, \Pi^t)] \leq \frac{\alpha}{K} \mathbb{E}[G_\mu(\mathbf{Z}^t, \Pi^{t-1})]$$

$$684 \quad (5.19) \quad + \frac{2\beta + c_0(K)}{K} \mathbb{E}[\text{dist}_V(\mathbf{Z}^t, \mathcal{Z}^*)^2] + \frac{4\beta}{K} \mathbb{E}[\text{dist}_V(\mathbf{Z}^{t-1}, \mathcal{Z}^*)^2].$$

685 By Assumption 5.3, for any saddle point $z^* \in \mathcal{Z}^*$,

$$686 \quad \text{dist}_V(\bar{z}, \mathcal{Z}^*)^2 \leq \frac{1}{\zeta} G_\mu(\bar{z}, z^*).$$

687 Applying this inequality with $(\bar{z}, z^*) = (\mathbf{Z}^t, \Pi^{t-1})$ and $(\bar{z}, z^*) = (\mathbf{Z}^{t-1}, \Pi^{t-2})$, we
 688 obtain, for all $t \geq 2$,

$$689 \quad \mathbb{E}[G_\mu(\mathbf{Z}^{t+1}, \Pi^t)] \leq \frac{4\beta}{K\zeta} \mathbb{E}[G_\mu(\mathbf{Z}^{t-1}, \Pi^{t-2})]$$

$$690 \quad (5.20) \quad + \frac{1}{K\zeta} (\alpha\zeta + 2\beta + c_0(K)) \mathbb{E}[G_\mu(\mathbf{Z}^t, \Pi^{t-1})].$$

691 Equivalently, using the definition of $c_0(K)$,

$$\begin{aligned}
 692 \quad \mathbb{E}[G_\mu(\mathbf{Z}^{t+1}, \Pi^t)] &\leq \frac{4\beta}{K\zeta} \mathbb{E}[G_\mu(\mathbf{Z}^{t-1}, \Pi^{t-2})] \\
 693 \quad (5.21) \quad &+ \frac{1}{K\zeta} \left(\alpha\zeta + 5\beta + 2 + \frac{1}{1-2\gamma_1^2} \right) \mathbb{E}[G_\mu(\mathbf{Z}^t, \Pi^{t-1})],
 \end{aligned}$$

694 for all $t \geq 2$. □

695 Now we are ready to prove theorem 5.5.

696 *Proof of Theorem 5.5.* Define

$$697 \quad a_t := \mathbb{E}[G_\mu(\mathbf{Z}^t, \Pi^{t-1})], \quad t \geq 1, \quad a_0 := \mathbb{E}[G_\mu(\mathbf{Z}^0, \Pi^0)].$$

698 By (5.14), the sequence $\{a_t\}_{t \geq 1}$ satisfies a second-order linear recursion. To simplify
699 the constants, define

$$700 \quad C := \frac{1}{\zeta} \max \left\{ \alpha\zeta + 5\beta + 2 + \frac{1}{1-2\gamma_1^2}, 4\beta \right\}.$$

701 The definition of C , together with the condition (5.3), implies $K \geq 100C$, and thus
702 both coefficients in (5.14) are bounded by $1/100$. Hence

$$703 \quad (5.22) \quad a_{t+1} \leq \frac{1}{100}(a_t + a_{t-1}), \quad t \geq 2.$$

704 Next, let $r := 1/5$, then $r^2 = \frac{1}{25} > \frac{1+r}{100}$. By choosing $M := \max\{a_0, 5a_1, 25a_2\}$, we
705 will show by induction that

$$706 \quad a_t \leq M r^{-t}, \quad \forall t \geq 1.$$

707 The claim holds for $t = 1, 2$ by the definition of M . Suppose it holds for $t-1$ and t .
708 Then

$$709 \quad a_{t+1} \leq \frac{1}{100}(a_t + a_{t-1}) \leq \frac{M}{100}(r^t + r^{t-1}) = M r^{t-1} \frac{1+r}{100} \leq M r^{t+1},$$

710 where the last inequality follows from $r^2 > (1+r)/100$. This completes the induction.

711 Since $\Pi^{t-1} \in \mathcal{Z}^*$, Assumption 5.3 gives

$$712 \quad \zeta \text{dist}_V(\mathbf{Z}^t, \mathcal{Z}^*)^2 \leq G_\mu(\mathbf{Z}^t, \Pi^{t-1}).$$

713 Taking expectations and using (5.4) yields (5.5). □

714 Finally, as a consequence of the linear convergence result, we can show that the
715 sequence $\{\mathbf{Z}^t\}_{t \geq 0}$ remains bounded in expectation within a ball centered at $\mathbf{Z}^0$. This
716 provides further support for the assumption $\mathbf{Z}^t \in \mathcal{R}$ for all $t \geq 0$ in Assumption 5.3,
717 and also yields an estimate of the radius of $\mathcal{R}$.

718 PROPOSITION 5.9. *Under the same conditions as theorem 5.5, we have*

$$719 \quad (5.23) \quad \mathbb{E} \|\mathbf{Z}^N - \mathbf{Z}^0\|_V^2 \leq \frac{20M}{\zeta}, \quad \forall N \geq 0.$$

Proof. According to Cauchy-Schwarz inequality, we have

$$\begin{aligned} \|\mathbf{Z}^N - \mathbf{Z}^0\|_V^2 &\leq \left(\sum_{t=1}^N \|\mathbf{Z}^t - \mathbf{Z}^{t-1}\|_V \right)^2 \leq \left(\sum_{t=1}^N 2^{-t} \right) \left(\sum_{t=1}^N 2^t \|\mathbf{Z}^t - \mathbf{Z}^{t-1}\|_V^2 \right) \\ &\leq \sum_{t=1}^N 2^t \|\mathbf{Z}^t - \mathbf{Z}^{t-1}\|_V^2. \end{aligned}$$

Besides, for any $t \geq 1$,

$$\begin{aligned} \mathbb{E} \|\mathbf{Z}^t - \mathbf{Z}^{t-1}\|_V^2 &\leq 2\mathbb{E} \|\mathbf{Z}^t - \Pi^{t-1}\|_V^2 + 2\mathbb{E} [\text{dist}_V(\mathbf{Z}^{t-1}, \mathbf{Z}^*)^2] \\ &\leq 6\mathbb{E} [\text{dist}_V(\mathbf{Z}^{t-1}, \mathbf{Z}^*)^2] \leq \frac{6M}{\zeta} 5^{-(t-1)}, \end{aligned} \quad (5.24)$$

where the second inequality comes from (5.17) and the third inequality comes from (5.5). Hence, we have

$$\mathbb{E} \|\mathbf{Z}^N - \mathbf{Z}^0\|_V^2 \leq \sum_{t=1}^N 2^t \mathbb{E} \|\mathbf{Z}^t - \mathbf{Z}^{t-1}\|_V^2 \leq \sum_{t=1}^N \frac{12M}{\zeta} \left(\frac{2}{5} \right)^{t-1} \leq \frac{20M}{\zeta}. \quad \square$$

6. Numerical Experiments. In this section, we illustrate the practical performance of DSPDHG and its restarted variant RDSPDHG on two classes of convex problems: soft-margin SVM under entrywise sampling and model predictive control (MPC) under blockwise sampling. Our experiments serve two purposes. First, they verify the sublinear convergence behavior predicted by the theory for general convex problems. Second, they demonstrate the effectiveness of restarting on problem instances that exhibit smoothed duality-gap quadratic growth.

Throughout this section, we compare deterministic PDHG, stochastic PDHG, doubly stochastic PDHG, and their restarted variants. We use uniform sampling on both primal and dual blocks. All methods are initialized from the same starting point. The step sizes are chosen according to the uniform rule

$$\sigma = 0.99 \frac{q}{\Lambda_{r,s}}, \quad \tau = 0.99 \frac{p}{\Lambda_{r,s}},$$

where $\Lambda_{r,s}$ is defined in (3.14) with $r = nq$ and $s = pm$. For SPDHG, we take $p = 1$, and for PDHG we take $p = q = 1$, in which case the rule reduces to the standard PDHG choice $\sigma = \tau = \frac{0.99}{\|A\|}$. For the restarted methods, we adopt an adaptive restart rule based on the relative KKT residual, following [22, 16]: a restart is triggered whenever

$$\text{relKKT}(z^k) \leq 0.8 \text{relKKT}(z^0). \quad (6.1)$$

This criterion is simple to implement and proved robust in our experiments. The relative error measures the gap between the objective value at the current iterate and the optimal objective value computed by the commercial solver COPT [13].

6.1. Soft-margin SVM. We first consider the soft-margin support vector machine problem

$$\min_{w,d} \frac{1}{2} \|w\|^2 + C \sum_{i=1}^n \max(0, 1 - b_i(w^\top a_i + d)), \quad (6.2)$$

where $w \in \mathbb{R}^m$, $d \in \mathbb{R}$, $a_i \in \mathbb{R}^m$ is the feature vector of the i th sample, and $b_i \in \{-1, +1\}$ is the corresponding label. The constant $C > 0$ is the penalty parameter.




Fig. 1: Relative error for soft-margin SVM. First column: colon-cancer ($n = 62, m = 2000$) with $p = 0.3, q = 0.15$; second column: mushrooms ($n = m = 100$) with $p = q = 0.1$; third column: w1a ($n = 2477, m = 300$) with $p = 0.2, q = 0.4$.

Reformulation. Problem (6.2) can be written in the form of (2.1) by introducing the variable $x = (w, d) \in \mathbb{R}^{m+1}$, and the matrix $A \in \mathbb{R}^{n \times (m+1)}$ whose i th row is $A_i = (b_i a_i^\top, b_i)$. We define

$$f_i(z) = C \max(0, 1 - z), \quad i = 1, \dots, n, \quad \text{and} \quad g_j(x_j) = \begin{cases} \frac{1}{2}x_j^2, & j = 1, \dots, m, \\ 0, & j = m + 1. \end{cases}$$

Then (6.2) takes the block-separable form (2.1). The Fenchel conjugate of f_i and its proximal mapping are

$$f_i^*(y) = \begin{cases} y, & y \in [-C, 0], \\ +\infty, & \text{otherwise,} \end{cases} \quad \text{and} \quad \text{prox}_{\sigma f_i^*}(x) = \begin{cases} -C, & x - \sigma < -C, \\ x - \sigma, & -C \leq x - \sigma \leq 0, \\ 0, & x - \sigma > 0. \end{cases}$$

Moreover,

$$\text{prox}_{\tau g_j}(x) = \begin{cases} \frac{x}{1 + \tau}, & j = 1, \dots, m, \\ x, & j = m + 1. \end{cases}$$

Experimental setup. We use several benchmark datasets from LIBSVM [8]. The primal variable is split coordinatewise, while the dual variable is naturally partitioned by samples. The penalty parameter C and the dataset statistics are reported together with the corresponding convergence plots in Figure 1.




Fig. 2: Constraint violation and relative error for MPC problems with $n_x = n_u = 20$ and $p = 0.1, q = 0.5$. First column: $T = 20$; second column: $T = 50$; third column: $T = 100$.

6.2. Model predictive control. We next consider the following model predictive control (MPC) problem:

$$\begin{aligned}
 \min_{\{x_t, u_t\}} \quad & x_T^\top H_T x_T + \sum_{t=0}^{T-1} (x_t^\top H_t x_t + u_t^\top R_t u_t) \\
 \text{s.t.} \quad & x_{t+1} = Ax_t + Bu_t, \quad t = 0, 1, \dots, T-1, \\
 & x_{t+1} \in \mathcal{X}, \quad u_t \in \mathcal{U}, \quad t = 0, 1, \dots, T-1, \\
 & x_0 = x_{\text{init}},
 \end{aligned}
 \tag{6.3}$$

where $x_t \in \mathbb{R}^{n_x}$ and $u_t \in \mathbb{R}^{n_u}$ denote the state and control variables, respectively. The box constraints are given by

$$\mathcal{X} = \{x \in \mathbb{R}^{n_x} : -\bar{x} \leq x \leq \bar{x}\}, \quad \mathcal{U} = \{u \in \mathbb{R}^{n_u} : -\bar{u} \leq u \leq \bar{u}\}.$$

The horizon length is T , and $x_{\text{init}} \in \mathbb{R}^{n_x}$ is the initial state. The matrices $H_t \in \mathbb{S}_+^{n_x}$, $R_t \in \mathbb{S}_+^{n_u}$, and $H_T \in \mathbb{S}_+^{n_x}$ define the stage and terminal costs.

Reformulation. Introducing dual variables y_t for the dynamics constraints $x_{t+1} - Ax_t - Bu_t = 0$, $t = 0, \dots, T-1$, yields the saddle-point formulation associated with the Lagrangian

$$\min_{x_t \in \mathcal{X}, u_t \in \mathcal{U}} x_T^\top H_T x_T + \sum_{t=0}^{T-1} (x_t^\top H_t x_t + u_t^\top R_t u_t) + \sum_{t=0}^{T-1} y_t^\top (x_{t+1} - Ax_t - Bu_t).$$

This problem is naturally block structured across time stages, making it well suited for stochastic primal-dual block updates.

Experimental setup. We generate synthetic instances with $n_x = n_u = 20$. The system matrix is chosen as $A = 0.5I + \Delta$, $\Delta_{ij} \sim \mathcal{N}(0, 0.1^2)$, and we keep only stable realizations whose eigenvalues lie strictly inside the unit circle. The input matrix is generated from $B_{ij} \sim \mathcal{N}(0, 1)$. The state cost matrix is $H_0 = \dots = H_T = \text{diag}(h)$,

where $h_i \sim \mathcal{U}(0, 10)$ and 70% of the entries of h are nonzero, and $R_0 = \dots = R_{T-1} = 0.1I$. The box constraints are generated from $\bar{x}_i \sim \mathcal{U}(1, 2)$, $\bar{u}_i \sim \mathcal{U}(0, 0.1)$, and the initial state is sampled as $x_{\text{init}} \sim \mathcal{U}(-0.5\bar{x}, 0.5\bar{x})$. We report results for horizons $T = 20, 50, 100$. The resulting convergence curves are reported in Figure 2.

Overall observations. As shown in both figures, the methods without restarting exhibit the expected sublinear convergence behavior, while restarting can significantly accelerate convergence when the local problem geometry is favorable. This is fully consistent with the theoretical results established in the previous sections. More importantly, across all tested instances, restarted DSPDHG delivers the best overall performance among the methods under comparison.

7. Conclusion. In this paper, we proposed DSPDHG, a doubly stochastic primal dual hybrid gradient method for block-structured convex saddle-point problems, together with its restarted variant RDSPDHG. By allowing randomized updates on both the primal and dual sides, the method extends classical PDHG and one-sided stochastic PDHG in a unified manner. For general convex problems, we established an $\mathcal{O}(1/K)$ ergodic convergence rate for the expected restricted primal-dual gap. Under a smoothed quadratic growth condition, we further proved linear convergence of the restarted scheme. Numerical experiments on soft-margin SVM and model predictive control demonstrate that the proposed methods are practically effective, and that restarting can substantially improve performance. These results suggest that doubly stochastic primal-dual updates provide a promising approach for solving large-scale structured convex optimization problems.

REFERENCES

- [1] J. M. AGUIRRE, D. CIFUENTES, V. GUIGUES, R. D. MONTEIRO, V. H. NASCIMENTO, AND A. SUJANANI, *cuhallr: A gpu accelerated low-rank augmented lagrangian method for large-scale semidefinite programming*, arXiv preprint arXiv:2505.13719, (2025).
- [2] A. ALACAOGLU, O. FERCOQ, AND V. CEVHER, *Random extrapolation for primal-dual coordinate descent*, in International conference on machine learning, PMLR, 2020, pp. 191–201.
- [3] A. ALACAOGLU, O. FERCOQ, AND V. CEVHER, *On the convergence of stochastic primal-dual hybrid gradient*, SIAM Journal on Optimization, 32 (2022), pp. 1288–1318.
- [4] D. APPLEGATE, M. DÍAZ, O. HINDER, H. LU, M. LUBIN, B. O'DONOGHUE, AND W. SCHUDY, *Practical large-scale linear programming using primal-dual hybrid gradient*, Advances in Neural Information Processing Systems, 34 (2021), pp. 20243–20257.
- [5] A. CHAMBOLLE, M. J. EHRHARDT, P. RICHTÁRIK, AND C.-B. SCHÖNLIEB, *Stochastic primal-dual hybrid gradient algorithm with arbitrary sampling and imaging applications*, SIAM Journal on Optimization, 28 (2018), pp. 2783–2808.
- [6] A. CHAMBOLLE AND T. POCK, *A first-order primal-dual algorithm for convex problems with applications to imaging*, Journal of mathematical imaging and vision, 40 (2011), pp. 120–145.
- [7] A. CHAMBOLLE AND T. POCK, *On the ergodic convergence rates of a first-order primal-dual algorithm*, Mathematical Programming, 159 (2016), pp. 253–287.
- [8] C.-C. CHANG AND C.-J. LIN, *Libsvm: A library for support vector machines*, ACM Trans. Intell. Syst. Technol., 2 (2011), <https://doi.org/10.1145/1961189.1961199>, <https://doi.org/10.1145/1961189.1961199>.
- [9] K. CHEN, D. SUN, Y. YUAN, G. ZHANG, AND X. ZHAO, *Hpr-lp: An implementation of an hpr method for solving linear programming*, arXiv preprint arXiv:2408.12179, (2024).
- [10] K. CHEN, D. SUN, Y. YUAN, G. ZHANG, AND X. ZHAO, *Hpr-qp: A dual halpern peaceman-rachford method for solving large-scale convex composite quadratic programming*, arXiv preprint arXiv:2507.02470, (2025).
- [11] L. DING, H. LU, AND J. YANG, *New understandings and computation on augmented lagrangian methods for low-rank semidefinite programming*, arXiv preprint arXiv:2505.15775, (2025).
- [12] O. FERCOQ, *Quadratic error bound of the smoothed gap and the restarted averaged primal-dual hybrid gradient*, arXiv preprint arXiv:2206.03041, (2022).

- [13] D. GE, Q. HUANGFU, Z. WANG, J. WU, AND Y. YE, *Cardinal optimizer (copt) user guide*, arXiv preprint arXiv:2208.14314, (2022).
- [14] Q. HAN, C. LI, Z. LIN, C. CHEN, Q. DENG, D. GE, H. LIU, AND Y. YE, *A low-rank admm splitting approach for semidefinite programming*, arXiv preprint arXiv:2403.09133, (2024).
- [15] Q. HAN, Z. LIN, H. LIU, C. CHEN, Q. DENG, D. GE, AND Y. YE, *Accelerating low-rank factorization-based semidefinite programming algorithms on gpu*, arXiv preprint arXiv:2407.15049, (2024).
- [16] Y. HUANG, W. ZHANG, H. LI, D. GE, H. LIU, AND Y. YE, *Restarted primal-dual hybrid conjugate gradient method for large-scale quadratic programming*, arXiv preprint arXiv:2405.16160, (2024).
- [17] Z. LIN, Z. XIONG, D. GE, AND Y. YE, *Pdcs: A primal-dual large-scale conic programming solver with gpu enhancements*, arXiv preprint arXiv:2505.00311, (2025).
- [18] H. LIU, Y. HUANG, H. LI, D. GE, AND Y. YE, *Pdhcg: A scalable first-order method for large-scale competitive market equilibrium computation*, 2025, <https://arxiv.org/abs/2506.06258>.
- [19] H. LU AND J. YANG, *cupdlp.jl: A gpu implementation of restarted primal-dual hybrid gradient for linear programming in julia*, arXiv preprint arXiv:2311.12180, (2023).
- [20] H. LU AND J. YANG, *Pdot: A practical primal-dual algorithm and a gpu-based solver for optimal transport*, arXiv preprint arXiv:2407.19689, (2024).
- [21] H. LU AND J. YANG, *Restarted halpern pdhg for linear programming*, arXiv preprint arXiv:2407.16144, (2024).
- [22] H. LU AND J. YANG, *A practical and optimal first-order method for large-scale convex quadratic programming*, Mathematical Programming, (2025), pp. 1–38.
- [23] C. MELCHER, A. JALILZADEH, AND E. Y. HAMEDANI, *Linear convergence of a unified primal-dual algorithm for convex-concave saddle point problems with quadratic growth*, 2025, <https://api.semanticscholar.org/CorpusID:282064933>.
- [24] A. W. YU, Q. LIN, AND T. YANG, *Doubly stochastic primal-dual coordinate method for bilinear saddle-point problem*, arXiv preprint arXiv:1508.03390, (2015).
- [25] F. ZHANG AND S. BOYD, *Solving large multicommodity network flow problems on gpus*, arXiv preprint arXiv:2501.17996, (2025).
- [26] G. ZHANG, Z. GU, Y. YUAN, AND D. SUN, *Hot: An efficient halpern accelerating algorithm for optimal transport problems*, IEEE Transactions on Pattern Analysis and Machine Intelligence, (2025).