

Electronic supplementary information

8,8'- μ -AMMONIUM DERIVATIVE OF COBALT BIS(DICARBOLLIDE) [8,8'- μ -NH₂-3,3'-Co(1,2-C₂B₉H₁₀)₂]

E. V. Bogdanova,*^a M. Yu. Stogniy,^{a,b} and S. A. Anufriev^a

^a Nesmeyanov Institute of Organoelement Compounds, Russian Academy of Sciences,
ul. Vavilova 28, str. 1, Moscow, 119334 Russia

^b Lomonosov Institute of Fine Chemical Technologies, MIREA–Russian Technological
University, pr. Vernadskogo 86, Moscow, 119571 Russia

Multinuclear NMR spectra of compound [8,8'- μ -NH₂-3,3'-Co(1,2-C₂B₉H₁₀)₂]




Figure S1. ¹H NMR spectrum of compound **2** (acetone-*d*₆).




Figure S2. $^{11}\text{B}\{^1\text{H}\}$ NMR spectrum of compound **2** (acetone- d_6).




Figure S3. ^{11}B NMR spectrum of compound **2** (acetone- d_6).




Figure S4. ^{13}C NMR spectrum of compound **2** (acetone- d_6).




Figure S5. ^1H - ^{13}C NMR HMQC spectrum of compound **2** (acetone- d_6).

Proposed reaction mechanism