

# Lecture Note 2

- Stationary Process
- Autocorrelation function
- Linear time series models
- Autoregressive of order one,  $AR(1)$ , process

## Stationary Process

**Definition 1** (Weak stationarity). A stochastic process  $\{y_t, t \in \mathcal{T}\}$  is said to be **weakly stationary** if it has a **constant mean** and **constant variance** and its **covariance function**,  $\gamma(t_1, t_2)$ , defined by

$$\gamma(t_1, t_2) = \text{cov}(y_{t_1}, y_{t_2}) = \mathbb{E}\{[y_{t_1} - \mathbb{E}(y_{t_1})][y_{t_2} - \mathbb{E}(y_{t_2})]\},$$

depends only on the **absolute difference**  $|t_1 - t_2|$ , namely

$$\gamma(t_1, t_2) = \gamma(|t_1 - t_2|).$$

**Definition 2** (Weakly trend stationary). A stochastic process  $\{y_t, t \in \mathcal{T}\}$  is said to be **weakly trend stationary** if  $y_t = x_t + d_t$ , where  $x_t$  is **weakly stationary** and  $d_t$  is **deterministic**.

**Definition 3.** (White noise process) The process  $\{\varepsilon_t, t \in \mathcal{T}\}$  is said to be a white noise process if it has **mean zero**, a **constant variance**, and  $\varepsilon_t$  and  $\varepsilon_s$  are **uncorrelated** for all  $s \neq t$ .

## Autocorrelation function

- In Statistics, the correlation coefficient between two random variables  $X$  and  $Y$  is defined as

$$\rho_{X,Y} = \frac{\text{cov}(X, Y)}{\sqrt{\text{var}(X)\text{var}(Y)}} = \frac{\mathbb{E}\{[X - \mathbb{E}(X)][Y - \mathbb{E}(Y)]\}}{\sqrt{\mathbb{E}\{[X - \mathbb{E}(X)]^2\}\mathbb{E}\{[Y - \mathbb{E}(Y)]^2\}}}$$

- When the sample  $\{(x_t, y_t) | t = 1, 2, \dots, T\}$  is available, the correlation can be consistency estimated by its sample counterpart

$$\hat{\rho}_{X,Y} = \frac{\frac{1}{T} \sum_{t=1}^T [(x_t - \bar{x})(y_t - \bar{y})]}{\sqrt{\frac{1}{T} \sum_{t=1}^T (x_t - \bar{x})^2 \frac{1}{T} \sum_{t=1}^T (y_t - \bar{y})^2}},$$

where  $\bar{x} = \frac{1}{T} \sum_{t=1}^T x_t$  and  $\bar{y} = \frac{1}{T} \sum_{t=1}^T y_t$ .

- The correlation coefficient is between  $-1$  and  $1$ ,  $-1 \leq \rho_{XY} \leq 1$ .

- Let  $x_t$  be a weakly stationary process. The correlation coefficient between  $x_t$  and  $x_{t-k}$  is called lag- $k$  autocorrelation of  $x_t$  and is commonly denoted by  $\rho(k)$ . Specifically, we define

$$\rho(k) = \frac{\text{cov}(x_t, x_{t-k})}{\sqrt{\text{var}(x_t)\text{var}(x_{t-k})}} = \frac{\gamma(t, t-k)}{\text{var}(x_t)} = \frac{\gamma(k)}{\gamma(0)}.$$

- $\rho(0) = 1$ ,  $\rho(k) = \rho(-k)$ , and  $-1 \leq \rho(k) \leq 1$ .
- $\hat{\rho}(k) = \frac{\frac{1}{T} \sum_{t=k+1}^T (x_t - \bar{x})(x_{t-k} - \bar{x})}{\frac{1}{T} \sum_{t=1}^T (x_t - \bar{x})^2}$  is a consistent estimator of  $\rho(k)$ . If  $\{x_t\}_{t=1}^T$  is a sequence of independently identically distributed (iid) random variables and  $\mathbb{E}(x_t)^2 < \infty$ , then  $\hat{\rho} \stackrel{a.s.}{\sim} \mathcal{N}(0, \frac{1}{T})$ .

Generally, if  $x_t$  is a weakly stationary process satisfying,  $x_t = \mu + \sum_{i=0}^q \delta_i \varepsilon_{t-i}$ , where  $\delta_0 = 1$  and  $\{\varepsilon\}_i$  is a sequence of iid random variables with mean 0, then  $\hat{\rho}(k) \stackrel{a.s.}{\sim} \mathcal{N}(\rho(k), \frac{1}{T} + \frac{2}{T} \sum_{i=1}^q \rho(i)^2)$ .

- Testing individual Autocorrelation Coefficient Function (ACF):

$$\mathcal{H}_0 : \rho(k) = 0, \text{ v.s. } \mathcal{H}_a : \rho(k) \neq 0 \longrightarrow \text{t-ratio} = \frac{\hat{\rho}(k)}{\sqrt{\frac{1}{T} + \frac{2}{T} \sum_{i=1}^q \hat{\rho}(i)^2}} \stackrel{a.s.}{\sim} \mathcal{N}(0, 1).$$

Or if we assume that  $\rho_i = 0$  for all  $i \neq k$  then:

$$\text{t-ratio} = \sqrt{T} \hat{\rho}(k) \stackrel{a.s.}{\sim} \mathcal{N}(0, 1).$$

- Joint test for several autocorrelations:

$$\mathcal{H}_0 : \rho(1) = \rho(2) = \dots = \rho(m) = 0, \text{ v.s. } \mathcal{H}_a : \rho(i) \neq 0 \text{ for at least one } i \in \{1, 2, \dots, m\},$$

Box and Pierce(1970):  $Q^*(m) = T \sum_{\ell=1}^m \hat{\rho}(\ell)^2 \stackrel{a.s.}{\sim} \chi^2(m)$ ,

Ljung and Pierce(1978):  $Q(m) = T(T+2) \sum_{\ell=1}^m \frac{\hat{\rho}(\ell)^2}{T-\ell} \stackrel{a.s.}{\sim} \chi^2(m)$ .

- Choice of  $m$ :  $m \approx T^{1/3}$  or  $m \approx \ln(T)$ .

## Linear time series models

**Definition 4.** A time series  $x_t$  is said to be linear if it can be written as:

$$x_t = \mu + \sum_{i=0}^{\infty} \delta_i \varepsilon_{t-i}, \text{ where } \varepsilon_t \stackrel{iid}{\sim} (0, \sigma^2). \quad (1)$$

**Q:** What is the expected value, variance, and autocovariance function of  $x_t$  generated by (1)?

**A:**  $\mathbb{E}(x_t) = \mu + \sum_{i=0}^{\infty} \delta_i \mathbb{E}(\varepsilon_{t-i}) = \mu.$

$$\text{var}(x_t) = \sum_{i=0}^{\infty} \delta_i^2 \text{var}(\varepsilon_{t-i}) = \sigma^2 \sum_{i=0}^{\infty} \delta_i^2.$$

$$\begin{aligned} \gamma(\ell) &= \text{cov}(x_t, x_{t-\ell}) = \mathbb{E}(x_t - \mu)(x_{t-\ell} - \mu) = \mathbb{E} \left[ \left( \sum_{i=0}^{\infty} \delta_i \varepsilon_{t-i} \right) \left( \sum_{j=0}^{\infty} \delta_j \varepsilon_{t-\ell-j} \right) \right] \\ &= \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \delta_i \delta_j \mathbb{E}(\varepsilon_{t-i} \varepsilon_{t-\ell-j}) = \sum_{i=0}^{\infty} \delta_i \delta_{i-\ell} \mathbb{E}(\varepsilon_{t-i}^2) = \sigma^2 \sum_{i=0}^{\infty} \delta_i \delta_{i-\ell} \end{aligned}$$

**Q:** Given the answer to the previous question, can we conclude that  $x_t$  is a weakly stationary process?

**A:** For  $x_t$  to be weakly stationary, we need its variance to exist, i.e.  $\gamma(0) = \text{var}(x_t) < \infty$ . So, we need  $\sum_{i=0}^{\infty} \delta_i^2 < \infty$ . We also need the autocovariance function to exist, i.e.  $\gamma(\ell) = \text{cov}(x_t, x_{t-\ell}) < \infty$  for all  $\ell$ . But note that,  $\sum_{i=0}^{\infty} \delta_i \delta_{i-\ell} \leq (\sum_{i=0}^{\infty} \delta_i^2)^{1/2} (\sum_{i=0}^{\infty} \delta_{i-\ell}^2)^{1/2}$ . So, if  $\sum_{i=0}^{\infty} \delta_i^2 < \infty$ , then  $\gamma(\ell) < \infty$  for all  $\ell$ . Assuming that  $\sum_{i=0}^{\infty} \delta_i^2 < \infty$ ,  $x_t$  have constant mean and variance, and its autocovariance function only depends on the absolute value of the time distance. So, we can conclude that  $x_t$  is a weakly stationary process.

• The autocorrelation function of  $x_t$  is as follows:

$$\rho(\ell) = \frac{\gamma(\ell)}{\gamma(0)} = \frac{\sum_{i=0}^{\infty} \delta_i \delta_{i-\ell}}{\sum_{i=0}^{\infty} \delta_i^2}$$

## Autoregressive of order one, AR(1), process

**Definition 5.** A time series process  $x_t$  is said to be Autoregressive of order one, AR(1), if it can be written as

$$x_t = \phi_0 + \phi_1 x_{t-1} + \varepsilon_t, \text{ where } \varepsilon_t \stackrel{iid}{\sim} (0, \sigma^2). \quad (2)$$

**Q:** Suppose that  $x_t$  given by (2) is a weakly stationary process. Compute its mean, variance and autocovariance function.

**A:**  $\mathbb{E}(x_t) = \phi_0 + \phi_1 \mathbb{E}(x_{t-1}) = \phi_0 + \phi_1 \mathbb{E}(x_t) \Rightarrow \mathbb{E}(x_t) = \frac{\phi_0}{1-\phi_1}.$

$$\text{var}(x_t) = \phi_1^2 \text{var}(x_{t-1}) + \text{var}(\varepsilon_t) = \phi_1^2 \text{var}(x_t) + \sigma^2 \Rightarrow \text{var}(x_t) = \frac{\sigma^2}{1-\phi_1^2}.$$

$\gamma(\ell) = \text{cov}(x_t, x_{t-\ell}) = \text{cov}(\phi_0 + \phi_1 x_{t-1} + \varepsilon_t, x_{t-\ell}) = \phi_1 \text{cov}(x_{t-1}, x_{t-\ell}) = \phi_1 \gamma(\ell-1)$ . By substituting,  $\gamma(\ell-1) = \phi_1 \gamma(\ell-2)$ , we can further write,  $\gamma(\ell) = \phi_1^2 \gamma(\ell-2)$ . By repeating

this for  $\ell$  times we can get  $\gamma(\ell) = \phi_1^\ell \gamma(0) = \phi_1^\ell \frac{\sigma^2}{1-\phi_1^2}$ .

**Q:** Find the condition(s) under which  $x_t$  given by (2) is a weakly stationary process.

**A:** By substituting  $x_{t-1} = \phi_0 + \phi_1 x_{t-2} + \varepsilon_{t-1}$  into the equation for  $x_t$ , we get

$$x_t = \phi_0 + \phi_0 \phi_1 + \phi_1^2 x_{t-2} + \varepsilon_t + \phi_1 \varepsilon_{t-1}.$$

By substituting  $x_{t-2} = \phi_0 + \phi_1 x_{t-3} + \varepsilon_{t-2}$ , we can further write

$$x_t = \phi_0(1 + \phi_1 + \phi_1^2) + \phi_1^3 x_{t-3} + \varepsilon_t + \phi_1 \varepsilon_{t-1} + \phi_1^2 \varepsilon_{t-2}.$$

Repeating this for  $n$  times, we get

$$x_t = \phi_0(1 + \phi_1 + \phi_1^2 + \cdots + \phi_1^n) + \phi_1^n x_{t-n} + \varepsilon_t + \phi_1 \varepsilon_{t-1} + \phi_1^2 \varepsilon_{t-2} + \cdots + \phi_1^n \varepsilon_{t-n}.$$

If  $|\phi_1| < 1$ , as  $n \rightarrow \infty$ , we get

$$x_t = \phi_0(1 + \phi_1 + \phi_1^2 + \cdots) + \varepsilon_t + \phi_1 \varepsilon_{t-1} + \phi_1^2 \varepsilon_{t-2} + \cdots.$$

Since  $|\phi_1| < 1$ ,  $1 + \phi_1 + \phi_1^2 + \cdots = \frac{1}{1-\phi_1}$ . Hence,

$$x_t = \frac{\phi_0}{1-\phi_1} + \sum_{i=0}^{\infty} \phi_1^i \varepsilon_{t-i}.$$

So, if  $|\phi_1| < 1$ , we can present  $AR(1)$  process as a linear time series process where  $\mu = \frac{\phi_0}{1-\phi_1}$  and  $\delta_i = \phi_1^i$ . We know that a linear time series process is weakly stationary if  $\sum_{i=0}^{\infty} \delta_i^2 < \infty$ . Here we have  $|\phi_1| < 1$ , therefore  $\sum_{i=0}^{\infty} \delta_i^2 = \sum_{i=0}^{\infty} (\phi_1^2)^i = \frac{1}{1-\phi_1^2} < \infty$ . So, we can conclude that a  $AR(1)$  process is weakly stationary if  $|\phi_1| < 1$ .