



---

# **MACHINE LEARNING**

## **CHAPTER 2: PROBABILITY DISTRIBUTIONS**

---

# Learning Objectives

---

- 1、 What are binary, multinomial and Gaussian distributions and their conjugate prior distributions?
  - 2、 What are the common properties of Gaussian distributions?
  - 3、 What are exponential families and their properties?
  - 4、 How to choose non-informative prior\*?
  - 5、 How to use non-parametric methods for learning?
  - 6、 What are KNN based methods?
-

# Outlines

---

- Binary Distributions
  - Multinomial Distributions
  - Gaussian Distributions
  - Exponential Families
  - Non-informative Priors
  - Non-parametric Methods
  - KNN
-


# Outlines

---


- Gaussian Distributions and Central Limit Theorem
  - Moments and Properties of Gaussians
  - Posterior of the Linear Gaussian Variable
  - Partitioned Gaussian Conditionals and Marginals
  - ML Learning of Gaussian Distributions
  - Sequential Learning of Gaussian Distributions
  - Bayesian Learning of Gaussian Distributions
  - Mixture of Gaussian Distributions
-

# The Gaussian Distribution

---



$$\mathcal{N}(x|\mu, \sigma^2) = \frac{1}{(2\pi\sigma^2)^{1/2}} \exp \left\{ -\frac{1}{2\sigma^2} (x - \mu)^2 \right\}$$



$$\mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{1}{(2\pi)^{D/2}} \frac{1}{|\boldsymbol{\Sigma}|^{1/2}} \exp \left\{ -\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right\}$$


---

# Central Limit Theorem

---

The distribution of the sum of  $N$  i.i.d. random variables becomes increasingly Gaussian as  $N$  grows.

Example:  $N$  uniform  $[0,1]$  random variables.



# Geometry of the Multivariate Gaussian


---

$$\Delta^2 = (\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu})$$

$$\boldsymbol{\Sigma}^{-1} = \sum_{i=1}^D \frac{1}{\lambda_i} \mathbf{u}_i \mathbf{u}_i^T$$

$$\Delta^2 = \sum_{i=1}^D \frac{y_i^2}{\lambda_i}$$

$$y_i = \mathbf{u}_i^T (\mathbf{x} - \boldsymbol{\mu})$$



# Moments of the Multivariate Gaussian (1)

---

$$\begin{aligned}\mathbb{E}[\mathbf{x}] &= \frac{1}{(2\pi)^{D/2}} \frac{1}{|\boldsymbol{\Sigma}|^{1/2}} \int \exp \left\{ -\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu}) \right\} \mathbf{x} \, d\mathbf{x} \\ &= \frac{1}{(2\pi)^{D/2}} \frac{1}{|\boldsymbol{\Sigma}|^{1/2}} \int \exp \left\{ -\frac{1}{2}\mathbf{z}^T \boldsymbol{\Sigma}^{-1}\mathbf{z} \right\} (\mathbf{z} + \boldsymbol{\mu}) \, d\mathbf{z}\end{aligned}$$

thanks to anti-symmetry of  $\mathbf{z}$

$$\mathbb{E}[\mathbf{x}] = \boldsymbol{\mu}$$

---






# Moments of the Multivariate Gaussian (2)

---

$$\mathbb{E}[\mathbf{x}\mathbf{x}^T] = \boldsymbol{\mu}\boldsymbol{\mu}^T + \boldsymbol{\Sigma}$$

$$\text{cov}[\mathbf{x}] = \mathbb{E} [(\mathbf{x} - \mathbb{E}[\mathbf{x}])(\mathbf{x} - \mathbb{E}[\mathbf{x}])^T] = \boldsymbol{\Sigma}$$

$$\text{cov}[A\mathbf{x}] = A\boldsymbol{\Sigma}A^T$$



# Properties of Gaussians

---

$$\left. \begin{array}{l} X \sim N(\mu, \sigma^2) \\ Y = aX + b \end{array} \right\} \Rightarrow Y \sim N(a\mu + b, a^2\sigma^2)$$

$$\left. \begin{array}{l} X_1 \sim N(\mu_1, \sigma_1^2) \\ X_2 \sim N(\mu_2, \sigma_2^2) \end{array} \right\} \Rightarrow p(X_1) \cdot p(X_2) \sim N\left(\frac{\sigma_2^2}{\sigma_1^2 + \sigma_2^2} \mu_1 + \frac{\sigma_1^2}{\sigma_1^2 + \sigma_2^2} \mu_2, \frac{1}{\sigma_1^{-2} + \sigma_2^{-2}}\right)$$



$$p(X) \sim N(\mu, \sigma^2)$$


Precision

$$\left[ \begin{array}{lcl} \sigma^{-2} & = & \sigma_1^{-2} + \sigma_2^{-2} \\ \sigma^{-2}\mu & = & \sigma_1^{-2}\mu_1 + \sigma_2^{-2}\mu_2 \end{array} \right.$$

# Properties of Gaussians

---

$$p_{X_1}(x)p_{X_2}(x) \propto e^{-\frac{(x-\mu_1)^2}{2\sigma_1^2}} e^{-\frac{(x-\mu_2)^2}{2\sigma_2^2}}$$



$$p_X(x) \propto e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

Quadratic terms of  $x$  ( $x^2$ ) are equal

1<sup>st</sup> order terms of  $x$  are also equal

$$\begin{cases} \sigma^{-2} &= \sigma_1^{-2} + \sigma_2^{-2} \\ \sigma^{-2}\mu &= \sigma_1^{-2}\mu_1 + \sigma_2^{-2}\mu_2 \end{cases}$$

# Properties of Multivariate Gaussians

---

$$\left. \begin{array}{l} X \sim N(\mu, \Sigma) \\ Y = AX + B \end{array} \right\} \Rightarrow Y \sim N(A\mu + B, A\Sigma A^T)$$

$$\left. \begin{array}{l} X_1 \sim N(\mu_1, \Sigma_1) \\ X_2 \sim N(\mu_2, \Sigma_2) \end{array} \right\} \Rightarrow p(X_1) \cdot p(X_2) \sim N\left( \frac{\Sigma_2}{\Sigma_1 + \Sigma_2} \mu_1 + \frac{\Sigma_1}{\Sigma_1 + \Sigma_2} \mu_2, \frac{1}{\Sigma_1^{-1} + \Sigma_2^{-1}} \right)$$

(where division "-" denotes matrix inversion)

- We **stay Gaussian** as long as we start with Gaussians and perform only **linear transformations**
-

# Properties of Multivariate Gaussians

---

$$p_X(x) \sim N(\mu, \Sigma) \iff \begin{aligned} p_{X_1}(x) &\sim N(\mu_1, \Sigma_1) \\ p_{X_2}(x) &\sim N(\mu_2, \Sigma_2) \end{aligned}$$

Precision

$$\left[ \begin{array}{l} \Sigma^{-1} \\ \Sigma^{-1} \mu \end{array} \right] = \begin{array}{l} \Sigma_1^{-1} + \Sigma_2^{-1} \\ \Sigma_1^{-1} \mu_1 + \Sigma_2^{-1} \mu_2 \end{array}$$

Mean

# Bayes' Theorem for Gaussian Variables (1)

---

Given  $y = Ax + v$

$$p(x) = \mathcal{N}(x|\mu, \Sigma) \quad p(v) = \mathcal{N}(v|0, Q)$$

we have  $p(y|x) = \mathcal{N}(y|Ax, Q)$

$$p(y) = \mathcal{N}(y|A\mu, A\Sigma A^T + Q)$$

Then what is  $p(x|y)$  ?

---

# Bayes' Theorem for Gaussian Variables (2)

---

Given  $x = m + u$

$$p(x|y) = \mathcal{N}(x|m, L) \quad p(u) = \mathcal{N}(u|0, L)$$

we have

$$p(x|y) \propto p(y|x)p(x) = \mathcal{N}(y|Ax, Q)\mathcal{N}(x|\mu, \Sigma)$$

$$-\frac{1}{2}(x - m)^T L^{-1}(x - m) \quad \propto \quad -\frac{1}{2}(y - Ax)^T Q^{-1}(y - Ax) \\ -\frac{1}{2}(x - \mu)^T \Sigma^{-1}(x - \mu)$$

---

# Bayes' Theorem for Gaussian Variables (3)

---

$$-\frac{1}{2}(x - m)^T L^{-1}(x - m) \propto -\frac{1}{2}(y - Ax)^T Q^{-1}(y - Ax) - \frac{1}{2}(x - \mu)^T \Sigma^{-1}(x - \mu)$$

Quadratic terms of  $x$  ( $x^T ** x$ ) are equal

$$\begin{cases} L^{-1} &= A^T Q^{-1} A + \Sigma^{-1} \\ L^{-1} m &= A^T Q^{-1} y + \Sigma^{-1} \mu \end{cases}$$

1<sup>st</sup> order terms of  $x$  ( $x^T **$ ) are also equal

---



## Bayes' Theorem for Gaussian Variables (4)

---

$$p(x|y) = \mathcal{N}(x|m, L)$$

where

$$\begin{cases} L^{-1} &= A^T Q^{-1} A + \Sigma^{-1} \\ m &= L \{ A^T Q^{-1} y + \Sigma^{-1} \mu \} \end{cases}$$

# Matrix Inversion Lemma

---

If  $A$ ,  $C$ ,  $BCD$  are non-singular square matrix (the inverse exists) then

$$[A + BCD]^{-1} = A^{-1} - A^{-1}B[C^{-1} + DA^{-1}B]^{-1}DA^{-1}$$

# Matrix Inversion Lemma Proof

---

$$\begin{aligned} & [A + BCD] [A^{-1} - A^{-1}B[C^{-1} + DA^{-1}B]^{-1}DA^{-1}] \\ &= I + BCDA^{-1} - B[C^{-1} + DA^{-1}B]^{-1}DA^{-1} \\ &\quad - BCDA^{-1}B[C^{-1} + DA^{-1}B]^{-1}DA^{-1} \\ &= I + BCDA^{-1} - B\{I + CDA^{-1}B\}[C^{-1} + DA^{-1}B]^{-1}DA^{-1} \\ &= I + BCDA^{-1} - BC\{C^{-1} + DA^{-1}B\}[C^{-1} + DA^{-1}B]^{-1}DA^{-1} \\ &= I \end{aligned}$$

---

# Bayes' Theorem for Gaussian Variables (5)

---

Then

$$L = \Sigma - \Sigma A^T (A^T \Sigma A + Q)^{-1} A \Sigma$$

$$\begin{cases} L &= (I - KA)\Sigma \\ m &= \mu + K(y - A\mu) \end{cases}$$

Kalman Gain  $\longrightarrow K = \Sigma A^T (A^T \Sigma A + Q)^{-1}$

$$p(x|y) = \mathcal{N}(x|\mu + K(y - A\mu), (I - KA)\Sigma)$$

---

# Bayes' Theorem for Gaussian Variables (6)\*

---

Given

$$\begin{aligned}p(\mathbf{x}) &= \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Lambda}^{-1}) \\p(\mathbf{y}|\mathbf{x}) &= \mathcal{N}(\mathbf{y}|\mathbf{Ax} + \mathbf{b}, \mathbf{L}^{-1})\end{aligned}$$

we have

$$\begin{aligned}p(\mathbf{y}) &= \mathcal{N}(\mathbf{y}|\mathbf{A}\boldsymbol{\mu} + \mathbf{b}, \mathbf{L}^{-1} + \mathbf{A}\boldsymbol{\Lambda}^{-1}\mathbf{A}^T) \\p(\mathbf{x}|\mathbf{y}) &= \mathcal{N}(\mathbf{x}|\boldsymbol{\Sigma}\{\mathbf{A}^T\mathbf{L}(\mathbf{y} - \mathbf{b}) + \boldsymbol{\Lambda}\boldsymbol{\mu}\}, \boldsymbol{\Sigma})\end{aligned}$$

where

$$\boldsymbol{\Sigma} = (\boldsymbol{\Lambda} + \mathbf{A}^T\mathbf{L}\mathbf{A})^{-1}$$

---

# Partitioned Gaussian Distributions (1)

---

$$p(\mathbf{x}) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) \quad \boldsymbol{\Lambda} \equiv \boldsymbol{\Sigma}^{-1} \quad \boldsymbol{\Lambda} = \begin{pmatrix} \boldsymbol{\Lambda}_{aa} & \boldsymbol{\Lambda}_{ab} \\ \boldsymbol{\Lambda}_{ba} & \boldsymbol{\Lambda}_{bb} \end{pmatrix}$$

$$\mathbf{x} = \begin{pmatrix} \mathbf{x}_a \\ \mathbf{x}_b \end{pmatrix} \quad \boldsymbol{\mu} = \begin{pmatrix} \boldsymbol{\mu}_a \\ \boldsymbol{\mu}_b \end{pmatrix} \quad \boldsymbol{\Sigma} = \begin{pmatrix} \boldsymbol{\Sigma}_{aa} & \boldsymbol{\Sigma}_{ab} \\ \boldsymbol{\Sigma}_{ba} & \boldsymbol{\Sigma}_{bb} \end{pmatrix}$$

$$\mathbf{x}_a = \mathbf{A}\mathbf{x}_b + \mathbf{w} \quad \boldsymbol{\Sigma}_{a|b} = \boldsymbol{\Sigma}_w$$

$$\mathbf{x}_a - \boldsymbol{\mu}_a = \mathbf{A}(\mathbf{x}_b - \boldsymbol{\mu}_b) + \mathbf{w} \Rightarrow \boldsymbol{\mu}_{a|b} - \boldsymbol{\mu}_a = \mathbf{A}(\mathbf{x}_b - \boldsymbol{\mu}_b)$$

$$\boldsymbol{\Sigma}_{ab} = \mathbf{A}\boldsymbol{\Sigma}_{bb} \Rightarrow \mathbf{A} = \boldsymbol{\Sigma}_{ab}\boldsymbol{\Sigma}_{bb}^{-1}$$

$$\boldsymbol{\Sigma}_{aa} = \mathbf{A}\boldsymbol{\Sigma}_{bb}\mathbf{A}^T + \boldsymbol{\Sigma}_w = \mathbf{A}\boldsymbol{\Sigma}_{bb}\mathbf{A}^T + \boldsymbol{\Sigma}_{a|b} = \boldsymbol{\Sigma}_{ab}\boldsymbol{\Sigma}_{bb}^{-1}\boldsymbol{\Sigma}_{ba} + \boldsymbol{\Sigma}_{a|b}$$

---

# Partitioned Gaussian Distributions (2)

---

$$p(\mathbf{x}) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

$$\mathbf{x} = \begin{pmatrix} \mathbf{x}_a \\ \mathbf{x}_b \end{pmatrix} \quad \boldsymbol{\mu} = \begin{pmatrix} \boldsymbol{\mu}_a \\ \boldsymbol{\mu}_b \end{pmatrix} \quad \boldsymbol{\Sigma} = \begin{pmatrix} \boldsymbol{\Sigma}_{aa} & \boldsymbol{\Sigma}_{ab} \\ \boldsymbol{\Sigma}_{ba} & \boldsymbol{\Sigma}_{bb} \end{pmatrix}$$

$$\boldsymbol{\mu}_{a|b} = \boldsymbol{\mu}_a + \boldsymbol{\Sigma}_{ab} \boldsymbol{\Sigma}_{bb}^{-1} (\mathbf{x}_b - \boldsymbol{\mu}_b)$$

$$\boldsymbol{\Sigma}_{a|b} = \boldsymbol{\Sigma}_{aa} - \boldsymbol{\Sigma}_{ab} \boldsymbol{\Sigma}_{bb}^{-1} \boldsymbol{\Sigma}_{ba}$$

---

# Inverse Covariance Matrix\*

---

$$\begin{aligned}
 -\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu}) = \\
 \boxed{-\frac{1}{2}(\mathbf{x}_a - \boldsymbol{\mu}_a)^T \boldsymbol{\Lambda}_{aa}(\mathbf{x}_a - \boldsymbol{\mu}_a)} - \frac{1}{2}(\mathbf{x}_a - \boldsymbol{\mu}_a)^T \boldsymbol{\Lambda}_{ab}(\mathbf{x}_b - \boldsymbol{\mu}_b) \\
 - \frac{1}{2}(\mathbf{x}_b - \boldsymbol{\mu}_b)^T \boldsymbol{\Lambda}_{ba}(\mathbf{x}_a - \boldsymbol{\mu}_a) - \frac{1}{2}(\mathbf{x}_b - \boldsymbol{\mu}_b)^T \boldsymbol{\Lambda}_{bb}(\mathbf{x}_b - \boldsymbol{\mu}_b).
 \end{aligned}$$

$$-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu}) = -\frac{1}{2}\mathbf{x}^T \boldsymbol{\Sigma}^{-1}\mathbf{x} + \mathbf{x}^T \boldsymbol{\Sigma}^{-1}\boldsymbol{\mu} + \text{const}$$

$$-\frac{1}{2}(\mathbf{x}_a - \boldsymbol{\mu}_{a|b})^T \boldsymbol{\Sigma}_{a|b}^{-1}(\mathbf{x}_a - \boldsymbol{\mu}_{a|b}) = \boxed{-\frac{1}{2}\mathbf{x}_a^T \boldsymbol{\Sigma}_{a|b}^{-1}\mathbf{x}_a} + \mathbf{x}_a^T \boldsymbol{\Sigma}_{a|b}^{-1}\boldsymbol{\mu}_{a|b} + \text{const.}$$

$$\Rightarrow \underbrace{\boldsymbol{\Sigma}_{a|b} = \boldsymbol{\Lambda}_{aa}^{-1}}_{-\frac{1}{2}\mathbf{x}_a^T * \mathbf{x}_a} \quad \underbrace{\boldsymbol{\Lambda}_{aa}\boldsymbol{\mu}_a - \boldsymbol{\Lambda}_{ab}(\mathbf{x}_b - \boldsymbol{\mu}_b)}_{\mathbf{x}_a^T *} = \boldsymbol{\Sigma}_{a|b}^{-1}\boldsymbol{\mu}_{a|b}$$


---



# Inverse Matrix Lemma\*

---

$$\begin{pmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{pmatrix}^{-1} = \begin{pmatrix} \mathbf{M} & -\mathbf{M}\mathbf{B}\mathbf{D}^{-1} \\ -\mathbf{D}^{-1}\mathbf{C}\mathbf{M} & \mathbf{D}^{-1} + \mathbf{D}^{-1}\mathbf{C}\mathbf{M}\mathbf{B}\mathbf{D}^{-1} \end{pmatrix}$$

$$\mathbf{M} = (\mathbf{A} - \mathbf{B}\mathbf{D}^{-1}\mathbf{C})^{-1}$$

---

# Inverse Covariance Matrix\*

---

$$\begin{pmatrix} \Sigma_{aa} & \Sigma_{ab} \\ \Sigma_{ba} & \Sigma_{bb} \end{pmatrix}^{-1} = \begin{pmatrix} \Lambda_{aa} & \Lambda_{ab} \\ \Lambda_{ba} & \Lambda_{bb} \end{pmatrix}$$

$$\Lambda_{aa} = (\Sigma_{aa} - \Sigma_{ab}\Sigma_{bb}^{-1}\Sigma_{ba})^{-1}$$

$$\Lambda_{ab} = -(\Sigma_{aa} - \Sigma_{ab}\Sigma_{bb}^{-1}\Sigma_{ba})^{-1}\Sigma_{ab}\Sigma_{bb}^{-1}$$

$$\Sigma_{a|b} = \Sigma_{aa} - \Sigma_{ab}\Sigma_{bb}^{-1}\Sigma_{ba}$$

---

# Partitioned Conditionals and Marginals\*

---

$$p(\mathbf{x}_a | \mathbf{x}_b) = \mathcal{N}(\mathbf{x}_a | \boldsymbol{\mu}_{a|b}, \boldsymbol{\Sigma}_{a|b})$$



$$\begin{aligned}\boldsymbol{\Sigma}_{a|b} &= \boldsymbol{\Lambda}_{aa}^{-1} = \boldsymbol{\Sigma}_{aa} - \boldsymbol{\Sigma}_{ab} \boldsymbol{\Sigma}_{bb}^{-1} \boldsymbol{\Sigma}_{ba} \\ \boldsymbol{\mu}_{a|b} &= \boldsymbol{\Sigma}_{a|b} \{ \boldsymbol{\Lambda}_{aa} \boldsymbol{\mu}_a - \boldsymbol{\Lambda}_{ab} (\mathbf{x}_b - \boldsymbol{\mu}_b) \} \\ &= \boldsymbol{\mu}_a - \boldsymbol{\Lambda}_{aa}^{-1} \boldsymbol{\Lambda}_{ab} (\mathbf{x}_b - \boldsymbol{\mu}_b) \\ &= \boldsymbol{\mu}_a + \boldsymbol{\Sigma}_{ab} \boldsymbol{\Sigma}_{bb}^{-1} (\mathbf{x}_b - \boldsymbol{\mu}_b)\end{aligned}$$

$$\begin{aligned}p(\mathbf{x}_a) &= \int p(\mathbf{x}_a, \mathbf{x}_b) d\mathbf{x}_b \\ &= \mathcal{N}(\mathbf{x}_a | \boldsymbol{\mu}_a, \boldsymbol{\Sigma}_{aa})\end{aligned}$$

---

# Partitioned Conditionals and Marginals

---



# Maximum Likelihood for the Gaussian (1)

---

Given i.i.d. data  $\mathbf{X} = (\mathbf{x}_1, \dots, \mathbf{x}_N)^T$ , the log likelihood function is given by

$$\ln p(\mathbf{X}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = -\frac{ND}{2} \ln(2\pi) - \frac{N}{2} \ln |\boldsymbol{\Sigma}| - \frac{1}{2} \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{x}_n - \boldsymbol{\mu})$$

Sufficient statistics

$$\sum_{n=1}^N \mathbf{x}_n$$

$$\sum_{n=1}^N \mathbf{x}_n \mathbf{x}_n^T$$

---

# Maximum Likelihood for the Gaussian (2)

---

Set the derivative of the log likelihood function to zero,

$$\frac{\partial}{\partial \boldsymbol{\mu}} \ln p(\mathbf{X} | \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \sum_{n=1}^N \boldsymbol{\Sigma}^{-1} (\mathbf{x}_n - \boldsymbol{\mu}) = 0$$

and solve to obtain

$$\boldsymbol{\mu}_{\text{ML}} = \frac{1}{N} \sum_{n=1}^N \mathbf{x}_n.$$

Similarly

$$\boldsymbol{\Sigma}_{\text{ML}} = \frac{1}{N} \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}})(\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}})^{\text{T}}.$$

---

# Maximum Likelihood for the Gaussian (3)

---

The trace is invariant under cyclic permutation of matrix product

$$\text{tr}\{ABC\} = \text{tr}\{CAB\} = \text{tr}\{BCA\}$$

as follows

$$x^T \Sigma^{-1} x = \text{tr}\{x^T \Sigma^{-1} x\} = \text{tr}\{x x^T \Sigma^{-1}\}$$

Besides

$$\frac{\partial \text{tr}\{AB\}}{\partial A} = B^T$$

then

$$\frac{\partial \text{tr}\{x x^T \Sigma^{-1}\}}{\partial \Sigma^{-1}} = x x^T$$

---

# Maximum Likelihood for the Gaussian (4)

---

Given  $\frac{\partial \ln |A|}{\partial A} = (A^{-1})^T = (A^T)^{-1}$

then  $\frac{\partial \ln |\Sigma^{-1}|}{\partial \Sigma^{-1}} = \Sigma$

As follows

$$\frac{\partial \ln p(X|\mu, \Sigma)}{\partial \Sigma^{-1}} = \frac{N}{2} \Sigma - \frac{1}{2} \sum_{n=1}^N (x_n - \mu)(x_n - \mu)^T = 0$$

---



# Maximum Likelihood for the Gaussian (5)

---

Under the true distribution

$$\begin{aligned}\mathbb{E}[\boldsymbol{\mu}_{\text{ML}}] &= \boldsymbol{\mu} \\ \mathbb{E}[\boldsymbol{\Sigma}_{\text{ML}}] &= \frac{N-1}{N} \boldsymbol{\Sigma}.\end{aligned}$$

Hence define

$$\tilde{\boldsymbol{\Sigma}} = \frac{1}{N-1} \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}})(\mathbf{x}_n - \boldsymbol{\mu}_{\text{ML}})^{\text{T}}.$$

# Sequential Estimation

---

Contribution of the  $N^{\text{th}}$  data point,  $\mathbf{x}_N$

$$\begin{aligned}\mu_{\text{ML}}^{(N)} &= \frac{1}{N} \sum_{n=1}^N \mathbf{x}_n \\&= \frac{1}{N} \mathbf{x}_N + \frac{1}{N} \sum_{n=1}^{N-1} \mathbf{x}_n \\&= \frac{1}{N} \mathbf{x}_N + \frac{N-1}{N} \mu_{\text{ML}}^{(N-1)} \\&= \underbrace{\mu_{\text{ML}}^{(N-1)}}_{\text{old estimate}} + \underbrace{\frac{1}{N}}_{\text{correction weight}} \underbrace{(\mathbf{x}_N - \mu_{\text{ML}}^{(N-1)})}_{\text{correction given } \mathbf{x}_N}\end{aligned}$$

---

# The Robbins-Monro Algorithm (1)\*

---

Consider  $\theta$  and  $z$  governed by  $p(z, \theta)$  and define the *regression function*


$$f(\theta) \equiv \mathbb{E}[z|\theta] = \int z p(z|\theta) \, dz$$

Seek  $\theta^*$  such that  $f(\theta^*) = 0$ .

---

# The Robbins-Monro Algorithm (2)\*

---



Assume we are given samples from  $p(z, \theta)$ , one at the time.

---

# The Robbins-Monro Algorithm (3)\*

---

Successive estimates of  $\theta^*$  are then given by

$$\theta^{(N)} = \theta^{(N-1)} - a_{N-1} z(\theta^{(N-1)}).$$

Conditions on  $a_N$  for convergence :

$$\lim_{N \rightarrow \infty} a_N = 0 \qquad \sum_{N=1}^{\infty} a_N = \infty \qquad \sum_{N=1}^{\infty} a_N^2 < \infty$$

# Robbins-Monro for Maximum Likelihood (1)\*

---

Regarding

$$-\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \frac{\partial}{\partial \theta} \ln p(x_n | \theta) = \mathbb{E}_x \left[ -\frac{\partial}{\partial \theta} \ln p(x | \theta) \right]$$

as a regression function, finding its root is equivalent to finding the maximum likelihood solution  $\theta_{\text{ML}}$ . Thus

$$\theta^{(N)} = \theta^{(N-1)} - a_{N-1} \frac{\partial}{\partial \theta^{(N-1)}} \left[ -\ln p(x_N | \theta^{(N-1)}) \right].$$

---

# Robbins-Monro for Maximum Likelihood (2)\*



---

Example: estimate the mean of a Gaussian.

$$\begin{aligned} z &= \frac{\partial}{\partial \mu_{\text{ML}}} [-\ln p(x|\mu_{\text{ML}}, \sigma^2)] \\ &= -\frac{1}{\sigma^2} (x - \mu_{\text{ML}}) \end{aligned}$$

The distribution of  $z$  is Gaussian with mean  $\mu - \mu_{\text{ML}}$ .

For the Robbins-Monro update equation,  $a_N = \sigma^2 / N$ .



SEQUENTIAL estimation

# Bayesian Inference for the Gaussian (1)

---

Assume  $\sigma^2$  is known. Given i.i.d. data

$\mathbf{x} = \{x_1, \dots, x_N\}$ , the likelihood function for  $\mu$  is given by

$$p(\mathbf{x}|\mu) = \prod_{n=1}^N p(x_n|\mu) = \frac{1}{(2\pi\sigma^2)^{N/2}} \exp \left\{ -\frac{1}{2\sigma^2} \sum_{n=1}^N (x_n - \mu)^2 \right\}.$$

This has a Gaussian shape as a function of  $\mu$   
(but it is *not* a distribution over  $\mu$ ).

---



# Bayesian Inference for the Gaussian (2)

---

Combined with a Gaussian prior over  $\mu$ ,

$$p(\mu) = \mathcal{N}(\mu | \mu_0, \sigma_0^2).$$

this gives the posterior

$$p(\mu | \mathbf{x}) \propto p(\mathbf{x} | \mu) p(\mu).$$

Completing the square over  $\mu$ , we see that

$$p(\mu | \mathbf{x}) = \mathcal{N}(\mu | \mu_N, \sigma_N^2)$$

---

# Bayesian Inference for the Gaussian (3)

---

... where

$$\mu_N = \frac{\sigma^2}{N\sigma_0^2 + \sigma^2}\mu_0 + \frac{N\sigma_0^2}{N\sigma_0^2 + \sigma^2}\mu_{\text{ML}}, \quad \mu_{\text{ML}} = \frac{1}{N} \sum_{n=1}^N x_n$$

$$\frac{1}{\sigma_N^2} = \frac{1}{\sigma_0^2} + \frac{N}{\sigma^2}.$$


Note:

	$N = 0$	$N \rightarrow \infty$
$\mu_N$	$\mu_0$	$\mu_{\text{ML}}$
$\sigma_N^2$	$\sigma_0^2$	0

# Bayesian Inference for the Gaussian (4)

---

Example:  $p(\mu|\mathbf{x}) = \mathcal{N}(\mu|\mu_N, \sigma_N^2)$  for  $N = 0, 1, 2$  and 10.



# Bayesian Inference for the Gaussian (5)

---

## Sequential Estimation

$$\begin{aligned} p(\mu|\mathbf{x}) &\propto p(\mu)p(\mathbf{x}|\mu) \\ &= \left[ p(\mu) \prod_{n=1}^{N-1} p(x_n|\mu) \right] p(x_N|\mu) \\ &\propto \mathcal{N}(\mu|\mu_{N-1}, \sigma_{N-1}^2) p(x_N|\mu) \end{aligned}$$

The posterior obtained after observing  $N - 1$  data points becomes the prior when we observe the  $N^{\text{th}}$  data point.

---

# Bayesian Inference for the Gaussian (6)

---

Now assume  $\mu$  is known. The likelihood function for  $\lambda = 1/\sigma^2$  is given by

$$p(\mathbf{x}|\lambda) = \prod_{n=1}^N \mathcal{N}(x_n|\mu, \lambda^{-1}) \propto \lambda^{N/2} \exp \left\{ -\frac{\lambda}{2} \sum_{n=1}^N (x_n - \mu)^2 \right\}.$$

This has a Gamma shape as a function of  $\lambda$ .

---




# Bayesian Inference for the Gaussian (7)

---

## The Gamma distribution

$$\text{Gam}(\lambda|a, b) = \frac{1}{\Gamma(a)} b^a \lambda^{a-1} \exp(-b\lambda)$$

$$\mathbb{E}[\lambda] = \frac{a}{b} \qquad \text{var}[\lambda] = \frac{a}{b^2}$$



# Bayesian Inference for the Gaussian (8)

---

Now we combine a Gamma prior,  $\text{Gam}(\lambda|a_0, b_0)$ , with the likelihood function for  $\lambda$  to obtain

$$p(\lambda|\mathbf{x}) \propto \lambda^{a_0-1} \lambda^{N/2} \exp \left\{ -b_0 \lambda - \frac{\lambda}{2} \sum_{n=1}^N (x_n - \mu)^2 \right\}$$

which we recognize as  $\text{Gam}(\lambda|a_N, b_N)$  with

$$\begin{aligned} a_N &= a_0 + \frac{N}{2} \\ b_N &= b_0 + \frac{1}{2} \sum_{n=1}^N (x_n - \mu)^2 = b_0 + \frac{N}{2} \sigma_{\text{ML}}^2. \end{aligned}$$

---

# Bayesian Inference for the Gaussian (9)

---

If both  $\mu$  and  $\lambda$  are unknown, the joint likelihood function is given by

$$\begin{aligned} p(\mathbf{x}|\mu, \lambda) &= \prod_{n=1}^N \left( \frac{\lambda}{2\pi} \right)^{1/2} \exp \left\{ -\frac{\lambda}{2} (x_n - \mu)^2 \right\} \\ &\propto \left[ \lambda^{1/2} \exp \left( -\frac{\lambda \mu^2}{2} \right) \right]^N \exp \left\{ \lambda \mu \sum_{n=1}^N x_n - \frac{\lambda}{2} \sum_{n=1}^N x_n^2 \right\}. \end{aligned}$$

We need a prior with the same functional dependence on  $\mu$  and  $\lambda$ .

---



# Bayesian Inference for the Gaussian (10)

---

The Gaussian-gamma distribution prior

$$\begin{aligned} p(\mu, \lambda) &\propto \left[ \lambda^{1/2} \exp \left( -\frac{\lambda \mu^2}{2} \right) \right]^\beta \exp \{ c \lambda \mu - d \lambda \} \\ &= \exp \left\{ -\frac{\beta \lambda}{2} (\mu - c/\beta)^2 \right\} \lambda^{\beta/2} \exp \left\{ - \left( d - \frac{c^2}{2\beta} \right) \lambda \right\} \end{aligned}$$

Then the posterior is given by

$$\beta_N = \beta + N \quad c_N = c + \sum_{n=1}^N x_N \quad d_N = d + \frac{1}{2} \sum_{n=1}^N x_N^2$$

---

# Bayesian Inference for the Gaussian (11)

---

## The Gaussian-gamma distribution

$$p(\mu, \lambda) = \mathcal{N}(\mu|\mu_0, (\beta\lambda)^{-1})\text{Gam}(\lambda|a, b)$$
$$\propto \underbrace{\exp\left\{-\frac{\beta\lambda}{2}(\mu - \mu_0)^2\right\}}_{\text{Quadratic in } \mu} \underbrace{\lambda^{a-1} \exp\{-b\lambda\}}_{\text{Gamma distribution over } \lambda}$$

- Quadratic in  $\mu$ .
- Linear in  $\lambda$ .
- Gamma distribution over  $\lambda$ .
- Independent of  $\mu$ .

$$\mu_0 = c/\beta$$

$$a = 1 + \beta/2$$


$$b = d - c^2/2\beta$$

---

# Bayesian Inference for the Gaussian (12)

---

## The Gaussian-gamma distribution



# Bayesian Inference for the Gaussian (13)\*

---

Multivariate conjugate priors

- $\boldsymbol{\mu}$  unknown,  $\boldsymbol{\Lambda}$  known:  $p(\boldsymbol{\mu})$  Gaussian.
- $\boldsymbol{\Lambda}$  unknown,  $\boldsymbol{\mu}$  known:  $p(\boldsymbol{\Lambda})$  Wishart,


$$\mathcal{W}(\boldsymbol{\Lambda}|\mathbf{W}, \nu) = B|\boldsymbol{\Lambda}|^{(\nu-D-1)/2} \exp\left(-\frac{1}{2}\text{Tr}(\mathbf{W}^{-1}\boldsymbol{\Lambda})\right).$$

- $\boldsymbol{\Lambda}$  and  $\boldsymbol{\mu}$  unknown:  $p(\boldsymbol{\mu}, \boldsymbol{\Lambda})$  Gaussian-Wishart,  $p(\boldsymbol{\mu}, \boldsymbol{\Lambda}|\boldsymbol{\mu}_0, \beta, \mathbf{W}, \nu) = \mathcal{N}(\boldsymbol{\mu}|\boldsymbol{\mu}_0, (\beta\boldsymbol{\Lambda})^{-1}) \mathcal{W}(\boldsymbol{\Lambda}|\mathbf{W}, \nu)$
-


# Mixtures of Gaussians (1)

---

Old Faithful data set



Single Gaussian



Mixture of two Gaussians

# Mixtures of Gaussians (2)


---

Combine simple models  
into a complex model:

$$p(\mathbf{x}) = \sum_{k=1}^K \pi_k \underbrace{\mathcal{N}(\mathbf{x} | \boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}_{\text{Component}}$$




Mixing coefficient

$$\forall k : \pi_k \geq 0 \quad \sum_{k=1}^K \pi_k = 1$$



# Mixtures of Gaussians (3)

---



# Mixtures of Gaussians (4)

---

Determining parameters  $\mu$ ,  $\Sigma$ , and  $\pi$  using maximum log likelihood

$$\ln p(\mathbf{X}|\pi, \mu, \Sigma) = \sum_{n=1}^N \ln \left\{ \underbrace{\sum_{k=1}^K \pi_k \mathcal{N}(\mathbf{x}_n | \mu_k, \Sigma_k)} \right\}$$

Log of a sum; no closed form maximum.

Solution: use standard, iterative, numeric optimization methods or the *expectation maximization* algorithm (Chapter 9).

---



# Mixtures of Gaussians (5)

---

The posterior probability of each data point being responsible for each cluster

$$\begin{aligned}\gamma_k(\mathbf{x}) &\equiv p(k|\mathbf{x}) \\ &= \frac{p(k)p(\mathbf{x}|k)}{\sum_l p(l)p(\mathbf{x}|l)} \\ &= \frac{\pi_k \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}_k, \boldsymbol{\Sigma}_k)}{\sum_l \pi_l \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}_l, \boldsymbol{\Sigma}_l)}\end{aligned}$$

---

# Student's t-Distribution\*

---

$$\begin{aligned} p(x|\mu, a, b) &= \int_0^\infty \mathcal{N}(x|\mu, \tau^{-1}) \text{Gam}(\tau|a, b) \, d\tau \\ &= \int_0^\infty \mathcal{N}(x|\mu, (\eta\lambda)^{-1}) \text{Gam}(\eta|\nu/2, \nu/2) \, d\eta \quad \leftarrow \\ &= \frac{\Gamma(\nu/2 + 1/2)}{\Gamma(\nu/2)} \left( \frac{\lambda}{\pi\nu} \right)^{1/2} \left[ 1 + \frac{\lambda(x - \mu)^2}{\nu} \right]^{-\nu/2 - 1/2} \\ &= \text{St}(x|\mu, \lambda, \nu) \end{aligned}$$

where


$$\lambda = a/b \qquad \eta = \tau b/a \qquad \nu = 2a.$$

Infinite mixture of Gaussians.

---

# Student's t-Distribution\*

---





	$\nu = 1$	$\nu \rightarrow \infty$
$\text{St}(x \mu, \lambda, \nu)$	Cauchy	$\mathcal{N}(x \mu, \lambda^{-1})$

---

# Student's t-Distribution\*

---

Robustness to outliers: Gaussian vs t-distribution.



# Student's t-Distribution\*

---

The  $D$ -variate case:

$$\begin{aligned}\text{St}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Lambda}, \nu) &= \int_0^\infty \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, (\eta\boldsymbol{\Lambda})^{-1}) \text{Gam}(\eta|\nu/2, \nu/2) \, d\eta \\ &= \frac{\Gamma(D/2 + \nu/2)}{\Gamma(\nu/2)} \frac{|\boldsymbol{\Lambda}|^{1/2}}{(\pi\nu)^{D/2}} \left[ 1 + \frac{\Delta^2}{\nu} \right]^{-D/2-\nu/2}\end{aligned}$$

where  $\Delta^2 = (\mathbf{x} - \boldsymbol{\mu})^\text{T} \boldsymbol{\Lambda} (\mathbf{x} - \boldsymbol{\mu})$ .

Properties:

$$\begin{aligned}\mathbb{E}[\mathbf{x}] &= \boldsymbol{\mu}, & \text{if } \nu > 1 \\ \text{cov}[\mathbf{x}] &= \frac{\nu}{(\nu - 2)} \boldsymbol{\Lambda}^{-1}, & \text{if } \nu > 2 \\ \text{mode}[\mathbf{x}] &= \boldsymbol{\mu}\end{aligned}$$

---

# Periodic variables\*

---

- Examples: calendar time, direction, ...
- We require

$$\begin{aligned}p(\theta) &\geq 0 \\ \int_0^{2\pi} p(\theta) \, d\theta &= 1 \\ p(\theta + 2\pi) &= p(\theta).\end{aligned}$$

# von Mises Distribution (1)\*

---

This requirement is satisfied by

$$p(\theta|\theta_0, m) = \frac{1}{2\pi I_0(m)} \exp \{m \cos(\theta - \theta_0)\}$$

where



$$I_0(m) = \frac{1}{2\pi} \int_0^{2\pi} \exp \{m \cos \theta\} d\theta$$

is the 0<sup>th</sup> order modified Bessel function of the 1<sup>st</sup> kind.

---

# von Mises Distribution (2)\*

---





# Maximum Likelihood for von Mises\*

---

Given a data set,  $\mathcal{D} = \{\theta_1, \dots, \theta_N\}$ , the log likelihood function is given by

$$\ln p(\mathcal{D}|\theta_0, m) = -N \ln(2\pi) - N \ln I_0(m) + m \sum_{n=1}^N \cos(\theta_n - \theta_0).$$

Maximizing with respect to  $\theta_0$  we directly obtain

$$\theta_0^{\text{ML}} = \tan^{-1} \left\{ \frac{\sum_n \sin \theta_n}{\sum_n \cos \theta_n} \right\}.$$

Similarly, maximizing with respect to  $m$  we get

$$\frac{I_1(m_{\text{ML}})}{I_0(m_{\text{ML}})} = \frac{1}{N} \sum_{n=1}^N \cos(\theta_n - \theta_0^{\text{ML}})$$

which can be solved numerically for  $m_{\text{ML}}$ .

---