



CS215 DISCRETE MATH

Dr. QI WANG

Department of Computer Science and Engineering

Office: Room413, CoE South Tower

Email: wangqi@sustech.edu.cn

Application of Number Theory

- G. H. Hardy (1877 - 1947)

In his 1940 autobiography *A Mathematician's Apology*, Hardy wrote “The great modern achievements of applied mathematics have been in relativity and quantum mechanics, and these subjects are, at present, **almost as ‘useless’ as the theory of numbers.**”



Application of Number Theory

- G. H. Hardy (1877 - 1947)

In his 1940 autobiography *A Mathematician's Apology*, Hardy wrote “The great modern achievements of applied mathematics have been in relativity and quantum mechanics, and these subjects are, at present, **almost as ‘useless’ as the theory of numbers.**”



If he could see the world now, Hardy would be spinning in his grave.

Number Theory

- *Number theory* is a branch of mathematics that explores integers and their properties, is the basis of **cryptography**, **coding theory**, **computer security**, **e-commerce**, etc.



Number Theory

- *Number theory* is a branch of mathematics that explores integers and their properties, is the basis of **cryptography**, **coding theory**, **computer security**, **e-commerce**, etc.
- At one point, the largest employer of mathematicians in the United States, and probably the world, was the **National Security Agency** (NSA). The NSA is the largest spy agency in the US (bigger than CIA, Central Intelligence Agency), and has the responsibility for code design and breaking.



Division

- If a and b are integers with $a \neq 0$, we say that a divides b if there is an integer k such that $b = ak$, or equivalently b/a is an integer. In this case, we say that a is a *factor* or *divisor* of b , and b is a *multiple* of a . (We use the notations $a|b$, $a \nmid b$)



Division

- If a and b are integers with $a \neq 0$, we say that a divides b if there is an integer k such that $b = ak$, or equivalently b/a is an integer. In this case, we say that a is a *factor* or *divisor* of b , and b is a *multiple* of a . (We use the notations $a|b$, $a \nmid b$)

Example

◇ $4 \mid 24$

◇ $3 \nmid 7$



Divisibility

- All integers divisible by $d > 0$ can be enumerated as:
 $\dots, -kd, \dots, -2d, -d, 0, d, 2d, \dots, kd, \dots$



Divisibility

- **All integers divisible by $d > 0$** can be **enumerated** as:
 $\dots, -kd, \dots, -2d, -d, 0, d, 2d, \dots, kd, \dots$
- **Question:** Let n and d be two positive integers. How many positive integers **not exceeding n** are divisible by d ?



Divisibility

- **All integers divisible by $d > 0$** can be **enumerated** as:
 $\dots, -kd, \dots, -2d, -d, 0, d, 2d, \dots, kd, \dots$
- **Question:** Let n and d be two positive integers. How many positive integers **not exceeding n** are divisible by d ?

Answer: Count the number of integers such that $0 < kd \leq n$. Therefore, there are $\lfloor n/d \rfloor$ such positive integers.



Divisibility

■ Properties

Let a, b, c be integers. Then the following hold:

- (i) if $a|b$ and $a|c$, then $a|(b + c)$
- (ii) if $a|b$ then $a|bc$ for all integers c
- iii) if $a|b$ and $b|c$, then $a|c$



Divisibility

■ Properties

Let a, b, c be integers. Then the following hold:

- (i) if $a|b$ and $a|c$, then $a|(b + c)$
- (ii) if $a|b$ then $a|bc$ for all integers c
- iii) if $a|b$ and $b|c$, then $a|c$

Proof.



Divisibility

- **Corollary** If a, b, c are integers, where $a \neq 0$, such that $a|b$ and $a|c$, then $a|(mb + nc)$ whenever m and n are integers.



Divisibility

- **Corollary** If a, b, c are integers, where $a \neq 0$, such that $a|b$ and $a|c$, then $a|(mb + nc)$ whenever m and n are integers.

Proof. By part (ii) and part (i) of Properties.

The Division Algorithm

- If a is an integer and d a positive integer, then there are **unique** integers q and r , with $0 \leq r < d$, such that $a = dq + r$. In this case, d is called the *divisor*, a is called the *dividend*, q is called the *quotient*, and r is called the *remainder*.



The Division Algorithm

- If a is an integer and d a positive integer, then there are **unique** integers q and r , with $0 \leq r < d$, such that $a = dq + r$. In this case, d is called the *divisor*, a is called the *dividend*, q is called the *quotient*, and r is called the *remainder*.

In this case, we use the notations $q = a \text{ div } d$ and $r = a \bmod d$.



Congruence Relation

- If a and b are integers and m is a positive integer, then a is *congruent to b modulo m if m divides $a - b$* , denoted by $a \equiv b \pmod{m}$. This is called *congruence* and m is its *modulus*.



Congruence Relation

- If a and b are integers and m is a positive integer, then a is *congruent to b modulo m if m divides $a - b$* , denoted by $a \equiv b \pmod{m}$. This is called *congruence* and m is its *modulus*.

Example

- ◇ $15 \equiv 3 \pmod{6}$
- ◇ $-1 \equiv 11 \pmod{6}$



More on Congruences

- Let m be a positive integer. The integers a and b are congruent modulo m if and only if there is an integer k such that $a = b + km$.



More on Congruences

- Let m be a positive integer. The integers a and b are congruent modulo m if and only if there is an integer k such that $a = b + km$.

Proof.

“only if” part

“if” part



$(\text{mod } m)$ and $\text{mod } m$ Notations

- $a \equiv b \pmod{m}$ and $a \bmod m = b$ are different.
 - ◇ $a \equiv b \pmod{m}$ is a **relation** on the set of integers
 - ◇ In $a \bmod m = b$, the notation **mod** denotes a function



$(\text{mod } m)$ and $\text{mod } m$ Notations

- $a \equiv b \pmod{m}$ and $a \bmod m = b$ are different.
 - ◇ $a \equiv b \pmod{m}$ is a **relation** on the set of integers
 - ◇ In $a \bmod m = b$, the notation **mod** denotes a function
- Let a and b be integers, and let m be a positive integer.
Then $a \equiv b \bmod m$ if and only if $a \bmod m = b \bmod m$



$(\bmod m)$ and $\bmod m$ Notations

- $a \equiv b \pmod{m}$ and $a \bmod m = b$ are different.
 - ◇ $a \equiv b \pmod{m}$ is a **relation** on the set of integers
 - ◇ In $a \bmod m = b$, the notation **mod** denotes a **function**
- Let a and b be integers, and let m be a positive integer.
Then $a \equiv b \bmod m$ if and only if $a \bmod m = b \bmod m$

Proof.



Congruences of Sums and Products

- Let m be a positive integer. If $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$, then $a + c \equiv b + d \pmod{m}$ and $ac \equiv bd \pmod{m}$



Congruences of Sums and Products

- Let m be a positive integer. If $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$, then $a + c \equiv b + d \pmod{m}$ and $ac \equiv bd \pmod{m}$

Proof.

Algebraic Manipulation of Congruences

- If $a \equiv b \pmod{m}$, then
 - $c \cdot a \equiv c \cdot b \pmod{m}$?
 - $c + a \equiv c + b \pmod{m}$?
 - $a/c \equiv b/c \pmod{m}$?



Algebraic Manipulation of Congruences

- If $a \equiv b \pmod{m}$, then
 - $c \cdot a \equiv c \cdot b \pmod{m}$?
 - $c + a \equiv c + b \pmod{m}$?
 - $a/c \equiv b/c \pmod{m}$?

$$14 \equiv 8 \pmod{6} \text{ but } 7 \not\equiv 4 \pmod{6}$$



Computing the mod Function

- **Corollary** Let m be a positive integer and let a and b be integers. Then

$$(a + b) \bmod m = ((a \bmod m) + (b \bmod m)) \bmod m$$

$$ab \bmod m = ((a \bmod m)(b \bmod m)) \bmod m$$



Computing the mod Function

- **Corollary** Let m be a positive integer and let a and b be integers. Then

$$(a + b) \bmod m = ((a \bmod m) + (b \bmod m)) \bmod m$$
$$ab \bmod m = ((a \bmod m)(b \bmod m)) \bmod m$$

Proof.



Arithmetic Modulo m

- Let \mathbf{Z}_m be the set of nonnegative integers less than m : $\{0, 1, \dots, m - 1\}$.



Arithmetic Modulo m

- Let \mathbf{Z}_m be the set of nonnegative integers less than m : $\{0, 1, \dots, m - 1\}$.

$$+_m : a +_m b = (a + b) \bmod m$$

$$\cdot_m : a \cdot_m b = ab \bmod m$$



Arithmetic Modulo m

- Let \mathbf{Z}_m be the set of nonnegative integers less than m : $\{0, 1, \dots, m - 1\}$.

$$+_m : a +_m b = (a + b) \bmod m$$

$$\cdot_m : a \cdot_m b = ab \bmod m$$

Example

$$\diamond 7 +_{11} 9 = ?$$

$$\diamond 7 \cdot_{11} 9 = ?$$



Arithmetic Modulo m

- **Closure:** if $a, b \in \mathbf{Z}_m$, then $a +_m b, a \cdot_m b \in \mathbf{Z}_m$



Arithmetic Modulo m

- **Closure**: if $a, b \in \mathbf{Z}_m$, then $a +_m b, a \cdot_m b \in \mathbf{Z}_m$
- **Associativity**: if $a, b, c \in \mathbf{Z}_m$, then
 $(a +_m b) +_m c = a +_m (b +_m c)$ and
 $(a \cdot_m b) \cdot_m c = a \cdot_m (b \cdot_m c)$



Arithmetic Modulo m

- **Closure:** if $a, b \in \mathbf{Z}_m$, then $a +_m b, a \cdot_m b \in \mathbf{Z}_m$
- **Associativity:** if $a, b, c \in \mathbf{Z}_m$, then
 $(a +_m b) +_m c = a +_m (b +_m c)$ and
 $(a \cdot_m b) \cdot_m c = a \cdot_m (b \cdot_m c)$
- **Identity elements:** $a +_m 0 = a$ and $a \cdot_m 1 = a$



Arithmetic Modulo m

- **Closure:** if $a, b \in \mathbf{Z}_m$, then $a +_m b, a \cdot_m b \in \mathbf{Z}_m$
- **Associativity:** if $a, b, c \in \mathbf{Z}_m$, then
 $(a +_m b) +_m c = a +_m (b +_m c)$ and
 $(a \cdot_m b) \cdot_m c = a \cdot_m (b \cdot_m c)$
- **Identity elements:** $a +_m 0 = a$ and $a \cdot_m 1 = a$
- **Additive inverses:** if $a \neq 0$ and $a \in \mathbf{Z}_m$, then $m - a$ is an additive inverse of a modulo m



Arithmetic Modulo m

- **Closure**: if $a, b \in \mathbf{Z}_m$, then $a +_m b, a \cdot_m b \in \mathbf{Z}_m$
- **Associativity**: if $a, b, c \in \mathbf{Z}_m$, then
 $(a +_m b) +_m c = a +_m (b +_m c)$ and
 $(a \cdot_m b) \cdot_m c = a \cdot_m (b \cdot_m c)$
- **Identity elements**: $a +_m 0 = a$ and $a \cdot_m 1 = a$
- **Additive inverses**: if $a \neq 0$ and $a \in \mathbf{Z}_m$, then $m - a$ is an additive inverse of a modulo m
- **Commutativity**: if $a, b \in \mathbf{Z}_m$, then $a +_m b = b +_m a$



Arithmetic Modulo m

- **Closure**: if $a, b \in \mathbf{Z}_m$, then $a +_m b, a \cdot_m b \in \mathbf{Z}_m$
- **Associativity**: if $a, b, c \in \mathbf{Z}_m$, then
 $(a +_m b) +_m c = a +_m (b +_m c)$ and
 $(a \cdot_m b) \cdot_m c = a \cdot_m (b \cdot_m c)$
- **Identity elements**: $a +_m 0 = a$ and $a \cdot_m 1 = a$
- **Additive inverses**: if $a \neq 0$ and $a \in \mathbf{Z}_m$, then $m - a$ is an additive inverse of a modulo m
- **Commutativity**: if $a, b \in \mathbf{Z}_m$, then $a +_m b = b +_m a$
- **Distributivity**: if $a, b, c \in \mathbf{Z}_m$, then
 $a \cdot_m (b +_m c) = (a \cdot_m b) +_m (a \cdot_m c)$ and
 $(a +_m b) \cdot_m c = (a \cdot_m c) +_m (b \cdot_m c)$



Group

- A set G of elements, along with a **binary operation** \star , must satisfy the following four properties to be called a *group*.



Group

- A set G of elements, along with a **binary operation** \star , must satisfy the following four properties to be called a **group**.

Closure: If $a, b \in G$, then $a \star b = c$ also belongs to G .

Associativity: $(a \star b) \star c = a \star (b \star c)$

Identity element: There is a **unique** element 1_e , such that for every $a \in G$, we have $a \star 1_e = a$.

Inverse: For every $a \in G$, there exists an element, denoted by a^{-1} , such that $a \star a^{-1} = 1_e$.



Group

- A set G of elements, along with a **binary operation** \star , must satisfy the following four properties to be called a **group**.

Closure: If $a, b \in G$, then $a \star b = c$ also belongs to G .

Associativity: $(a \star b) \star c = a \star (b \star c)$

Identity element: There is a **unique** element 1_e , such that for every $a \in G$, we have $a \star 1_e = a$.

Inverse: For every $a \in G$, there exists an element, denoted by a^{-1} , such that $a \star a^{-1} = 1_e$.

Example:

$(\mathbb{Z}, +)$, $(\mathbb{Q}, +)$, $(\mathbb{R}, +)$, $(\mathbb{M}_{n \times n}, +)$?

(\mathbb{Z}^*, \times) , (\mathbb{Q}^*, \times) , (\mathbb{R}^*, \times) , $(\mathbb{M}_{n \times n}^*, \cdot)$?



Permutation Group

- Let $s_n = \langle 1, 2, \dots, n \rangle$ denote a *sequence* of integers 1 through n . Denote by P_n the set of all *permutations* of the sequence s_n .



Permutation Group

- Let $s_n = \langle 1, 2, \dots, n \rangle$ denote a *sequence* of integers 1 through n . Denote by P_n the set of all *permutations* of the sequence s_n .

For example, $s_3 = \langle 1, 2, 3 \rangle$

$$P_3 = \{ \langle 1, 2, 3 \rangle, \langle 1, 3, 2 \rangle, \langle 2, 1, 3 \rangle, \langle 2, 3, 1 \rangle, \langle 3, 1, 2 \rangle, \langle 3, 2, 1 \rangle \}$$



Permutation Group

- Let $s_n = \langle 1, 2, \dots, n \rangle$ denote a *sequence* of integers 1 through n . Denote by P_n the set of all *permutations* of the sequence s_n .

For example, $s_3 = \langle 1, 2, 3 \rangle$

$$P_3 = \{ \langle 1, 2, 3 \rangle, \langle 1, 3, 2 \rangle, \langle 2, 1, 3 \rangle, \langle 2, 3, 1 \rangle, \langle 3, 1, 2 \rangle, \langle 3, 2, 1 \rangle \}$$

- Define a binary operation \circ on the elements of P_n :
for $\rho, \pi \in P_n$, $\pi \circ \rho$ denotes a *re-permutation* of the elements of ρ according to the elements of π .



Permutation Group

- Consider $s_3 = \langle 1, 2, 3 \rangle$, and
 $P_3 = \{ \langle p_1, p_2, p_3 \rangle \mid p_1, p_2, p_3 \in s_3 \text{ with } p_1 \neq p_2 \neq p_3 \}$.



Permutation Group

- Consider $s_3 = \langle 1, 2, 3 \rangle$, and
 $P_3 = \{ \langle p_1, p_2, p_3 \rangle \mid p_1, p_2, p_3 \in s_3 \text{ with } p_1 \neq p_2 \neq p_3 \}$.
- $\pi = \langle 3, 2, 1 \rangle$, $\rho = \langle 1, 3, 2 \rangle$, what is $\pi \circ \rho$?



Permutation Group

- Consider $s_3 = \langle 1, 2, 3 \rangle$, and
 $P_3 = \{ \langle p_1, p_2, p_3 \rangle \mid p_1, p_2, p_3 \in s_3 \text{ with } p_1 \neq p_2 \neq p_3 \}$.
- $\pi = \langle 3, 2, 1 \rangle$, $\rho = \langle 1, 3, 2 \rangle$, what is $\pi \circ \rho$?
 $\pi \circ \rho = \langle 2, 3, 1 \rangle \in P_3$



Permutation Group

- Consider $s_3 = \langle 1, 2, 3 \rangle$, and $P_3 = \{ \langle p_1, p_2, p_3 \rangle \mid p_1, p_2, p_3 \in s_3 \text{ with } p_1 \neq p_2 \neq p_3 \}$.
- $\pi = \langle 3, 2, 1 \rangle$, $\rho = \langle 1, 3, 2 \rangle$, what is $\pi \circ \rho$?
 $\pi \circ \rho = \langle 2, 3, 1 \rangle \in P_3$
- We can verify the other three properties.
$$\rho_1 \circ (\rho_2 \circ \rho_3) = (\rho_1 \circ \rho_2) \circ \rho_3$$
$$\langle 1, 2, 3 \rangle \circ \rho = \rho \circ \langle 1, 2, 3 \rangle = \rho$$

For each $\rho \in P_3$, there exists another unique $\pi \in P_3$ such that

$$\rho \circ \pi = \pi \circ \rho = \langle 1, 2, 3 \rangle$$



Permutation Group

- Consider $s_3 = \langle 1, 2, 3 \rangle$, and
 $P_3 = \{ \langle p_1, p_2, p_3 \rangle \mid p_1, p_2, p_3 \in s_3 \text{ with } p_1 \neq p_2 \neq p_3 \}$.
- $\pi = \langle 3, 2, 1 \rangle$, $\rho = \langle 1, 3, 2 \rangle$, what is $\pi \circ \rho$?
 $\pi \circ \rho = \langle 2, 3, 1 \rangle \in P_3$
- We can verify the other three properties.
$$\rho_1 \circ (\rho_2 \circ \rho_3) = (\rho_1 \circ \rho_2) \circ \rho_3$$
$$\langle 1, 2, 3 \rangle \circ \rho = \rho \circ \langle 1, 2, 3 \rangle = \rho$$

For each $\rho \in P_3$, there exists another unique $\pi \in P_3$ such that

$$\rho \circ \pi = \pi \circ \rho = \langle 1, 2, 3 \rangle$$

(P_n, \circ) is called a *permutation group*.



Abelian Group

- If the operation on the set elements is *commutative*, the group is called an *abelian group*. ($a \star b = b \star a$)



Abelian Group

- If the operation on the set elements is *commutative*, the group is called an *abelian group*. ($a \star b = b \star a$)

- **Example**

$(\mathbb{Z}, +)$, $(\mathbb{Q}, +)$, $(\mathbb{R}, +)$, $(\mathbb{M}_{n \times n}, +)$?

$(GL(n), \cdot)$, (P_n, \circ) ?



Abelian Group

- If the operation on the set elements is *commutative*, the group is called an *abelian group*. ($a \star b = b \star a$)

- **Example**

$$(\mathbb{Z}, +), (\mathbb{Q}, +), (\mathbb{R}, +), (\mathbb{M}_{n \times n}, +) ?$$

$$(GL(n), \cdot), (P_n, \circ) ?$$

- If the group operation is referred to as *addition* (*multiplication*), then the group also allows for *subtraction* (*division*).

$$a - b = a + (-b)$$

$$a/b = a \cdot b^{-1}$$



Ring

- If $(R, +)$ is an *abelian group*, we define one more operation (denoted as *multiplication* \times for convenience) to have a *ring* $(R, +, \times)$ satisfying the following properties.



Ring

- If $(R, +)$ is an *abelian group*, we define one more operation (denoted as *multiplication* \times for convenience) to have a *ring* $(R, +, \times)$ satisfying the following properties.

Closure: R must be closed w.r.t. \times

Associativity: $(a \times b) \times c = a \times (b \times c)$

Distributivity: $a \times (b + c) = a \times b + a \times c$
 $(a + b) \times c = a \times c + b \times c$



Ring

- If $(R, +)$ is an *abelian group*, we define one more operation (denoted as *multiplication* \times for convenience) to have a *ring* $(R, +, \times)$ satisfying the following properties.

Closure: R must be closed w.r.t. \times

Associativity: $(a \times b) \times c = a \times (b \times c)$

Distributivity: $a \times (b + c) = a \times b + a \times c$
 $(a + b) \times c = a \times c + b \times c$

Example:

$(\mathbb{Z}, +, \times)$, $(\mathbb{Q}, +, \times)$, $(\mathbb{R}, +, \times)$, $(\mathbb{M}_{n \times n}, +, \cdot)$?



Commutative Ring, Integral Domain

- A *ring* is *commutative* if the **multiplication operation** is *commutative* for all elements in the ring. ($ab = ba$)



Commutative Ring, Integral Domain

- A *ring* is *commutative* if the **multiplication operation** is *commutative* for all elements in the ring. ($ab = ba$)
- An *integral domain* $(R, +, \times)$ is a *commutative ring* that satisfies the following two additional properties.

Identity element for multiplication: $a1 = 1a = a$

Nonzero product for any two nonzero elements:
if $ab = 0$, then either a or b **must be** 0.



Commutative Ring, Integral Domain

- A *ring* is *commutative* if the **multiplication operation** is *commutative* for all elements in the ring. ($ab = ba$)
- An *integral domain* $(R, +, \times)$ is a *commutative ring* that satisfies the following two additional properties.

Identity element for multiplication: $a1 = 1a = a$

Nonzero product for any two nonzero elements:

if $ab = 0$, then either a or b **must be** 0.

Example:

$(\mathbb{Z}, +, \times)$, $(\mathbb{Q}, +, \times)$, $(\mathbb{R}, +, \times)$?

$(\mathbb{Z}_m, +, \times)$, $(M_{n \times n}, +, \cdot)$?



Field

- A *field*, denoted by $(F, +, \times)$, is an *integral domain* whose elements satisfy the following additional property.

Inverse for multiplication: For every $a \in F$, there exists an element b , denoted by a^{-1} , such that $ab = ba = 1$.

Field

- A *field*, denoted by $(F, +, \times)$, is an *integral domain* whose elements satisfy the following additional property.

Inverse for multiplication: For every $a \in F$, there exists an element b , denoted by a^{-1} , such that $ab = ba = 1$.

- **Example:**

$(\mathbb{Z}, +, \times)$, $(\mathbb{Q}, +, \times)$, $(\mathbb{R}, +, \times)$?

$(\mathbb{Z}_p, +, \times)$?



Representations of Integers

- We may use *decimal* (*base 10*) or *binary* or *octal* or *hexadecimal* or other notations to represent integers.



Representations of Integers

- We may use *decimal* (*base 10*) or *binary* or *octal* or *hexadecimal* or other notations to represent integers.
- Let $b > 1$ be an integer. Then if n is a positive integer, it can be expressed **uniquely in the form**
$$n = a_k b^k + a_{k-1} b^{k-1} + \dots + a_1 b + a_0,$$
 where k is nonnegative, a_i 's are nonnegative integers less than b . The representation of n is called *the base- b expansion of n* and is denoted by $(a_k a_{k-1} \dots a_1 a_0)_b$.



Base- b Expansions

- To get the decimal expansion is easy.



Base- b Expansions

- To get the decimal expansion is easy.

Example

- ◇ $(10101111)_2 = 2^8 + 2^6 + 2^4 + 2^3 + 2^2 + 2^1 + 2^0 = 351$
- ◇ $(7016)_8 = 7 \cdot 8^3 + 1 \cdot 8 + 6 = 3598$



Base- b Expansions

- To get the decimal expansion is easy.

Example

- ◇ $(10101111)_2 = 2^8 + 2^6 + 2^4 + 2^3 + 2^2 + 2^1 + 2^0 = 351$
- ◇ $(7016)_8 = 7 \cdot 8^3 + 1 \cdot 8 + 6 = 3598$

- Conversions between binary, octal, hexadecimal expansions are easy.



Base- b Expansions

- To get the decimal expansion is easy.

Example

- ◇ $(101011111)_2 = 2^8 + 2^6 + 2^4 + 2^3 + 2^2 + 2^1 + 2^0 = 351$
- ◇ $(7016)_8 = 7 \cdot 8^3 + 1 \cdot 8 + 6 = 3598$

- Conversions between binary, octal, hexadecimal expansions are easy.

Example

- ◇ $(101011111)_2 = (\underline{101}\overline{011}\underline{111}) = (537)_8$
- ◇ $(7016)_8 = (\underline{111}\overline{000}\underline{001}\overline{110})_2$
 $= (\underline{111}\overline{000}\underline{001}\overline{110})_2 = (E0E)_{16}$

Base- b Expansions

$$\begin{aligned}n &= a_k b^k + a_{k-1} b^{k-1} + a_{k-2} b^{k-2} + \cdots + a_2 b^2 + a_1 b + a_0 \\&= b(a_k b^{k-1} + a_{k-1} b^{k-2} + a_{k-2} b^{k-3} + \cdots + a_2 b + a_1) + \textcolor{red}{a_0} \\&= b(b(a_k b^{k-2} + a_{k-1} b^{k-3} + a_{k-2} b^{k-4} + \cdots + a_2) + \textcolor{red}{a_1}) + \textcolor{blue}{a_0} \\&= \cdots\end{aligned}$$



Base- b Expansions

$$\begin{aligned}n &= a_k b^k + a_{k-1} b^{k-1} + a_{k-2} b^{k-2} + \cdots + a_2 b^2 + a_1 b + a_0 \\&= b(a_k b^{k-1} + a_{k-1} b^{k-2} + a_{k-2} b^{k-3} + \cdots + a_2 b + a_1) + \textcolor{red}{a_0} \\&= b(b(a_k b^{k-2} + a_{k-1} b^{k-3} + a_{k-2} b^{k-4} + \cdots + a_2) + \textcolor{red}{a_1}) + \textcolor{blue}{a_0} \\&= \cdots\end{aligned}$$

To construct the base- b expansion of an integer n ,

- Divide n by b to obtain $\textcolor{blue}{n = bq_0 + a_0}$, with $0 \leq a_0 < b$
- The remainder a_0 is the rightmost digit in the base- b expansion of n . Then divide q_0 by b to get $\textcolor{blue}{q_0 = bq_1 + a_1}$ with $0 \leq a_1 < b$
- a_1 is the second digit from the right. Continue by successively dividing the quotients by b until **the quotient is 0**



Algorithm: Constructing Base- b Expansions

```
procedure base  $b$  expansion( $n, b$ : positive integers with  $b > 1$ )  
   $q := n$   
   $k := 0$   
  while ( $q \neq 0$ )  
     $a_k := q \bmod b$   
     $q := q \operatorname{div} b$   
     $k := k + 1$   
  return( $a_{k-1}, \dots, a_1, a_0$ ) { ( $a_{k-1} \dots a_1 a_0$ ) $_b$  is base  $b$  expansion of  $n$  }
```

Example

- $(12345)_{10} = (30071)_8$



Example

■ $(12345)_{10} = (30071)_8$

$$12345 = 8 \cdot 1543 + 1$$

$$1543 = 8 \cdot 192 + 7$$

$$192 = 8 \cdot 24 + 0$$

$$24 = 8 \cdot 3 + 0$$

$$3 = 8 \cdot 0 + 3$$



Binary Addition of Integers

$$a = (a_{n-1}a_{n-2} \dots a_1a_0), \quad b = (b_{n-1}b_{n-2} \dots b_1b_0)$$

procedure *add*(*a, b*: positive integers)

{the binary expansions of *a* and *b* are $(a_{n-1}, a_{n-2}, \dots, a_0)_2$ and $(b_{n-1}, b_{n-2}, \dots, b_0)_2$, respectively}

c := 0

for *j* := 0 to *n* − 1

d := $\lfloor (a_j + b_j + c) / 2 \rfloor$

*s*_{*j*} := $a_j + b_j + c - 2d$

c := *d*

*s*_{*n*} := *c*

return(*s*₀, *s*₁, ..., *s*_{*n*}) {the binary expansion of the sum is $(s_n, s_{n-1}, \dots, s_0)_2$ }

Binary Addition of Integers

$$a = (a_{n-1}a_{n-2} \dots a_1a_0), \quad b = (b_{n-1}b_{n-2} \dots b_1b_0)$$

```
procedure add(a, b: positive integers)
{the binary expansions of a and b are  $(a_{n-1}, a_{n-2}, \dots, a_0)_2$  and  $(b_{n-1}, b_{n-2}, \dots, b_0)_2$ , respectively}
c := 0
for j := 0 to n − 1
    d :=  $\lfloor (a_j + b_j + c) / 2 \rfloor$ 
    sj :=  $a_j + b_j + c - 2d$ 
    c := d
sn := c
return(s0, s1, ..., sn) {the binary expansion of the sum is  $(s_n, s_{n-1}, \dots, s_0)_2$ }
```

$O(n)$ bit additions

Algorithm: Binary Multiplication of Integers

$$a = (a_{n-1}a_{n-2} \dots a_1a_0)_2, \quad b = (b_{n-1}b_{n-2} \dots b_1b_0)_2$$

$$\begin{aligned} ab &= a(b_02^0 + b_12^1 + \dots + b_{n-1}2^{n-1}) \\ &= a(b_02^0) + a(b_12^1) + \dots + a(b_{n-1}2^{n-1}) \end{aligned}$$

```
procedure multiply(a, b: positive integers)
{the binary expansions of a and b are  $(a_{n-1}, a_{n-2}, \dots, a_0)_2$  and  $(b_{n-1}, b_{n-2}, \dots, b_0)_2$ , respectively}
for j := 0 to n - 1
    if  $b_j = 1$  then  $c_j = a$  shifted j places
    else  $c_j := 0$ 
{ $c_0, c_1, \dots, c_{n-1}$  are the partial products}
p := 0
for j := 0 to n - 1
    p := p +  $c_j$ 
return p {p is the value of ab}
```

Algorithm: Binary Multiplication of Integers

$$a = (a_{n-1}a_{n-2} \dots a_1a_0)_2, \quad b = (b_{n-1}b_{n-2} \dots b_1b_0)_2$$

$$\begin{aligned} ab &= a(b_02^0 + b_12^1 + \dots + b_{n-1}2^{n-1}) \\ &= a(b_02^0) + a(b_12^1) + \dots + a(b_{n-1}2^{n-1}) \end{aligned}$$

```
procedure multiply(a, b: positive integers)
{the binary expansions of a and b are  $(a_{n-1}, a_{n-2}, \dots, a_0)_2$  and  $(b_{n-1}, b_{n-2}, \dots, b_0)_2$ , respectively}
for j := 0 to n - 1
    if  $b_j = 1$  then  $c_j = a$  shifted j places
    else  $c_j := 0$ 
{ $c_0, c_1, \dots, c_{n-1}$  are the partial products}
p := 0
for j := 0 to n - 1
    p := p +  $c_j$ 
return p {p is the value of  $ab$ }
```

$O(n^2)$ shifts and $O(n^2)$ bit additions

Algorithm: Computing div and mod

```
procedure division algorithm (a: integer, d: positive integer)
  q := 0
  r := |a|
  while r ≥ d
    r := r - d
    q := q + 1
  if a < 0 and r > 0 then
    r := d - r
    q := -(q+1)
  return (q, r) {q = a div d is the quotient, r = a mod d is the
  remainder }
```



Algorithm: Computing div and mod

```
procedure division algorithm (a: integer, d: positive integer)
  q := 0
  r := |a|
  while r ≥ d
    r := r - d
    q := q + 1
  if a < 0 and r > 0 then
    r := d - r
    q := -(q+1)
  return (q, r) {q = a div d is the quotient, r = a mod d is the
  remainder }
```

$O(q \log a)$ bit operations. But there exist more efficient algorithms with complexity $O(n^2)$, where $n = \max(\log a, \log d)$

Algorithm: Computing div and mod (cont)

```
■ procedure division2 ( $a, d \in \mathbb{N}, d \geq 1$ )  
  if  $a < d$   
    return  $(q, r) = (0, a)$   
   $(q, r) = \text{division2}(\lfloor a/2 \rfloor, d)$   
   $q = 2q, r = 2r$   
  if  $a$  is odd  
     $r = r + 1$   
  if  $r \geq d$   
     $r = r - d$   
     $q = q + 1$   
  return  $(q, r)$ 
```

Algorithm: Computing div and mod (cont)

```
■ procedure division2 ( $a, d \in \mathbb{N}, d \geq 1$ )  
  if  $a < d$   
    return  $(q, r) = (0, a)$   
   $(q, r) = \text{division2}(\lfloor a/2 \rfloor, d)$   
   $q = 2q, r = 2r$   
  if  $a$  is odd  
     $r = r + 1$   
  if  $r \geq d$   
     $r = r - d$   
     $q = q + 1$   
  return  $(q, r)$ 
```

$O(\log q \log a)$ bit operations.

Algorithm: Binary Modular Exponentiation

$$b^n = b^{a_{k-1} \cdot 2^{k-1} + \dots + a_1 \cdot 2 + a_0} = b^{a_{k-1} \cdot 2^{k-1}} \dots b^{a_1 \cdot 2} \cdot b^{a_0}$$

Successively finds $b \bmod m$, $b^2 \bmod m$, $b^4 \bmod m$, \dots , $b^{2^{k-1}} \bmod m$, and multiplies together the terms $b^{2^j} \bmod m$ where $a_j = 1$.

```
procedure modular_exponentiation( $b$ : integer,  $n = (a_{k-1}a_{k-2}\dots a_1a_0)_2$ ,  $m$ : positive integers)
   $x := 1$ 
   $power := b \bmod m$ 
  for  $i := 0$  to  $k - 1$ 
    if  $a_i = 1$  then  $x := (x \cdot power) \bmod m$ 
     $power := (power \cdot power) \bmod m$ 
  return  $x$  { $x$  equals  $b^n \bmod m$ }
```


Algorithm: Binary Modular Exponentiation

$$b^n = b^{a_{k-1} \cdot 2^{k-1} + \dots + a_1 \cdot 2 + a_0} = b^{a_{k-1} \cdot 2^{k-1}} \dots b^{a_1 \cdot 2} \cdot b^{a_0}$$

Successively finds $b \bmod m$, $b^2 \bmod m$, $b^4 \bmod m$, \dots , $b^{2^{k-1}} \bmod m$, and multiplies together the terms $b^{2^j} \bmod m$ where $a_j = 1$.

```
procedure modular_exponentiation( $b$ : integer,  $n = (a_{k-1}a_{k-2}\dots a_1a_0)_2$ ,  $m$ : positive integers)
   $x := 1$ 
   $power := b \bmod m$ 
  for  $i := 0$  to  $k - 1$ 
    if  $a_i = 1$  then  $x := (x \cdot power) \bmod m$ 
     $power := (power \cdot power) \bmod m$ 
  return  $x$  { $x$  equals  $b^n \bmod m$ }
```

$O((\log m)^2 \log n)$ bit operations



Next Lecture

- number theory, cryptography ...

