



MACHINE LEARNING

CHAPTER 1: PRELIMINARY

Learning Objectives









- 1、 What are pattern recognition and machine learning?
 - 2、 What are curve fitting and regularization?
 - 3、 What are ML and MAP Bayesian inferences?
 - 4、 How to deal with the curse of dimensionality?
 - 5、 What is the relationship between decision theory and machine learning?
 - 6、 What are generative and discriminative models?
 - 7、 How to use entropy、 KL divergence and mutual information for machine learning?
-

Outlines

- Pattern Recognition and Machine Learning
 - Curve Fitting and Regularization
 - Probabilities and Gaussian Distributions
 - Bayesian Inferences (ML and MAP)
 - Curse of Dimensionality
 - Decision Theory
 - Entropy and Information
-

Example

Handwritten Digit Recognition



Machine Learning Pipeline

Dimension Reduction

Feature Modeling

Decision

Data Set

Vector Space

Feature Space




Function Space

Decision Space

Feature Selection


Boundary Modeling

Prediction



Feature Model

Boundary Model



Deterministic Machine Learning

Process of Machine Learning:

$$\theta^* = \min_{\theta} E(f_{\theta}(x), y) + \lambda \Omega(\theta)$$

θ : model parameters
 x : input data
 y : output data
 λ : weights
 $f_{\theta}(\bullet)$: model
 $E(\bullet)$: cost function
 $\Omega(\theta)$: model constraints

Process of Prediction:

$$y' = f_{\theta^*}(x')$$

x' : new input data
 y' : new output data
 $f_{\theta^*}(\bullet)$: learned model

Probabilistic Machine Learning

Goal of Machine Learning: $p(\text{real data}) \approx p(\text{generated data} / \text{model}, \theta)$

we assume: $p(\text{training data}) \approx p(\text{real data})$

Operation of Machine Learning: $p(\text{training data}) \approx p(\text{generated data} / \text{model}, \theta)$

Process of Machine Learning: $\theta^* \approx \max_{\theta} p(\text{training data} / \text{model}, \theta)$

Process of Machine Learning: $\theta^* \approx \max_{\theta} p(\theta / \text{training data}, \text{model})$

Process of prediction: $p(\text{testing data} / \text{model}, \theta^*)$

Bayesian Machine Learning

Process of Machine Learning:

$$p(\theta | \text{training data, model}) \propto p(\text{training data} / \text{model}, \theta) p_0(\theta)$$

posterior

likelihood

prior

Process of Prediction:


$$p(\text{testing data} | \text{training data, model}) =$$

$$\int p(\text{testing data} / \text{model}, \theta) p(\theta | \text{training data, model}) d\theta$$

Outlines


- Pattern Recognition and Machine Learning
 - Curve Fitting and Regularization
 - Probabilities and Gaussian Distributions
 - Bayesian Inferences (ML and MAP)
 - Curse of Dimensionality
 - Decision Theory
 - Entropy and Information
-

Polynomial Curve Fitting




$$y(x, \mathbf{w}) = w_0 + w_1x + w_2x^2 + \dots + w_Mx^M = \sum_{j=0}^M w_jx^j$$

Sum-of-Squares Error Function




$$E(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2$$


0th Order Polynomial




1st Order Polynomial




3rd Order Polynomial



9th Order Polynomial



Over-fitting




Root-Mean-Square (RMS) Error: $E_{\text{RMS}} = \sqrt{2E(\mathbf{w}^*)/N}$

Polynomial Coefficients

	$M = 0$	$M = 1$	$M = 3$	$M = 9$
w_0^*	0.19	0.82	0.31	0.35
w_1^*		-1.27	7.99	232.37
w_2^*			-25.43	-5321.83
w_3^*			17.37	48568.31
w_4^*				-231639.30
w_5^*				640042.26
w_6^*				-1061800.52
w_7^*				1042400.18
w_8^*				-557682.99
w_9^*				125201.43


Data Set Size: $N = 15$

9th Order Polynomial



Data Set Size: $N = 100$

9th Order Polynomial




Regularization


Penalize large coefficient values

$$\tilde{E}(\mathbf{w}) = \frac{1}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2$$


Regularization: $\ln \lambda = -18$



Regularization: $\ln \lambda = 0$



Regularization: E_{RMS} vs. $\ln \lambda$



Polynomial Coefficients


	$\ln \lambda = -\infty$	$\ln \lambda = -18$	$\ln \lambda = 0$
w_0^*	0.35	0.35	0.13
w_1^*	232.37	4.74	-0.05
w_2^*	-5321.83	-0.77	-0.06
w_3^*	48568.31	-31.97	-0.05
w_4^*	-231639.30	-3.89	-0.03
w_5^*	640042.26	55.28	-0.02
w_6^*	-1061800.52	41.32	-0.01
w_7^*	1042400.18	-45.95	-0.00
w_8^*	-557682.99	-91.53	0.00
w_9^*	125201.43	72.68	0.01

Outlines


- Pattern Recognition and Machine Learning
 - Curve Fitting and Regularization
 - Probabilities and Gaussian Distributions
 - Bayesian Inferences (ML and MAP)
 - Curse of Dimensionality
 - Decision Theories
 - Entropy and Information
-

Probability Theory

Apples and Oranges



Probability Theory



Marginal Probability

$$p(X = x_i) = \frac{c_i}{N}.$$


Joint Probability

$$p(X = x_i, Y = y_j) = \frac{n_{ij}}{N}$$

Conditional Probability

$$p(Y = y_j | X = x_i) = \frac{n_{ij}}{c_i}$$

Probability Theory



Sum Rule

$$\begin{aligned} p(X = x_i) &= \frac{c_i}{N} = \frac{1}{N} \sum_{j=1}^L n_{ij} \\ &= \sum_{j=1}^L p(X = x_i, Y = y_j) \end{aligned}$$

Product Rule

$$\begin{aligned} p(X = x_i, Y = y_j) &= \frac{n_{ij}}{N} = \frac{n_{ij}}{c_i} \cdot \frac{c_i}{N} \\ &= p(Y = y_j | X = x_i) p(X = x_i) \end{aligned}$$

The Rules of Probability

Sum Rule

$$p(X) = \sum_Y p(X, Y)$$

Product Rule

$$p(X, Y) = p(Y|X)p(X)$$

Bayes' Theorem


$$p(Y|X) = \frac{p(X|Y)p(Y)}{p(X)}$$

$$p(X) = \sum_Y p(X|Y)p(Y) \quad : \text{normalization}$$

posterior \propto likelihood \times prior

$$p(Y/X) \qquad p(X/Y) \qquad p(Y)$$

Probability Densities




$$p(x \in (a, b)) = \int_a^b p(x) dx$$

$$P(z) = \int_{-\infty}^z p(x) dx$$

$$p(x) \geq 0$$

$$\int_{-\infty}^{\infty} p(x) dx = 1$$

Transformed Densities




$$\begin{aligned} p_y(y) &= p_x(x) \left| \frac{dx}{dy} \right| \\ &= p_x(g(y)) |g'(y)| \end{aligned}$$

$$x = g(y)$$

Expectations

$$\mathbb{E}[f] = \sum_x p(x) f(x)$$

$$\mathbb{E}[f] = \int p(x) f(x) \, dx$$

$$\mathbb{E}_x[f|y] = \sum_x p(x|y) f(x)$$


Conditional Expectation
(discrete)

$$\mathbb{E}[f] \simeq \frac{1}{N} \sum_{n=1}^N f(x_n)$$

Approximate Expectation
(discrete and continuous)

Variances and Covariances


$$\text{var}[f] = \mathbb{E} \left[(f(x) - \mathbb{E}[f(x)])^2 \right] = \mathbb{E}[f(x)^2] - \mathbb{E}[f(x)]^2$$

$$\begin{aligned} \text{cov}[x, y] &= \mathbb{E}_{x,y} [\{x - \mathbb{E}[x]\} \{y - \mathbb{E}[y]\}] \\ &= \mathbb{E}_{x,y} [xy] - \mathbb{E}[x]\mathbb{E}[y] \end{aligned}$$

$$\begin{aligned} \text{cov}[\mathbf{x}, \mathbf{y}] &= \mathbb{E}_{\mathbf{x}, \mathbf{y}} [\{\mathbf{x} - \mathbb{E}[\mathbf{x}]\} \{\mathbf{y}^T - \mathbb{E}[\mathbf{y}^T]\}] \\ &= \mathbb{E}_{\mathbf{x}, \mathbf{y}} [\mathbf{x} \mathbf{y}^T] - \mathbb{E}[\mathbf{x}] \mathbb{E}[\mathbf{y}^T] \end{aligned}$$

The Gaussian Distribution

$$\mathcal{N}(x|\mu, \sigma^2) = \frac{1}{(2\pi\sigma^2)^{1/2}} \exp \left\{ -\frac{1}{2\sigma^2} (x - \mu)^2 \right\}$$



$$\mathcal{N}(x|\mu, \sigma^2) > 0$$

$$\int_{-\infty}^{\infty} \mathcal{N}(x|\mu, \sigma^2) dx = 1$$

Gaussian Mean and Variance


$$\mathbb{E}[x] = \int_{-\infty}^{\infty} \mathcal{N}(x|\mu, \sigma^2) x \, dx = \mu$$

$$\mathbb{E}[x^2] = \int_{-\infty}^{\infty} \mathcal{N}(x|\mu, \sigma^2) x^2 \, dx = \mu^2 + \sigma^2$$

$$\text{var}[x] = \mathbb{E}[x^2] - \mathbb{E}[x]^2 = \sigma^2$$

The Multivariate Gaussian


$$\mathcal{N}(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{1}{(2\pi)^{D/2}} \frac{1}{|\boldsymbol{\Sigma}|^{1/2}} \exp \left\{ -\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu}) \right\}$$



Outlines

- Pattern Recognition and Machine Learning
 - Curve Fitting and Regularization
 - Probabilities and Gaussian Distributions
 - Bayesian Inferences (ML and MAP)
 - Curse of Dimensionality
 - Decision Theories
 - Entropy and Information
-

Gaussian Parameter Estimation



$$p(\mathbf{x} | \mu, \sigma^2) = \prod_{n=1}^N \mathcal{N}(x_n | \mu, \sigma^2)$$

Maximum (Log) Likelihood

$$\ln p(\mathbf{x}|\mu, \sigma^2) = -\frac{1}{2\sigma^2} \sum_{n=1}^N (x_n - \mu)^2 - \frac{N}{2} \ln \sigma^2 - \frac{N}{2} \ln(2\pi)$$


$$\mu_{\text{ML}} = \frac{1}{N} \sum_{n=1}^N x_n \qquad \sigma_{\text{ML}}^2 = \frac{1}{N} \sum_{n=1}^N (x_n - \mu_{\text{ML}})^2$$

Properties of μ_{ML} and σ_{ML}^2


$$\mathbb{E}[\mu_{\text{ML}}] = \mu$$

$$\mathbb{E}[\sigma_{\text{ML}}^2] = \left(\frac{N-1}{N}\right) \sigma^2$$

$$\begin{aligned}\tilde{\sigma}^2 &= \frac{N}{N-1} \sigma_{\text{ML}}^2 \\ &= \frac{1}{N-1} \sum_{n=1}^N (x_n - \mu_{\text{ML}})^2\end{aligned}$$



Curve Fitting Re-visited



(t, x) : training data $\Rightarrow \mathbf{w}, \beta$ (\mathbf{w}, β, x_0) : $\Rightarrow p(t|x_0, \mathbf{w}, \beta)$

Maximum Likelihood

$$p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \beta) = \prod_{n=1}^N \mathcal{N}(t_n|y(x_n, \mathbf{w}), \beta^{-1})$$


$$\ln p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \beta) = - \underbrace{\frac{\beta}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2}_{\beta E(\mathbf{w})} + \frac{N}{2} \ln \beta - \frac{N}{2} \ln(2\pi)$$

Determine \mathbf{w}_{ML} by minimizing sum-of-squares error, $E(\mathbf{w})$.

$$\frac{1}{\beta_{\text{ML}}} = \frac{1}{N} \sum_{n=1}^N \{y(x_n, \mathbf{w}_{\text{ML}}) - t_n\}^2$$

Predictive Distribution

$$p(t|x, \mathbf{w}_{\text{ML}}, \beta_{\text{ML}}) = \mathcal{N}(t|y(x, \mathbf{w}_{\text{ML}}), \beta_{\text{ML}}^{-1})$$



MAP: A Step towards Bayes

MAP: Maximum *A Posteriori*

$$p(\mathbf{w}|\alpha) = \mathcal{N}(\mathbf{w}|\mathbf{0}, \alpha^{-1}\mathbf{I}) = \left(\frac{\alpha}{2\pi}\right)^{(M+1)/2} \exp\left\{-\frac{\alpha}{2}\mathbf{w}^T\mathbf{w}\right\}$$

$$\boxed{\textit{posteriori}} \longrightarrow p(\mathbf{w}|\mathbf{x}, \mathbf{t}, \alpha, \beta) \propto \underset{\substack{\uparrow \\ \boxed{\textit{likelihood}}}}{p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \beta)} p(\mathbf{w}|\alpha) \longleftarrow \boxed{\textit{priori}}$$

$$\beta\tilde{E}(\mathbf{w}) = \frac{\beta}{2} \sum_{n=1}^N \{y(x_n, \mathbf{w}) - t_n\}^2 + \frac{\alpha}{2} \mathbf{w}^T \mathbf{w}$$

Determine \mathbf{w}_{MAP} by minimizing regularized sum-of-squares error, $\tilde{E}(\mathbf{w})$.

Bayesian Curve Fitting

$$p(t|x, \mathbf{x}, \mathbf{t}) = \int p(t|x, \mathbf{w})p(\mathbf{w}|\mathbf{x}, \mathbf{t}) d\mathbf{w} = \mathcal{N}(t|m(x), s^2(x))$$


$$m(x) = \beta \phi(x)^T \mathbf{S} \sum_{n=1}^N \phi(x_n) t_n \quad s^2(x) = \beta^{-1} + \phi(x)^T \mathbf{S} \phi(x)$$

$$\mathbf{S}^{-1} = \alpha \mathbf{I} + \beta \sum_{n=1}^N \phi(x_n) \phi(x_n)^T \quad \phi(x_n) = (x_n^0, \dots, x_n^M)^T$$

We will go through more details in a later lecture.

Bayesian Predictive Distribution

$$p(t|x, \mathbf{x}, \mathbf{t}) = \mathcal{N}(t|m(x), s^2(x))$$



Model Selection and Evaluation

Cross-Validation

