

CS215: Discrete Math (H)
2023 Fall Semester Written Assignment #1
Due: Oct. 9th, 2023, please submit at the beginning of class

Q.1 Let p, q be the propositions

p : You get 100 marks on the final.

q : You get an A in this course.

Write these propositions using p and q and logical connectives (including negations).

- (a) You get 100 marks on the final, but you do not get an A in this course.
- (b) You will get an A in this course if you get 100 marks on the final.
- (c) If you do not get 100 marks on the final, then you will not get an A in this course.
- (d) Getting 100 marks on the final is sufficient for getting an A in this course.
- (e) Whenever you get an A in this course, you got 100 marks on the final.

Q.2 Construct a truth table for each of these compound propositions.

- (a) $(p \leftrightarrow q) \oplus (\neg p \leftrightarrow q)$
- (b) $(p \oplus q) \wedge (p \oplus \neg q)$

Q.3 Use truth tables to decide whether or not the following two propositions are equivalent.

- (a) $(p \rightarrow q) \vee (p \rightarrow r)$ and $p \rightarrow (q \vee r)$
- (b) $(p \rightarrow q) \rightarrow r$ and $p \rightarrow (q \rightarrow r)$
- (c) $(p \vee q) \rightarrow r$ and $(p \rightarrow r) \wedge (q \rightarrow r)$

Q.4 Use logical equivalences to prove the following statements.

- (a) $\neg p \rightarrow (q \rightarrow r)$ and $q \rightarrow (p \vee r)$ are equivalent.
- (b) $(p \rightarrow q) \rightarrow ((r \rightarrow p) \rightarrow (r \rightarrow q))$ is a tautology.

Q.5 Suppose that p, q, r, s are all logical propositions. You are given the following statement

$$(q \rightarrow (r \vee p)) \rightarrow ((\neg r \vee s) \wedge \neg s).$$

Prove that this implies $\neg r$ using logical equivalences and inference rules.

Q.6 Let $F(x, y)$ be the statement “ x can fool y ”, where the domain consists of all people in the world. Use quantifiers to express each of these statement.

- (a) Everybody can fool Fred.
- (b) Everybody can fool somebody.
- (c) There is no one who can fool everybody.
- (d) Everyone can be fooled by somebody.
- (e) No one can fool both Fred and Jerry.
- (f) Nancy can fool exactly two people.
- (g) There is exactly one person whom everybody can fool.
- (h) There is someone who can fool exactly one person besides himself or herself.

Q.7 Given the following predicate on the set P of all people who ever lived: Parent(x, y) means x is the parent of y .

- (1) Rewrite the following in the language of mathematical logic (you may use the equality/inequality operators):

“All people have two parents.”

- (2) Based on the definition of Parent, we will *recursively* define the concept of *ancestor*: “An ancestor of a person is one of the person’s parents or the ancestor of (at least) one of the person’s parents.”

Rewrite this definition using the language of mathematical logic. Specifically, you need to provide a necessary and sufficient condition for the predicate Ancestor(x, y) to be true. Note that you can *recursively* use the predicate Ancestor(x, y) in the condition itself.

Q.8 Express the negations of each of these statements so that all negation symbols immediately precede predicates.

(a) $\exists x \exists y P(x, y) \wedge \forall x \forall y Q(x, y)$

(b) $\forall x \exists y P(x, y) \vee \forall x \exists y Q(x, y)$

(c) $\forall x \exists y (P(x, y) \rightarrow Q(x, y))$

Q.9 Let $P(x, y)$ be a propositional function. Prove or disprove that $\exists x \forall y P(x, y) \rightarrow \forall y \exists x P(x, y)$ is a tautology.

Q.10 For the following argument, explain which rules of inference are used for each step.

“Each of five roommates, A, B, C, D, and E, has taken a course in discrete mathematics. Every student who has taken a course in discrete mathematics can take a course in algorithms. Therefore, all five roommates can take a course in algorithms next year.”

Q.11

- (a) Let P be a proposition in atomic propositions p and q . If $P = \neg(p \leftrightarrow (q \vee \neg p))$, prove that P is equivalent to $\neg p \vee \neg q$.
- (b) If P is of any length, using any of the logical connectives $\neg, \wedge, \vee, \rightarrow, \leftrightarrow$, prove that P is logically equivalent to a proposition of the form

$$A \square B,$$

where \square is one of $\wedge, \vee, \leftrightarrow$, and A and B are chosen from $\{p, \neg p, q, \neg q\}$.

Q.12 Prove or disprove that if a and b are rational numbers, then a^b is also rational. .

Q.13

Prove or disprove the following.

- (1) For two irrational numbers a and b , a^b is also irrational.
- (2) For an irrational number a , \sqrt{a} is also irrational.

Q.14 Prove that $\sqrt[3]{2}$ is irrational.

Q.15 Prove that between every two rational numbers there is an irrational number.

Q.16 Give a direct proof that: Let a and b be integers. If $a^2 + b^2$ is even, then $a + b$ is even.

Q.17 Let the coefficients of the polynomial $f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + x^n$ be integers. Show that any *real* root of the equation $f(x) = 0$ is either integral or irrational. Note that in your proof, you may direct use the following result without a proof. “**Fact.** If a prime p is a factor of some power of an integer, then it is a factor of that integer.”