## AI1110 Assignment 11

# Dondapati Chandrahas Reddy AI21BTECH11010

June 20, 2022

### Question(Papoulis Exercise 15.3)

Find the stationary distribution  $q_0, q_1, ...$  for the Markov chain whose only nonzero stationary probabilities are

$$p_{i,1} = \frac{i}{i+1}$$
  $p_{i,i+1} = \frac{1}{i+1}$   $i = 1, 2, ...$ 

#### Solution

$$q_{i+1} = p_{i,i+1} \ q_i = \frac{1}{i+1} q_i \tag{1}$$

$$q_i = p_{i-1,i} \ q_{i-1} = \frac{1}{i} q_{i-1}$$
 (2)

:

$$q_1 = p_{0,1} \ q_0 = \frac{1}{1} q_0 \tag{3}$$

$$\implies q_{i+1} = \frac{q_0}{(i+1)!} \tag{4}$$

### Solution(Contd.)

Summation of all probabilities equals one.

$$\sum_{k=1}^{\infty} q_k = q_0 \sum_{k=1}^{\infty} \frac{1}{k!} = 1 \tag{5}$$

$$q_0e = 1 \tag{6}$$

$$\implies q_0 = \frac{1}{e} \tag{7}$$

Therefore, the steady state probabilities are given by,

$$q_k = \frac{1}{e^{k!}} \tag{8}$$