Statistics Mcq Questions

Q1._____is not a common measure of central tendency.

a)Mean

b)Median

c)Mode

d)Range

Q2. _____ is not a measure of variability

a)Median

b)Standard Variance

c)Variance

d)Range

Q3.Which of these is a type of statistics?

(A) Descriptive statistics

(B) Inferential statistics

(C) Industry statistics

(D)Both A and B

Q4.Variables which can be measured in terms of height , weight and length are ___?

A) continuous variables

B) measuring variables

C) flowchart variables

D) discrete variables

Q5. The discrete variables and continuous variables are the types of

A. open end classification

B. time series classification

C. qualitative classification

D. quantitative classification

Q6. In the systematic sampling, the value of k is __?

A. sampling interval

B. sub stage interval

C. secondary stage interval

D. multistage interval

Q7. ____ involve methods of organizing ,picturing, and summarizing information from samples and populations.

a) Descriptive statistics

b) Inferential statistics

c) Analytical statistics

d) All of the above

Q8._____ is a descriptive measurement computed from an entire population of data.

a) Statistics
b) Parameter
c) Clustering
d) None of these

Q9. Score Of computer exam is an example of ___?

a) Ratio data

b) Interval Data
c) Ordinal Data
d) Nominal Data

Q10.Pin Code is an example of ___?

a) Ratio data
b) Interval Data
c) Ordinal Data
d) Nominal data

Q11. Sample is regared as a subset of___?

(a) Data

(b) Set

(c) Distribution

(d) Population

Q12. _____technique where every item in the population has equal probability of being selected?

(a) Simple random sampling

(b) Stratified random sampling

(c) Systematic sampling

(d) Probability sampling

Q13. A number in a data-set which is very different from others is called _____?

a) Error
b) Mode
c) Outliers
d) Median

Q14. The number of observations in a particular class is called:

a) Width of the class

b) Class mark

c) Frequency

d) None of the above

Q15. The square root of the variance is called the _____ deviation.

a) empirical
b) mean

c) continuous
d) standard Deviation

Q16. When the correlation coefficient, r, is close to one:
a) There is no relationship between the two variables

b) There is a strong linear relationship between the two variables

c) It is impossible to tell if there is a relationship between the two variables

d) The slope of the regression line will be close to one

Q17. If the assumed hypothesis is tested for rejection considering it to be true is called?
a) Null Hypothesis
b) Statistical Hypothesis
c) Simple Hypothesis
d) Composite Hypothesis

Q18. The values of mean, median and mode can be

a) Some time equal
b) Never equal

c) Always equal
d) None of these

Q19.Cumulative frequency is____ frequency?

a)Decreasing

b)Increasing

c)Different

d)None of these

Q20. If the mean of the sampling distribution is equal to the parameters then estimators will be

A)Biased

B)Consistent

C)Sufficient

D)Unbiased

MACHINE LEARNING QUESTIONS

Q1.Which of the following is not a supervised learning?

a)Naive Bayes

b)PCA

c)Linear Regression

d)Decision Tree Answer

Q2. Which of the following is not type of learning?

a)Unsupervised Learning

b)Supervised Learning

c)Semi-unsupervised Learning

d)Reinforcement Learning

Q3. Fraud Detection, Image Classification, Diagnostic, and Customer Retention are applications in which of the following

a)Unsupervised Learning: Regression

b)Supervised Learning: Classification

c)Unsupervised Learning: Clustering

d)Reinforcement Learning

Q4. Which of the following is a widely used and effective machine learning algorithm based on the idea of bagging?

A. Decision Tree

B. Random Forest

C. Regression

D. Classification

Q5. A model can learn based on the rewards it received for its previous action is known as:

A. Supervised learning

B. Unsupervised learning

C. Reinforcement learning

D. Concept learning

Q6. Which of the following is an application of reinforcement learning?

A) Topic modeling

B) Recommendation system

C)Pattern recognition

D)Image classification

Q6. What would be the best regression model for more than one independent variable?

A) Simple Linear Regression

B) Multiple Linear Regression

C) Logistic Regression

D) All of the Above

Q7. Regression is a __?

A. Supervised Learning Algorithm

B. Unsupervised Learning Algorithm

C. Reinforcement Learning Algorithm

D. None

Q8. Linear regression is___ to Outliers

A) sensitive

B) not sensitive

C) not affected by outliers

D) None

Q9.What does k stand for in the KNN algorithm?

A. Number of neighbors

B. Number of output classes

C. Number of input features

D. None

Q10. For clustering, we do not require-

A. Labeled data

B. Unlabeled data

C. Numerical data

D. Categorical data

Q11. The average squared difference between classifier predicted output and actual output.

a) mean squared error

b) root mean squared error

c) mean absolute error

d) mean relative error

Q12. Regression trees are often used to model _____ data.

a) Linear

b) Nonlinear

c) Categorical

d) Symmetrical

Q13 What device below is not an example of Machine Learning?

A)Speech to text

B)Google Search

C)Google Assistant

D)None Of The Above

Q14. How many dependent variables are used in multiple regression?

a) One

b) One or more

c) Two or more

d) Two

Q15. If a distribution is skewed to the left, then it is .

a) Negatively skewed

b) Positively skewed

c) Symmetrically skewed

d) Symmetrical

Q16. A Type I error is also known as a _____.

a. False positive

b. False negative

c. Double negative

d. Positive negative

Q17. A Type II error is also known as a _____.

a. False positive

b. False negative

c. Double negative

d. Positive negative

Q18. High Entropy Means That the partitions in Classification are?

A)Useful

B)Useless

C)Not Pure

D)Pure

Q19.Probability Of Occurrence Of A single event?

A. conditional probability

B. marginal probability

C. non conditional probability

D. occurrence probability

Q20 For two events, the probability of occurrence of both events at same time or occurrence in series is Classified as

A. joint probability

B. dependent probability

C. series probability

D. conditional probability