

# Numerical Linear Algebra

Ramaz Botchorishvili

Kutaisi International University

November 8, 2023

# Numerical Linear Algebra

Ramaz Botchorishvili

Kutaisi International University

November 8, 2023

# LU factorization, pivoting and stability of Gaussian elimination

- ▶ Recap of Previous Lecture
- ▶ Examples of LU factorization
- ▶ Existence of LU factorization
- ▶ Gaussian elimination(GE) and LU factorization
- ▶ Pivoting
- ▶ Stability of GE
- ▶ Q & A

# Recap of Previous Lecture

- ▶ Perturbations in right hand side and coefficients
- ▶ Round-off errors
- ▶ How to avoid cancellation and recursion errors
- ▶ Computational template for numerical linear algebra
- ▶ Thomas algorithm

# LU Decomposition Examples 1

## Example 8.1

General  $LU$  decomposition

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix} = \begin{pmatrix} l_{11} & 0 & \dots & 0 \\ l_{21} & l_{22} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ l_{n1} & l_{n2} & \dots & l_{nn} \end{pmatrix} \begin{pmatrix} u_{11} & u_{12} & \dots & u_{1n} \\ 0 & u_{22} & \dots & u_{2n} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & u_{nn} \end{pmatrix}$$

# LU Decomposition Examples 1

## Example 8.1

General  $LU$  decomposition

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix} = \begin{pmatrix} l_{11} & 0 & \dots & 0 \\ l_{21} & l_{22} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ l_{n1} & l_{n2} & \dots & l_{nn} \end{pmatrix} \begin{pmatrix} u_{11} & u_{12} & \dots & u_{1n} \\ 0 & u_{22} & \dots & u_{2n} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & u_{nn} \end{pmatrix}$$

## Example 8.2

Doolittle  $LU$  decomposition

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix} = \begin{pmatrix} 1 & 0 & \dots & 0 \\ l_{21} & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ l_{n1} & l_{n2} & \dots & 1 \end{pmatrix} \begin{pmatrix} u_{11} & u_{12} & \dots & u_{1n} \\ 0 & u_{22} & \dots & u_{2n} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & u_{nn} \end{pmatrix}$$

# LU Decomposition Examples 1

## Example 8.1

General  $LU$  decomposition

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix} = \begin{pmatrix} l_{11} & 0 & \dots & 0 \\ l_{21} & l_{22} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ l_{n1} & l_{n2} & \dots & l_{nn} \end{pmatrix} \begin{pmatrix} u_{11} & u_{12} & \dots & u_{1n} \\ 0 & u_{22} & \dots & u_{2n} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & u_{nn} \end{pmatrix}$$

## Example 8.2

Doolittle  $LU$  decomposition

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix} = \begin{pmatrix} 1 & 0 & \dots & 0 \\ l_{21} & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ l_{n1} & l_{n2} & \dots & 1 \end{pmatrix} \begin{pmatrix} u_{11} & u_{12} & \dots & u_{1n} \\ 0 & u_{22} & \dots & u_{2n} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & u_{nn} \end{pmatrix}$$

## Example 8.3

Crout  $LU$  decomposition

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix} = \begin{pmatrix} l_{11} & 0 & \dots & 0 \\ l_{21} & l_{22} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ l_{n1} & l_{n2} & \dots & l_{nn} \end{pmatrix} \begin{pmatrix} 1 & u_{12} & \dots & u_{1n} \\ 0 & 1 & \dots & u_{2n} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 \end{pmatrix}$$

## LU Decomposition Examples 2

### Example 8.4

General  $LDU$  decomposition

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix} = \begin{pmatrix} 1 & 0 & \dots & 0 \\ l_{21} & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ l_{n1} & l_{n2} & \dots & 1 \end{pmatrix} \begin{pmatrix} d_{11} & 0 & \dots & 0 \\ 0 & d_{22} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & d_{nn} \end{pmatrix} \begin{pmatrix} 1 & u_{12} & \dots & u_{1n} \\ 0 & 1 & \dots & u_{2n} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 \end{pmatrix}$$



## LU Decomposition Examples 2

### Example 8.4

General  $LDU$  decomposition

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix} = \begin{pmatrix} 1 & 0 & \dots & 0 \\ l_{21} & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ l_{n1} & l_{n2} & \dots & 1 \end{pmatrix} \begin{pmatrix} d_{11} & 0 & \dots & 0 \\ 0 & d_{22} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & d_{nn} \end{pmatrix} \begin{pmatrix} 1 & u_{12} & \dots & u_{1n} \\ 0 & 1 & \dots & u_{2n} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 \end{pmatrix}$$

### Example 8.5

- Doolittle decomposition  $A = LDU \Rightarrow A = (LD)U$

## LU Decomposition Examples 2

### Example 8.4

General  $LDU$  decomposition

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix} = \begin{pmatrix} 1 & 0 & \dots & 0 \\ l_{21} & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ l_{n1} & l_{n2} & \dots & 1 \end{pmatrix} \begin{pmatrix} d_{11} & 0 & \dots & 0 \\ 0 & d_{22} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & d_{nn} \end{pmatrix} \begin{pmatrix} 1 & u_{12} & \dots & u_{1n} \\ 0 & 1 & \dots & u_{2n} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 \end{pmatrix}$$

### Example 8.5

- ▶ Doolittle decomposition  $A = LDU \Rightarrow A = (LD)U$
- ▶ Crout decomposition  $A = LDU \Rightarrow A = L(DU)$

## LU Decomposition Examples 2

### Example 8.4

General  $LDU$  decomposition

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{pmatrix} = \begin{pmatrix} 1 & 0 & \dots & 0 \\ l_{21} & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ l_{n1} & l_{n2} & \dots & 1 \end{pmatrix} \begin{pmatrix} d_{11} & 0 & \dots & 0 \\ 0 & d_{22} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & d_{nn} \end{pmatrix} \begin{pmatrix} 1 & u_{12} & \dots & u_{1n} \\ 0 & 1 & \dots & u_{2n} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 \end{pmatrix}$$

### Example 8.5

- ▶ Doolittle decomposition  $A = LDU \Rightarrow A = (LD)U$
- ▶ Crout decomposition  $A = LDU \Rightarrow A = L(DU)$
- ▶ Exchanging rows or columns may be needed for implementation

## LU Decomposition Examples 3

### Example 8.6

►  $B = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix}$ ,  $\det(B) = 0$ ,  $\det(B_1) = 1 \neq 0$

## LU Decomposition Examples 3

### Example 8.6

- ▶  $B = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix}$ ,  $\det(B) = 0$ ,  $\det(B_1) = 1 \neq 0$
- ▶ Unique  $LU$  factorization exists

## LU Decomposition Examples 3

### Example 8.6

- ▶  $B = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix}$ ,  $\det(B) = 0$ ,  $\det(B_1) = 1 \neq 0$
- ▶ Unique  $LU$  factorization exists

### Example 8.7

- ▶  $D = \begin{pmatrix} 0 & 1 \\ 0 & 2 \end{pmatrix}$ ,  $\det(D) = 0$ ,  $\det(D_1) = 0$

## LU Decomposition Examples 3

### Example 8.6

- ▶  $B = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix}$ ,  $\det(B) = 0$ ,  $\det(B_1) = 1 \neq 0$
- ▶ Unique  $LU$  factorization exists

### Example 8.7

- ▶  $D = \begin{pmatrix} 0 & 1 \\ 0 & 2 \end{pmatrix}$ ,  $\det(D) = 0$ ,  $\det(D_1) = 0$
- ▶ Infinitely many  $LU$  factorizations exist

## LU Decomposition Examples 3

### Example 8.6

- ▶  $B = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix}$ ,  $\det(B) = 0$ ,  $\det(B_1) = 1 \neq 0$
- ▶ Unique  $LU$  factorization exists

### Example 8.7

- ▶  $D = \begin{pmatrix} 0 & 1 \\ 0 & 2 \end{pmatrix}$ ,  $\det(D) = 0$ ,  $\det(D_1) = 0$
- ▶ Infinitely many  $LU$  factorizations exist
- ▶  $D = L_\beta U_\beta = \begin{pmatrix} 1 & 0 \\ \beta & 1 \end{pmatrix} \cdot \begin{pmatrix} 0 & 1 \\ 0 & 2 - \beta \end{pmatrix}$



## LU Decomposition Examples 3

### Example 8.6

- ▶  $B = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix}$ ,  $\det(B) = 0$ ,  $\det(B_1) = 1 \neq 0$
- ▶ Unique  $LU$  factorization exists

### Example 8.7

- ▶  $D = \begin{pmatrix} 0 & 1 \\ 0 & 2 \end{pmatrix}$ ,  $\det(D) = 0$ ,  $\det(D_1) = 0$
- ▶ Infinitely many  $LU$  factorizations exist
- ▶  $D = L_\beta U_\beta = \begin{pmatrix} 1 & 0 \\ \beta & 1 \end{pmatrix} \cdot \begin{pmatrix} 0 & 1 \\ 0 & 2 - \beta \end{pmatrix}$

### Example 8.8

- ▶  $C = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ ,  $\det(C) = -1$ ,  $\det(C_1) = 0$ : no  $LU$  factorization exists

## LU Decomposition Examples 3

### Example 8.6

- ▶  $B = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix}$ ,  $\det(B) = 0$ ,  $\det(B_1) = 1 \neq 0$
- ▶ Unique  $LU$  factorization exists

### Example 8.7

- ▶  $D = \begin{pmatrix} 0 & 1 \\ 0 & 2 \end{pmatrix}$ ,  $\det(D) = 0$ ,  $\det(D_1) = 0$
- ▶ Infinitely many  $LU$  factorizations exist
- ▶  $D = L_\beta U_\beta = \begin{pmatrix} 1 & 0 \\ \beta & 1 \end{pmatrix} \cdot \begin{pmatrix} 0 & 1 \\ 0 & 2 - \beta \end{pmatrix}$

### Example 8.8

- ▶  $C = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ ,  $\det(C) = -1$ ,  $\det(C_1) = 0$ : no  $LU$  factorization exists
- ▶  $\begin{pmatrix} 1 & 0 \\ l_{21} & 1 \end{pmatrix} \cdot \begin{pmatrix} u_{11} & u_{12} \\ 0 & u_{22} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ : no solution exists

# Existence and uniqueness of LU factorization 1

## Theorem 8.9

*Unique LU factorization(Doolittle) of  $A \in \mathbb{R}^{n \times n}$  exists iff*

# Existence and uniqueness of LU factorization 1

## Theorem 8.9

*Unique LU factorization (Doolittle) of  $A \in \mathbb{R}^{n \times n}$  exists iff*

- ▶  $\det(A_i) \neq 0, i = 1, 2, \dots, n - 1$

# Existence and uniqueness of LU factorization 1

## Theorem 8.9

*Unique LU factorization(Doolittle) of  $A \in \mathbb{R}^{n \times n}$  exists iff*

- ▶  $\det(A_i) \neq 0, i = 1, 2, \dots, n - 1$
- ▶  $A_i$  -main submatrices,  $A \in \mathbb{R}^{i \times i}$

# Existence and uniqueness of LU factorization 1

## Theorem 8.9

*Unique LU factorization(Doolittle) of  $A \in \mathbb{R}^{n \times n}$  exists iff*

- ▶  $\det(A_i) \neq 0, i = 1, 2, \dots, n - 1$
- ▶  $A_i$  -main submatrices,  $A \in \mathbb{R}^{i \times i}$

## Proof.

Proof for sufficient condition, p1

# Existence and uniqueness of LU factorization 1

## Theorem 8.9

*Unique LU factorization(Doolittle) of  $A \in \mathbb{R}^{n \times n}$  exists iff*

- ▶  $\det(A_i) \neq 0, i = 1, 2, \dots, n - 1$
- ▶  $A_i$  -main submatrices,  $A \in \mathbb{R}^{i \times i}$

## Proof.

Proof for sufficient condition, p1

- ▶ proof by mathematical induction

# Existence and uniqueness of LU factorization 1

## Theorem 8.9

*Unique LU factorization(Doolittle) of  $A \in \mathbb{R}^{n \times n}$  exists iff*

- ▶  $\det(A_i) \neq 0, i = 1, 2, \dots, n - 1$
- ▶  $A_i$  -main submatrices,  $A \in \mathbb{R}^{i \times i}$

## Proof.

Proof for sufficient condition, p1

- ▶ proof by mathematical induction
- ▶  $i = 1, A_1 = a_{11}, \det(A_1) \neq 0 \Rightarrow A_1 = L_1 U_1 = 1 \cdot a_{11}$



# Existence and uniqueness of LU factorization 1

## Theorem 8.9

*Unique LU factorization (Doolittle) of  $A \in \mathbb{R}^{n \times n}$  exists iff*

- ▶  $\det(A_i) \neq 0, i = 1, 2, \dots, n - 1$
- ▶  $A_i$  -main submatrices,  $A \in \mathbb{R}^{i \times i}$

## Proof.

Proof for sufficient condition, p1

- ▶ proof by mathematical induction
- ▶  $i = 1, A_1 = a_{11}, \det(A_1) \neq 0 \Rightarrow A_1 = L_1 U_1 = 1 \cdot a_{11}$
- ▶ Assume  $A_{i-1} = L_{i-1} U_{i-1}$

# Existence and uniqueness of LU factorization 1

## Theorem 8.9

Unique LU factorization (Doolittle) of  $A \in \mathbb{R}^{n \times n}$  exists iff

- ▶  $\det(A_i) \neq 0, i = 1, 2, \dots, n - 1$
- ▶  $A_i$  -main submatrices,  $A \in \mathbb{R}^{i \times i}$

## Proof.

Proof for sufficient condition, p1

- ▶ proof by mathematical induction
- ▶  $i = 1, A_1 = a_{11}, \det(A_1) \neq 0 \Rightarrow A_1 = L_1 U_1 = 1 \cdot a_{11}$
- ▶ Assume  $A_{i-1} = L_{i-1} U_{i-1}$

- ▶ Denote  $A_i = \begin{pmatrix} A_{i-1} & c_i \\ d_i & a_{ii} \end{pmatrix}, c_i = \begin{pmatrix} a_{1i} \\ \vdots \\ a_{i-1,i} \end{pmatrix}, d_i = \begin{pmatrix} a_{i1} & \dots & a_{ii-1} \end{pmatrix}$

$$L_i = \begin{pmatrix} L_{i-1} & 0 \\ l_i & 1 \end{pmatrix}, l_i = \begin{pmatrix} l_{i1} & \dots & l_{ii-1} \end{pmatrix}, U_i = \begin{pmatrix} U_{i-1} & u_i \\ 0 & u_{ii} \end{pmatrix}, u_i = \begin{pmatrix} u_{1i} \\ \vdots \\ u_{i-1,i} \end{pmatrix}$$

Proof.

Proof for sufficient condition, p2

## Proof.

Proof for sufficient condition, p2

- ▶ Assume  $A_{i-1} = L_{i-1}U_{i-1}$

## Proof.

Proof for sufficient condition, p2

► Assume  $A_{i-1} = L_{i-1} U_{i-1}$

► Denote  $A_i = \begin{pmatrix} A_{i-1} & c_i \\ d_i & a_{ii} \end{pmatrix}$ ,  $c_i = \begin{pmatrix} a_{1i} \\ \vdots \\ a_{i-1i} \end{pmatrix}$ ,  $d_i = (a_{i1} \quad \dots \quad a_{ii-1})$

$$L_i = \begin{pmatrix} L_{i-1} & 0 \\ l_i & 1 \end{pmatrix}, l_i = (l_{i1} \quad \dots \quad l_{ii-1}), U_i = \begin{pmatrix} U_{i-1} & u_i \\ 0 & u_{ii} \end{pmatrix}, u_i = \begin{pmatrix} u_{1i} \\ \vdots \\ u_{i-1i} \end{pmatrix}$$

## Proof.

Proof for sufficient condition, p2

► Assume  $A_{i-1} = L_{i-1}U_{i-1}$

► Denote  $A_i = \begin{pmatrix} A_{i-1} & c_i \\ d_i & a_{ii} \end{pmatrix}$ ,  $c_i = \begin{pmatrix} a_{1i} \\ \vdots \\ a_{i-1i} \end{pmatrix}$ ,  $d_i = (a_{i1} \quad \dots \quad a_{ii-1})$

$$L_i = \begin{pmatrix} L_{i-1} & 0 \\ l_i & 1 \end{pmatrix}, l_i = (l_{i1} \quad \dots \quad l_{ii-1}), U_i = \begin{pmatrix} U_{i-1} & u_i \\ 0 & u_{ii} \end{pmatrix}, u_i = \begin{pmatrix} u_{1i} \\ \vdots \\ u_{i-1i} \end{pmatrix}$$

► Formula to prove:  $A_i = L_iU_i$

## Proof.

Proof for sufficient condition, p2

► Assume  $A_{i-1} = L_{i-1}U_{i-1}$

► Denote  $A_i = \begin{pmatrix} A_{i-1} & c_i \\ d_i & a_{ii} \end{pmatrix}$ ,  $c_i = \begin{pmatrix} a_{1i} \\ \vdots \\ a_{i-1i} \end{pmatrix}$ ,  $d_i = (a_{i1} \quad \dots \quad a_{ii-1})$

$$L_i = \begin{pmatrix} L_{i-1} & 0 \\ l_i & 1 \end{pmatrix}, l_i = (l_{i1} \quad \dots \quad l_{ii-1}), U_i = \begin{pmatrix} U_{i-1} & u_i \\ 0 & u_{ii} \end{pmatrix}, u_i = \begin{pmatrix} u_{1i} \\ \vdots \\ u_{i-1i} \end{pmatrix}$$

► Formula to prove:  $A_i = L_i U_i$

$$\text{► } \begin{pmatrix} L_{i-1} & 0 \\ l_i & 1 \end{pmatrix} \begin{pmatrix} U_{i-1} & u_i \\ 0 & u_{ii} \end{pmatrix} = \begin{pmatrix} A_{i-1} & c_i \\ d_i & a_{ii} \end{pmatrix} \Rightarrow \begin{cases} L_{i-1}U_{i-1} = A_{i-1} \\ L_{i-1}u_i = c_i \\ l_i U_{i-1} = d_i \\ l_i \cdot u_i + u_{ii} = a_{ii} \end{cases}$$

## Proof.

Proof for sufficient condition, p2

► Assume  $A_{i-1} = L_{i-1}U_{i-1}$

► Denote  $A_i = \begin{pmatrix} A_{i-1} & c_i \\ d_i & a_{ii} \end{pmatrix}$ ,  $c_i = \begin{pmatrix} a_{1i} \\ \vdots \\ a_{i-1i} \end{pmatrix}$ ,  $d_i = (a_{i1} \quad \dots \quad a_{ii-1})$

$$L_i = \begin{pmatrix} L_{i-1} & 0 \\ l_i & 1 \end{pmatrix}, l_i = (l_{i1} \quad \dots \quad l_{ii-1}), U_i = \begin{pmatrix} U_{i-1} & u_i \\ 0 & u_{ii} \end{pmatrix}, u_i = \begin{pmatrix} u_{1i} \\ \vdots \\ u_{i-1i} \end{pmatrix}$$

► Formula to prove:  $A_i = L_i U_i$

$$\text{► } \begin{pmatrix} L_{i-1} & 0 \\ l_i & 1 \end{pmatrix} \begin{pmatrix} U_{i-1} & u_i \\ 0 & u_{ii} \end{pmatrix} = \begin{pmatrix} A_{i-1} & c_i \\ d_i & a_{ii} \end{pmatrix} \Rightarrow \begin{cases} L_{i-1}U_{i-1} = A_{i-1} \\ L_{i-1}u_i = c_i \\ l_i U_{i-1} = d_i \\ l_i \cdot u_i + u_{ii} = a_{ii} \end{cases}$$

►  $\det(L_{i-1}) \neq 0, \det(U_{i-1}) \neq 0 \Rightarrow$  proof is complete





# Existence and uniqueness of LU factorization 3

## Theorem 8.10

*Unique LU factorization(Doolittle) of  $A \in \mathbb{R}^{n \times n}$  exists iff*

- ▶  $\det(A_i) \neq 0, i = 1, 2, \dots, n - 1$
- ▶  $A_i$  -main submatrices,  $A \in \mathbb{R}^{i \times i}$

# Existence and uniqueness of LU factorization 3

## Theorem 8.10

*Unique LU factorization(Doolittle) of  $A \in \mathbb{R}^{n \times n}$  exists iff*

- ▶  $\det(A_i) \neq 0, i = 1, 2, \dots, n - 1$
- ▶  $A_i$  -main submatrices,  $A \in \mathbb{R}^{i \times i}$

## Proof.

Proof for necessary condition, p3:

# Existence and uniqueness of LU factorization 3

## Theorem 8.10

*Unique LU factorization (Doolittle) of  $A \in \mathbb{R}^{n \times n}$  exists iff*

- ▶  $\det(A_i) \neq 0, i = 1, 2, \dots, n - 1$
- ▶  $A_i$  -main submatrices,  $A \in \mathbb{R}^{i \times i}$

## Proof.

Proof for necessary condition, p3:

If unique LU factorization exists then first  $n - 1$  main submatrices are not degenerated

# Existence and uniqueness of LU factorization 3

## Theorem 8.10

*Unique LU factorization (Doolittle) of  $A \in \mathbb{R}^{n \times n}$  exists iff*

- ▶  $\det(A_i) \neq 0, i = 1, 2, \dots, n - 1$
- ▶  $A_i$  -main submatrices,  $A \in \mathbb{R}^{i \times i}$

## Proof.

Proof for necessary condition, p3:

If unique LU factorization exists then first  $n - 1$  main submatrices are not degenerated

- ▶ proof by contradiction

# Existence and uniqueness of LU factorization 3

## Theorem 8.10

*Unique LU factorization (Doolittle) of  $A \in \mathbb{R}^{n \times n}$  exists iff*

- ▶  $\det(A_i) \neq 0, i = 1, 2, \dots, n - 1$
- ▶  $A_i$  -main submatrices,  $A \in \mathbb{R}^{i \times i}$

## Proof.

Proof for necessary condition, p3:

If unique LU factorization exists then first  $n - 1$  main submatrices are not degenerated

- ▶ proof by contradiction
- ▶ Assume unique LU factorization exists and for some  $k < n$  main submatrix  $A_k, 1 \leq k < n$  is degenerated and  $A_{k-1}, 2 \leq k < n$  is not degenerated

# Existence and uniqueness of LU factorization 3

## Theorem 8.10

Unique  $LU$  factorization (Doolittle) of  $A \in \mathbb{R}^{n \times n}$  exists iff

- ▶  $\det(A_i) \neq 0, i = 1, 2, \dots, n - 1$
- ▶  $A_i$  -main submatrices,  $A \in \mathbb{R}^{i \times i}$

## Proof.

Proof for necessary condition, p3:

If unique  $LU$  factorization exists then first  $n - 1$  main submatrices are not degenerated

- ▶ proof by contradiction
- ▶ Assume unique  $LU$  factorization exists and for some  $k < n$  main submatrix  $A_k, 1 \leq k < n$  is degenerated and  $A_{k-1}, 2 \leq k < n$  is not degenerated

$$\text{▶ } A = \begin{pmatrix} 1 & 0 & \dots & 0 \\ l_{21} & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ l_{n1} & l_{n2} & \dots & 1 \end{pmatrix} \begin{pmatrix} u_{11} & u_{12} & \dots & u_{1n} \\ 0 & u_{22} & \dots & u_{2n} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & u_{nn} \end{pmatrix} \Rightarrow \det(A) = \prod_{i=1}^n u_{ii}$$

# Existence and uniqueness of LU factorization 4

Proof.

Proof for necessary condition, p4:

## Existence and uniqueness of LU factorization 4

### Proof.

Proof for necessary condition, p4:

$$\blacktriangleright A = \begin{pmatrix} 1 & 0 & \dots & 0 \\ l_{21} & 1 & \dots & 0 \\ \vdots & \vdots & \dots & \vdots \\ l_{n1} & l_{n2} & \dots & 1 \end{pmatrix} \begin{pmatrix} u_{11} & u_{12} & \dots & u_{1n} \\ 0 & u_{22} & \dots & u_{2n} \\ \vdots & \vdots & \dots & \vdots \\ 0 & 0 & \dots & u_{nn} \end{pmatrix} \Rightarrow \det(A) = \prod_{i=1}^n u_{ii}$$



## Existence and uniqueness of LU factorization 4

### Proof.

Proof for necessary condition, p4:

$$\blacktriangleright A = \begin{pmatrix} 1 & 0 & \dots & 0 \\ l_{21} & 1 & \dots & 0 \\ \vdots & \vdots & \dots & \vdots \\ l_{n1} & l_{n2} & \dots & 1 \end{pmatrix} \begin{pmatrix} u_{11} & u_{12} & \dots & u_{1n} \\ 0 & u_{22} & \dots & u_{2n} \\ \vdots & \vdots & \dots & \vdots \\ 0 & 0 & \dots & u_{nn} \end{pmatrix} \Rightarrow \det(A) = \prod_{i=1}^n u_{ii}$$

$$\blacktriangleright \det(A_{n-1}) = 0, \det(A_{n-2}) \neq 0 \Rightarrow u_{n-1,n-1} = 0$$

# Existence and uniqueness of LU factorization 4

## Proof.

Proof for necessary condition, p4:

$$\blacktriangleright A = \begin{pmatrix} 1 & 0 & \dots & 0 \\ l_{21} & 1 & \dots & 0 \\ \vdots & \vdots & \dots & \vdots \\ l_{n1} & l_{n2} & \dots & 1 \end{pmatrix} \begin{pmatrix} u_{11} & u_{12} & \dots & u_{1n} \\ 0 & u_{22} & \dots & u_{2n} \\ \vdots & \vdots & \dots & \vdots \\ 0 & 0 & \dots & u_{nn} \end{pmatrix} \Rightarrow \det(A) = \prod_{i=1}^n u_{ii}$$

$$\blacktriangleright \det(A_{n-1}) = 0, \det(A_{n-2}) \neq 0 \Rightarrow u_{n-1,n-1} = 0$$

$$\blacktriangleright \begin{pmatrix} L_{n-1} & 0 \\ l_n & 1 \end{pmatrix} \begin{pmatrix} U_{n-1} & u_n \\ 0 & u_{nn} \end{pmatrix} = \begin{pmatrix} A_{n-1} & c_n \\ d_n & a_{nn} \end{pmatrix} \Rightarrow \begin{cases} L_{n-1} U_{n-1} = A_{n-1} \\ L_{n-1} u_n = A_{n-1} c_n \\ l_n U_{n-1} = d_n \\ l_n \cdot u_n + u_{nn} = a_{nn} \end{cases}$$

# Existence and uniqueness of LU factorization 4

## Proof.

Proof for necessary condition, p4:

$$\blacktriangleright A = \begin{pmatrix} 1 & 0 & \dots & 0 \\ l_{21} & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ l_{n1} & l_{n2} & \dots & 1 \end{pmatrix} \begin{pmatrix} u_{11} & u_{12} & \dots & u_{1n} \\ 0 & u_{22} & \dots & u_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & u_{nn} \end{pmatrix} \Rightarrow \det(A) = \prod_{i=1}^n u_{ii}$$

$$\blacktriangleright \det(A_{n-1}) = 0, \det(A_{n-2}) \neq 0 \Rightarrow u_{n-1,n-1} = 0$$

$$\blacktriangleright \begin{pmatrix} L_{n-1} & 0 \\ l_n & 1 \end{pmatrix} \begin{pmatrix} U_{n-1} & u_n \\ 0 & u_{nn} \end{pmatrix} = \begin{pmatrix} A_{n-1} & c_n \\ d_n & a_{nn} \end{pmatrix} \Rightarrow \begin{cases} L_{n-1} U_{n-1} = A_{n-1} \\ L_{n-1} u_n = A_{n-1} c_n \\ l_n U_{n-1} = d_n \\ l_n \cdot u_n + u_{nn} = a_{nn} \end{cases}$$

$$\blacktriangleright \det(U_{n-1}) = 0 \Rightarrow l_n U_{n-1} = d_n \text{ has no unique solution} \Rightarrow \text{contradiction, proof is complete}$$



# LU factorization from Gaussian elimination 1

## Example 8.11

$$\blacktriangleright \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$$

# LU factorization from Gaussian elimination 1

## Example 8.11

$$\blacktriangleright \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$$

$$\blacktriangleright \begin{pmatrix} 1 & 0 & 0 \\ x & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = \begin{pmatrix} y_1 \\ xy_1 + y_2 \\ y_3 \end{pmatrix}$$

# LU factorization from Gaussian elimination 1

## Example 8.11

$$\blacktriangleright \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$$

$$\blacktriangleright \begin{pmatrix} 1 & 0 & 0 \\ x & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = \begin{pmatrix} y_1 \\ xy_1 + y_2 \\ y_3 \end{pmatrix}$$

$$\blacktriangleright \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ x & 0 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = \begin{pmatrix} y_1 \\ y_2 \\ xy_1 + y_3 \end{pmatrix}$$

# LU factorization from Gaussian elimination 1

## Example 8.11

$$\blacktriangleright \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$$

$$\blacktriangleright \begin{pmatrix} 1 & 0 & 0 \\ x & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = \begin{pmatrix} y_1 \\ xy_1 + y_2 \\ y_3 \end{pmatrix}$$

$$\blacktriangleright \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ x & 0 & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = \begin{pmatrix} y_1 \\ y_2 \\ xy_1 + y_3 \end{pmatrix}$$

$$\blacktriangleright \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & x & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} y_0 \\ y_1 \\ y_2 \\ y_3 \\ y_4 \end{pmatrix} = \begin{pmatrix} y_0 \\ y_1 \\ y_2 \\ xy_1 + y_3 \\ y_4 \end{pmatrix}$$

# LU factorization from Gaussian elimination 2

## Example 8.12

$$\blacktriangleright \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ x_2 & 1 & 0 & \dots & 0 \\ x_3 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ x_k & 0 & 0 & & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \\ \dots \\ y_k \end{pmatrix} = \begin{pmatrix} y_1 \\ x_2 y_1 + y_2 \\ x_3 y_1 + y_3 \\ \dots \\ x_k y_1 + y_k \end{pmatrix}$$



# LU factorization from Gaussian elimination 2

## Example 8.12

$$\blacktriangleright \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ x_2 & 1 & 0 & \dots & 0 \\ x_3 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ x_k & 0 & 0 & \dots & 1 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \\ \vdots \\ y_k \end{pmatrix} = \begin{pmatrix} y_1 \\ x_2 y_1 + y_2 \\ x_3 y_1 + y_3 \\ \vdots \\ x_k y_1 + y_k \end{pmatrix}$$

$$\blacktriangleright \begin{pmatrix} y_1 \\ y_2 \\ y_3 \\ \vdots \\ y_k \end{pmatrix} = \begin{pmatrix} y_1 \\ 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix} \Rightarrow \begin{cases} x_2 = -\frac{y_2}{y_1} \\ x_3 = -\frac{y_3}{y_1} \\ \vdots \\ x_k = -\frac{y_k}{y_1} \end{cases} \Rightarrow \begin{cases} \text{Gaussian elimination} \\ \text{multipliers} = x_i, i = 2, \dots, n \end{cases}$$

# LU factorization from Gaussian elimination 3

►  $A \equiv A^{(0)}$

## LU factorization from Gaussian elimination 3

►  $A \equiv A^{(0)}$

►  $A^{(1)} = M_1 A^{(0)} = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ -\frac{a_{21}}{a_{11}} & 1 & 0 & \dots & 0 \\ -\frac{a_{31}}{a_{11}} & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -\frac{a_{n1}}{a_{11}} & 0 & 0 & \dots & 1 \end{pmatrix} \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \dots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{pmatrix}$

# LU factorization from Gaussian elimination 3

►  $A \equiv A^{(0)}$

►  $A^{(1)} = M_1 A^{(0)} = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ -\frac{a_{21}}{a_{11}} & 1 & 0 & \dots & 0 \\ -\frac{a_{31}}{a_{11}} & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ -\frac{a_{n1}}{a_{11}} & 0 & 0 & \dots & 1 \end{pmatrix} \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \dots & a_{3n} \\ \dots & \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{pmatrix}$

►  $A^{(1)} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ 0 & a_{22}^{(1)} & a_{23}^{(1)} & \dots & a_{2n}^{(1)} \\ 0 & a_{32}^{(1)} & a_{33}^{(1)} & \dots & a_{3n}^{(1)} \\ \dots & \dots & \dots & \dots & \dots \\ 0 & a_{n2}^{(1)} & a_{n3}^{(1)} & \dots & a_{nn}^{(1)} \end{pmatrix}, A^{(2)} = M_2 A^{(1)}$

# LU factorization from Gaussian elimination 3

►  $A \equiv A^{(0)}$

►  $A^{(1)} = M_1 A^{(0)} = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ -\frac{a_{21}}{a_{11}} & 1 & 0 & \dots & 0 \\ -\frac{a_{31}}{a_{11}} & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ -\frac{a_{n1}}{a_{11}} & 0 & 0 & \dots & 1 \end{pmatrix} \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \dots & a_{3n} \\ \dots & \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{pmatrix}$

►  $A^{(1)} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ 0 & a_{22}^{(1)} & a_{23}^{(1)} & \dots & a_{2n}^{(1)} \\ 0 & a_{32}^{(1)} & a_{33}^{(1)} & \dots & a_{3n}^{(1)} \\ \dots & \dots & \dots & \dots & \dots \\ 0 & a_{n2}^{(1)} & a_{n3}^{(1)} & \dots & a_{nn}^{(1)} \end{pmatrix}, A^{(2)} = M_2 A^{(1)}$

►  $M_2 A^{(1)} = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & -\frac{a_{32}^{(1)}}{a_{22}^{(1)}} & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & -\frac{a_{n2}^{(1)}}{a_{22}^{(1)}} & 0 & \dots & 1 \end{pmatrix} \cdot A^{(2)} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ 0 & a_{22}^{(1)} & a_{23}^{(2)} & \dots & a_{2n}^{(2)} \\ 0 & 0 & a_{33}^{(2)} & \dots & a_{3n}^{(2)} \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & a_{n3}^{(2)} & \dots & a_{nn}^{(2)} \end{pmatrix}$

## LU factorization from Gaussian elimination 4

$$\blacktriangleright A^{(k-1)} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & \dots & a_{1n} \\ 0 & a_{22}^{(1)} & a_{23}^{(2)} & \dots & \dots & a_{2n}^{(2)} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & a_{kk}^{(k-1)} & \dots & \dots & a_{kn}^{(k-1)} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & a_{nk}^{(k-1)} & \dots & \dots & a_{nn}^{(k-1)} \end{pmatrix}$$

# LU factorization from Gaussian elimination 4

$$\blacktriangleright A^{(k-1)} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & \dots & a_{1n} \\ 0 & a_{22}^{(1)} & a_{23}^{(2)} & \dots & \dots & a_{2n}^{(2)} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & a_{kk}^{(k-1)} & \dots & \dots & a_{kn}^{(k-1)} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & a_{nk}^{(k-1)} & \dots & \dots & a_{nn}^{(k-1)} \end{pmatrix}$$

$$A^{(k)} = M_k \cdot A^{(k-1)}$$

## LU factorization from Gaussian elimination 4

$$\blacktriangleright A^{(k-1)} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & \dots & a_{1n} \\ 0 & a_{22}^{(1)} & a_{23}^{(2)} & \dots & \dots & a_{2n}^{(2)} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & a_{kk}^{(k-1)} & \dots & \dots & a_{kn}^{(k-1)} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & a_{nk}^{(k-1)} & \dots & \dots & a_{nn}^{(k-1)} \end{pmatrix}$$

$$A^{(k)} = M_k \cdot A^{(k-1)}$$

- $\blacktriangleright$  Zero entries under main diagonal in the first  $k$  columns of  $A^{(k)}$



# LU factorization from Gaussian elimination 4

$$\blacktriangleright A^{(k-1)} = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & \dots & a_{1n} \\ 0 & a_{22}^{(1)} & a_{23}^{(2)} & \dots & \dots & a_{2n}^{(2)} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & a_{kk}^{(k-1)} & \dots & \dots & a_{kn}^{(k-1)} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & a_{nk}^{(k-1)} & \dots & \dots & a_{nn}^{(k-1)} \end{pmatrix}$$

$$A^{(k)} = M_k \cdot A^{(k-1)}$$

- $\blacktriangleright$  Zero entries under main diagonal in the first  $k$  columns of  $A^{(k)}$

$$\blacktriangleright M_k = \begin{pmatrix} 1 & 0 & 0 & \dots & \dots & \dots & 0 \\ 0 & 1 & 0 & \dots & \dots & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & 1 & \dots & \dots & \dots \\ \dots & \dots & \dots & -\frac{a_{k+1,k}^{(k-1)}}{a_{kk}^{(k-1)}} & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & -\frac{a_{nk}^{(k-1)}}{a_{kk}^{(k-1)}} & \dots & \dots & 1 \end{pmatrix}, \quad \left\{ \begin{array}{l} \text{multipliers :} \\ m_{k+1,k} = -\frac{a_{k+1,k}^{(k-1)}}{a_{kk}^{(k-1)}}, \\ \dots \\ m_{n,k} = -\frac{a_{nk}^{(k-1)}}{a_{kk}^{(k-1)}} \end{array} \right.$$

# LU factorization from Gaussian elimination 5



$$A^{(0)} = A, A^{(1)} = M_1 \cdot A^{(0)}, \dots, A^{(k)} = M_k \cdot A^{(k-1)}$$

## LU factorization from Gaussian elimination 5



$$A^{(0)} = A, A^{(1)} = M_1 \cdot A^{(0)}, \dots, A^{(k)} = M_k \cdot A^{(k-1)}$$



$$A^{(n-1)} = M_{n-1} \cdot A^{(n-2)} \equiv U$$

# LU factorization from Gaussian elimination 5



$$A^{(0)} = A, A^{(1)} = M_1 \cdot A^{(0)}, \dots, A^{(k)} = M_k \cdot A^{(k-1)}$$



$$A^{(n-1)} = M_{n-1} \cdot A^{(n-2)} \equiv U$$

►  $U = \begin{pmatrix} a_{11}^{(0)} & a_{11}^{(0)} & a_{13}^{(0)} & \dots & \dots & \dots & a_{1n}^{(0)} \\ 0 & a_{22}^{(1)} & a_{23}^{(1)} & \dots & \dots & \dots & a_{2n}^{(1)} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & a_{kk}^{(k-1)} & \dots & \dots & a_{kn}^{(k-1)} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & a_{n-1n-1}^{(n-2)} & a_{n-2n}^{(n-1)} \\ 0 & \dots & \dots & \dots & \dots & 0 & a_{nn}^{(n-1)} \end{pmatrix}$

# LU factorization from Gaussian elimination 5

$$A^{(0)} = A, A^{(1)} = M_1 \cdot A^{(0)}, \dots, A^{(k)} = M_k \cdot A^{(k-1)}$$

$$A^{(n-1)} = M_{n-1} \cdot A^{(n-2)} \equiv U$$

$$U = \begin{pmatrix} a_{11}^{(0)} & a_{11}^{(0)} & a_{13}^{(0)} & \dots & \dots & \dots & a_{1n}^{(0)} \\ 0 & a_{22}^{(1)} & a_{23}^{(1)} & \dots & \dots & \dots & a_{2n}^{(1)} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & a_{kk}^{(k-1)} & \dots & \dots & a_{kn}^{(k-1)} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & a_{n-1n-1}^{(n-2)} & a_{n-2n}^{(n-1)} \\ 0 & \dots & \dots & \dots & \dots & 0 & a_{nn}^{(n-1)} \end{pmatrix}$$

$$U = M_{n-1} \cdot A^{(n-2)} = \dots = M_{n-1} M_{n-2} \dots M_1 \cdot A^{(0)}$$

# LU factorization from Gaussian elimination 5

$$A^{(0)} = A, A^{(1)} = M_1 \cdot A^{(0)}, \dots, A^{(k)} = M_k \cdot A^{(k-1)}$$

$$A^{(n-1)} = M_{n-1} \cdot A^{(n-2)} \equiv U$$

$$U = \begin{pmatrix} a_{11}^{(0)} & a_{11}^{(0)} & a_{13}^{(0)} & \dots & \dots & \dots & a_{1n}^{(0)} \\ 0 & a_{22}^{(1)} & a_{23}^{(1)} & \dots & \dots & \dots & a_{2n}^{(1)} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & a_{kk}^{(k-1)} & \dots & \dots & a_{kn}^{(k-1)} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & a_{n-1n-1}^{(n-2)} & a_{n-2n}^{(n-1)} \\ 0 & \dots & \dots & \dots & \dots & 0 & a_{nn}^{(n-1)} \end{pmatrix}$$

$$U = M_{n-1} \cdot A^{(n-2)} = \dots = M_{n-1} M_{n-2} \dots M_1 \cdot A^{(0)}$$

$$M_{n-1} M_{n-2} \dots M_1 \cdot A = U \Rightarrow A = (M_{n-1} M_{n-2} \dots M_1)^{-1} U$$

# LU factorization from Gaussian elimination 5

$$A^{(0)} = A, A^{(1)} = M_1 \cdot A^{(0)}, \dots, A^{(k)} = M_k \cdot A^{(k-1)}$$

$$A^{(n-1)} = M_{n-1} \cdot A^{(n-2)} \equiv U$$

$$U = \begin{pmatrix} a_{11}^{(0)} & a_{11}^{(0)} & a_{13}^{(0)} & \dots & \dots & \dots & a_{1n}^{(0)} \\ 0 & a_{22}^{(1)} & a_{23}^{(1)} & \dots & \dots & \dots & a_{2n}^{(1)} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & a_{kk}^{(k-1)} & \dots & \dots & a_{kn}^{(k-1)} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & a_{n-1n-1}^{(n-2)} & a_{n-2n}^{(n-1)} \\ 0 & \dots & \dots & \dots & \dots & 0 & a_{nn}^{(n-1)} \end{pmatrix}$$

$$U = M_{n-1} \cdot A^{(n-2)} = \dots = M_{n-1} M_{n-2} \dots M_1 \cdot A^{(0)}$$

$$M_{n-1} M_{n-2} \dots M_1 \cdot A = U \Rightarrow A = (M_{n-1} M_{n-2} \dots M_1)^{-1} U$$

$$(CB)^{-1} = B^{-1} C^{-1} \Rightarrow A = M_1^{-1} M_2^{-1} \dots M_{n-2}^{-1} M_{n-1}^{-1} U$$

# LU factorization from Gaussian elimination 6



$$A = M_1^{-1} M_2^{-1} \dots M_{n-2}^{-1} M_{n-1}^{-1} U$$



# LU factorization from Gaussian elimination 6



$$A = M_1^{-1} M_2^{-1} \dots M_{n-2}^{-1} M_{n-1}^{-1} U$$



$$\begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ x_2 & 1 & 0 & \dots & 0 \\ x_3 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ x_k & 0 & 0 & & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ -x_2 & 1 & 0 & \dots & 0 \\ -x_3 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ -x_k & 0 & 0 & & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & & 1 \end{pmatrix}$$

# LU factorization from Gaussian elimination 6



$$A = M_1^{-1} M_2^{-1} \dots M_{n-2}^{-1} M_{n-1}^{-1} U$$



$$\begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ x_2 & 1 & 0 & \dots & 0 \\ x_3 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ x_k & 0 & 0 & \dots & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ -x_2 & 1 & 0 & \dots & 0 \\ -x_3 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ -x_k & 0 & 0 & \dots & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 1 \end{pmatrix}$$



$$M_k = \begin{pmatrix} 1 & 0 & 0 & \dots & \dots & \dots & 0 \\ 0 & 1 & 0 & \dots & \dots & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & 1 & \dots & \dots & \dots \\ \dots & \dots & \dots & m_{k+1k}^{(k-1)} & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & m_{nk}^{(k-1)} & \dots & \dots & 1 \end{pmatrix}, M_k^{-1} = \begin{pmatrix} 1 & 0 & 0 & \dots & \dots & \dots & \dots \\ 0 & 1 & 0 & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & 1 & \dots & \dots & \dots \\ \dots & \dots & \dots & -m_{k+1k}^{(k-1)} & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & -m_{nk}^{(k-1)} & \dots & \dots & \dots \end{pmatrix}$$

## LU factorization from Gaussian elimination 7

$$M_k = \begin{pmatrix} 1 & 0 & 0 & .. & .. & .. & 0 \\ 0 & 1 & 0 & .. & .. & .. & 0 \\ .. & .. & .. & .. & .. & .. & .. \\ .. & .. & .. & 1 & .. & .. & .. \\ .. & .. & .. & -\frac{a_{k+1k}^{(k-1)}}{a_{kk}^{(k-1)}} & .. & .. & .. \\ .. & .. & .. & .. & .. & .. & .. \\ 0 & .. & .. & -\frac{a_{nk}^{(k-1)}}{a_{kk}^{(k-1)}} & .. & .. & 1 \end{pmatrix}$$

# LU factorization from Gaussian elimination 7

$$M_k = \begin{pmatrix} 1 & 0 & 0 & .. & .. & .. & 0 \\ 0 & 1 & 0 & .. & .. & .. & 0 \\ .. & .. & .. & .. & .. & .. & .. \\ .. & .. & .. & 1 & .. & .. & .. \\ .. & .. & .. & -\frac{a_{k+1k}^{(k-1)}}{a_{kk}^{(k-1)}} & .. & .. & .. \\ .. & .. & .. & .. & .. & .. & .. \\ 0 & .. & .. & -\frac{a_{nk}^{(k-1)}}{a_{kk}^{(k-1)}} & .. & .. & 1 \end{pmatrix}$$

$$M_k^{-1} = \begin{pmatrix} 1 & 0 & 0 & .. & .. & .. & 0 \\ 0 & 1 & 0 & .. & .. & .. & 0 \\ .. & .. & .. & .. & .. & .. & .. \\ .. & .. & .. & 1 & .. & .. & .. \\ .. & .. & .. & \frac{a_{k+1k}^{(k-1)}}{a_{kk}^{(k-1)}} & .. & .. & .. \\ .. & .. & .. & .. & .. & .. & .. \\ 0 & .. & .. & \frac{a_{nk}^{(k-1)}}{a_{kk}^{(k-1)}} & .. & .. & 1 \end{pmatrix}$$

# LU factorization from Gaussian elimination 8

- ▶ Product of lower triangular matrices is lower triangular matrix

# LU factorization from Gaussian elimination 8

- ▶ Product of lower triangular matrices is lower triangular matrix
- ▶ Matrices  $M_k^{-1}, k = 1, 2, \dots, n - 1$  are lower triangular matrices

# LU factorization from Gaussian elimination 8

- ▶ Product of lower triangular matrices is lower triangular matrix
- ▶ Matrices  $M_k^{-1}, k = 1, 2, \dots, n - 1$  are lower triangular matrices
- ▶  $L = M_1^{-1} M_2^{-1} \dots M_{n-2}^{-1} M_{n-1}^{-1}$  is lower triangular matrix

# LU factorization from Gaussian elimination 8

- ▶ Product of lower triangular matrices is lower triangular matrix
- ▶ Matrices  $M_k^{-1}, k = 1, 2, \dots, n - 1$  are lower triangular matrices
- ▶  $L = M_1^{-1}M_2^{-1}\dots M_{n-2}^{-1}M_{n-1}^{-1}$  is lower triangular matrix

▶

$$A = M_1^{-1}M_2^{-1}\dots M_{n-2}^{-1}M_{n-1}^{-1}U$$



# LU factorization from Gaussian elimination 8

- ▶ Product of lower triangular matrices is lower triangular matrix
- ▶ Matrices  $M_k^{-1}, k = 1, 2, \dots, n - 1$  are lower triangular matrices
- ▶  $L = M_1^{-1}M_2^{-1}\dots M_{n-2}^{-1}M_{n-1}^{-1}$  is lower triangular matrix



$$A = M_1^{-1}M_2^{-1}\dots M_{n-2}^{-1}M_{n-1}^{-1}U$$



$$A = LU$$

# LU factorization from Gaussian elimination 8

- ▶ Product of lower triangular matrices is lower triangular matrix
- ▶ Matrices  $M_k^{-1}, k = 1, 2, \dots, n - 1$  are lower triangular matrices
- ▶  $L = M_1^{-1}M_2^{-1}\dots M_{n-2}^{-1}M_{n-1}^{-1}$  is lower triangular matrix



$$A = M_1^{-1}M_2^{-1}\dots M_{n-2}^{-1}M_{n-1}^{-1}U$$



$$A = LU$$

- ▶ Explicit expression for  $L$ ?

# LU factorization from Gaussian elimination 9

- Explicit expression for  $L = M_1^{-1}M_2^{-1}\dots M_{n-2}^{-1}M_{n-1}^{-1}$  ?

# LU factorization from Gaussian elimination 9

- Explicit expression for  $L = M_1^{-1}M_2^{-1}...M_{n-2}^{-1}M_{n-1}^{-1}$  ?

► 
$$\begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ x_2 & 1 & 0 & \dots & 0 \\ x_3 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ x_n & 0 & 0 & & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & y_3 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & y_n & 0 & & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ x_2 & 1 & 0 & \dots & 0 \\ x_3 & y_3 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ x_n & y_n & 0 & & 1 \end{pmatrix}$$

# LU factorization from Gaussian elimination 9

- Explicit expression for  $L = M_1^{-1}M_2^{-1}...M_{n-2}^{-1}M_{n-1}^{-1}$  ?

► 
$$\begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ x_2 & 1 & 0 & \dots & 0 \\ x_3 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ x_n & 0 & 0 & & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & y_3 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & y_n & 0 & & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ x_2 & 1 & 0 & \dots & 0 \\ x_3 & y_3 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ x_n & y_n & 0 & & 1 \end{pmatrix}$$

► 
$$M_k^{-1} = \begin{pmatrix} 1 & 0 & 0 & \dots & \dots & \dots & 0 \\ 0 & 1 & 0 & \dots & \dots & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & 1 & \dots & \dots & \dots \\ \dots & \dots & \dots & -m_{k+1k}^{(k-1)} & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & -m_{nk}^{(k-1)} & \dots & \dots & 1 \end{pmatrix}$$

# LU factorization from Gaussian elimination 9

- Explicit expression for  $L = M_1^{-1}M_2^{-1}\dots M_{n-2}^{-1}M_{n-1}^{-1}$

$$\begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ x_2 & 1 & 0 & \dots & 0 \\ x_3 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ x_n & 0 & 0 & \dots & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & y_3 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & y_n & 0 & \dots & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ x_2 & 1 & 0 & \dots & 0 \\ x_3 & y_3 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ x_n & y_n & 0 & \dots & 1 \end{pmatrix}$$

# LU factorization from Gaussian elimination 9

► Explicit expression for  $L = M_1^{-1}M_2^{-1}...M_{n-2}^{-1}M_{n-1}^{-1}$

► 
$$\begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ x_2 & 1 & 0 & \dots & 0 \\ x_3 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ x_n & 0 & 0 & & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & y_3 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & y_n & 0 & & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ x_2 & 1 & 0 & \dots & 0 \\ x_3 & y_3 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ x_n & y_n & 0 & & 1 \end{pmatrix}$$

► 
$$L = \begin{pmatrix} 1 & 0 & 0 & \dots & \dots & \dots & 0 \\ -m_{21}^{(1)} & 1 & 0 & \dots & \dots & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & 1 & \dots & \dots & \dots \\ -m_{k1}^{(1)} & \dots & \dots & -m_{k+1k}^{(k-1)} & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ -m_{n1}^{(1)} & \dots & \dots & -m_{nk}^{(k-1)} & \dots & -m_{nn-1}^{(n-1)} & 1 \end{pmatrix}$$

# LU factorization from Gaussian elimination 10

- ▶ Efficient storage scheme for  $LU$  factorization
  - ▶ store entries of  $U$  in upper triangular part of  $A$  including diagonal
  - ▶ store entries of  $L$  in lower triangular part of  $A$  excluding diagonal



# LU factorization from Gaussian elimination 10

- ▶ Efficient storage scheme for  $LU$  factorization

- ▶ store entries of  $U$  in upper triangular part of  $A$  including diagonal
- ▶ store entries of  $L$  in lower triangular part of  $A$  excluding diagonal

- ▶ 
$$\begin{pmatrix} a_{11}^{(0)} & a_{12}^{(0)} & a_{13}^{(0)} & \dots & \dots & \dots & a_{1n}^{(0)} \\ -m_{21} & a_{22}^{(1)} & a_{23}^{(1)} & \dots & \dots & \dots & a_{2n}^{(1)} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & a_{kk}^{(k-1)} & \dots & \dots & \dots \\ -m_{k+11} & \dots & \dots & -m_{k+1k} & \dots & \dots & a_{k+1n}^{(k)} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ -m_{n1} & \dots & \dots & -m_{nk} & \dots & \dots & a_{nn}^{(n-1)} \end{pmatrix}$$

# LU factorization from Gaussian elimination 11

1. Given input matrix  $A$ .

# LU factorization from Gaussian elimination 11

1. Given input matrix  $A$ .
2. Algorithm: for  $k = 1, 2, \dots, n - 1$

# LU factorization from Gaussian elimination 11

1. Given input matrix  $A$ .
2. Algorithm: for  $k = 1, 2, \dots, n - 1$ 
  - Compute multipliers and update some entries of  $k$ -th column

$$m_{ik} = \frac{a_{ik}}{a_{kk}}, a_{ik} \equiv m_{ik}, i = k + 1, \dots, n$$

# LU factorization from Gaussian elimination 11

1. Given input matrix  $A$ .
2. Algorithm: for  $k = 1, 2, \dots, n - 1$ 
  - Compute multipliers and update some entries of  $k$ -th column

$$m_{ik} = \frac{a_{ik}}{a_{kk}}, a_{ik} \equiv m_{ik}, i = k + 1, \dots, n$$

- Elimination step = update entries of submatrix  $(k + 1 : n, k + 1 : n)$

$$a_{ij} = a_{ij} + m_{ik} a_{kj}, i = k + 1, \dots, n, j = k + 1, \dots, n$$

# LU factorization from Gaussian elimination 11

1. Given input matrix  $A$ .

2. Algorithm: for  $k = 1, 2, \dots, n - 1$

► Compute multipliers and update some entries of  $k$ -th column

$$m_{ik} = \frac{a_{ik}}{a_{kk}}, a_{ik} \equiv m_{ik}, i = k + 1, \dots, n$$

► Elimination step = update entries of submatrix  $(k + 1 : n, k + 1 : n)$

$$a_{ij} = a_{ij} + m_{ik} a_{kj}, i = k + 1, \dots, n, j = k + 1, \dots, n$$

3. Output matrix  $A =$

$$\begin{pmatrix} a_{11}^{(0)} & a_{12}^{(0)} & a_{13}^{(0)} & \dots & \dots & \dots & a_{1n}^{(0)} \\ -m_{21} & a_{22}^{(1)} & a_{23}^{(1)} & \dots & \dots & \dots & a_{2n}^{(1)} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & a_{kk}^{(k-1)} & \dots & \dots & \dots \\ -m_{k+11} & \dots & \dots & -m_{k+1k} & \dots & \dots & a_{k+1n}^{(k)} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ -m_{n1} & \dots & \dots & -m_{nk} & \dots & \dots & a_{nn}^{(n-1)} \end{pmatrix}$$

# Difficulties with Gaussian elimination

## Example 8.13

Forsythe, Moler (1967)

- Impact of Finite Digit Arithmetic(case: 3 digit arithmetic)

# Difficulties with Gaussian elimination

## Example 8.13

Forsythe, Moler (1967)

- ▶ Impact of Finite Digit Arithmetic(case: 3 digit arithmetic)



$$A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$$



# Difficulties with Gaussian elimination

## Example 8.13

Forsythe, Moler (1967)

- ▶ Impact of Finite Digit Arithmetic(case: 3 digit arithmetic)



$$A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$$

- ▶ Exact  $LU$ :

# Difficulties with Gaussian elimination

## Example 8.13

Forsythe, Moler (1967)

- ▶ Impact of Finite Digit Arithmetic(case: 3 digit arithmetic)



$$A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$$

- ▶ Exact  $LU$ :  $m_{21} = -10^4$ ,  $L = \begin{pmatrix} 1 & 0 \\ 10^4 & 1 \end{pmatrix}$

# Difficulties with Gaussian elimination

## Example 8.13

Forsythe, Moler (1967)

- Impact of Finite Digit Arithmetic(case: 3 digit arithmetic)



$$A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$$

- Exact  $LU$ :  $m_{21} = -10^4$ ,  $L = \begin{pmatrix} 1 & 0 \\ 10^4 & 1 \end{pmatrix}$

$$U = A^{(1)} = \begin{pmatrix} 10^{-4} & 1 \\ 0 & 1 - 10^4 \end{pmatrix}, LU = A$$

# Difficulties with Gaussian elimination

## Example 8.13

Forsythe, Moler (1967)

- ▶ Impact of Finite Digit Arithmetic(case: 3 digit arithmetic)



$$A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$$

- ▶ Exact  $LU$ :  $m_{21} = -10^4$ ,  $L = \begin{pmatrix} 1 & 0 \\ 10^4 & 1 \end{pmatrix}$

$$U = A^{(1)} = \begin{pmatrix} 10^{-4} & 1 \\ 0 & 1 - 10^4 \end{pmatrix}, LU = A$$

- ▶ In three digit arithmetic with rounding

# Difficulties with Gaussian elimination

## Example 8.13

Forsythe, Moler (1967)

- Impact of Finite Digit Arithmetic(case: 3 digit arithmetic)



$$A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$$

- Exact  $LU$ :  $m_{21} = -10^4$ ,  $L = \begin{pmatrix} 1 & 0 \\ 10^4 & 1 \end{pmatrix}$

$$U = A^{(1)} = \begin{pmatrix} 10^{-4} & 1 \\ 0 & 1 - 10^4 \end{pmatrix}, LU = A$$

- In three digit arithmetic with rounding

$$fl(1 - 10^{-4}) = fl(-9999) = fl(-0.9999 \cdot 10^4) = -10^4$$

# Difficulties with Gaussian elimination

## Example 8.13

Forsythe, Moler (1967)

- Impact of Finite Digit Arithmetic(case: 3 digit arithmetic)



$$A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$$

- Exact  $LU$ :  $m_{21} = -10^4$ ,  $L = \begin{pmatrix} 1 & 0 \\ 10^4 & 1 \end{pmatrix}$

$$U = A^{(1)} = \begin{pmatrix} 10^{-4} & 1 \\ 0 & 1 - 10^4 \end{pmatrix}, LU = A$$

- In three digit arithmetic with rounding

$$fl(1 - 10^{-4}) = fl(-9999) = fl(-0.9999 \cdot 10^4) = -10^4 \quad L = \begin{pmatrix} 1 & 0 \\ 10^4 & 1 \end{pmatrix},$$

# Difficulties with Gaussian elimination

## Example 8.13

Forsythe, Moler (1967)

- Impact of Finite Digit Arithmetic(case: 3 digit arithmetic)



$$A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$$

- Exact  $LU$ :  $m_{21} = -10^4$ ,  $L = \begin{pmatrix} 1 & 0 \\ 10^4 & 1 \end{pmatrix}$

$$U = A^{(1)} = \begin{pmatrix} 10^{-4} & 1 \\ 0 & 1 - 10^4 \end{pmatrix}, LU = A$$

- In three digit arithmetic with rounding

$$fl(1 - 10^{-4}) = fl(-9999) = fl(-0.9999 \cdot 10^4) = -10^4 \quad L = \begin{pmatrix} 1 & 0 \\ 10^4 & 1 \end{pmatrix},$$

$$U = A^{(1)} = \begin{pmatrix} 10^{-4} & 1 \\ 0 & -10^4 \end{pmatrix},$$

# Difficulties with Gaussian elimination

## Example 8.13

Forsythe, Moler (1967)

- Impact of Finite Digit Arithmetic(case: 3 digit arithmetic)



$$A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$$

- Exact  $LU$ :  $m_{21} = -10^4$ ,  $L = \begin{pmatrix} 1 & 0 \\ 10^4 & 1 \end{pmatrix}$

$$U = A^{(1)} = \begin{pmatrix} 10^{-4} & 1 \\ 0 & 1 - 10^4 \end{pmatrix}, LU = A$$

- In three digit arithmetic with rounding

$$fl(1 - 10^{-4}) = fl(-9999) = fl(-0.9999 \cdot 10^4) = -10^4 \quad L = \begin{pmatrix} 1 & 0 \\ 10^4 & 1 \end{pmatrix},$$

$$U = A^{(1)} = \begin{pmatrix} 10^{-4} & 1 \\ 0 & -10^4 \end{pmatrix}, LU = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 0 \end{pmatrix} \neq A$$



# Pivoting and Gaussian elimination 1

## Example 8.14

Forsythe, Moler (1967)

- Impact of pivoting on finite precision arithmetic (3 digit example)

# Pivoting and Gaussian elimination 1

## Example 8.14

Forsythe, Moler (1967)

- ▶ Impact of pivoting on finite precision arithmetic (3 digit example)

- ▶  $A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$

# Pivoting and Gaussian elimination 1

## Example 8.14

Forsythe, Moler (1967)

- ▶ Impact of pivoting on finite precision arithmetic (3 digit example)
- ▶  $A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$
- ▶ Pivoting:  $A' = P_{12}A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 10^{-4} & 1 \end{pmatrix}, L'U' = A'$

# Pivoting and Gaussian elimination 1

## Example 8.14

Forsythe, Moler (1967)

- ▶ Impact of pivoting on finite precision arithmetic (3 digit example)
- ▶  $A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$
- ▶ Pivoting:  $A' = P_{12}A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 10^{-4} & 1 \end{pmatrix}, L'U' = A'$
- ▶  $L' = \begin{pmatrix} 1 & 0 \\ 10^{-4} & 1 \end{pmatrix}, U' = A^{(1)} = \begin{pmatrix} 1 & 1 \\ 0 & 1 - 10^{-4} \end{pmatrix},$

# Pivoting and Gaussian elimination 1

## Example 8.14

Forsythe, Moler (1967)

- ▶ Impact of pivoting on finite precision arithmetic (3 digit example)
- ▶  $A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$
- ▶ Pivoting:  $A' = P_{12}A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 10^{-4} & 1 \end{pmatrix}, L'U' = A'$
- ▶  $L' = \begin{pmatrix} 1 & 0 \\ 10^{-4} & 1 \end{pmatrix}, U' = A^{(1)} = \begin{pmatrix} 1 & 1 \\ 0 & 1 - 10^{-4} \end{pmatrix}, fl(1 - 10^{-4}) = 1$

# Pivoting and Gaussian elimination 1

## Example 8.14

Forsythe, Moler (1967)

- ▶ Impact of pivoting on finite precision arithmetic (3 digit example)
- ▶  $A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$
- ▶ Pivoting:  $A' = P_{12}A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 10^{-4} & 1 \end{pmatrix}, L'U' = A'$
- ▶  $L' = \begin{pmatrix} 1 & 0 \\ 10^{-4} & 1 \end{pmatrix}, U' = A^{(1)} = \begin{pmatrix} 1 & 1 \\ 0 & 1 - 10^{-4} \end{pmatrix}, fl(1 - 10^{-4}) = 1$
- ▶  $fl(L') = L',$

# Pivoting and Gaussian elimination 1

## Example 8.14

Forsythe, Moler (1967)

- ▶ Impact of pivoting on finite precision arithmetic (3 digit example)
- ▶  $A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$
- ▶ Pivoting:  $A' = P_{12}A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 10^{-4} & 1 \end{pmatrix}, L'U' = A'$
- ▶  $L' = \begin{pmatrix} 1 & 0 \\ 10^{-4} & 1 \end{pmatrix}, U' = A^{(1)} = \begin{pmatrix} 1 & 1 \\ 0 & 1 - 10^{-4} \end{pmatrix}, fl(1 - 10^{-4}) = 1$
- ▶  $fl(L') = L', fl(U') = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix},$

# Pivoting and Gaussian elimination 1

## Example 8.14

Forsythe, Moler (1967)

- ▶ Impact of pivoting on finite precision arithmetic (3 digit example)
- ▶  $A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$
- ▶ Pivoting:  $A' = P_{12}A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 10^{-4} & 1 \end{pmatrix}, L'U' = A'$
- ▶  $L' = \begin{pmatrix} 1 & 0 \\ 10^{-4} & 1 \end{pmatrix}, U' = A^{(1)} = \begin{pmatrix} 1 & 1 \\ 0 & 1 - 10^{-4} \end{pmatrix}, fl(1 - 10^{-4}) = 1$
- ▶  $fl(L') = L', fl(U') = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, fl(L')fl(U') = \begin{pmatrix} 1 & 1 \\ 10^{-4} & 1.0001 \end{pmatrix}$



# Pivoting and Gaussian elimination 1

## Example 8.14

Forsythe, Moler (1967)

- ▶ Impact of pivoting on finite precision arithmetic (3 digit example)

- ▶  $A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$

- ▶ Pivoting:  $A' = P_{12}A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 10^{-4} & 1 \end{pmatrix}, L'U' = A'$

- ▶  $L' = \begin{pmatrix} 1 & 0 \\ 10^{-4} & 1 \end{pmatrix}, U' = A^{(1)} = \begin{pmatrix} 1 & 1 \\ 0 & 1 - 10^{-4} \end{pmatrix}, fl(1 - 10^{-4}) = 1$

- ▶  $fl(L') = L', fl(U') = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, fl(L')fl(U') = \begin{pmatrix} 1 & 1 \\ 10^{-4} & 1.0001 \end{pmatrix}$

- ▶  $fl(fl(L')fl(U')) = \begin{pmatrix} 1 & 1 \\ 10^{-4} & 1 \end{pmatrix} = A'$

# Pivoting and Gaussian elimination 1

## Example 8.14

Forsythe, Moler (1967)

- ▶ Impact of pivoting on finite precision arithmetic (3 digit example)
- ▶  $A = \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix}$
- ▶ Pivoting:  $A' = P_{12}A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 10^{-4} & 1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 10^{-4} & 1 \end{pmatrix}, L'U' = A'$
- ▶  $L' = \begin{pmatrix} 1 & 0 \\ 10^{-4} & 1 \end{pmatrix}, U' = A^{(1)} = \begin{pmatrix} 1 & 1 \\ 0 & 1 - 10^{-4} \end{pmatrix}, fl(1 - 10^{-4}) = 1$
- ▶  $fl(L') = L', fl(U') = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, fl(L')fl(U') = \begin{pmatrix} 1 & 1 \\ 10^{-4} & 1.0001 \end{pmatrix}$
- ▶  $fl(fl(L')fl(U')) = \begin{pmatrix} 1 & 1 \\ 10^{-4} & 1 \end{pmatrix} = A'$

# Pivoting and Gaussian elimination 2

## Definition 8.15

$P$  is permutation matrix if in each row and column exactly one entry is 1 and other entries are 0.

# Pivoting and Gaussian elimination 2

## Definition 8.15

$P$  is permutation matrix if in each row and column exactly one entry is 1 and other entries are 0.

## Example 8.16

- ▶ Unit vector  $e_1, e_2, \dots, e_n$

# Pivoting and Gaussian elimination 2

## Definition 8.15

$P$  is permutation matrix if in each row and column exactly one entry is 1 and other entries are 0.

## Example 8.16

► Unit vector  $e_1, e_2, \dots, e_n$

►

$$P_{12} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} e_2^T \\ e_1^T \end{pmatrix}, P_{12}P_{12}^T = I$$

# Pivoting and Gaussian elimination 2

## Definition 8.15

$P$  is permutation matrix if in each row and column exactly one entry is 1 and other entries are 0.

## Example 8.16

► Unit vector  $e_1, e_2, \dots, e_n$

►

$$P_{12} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} e_2^T \\ e_1^T \end{pmatrix}, P_{12}P_{12}^T = I$$

► General permutation matrix

$$P = \begin{pmatrix} e_{\alpha_1}^T \\ e_{\alpha_2}^T \\ \vdots \\ e_{\alpha_n}^T \end{pmatrix}, 1 \leq i, \alpha_i \leq n, PP^T = I, PA = \begin{pmatrix} \alpha_1 - \text{th row of } A \\ \alpha_2 - \text{th row of } A \\ \vdots \\ \alpha_n - \text{th row of } A \end{pmatrix}$$

## Pivoting and Gaussian elimination 3

- Premultiplication:  $P_{left}A$

## Pivoting and Gaussian elimination 3

- ▶ Premultiplication:  $P_{left}A$
- ▶ Postmultiplication:  $AP_{right}$



## Pivoting and Gaussian elimination 3

- ▶ Premultiplication:  $P_{left}A$
- ▶ Postmultiplication:  $AP_{right}$
- ▶ General permutation matrix  $P_{left}$

$$P_{left} = \begin{pmatrix} e_{\alpha_1}^T \\ e_{\alpha_2}^T \\ \cdot \\ \cdot \\ e_{\alpha_n}^T \end{pmatrix}$$

## Pivoting and Gaussian elimination 3

- ▶ Premultiplication:  $P_{left}A$
- ▶ Postmultiplication:  $AP_{right}$
- ▶ General permutation matrix  $P_{left}$

$$P_{left} = \begin{pmatrix} e_{\alpha_1}^T \\ e_{\alpha_2}^T \\ \vdots \\ e_{\alpha_n}^T \end{pmatrix}$$

$$P_{left}A = \begin{pmatrix} \alpha_1 - \text{th row of matrix } A \\ \alpha_2 - \text{th row of matrix } A \\ \vdots \\ \alpha_n - \text{th row of matrix } A \end{pmatrix}$$

## Pivoting and Gaussian elimination 3

- ▶ Premultiplication:  $P_{left}A$
- ▶ Postmultiplication:  $AP_{right}$
- ▶ General permutation matrix  $P_{left}$

$$P_{left} = \begin{pmatrix} e_{\alpha_1}^T \\ e_{\alpha_2}^T \\ \vdots \\ e_{\alpha_n}^T \end{pmatrix}$$

$$P_{left}A = \begin{pmatrix} \alpha_1 - \text{th row of matrix } A \\ \alpha_2 - \text{th row of matrix } A \\ \vdots \\ \alpha_n - \text{th row of matrix } A \end{pmatrix}$$

- ▶ General permutation matrix  $P_{right} = (e_{\alpha_1}, e_{\alpha_2}, \dots, e_{\alpha_n})$

## Pivoting and Gaussian elimination 3

- ▶ Premultiplication:  $P_{left}A$
- ▶ Postmultiplication:  $AP_{right}$
- ▶ General permutation matrix  $P_{left}$

$$P_{left} = \begin{pmatrix} e_{\alpha_1}^T \\ e_{\alpha_2}^T \\ \vdots \\ e_{\alpha_n}^T \end{pmatrix}$$

$$P_{left}A = \begin{pmatrix} \alpha_1 - \text{th row of matrix } A \\ \alpha_2 - \text{th row of matrix } A \\ \vdots \\ \alpha_n - \text{th row of matrix } A \end{pmatrix}$$

- ▶ General permutation matrix  $P_{right} = (e_{\alpha_1}, e_{\alpha_2}, \dots, e_{\alpha_n})$

$$AP_{right} = (\alpha_1 - \text{th column}, \alpha_2 - \text{th column}, \dots, \alpha_n - \text{th column of } A)$$

# Pivoting and Gaussian elimination 4

- ▶ Gaussian Elimination with Partial Pivoting (GEPP)

## Pivoting and Gaussian elimination 4

- ▶ Gaussian Elimination with Partial Pivoting (GEPP)

$$A^{(1)} = M_1 P_1 A$$

$$A^{(2)} = M_2 P_2 A^{(1)} = M_2 P_2 M_1 P_1 A$$

..

$$A^{(n-1)} = M_{n-1} P_{n-1} \cdots M_2 P_2 M_1 P_1 A$$

## Pivoting and Gaussian elimination 4

- ▶ Gaussian Elimination with Partial Pivoting (GEPP)

$$A^{(1)} = M_1 P_1 A$$

$$A^{(2)} = M_2 P_2 A^{(1)} = M_2 P_2 M_1 P_1 A$$

..

$$A^{(n-1)} = M_{n-1} P_{n-1} \cdots M_2 P_2 M_1 P_1 A$$

- ▶ GEPP and  $LU$  decomposition

## Pivoting and Gaussian elimination 4

### ► Gaussian Elimination with Partial Pivoting (GEPP)

$$A^{(1)} = M_1 P_1 A$$

$$A^{(2)} = M_2 P_2 A^{(1)} = M_2 P_2 M_1 P_1 A$$

..

$$A^{(n-1)} = M_{n-1} P_{n-1} \cdots M_2 P_2 M_1 P_1 A$$

### ► GEPP and $LU$ decomposition

$$\begin{aligned} A^{(n-1)} &= M_{n-1} P_{n-1} \cdots M_2 P_2 M_1 P_1 A = \\ &= M_{n-1} P_{n-1} \cdots M_2 P_2 M_1 (P_2^T \cdots P_{n-1}^T P_{n-1} \cdots P_2) P_1 A = \\ &= M_{n-1} P_{n-1} \cdots M_2 P_2 M_1 (P_2^T \cdots P_{n-1}^T) P_{n-1} \cdots P_2 P_1 A = \\ &= M_{n-1} P_{n-1} \cdots M_2 (P_3^T \cdots P_{n-1}^T P_{n-1} \cdots P_3) P_2 M_1 (P_2^T \cdots P_{n-1}^T) P A = \\ &= M_{n-1} P_{n-1} \cdots M_2 (P_3^T \cdots P_{n-1}^T (P_{n-1} \cdots P_3 P_2 M_1 P_2^T \cdots P_{n-1}^T) P A = \\ &= M_{n-1} P_{n-1} \cdots M_2 (P_3^T \cdots P_{n-1}^T) M'_1 P A = \\ &= M'_{n-1} \cdots M'_2 M'_1 P A \end{aligned}$$



# Pivoting and Gaussian elimination 5

## ► GEPP and $LU$ decomposition

$$A^{(n-1)} = M'_{n-1} \cdots M'_2 M'_1 P A$$

$$P = P_{n-1} \cdots P_2 P_1$$

$$M'_1 = P_{n-1} \cdots P_3 P_2 M_1 P_2^T \cdots P_{n-1}^T$$

$$M'_2 = P_{n-1} \cdots P_3 M_1 P_3^T \cdots P_{n-1}^T$$

$$M'_3 = P_{n-1} \cdots P_4 M_1 P_4^T \cdots P_{n-1}^T$$

..

$$M'_{n-1} = M_{n-1}$$

# Pivoting and Gaussian elimination 5

## ► GEPP and $LU$ decomposition

$$A^{(n-1)} = M'_{n-1} \cdots M'_2 M'_1 P A$$

$$P = P_{n-1} \cdots P_2 P_1$$

$$M'_1 = P_{n-1} \cdots P_3 P_2 M_1 P_2^T \cdots P_{n-1}^T$$

$$M'_2 = P_{n-1} \cdots P_3 M_1 P_3^T \cdots P_{n-1}^T$$

$$M'_3 = P_{n-1} \cdots P_4 M_1 P_4^T \cdots P_{n-1}^T$$

..

$$M'_{n-1} = M_{n-1}$$



$$A^{(n-1)} = U$$

$$P A = L U$$

$$L = (M'_{n-1} \cdots M'_2 M'_1)^{-1}$$

# Pivoting and Gaussian elimination 5

- ▶ Gaussian elimination with complete pivoting (GECP)

# Pivoting and Gaussian elimination 5

- ▶ Gaussian elimination with complete pivoting (GECP)



$$A^{(k)} = M_k P_k A^{(k-1)} Q_k$$

# Pivoting and Gaussian elimination 5

- ▶ Gaussian elimination with complete pivoting (GECP)



$$A^{(k)} = M_k P_k A^{(k-1)} Q_k$$

- ▶  $P_k$  - for row exchange

# Pivoting and Gaussian elimination 5

- ▶ Gaussian elimination with complete pivoting (GECP)



$$A^{(k)} = M_k P_k A^{(k-1)} Q_k$$

- ▶  $P_k$  - for row exchange
- ▶  $Q_k$  - for column exchange

# Pivoting and Gaussian elimination 5

- ▶ Gaussian elimination with complete pivoting (GECP)



$$A^{(k)} = M_k P_k A^{(k-1)} Q_k$$

- ▶  $P_k$  - for row exchange
- ▶  $Q_k$  - for column exchange
- ▶  $M_k$  - for elimination step in column  $k$

# Pivoting and Gaussian elimination 5

- ▶ Gaussian elimination with complete pivoting (GECP)



$$A^{(k)} = M_k P_k A^{(k-1)} Q_k$$

- ▶  $P_k$  - for row exchange
- ▶  $Q_k$  - for column exchange
- ▶  $M_k$  - for elimination step in column  $k$



$$A^{(n-1)} = U$$

$$PAQ = LU$$

$$P = P_{n-1} \dots P_2 P_1$$

$$Q = Q_1 Q_2 \dots Q_{n-1}$$



# Stability of Gaussian elimination 1

## Definition 8.17

► The **growth factor**  $\rho$

►

$$\rho = \frac{\max\{\alpha_0, \alpha_1, \dots, \alpha_{n-1}\}}{\alpha_0}$$

$$\alpha_k = \max_{i,j} |a_{ij}^{(k)}|, k = 0, 1, \dots, n-1$$

►  $A^{(0)} = A$ ,  $A^{(k)}$  - matrix obtained after  $k$  steps of Gaussian elimination

# Stability of Gaussian elimination 1

## Definition 8.17

- ▶ The **growth factor**  $\rho$



$$\rho = \frac{\max\{\alpha_0, \alpha_1, \dots, \alpha_{n-1}\}}{\alpha_0}$$

$$\alpha_k = \max_{i,j} |a_{ij}^{(k)}|, k = 0, 1, \dots, n-1$$

- ▶  $A^{(0)} = A$ ,  $A^{(k)}$  - matrix obtained after  $k$  steps of Gaussian elimination
- ▶ Growth factor of GEPP  $\rho \leq 2^{n-1}$

# Stability of Gaussian elimination 1

## Definition 8.17

- ▶ The **growth factor**  $\rho$



$$\rho = \frac{\max\{\alpha_0, \alpha_1, \dots, \alpha_{n-1}\}}{\alpha_0}$$

$$\alpha_k = \max_{i,j} |a_{ij}^{(k)}|, k = 0, 1, \dots, n-1$$

- ▶  $A^{(0)} = A$ ,  $A^{(k)}$  - matrix obtained after  $k$  steps of Gaussian elimination
- ▶ Growth factor of GEPP  $\rho \leq 2^{n-1}$
- ▶ Growth factor of GECP  $\rho \leq (n \cdot 2^1 \cdot 3^{\frac{1}{2}} \cdot 4^{\frac{1}{3}} \dots n^{\frac{1}{n-1}})^{\frac{1}{2}}$

# Stability of Gaussian elimination 1

## Definition 8.17

- ▶ The **growth factor**  $\rho$



$$\rho = \frac{\max\{\alpha_0, \alpha_1, \dots, \alpha_{n-1}\}}{\alpha_0}$$

$$\alpha_k = \max_{i,j} |a_{ij}^{(k)}|, k = 0, 1, \dots, n-1$$

- ▶  $A^{(0)} = A$ ,  $A^{(k)}$  - matrix obtained after  $k$  steps of Gaussian elimination
- ▶ Growth factor of GEPP  $\rho \leq 2^{n-1}$
- ▶ Growth factor of GECP  $\rho \leq (n \cdot 2^1 \cdot 3^{\frac{1}{2}} \cdot 4^{\frac{1}{3}} \dots n^{\frac{1}{n-1}})^{\frac{1}{2}}$
- ▶ The textbook, chapter 5, pp.108,109

## Stability of Gaussian elimination 2

- ▶ A priori estimates: theorems 5.16 and 5.18 in the textbook

## Stability of Gaussian elimination 2

- ▶ A priori estimates: theorems 5.16 and 5.18 in the textbook

### Theorem 8.18

- ▶  $LU = A + E$ ,  $E = \text{error}$

## Stability of Gaussian elimination 2

- ▶ A priori estimates: theorems 5.16 and 5.18 in the textbook

### Theorem 8.18

- ▶  $LU = A + E$ ,  $E = \text{error}$
- ▶ If  $L$  and  $U$  are computed by Gaussian elimination without pivoting then

$$\|E\| \leq n\mu \|L\| \cdot \|U\|$$

# Stability of Gaussian elimination 2

- ▶ A priori estimates: theorems 5.16 and 5.18 in the textbook

## Theorem 8.18

- ▶  $LU = A + E$ ,  $E = \text{error}$
- ▶ If  $L$  and  $U$  are computed by Gaussian elimination without pivoting then

$$\|E\| \leq n\mu \|L\| \cdot \|U\|$$

- ▶ If  $L$  and  $U$  are computed with GEPP then

$$\|E\|_{\infty} \leq n^3 \mu \rho \|A\|_{\infty}$$



# Stability of Gaussian elimination 2

- ▶ A prior estimates: theorems 5.16 and 5.18 in the textbook

## Theorem 8.18

- ▶  $LU = A + E$ ,  $E = \text{error}$
- ▶ If  $L$  and  $U$  are computed by Gaussian elimination without pivoting then

$$\|E\| \leq n\mu \|L\| \cdot \|U\|$$

- ▶ If  $L$  and  $U$  are computed with GEPP then

$$\|E\|_{\infty} \leq n^3 \mu \rho \|A\|_{\infty}$$

where  $\rho$  is growth factor,  $|L|, |U|$  are matrices where entries are replaced by their absolute values

- ▶ Q: Which estimate is better?

# Stability of Gaussian elimination 2

- ▶ A prior estimates: theorems 5.16 and 5.18 in the textbook

## Theorem 8.18

- ▶  $LU = A + E$ ,  $E = \text{error}$
- ▶ If  $L$  and  $U$  are computed by Gaussian elimination without pivoting then

$$\|E\| \leq n\mu \|L\| \cdot \|U\|$$

- ▶ If  $L$  and  $U$  are computed with GEPP then

$$\|E\|_{\infty} \leq n^3 \mu \rho \|A\|_{\infty}$$

where  $\rho$  is growth factor,  $|L|, |U|$  are matrices where entries are replaced by their absolute values

- ▶ Q: Which estimate is better?
- ▶ A posterior estimate:  $\|LU - A\|$  - extra FLOPS needed

# Residual theorem

- ▶ A posterior estimate: theorems 6.12 in the textbook

## Theorem 8.19

- ▶  $Ax = b, \tilde{r} = A\tilde{x} - b$



$$\frac{\|\tilde{x} - x\|}{\|x\|} \leq \text{cond}(A) \frac{\|\tilde{r}\|}{\|b\|}$$

# Residual theorem

- ▶ A posterior estimate: theorems 6.12 in the textbook

## Theorem 8.19

- ▶  $Ax = b, \tilde{r} = A\tilde{x} - b$



$$\frac{\|\tilde{x} - x\|}{\|x\|} \leq \text{cond}(A) \frac{\|\tilde{r}\|}{\|b\|}$$

- ▶ Symptoms of ill conditioning

# Residual theorem

- ▶ A posterior estimate: theorems 6.12 in the textbook

## Theorem 8.19

- ▶  $Ax = b, \tilde{r} = A\tilde{x} - b$



$$\frac{\|\tilde{x} - x\|}{\|x\|} \leq \text{cond}(A) \frac{\|\tilde{r}\|}{\|b\|}$$

- ▶ Symptoms of ill conditioning
  - ▶ A large computed solution (e.g. large solution and "not large" RHS  $b$ )

# Residual theorem

- ▶ A posterior estimate: theorems 6.12 in the textbook

## Theorem 8.19

- ▶  $Ax = b, \tilde{r} = A\tilde{x} - b$



$$\frac{\|\tilde{x} - x\|}{\|x\|} \leq \text{cond}(A) \frac{\|\tilde{r}\|}{\|b\|}$$

- ▶ Symptoms of ill conditioning
  - ▶ A large computed solution (e.g. large solution and "not large" RHS  $b$ )
  - ▶ A small pivot

# Residual theorem

- ▶ A posterior estimate: theorems 6.12 in the textbook

## Theorem 8.19

- ▶  $Ax = b, \tilde{r} = A\tilde{x} - b$



$$\frac{\|\tilde{x} - x\|}{\|x\|} \leq \text{cond}(A) \frac{\|\tilde{r}\|}{\|b\|}$$

- ▶ Symptoms of ill conditioning
  - ▶ A large computed solution (e.g. large solution and "not large" RHS  $b$ )
  - ▶ A small pivot
  - ▶ A big residual vector

# Scaling

## Definition 8.20

►  $Ax = b$



# Scaling

## Definition 8.20

- ▶  $Ax = b$
- ▶  $(D, F)$  - ordered pair of non-singular diagonal matrices

# Scaling

## Definition 8.20

- ▶  $Ax = b$
- ▶  $(D, F)$  - ordered pair of non-singular diagonal matrices
- ▶  $A_s = DAF$  - scaled matrix

# Scaling

## Definition 8.20

- ▶  $Ax = b$
- ▶  $(D, F)$  - ordered pair of non-singular diagonal matrices
- ▶  $A_s = DAF$  - scaled matrix
- ▶ scaled system  $DAFy = Db, y = F^{-1}x$

# Scaling

## Definition 8.20

- ▶  $Ax = b$
- ▶  $(D, F)$  - ordered pair of non-singular diagonal matrices
- ▶  $A_s = DAF$  - scaled matrix
- ▶ scaled system  $DAFy = Db$ ,  $y = F^{-1}x$
- ▶  $(D, I)$  defines row scaling

# Scaling

## Definition 8.20

- ▶  $Ax = b$
- ▶  $(D, F)$  - ordered pair of non-singular diagonal matrices
- ▶  $A_s = DAF$  - scaled matrix
- ▶ scaled system  $DAFy = Db, y = F^{-1}x$
- ▶  $(D, I)$  defines row scaling
- ▶  $(I, F)$  defines column scaling

# Scaling

## Definition 8.20

- ▶  $Ax = b$
- ▶  $(D, F)$  - ordered pair of non-singular diagonal matrices
- ▶  $A_s = DAF$  - scaled matrix
- ▶ scaled system  $DAFy = Db, y = F^{-1}x$
- ▶  $(D, I)$  defines row scaling
- ▶  $(I, F)$  defines column scaling
- ▶ Why to apply scaling

# Scaling

## Definition 8.20

- ▶  $Ax = b$
- ▶  $(D, F)$  - ordered pair of non-singular diagonal matrices
- ▶  $A_s = DAF$  - scaled matrix
- ▶ scaled system  $DAFy = Db, y = F^{-1}x$
- ▶  $(D, I)$  defines row scaling
- ▶  $(I, F)$  defines column scaling
- ▶ Why to apply scaling
  - ▶ Make  $\text{cond}(A_s)$  smaller (Golub, VanLoan, 1989)

# Scaling

## Definition 8.20

- ▶  $Ax = b$
- ▶  $(D, F)$  - ordered pair of non-singular diagonal matrices
- ▶  $A_s = DAF$  - scaled matrix
- ▶ scaled system  $DAFy = Db, y = F^{-1}x$
- ▶  $(D, I)$  defines row scaling
- ▶  $(I, F)$  defines column scaling
- ▶ Why to apply scaling
  - ▶ Make  $\text{cond}(A_s)$  smaller (Golub, VanLoan, 1989)
  - ▶ Get more balanced matrix = eliminate disparity in the size of entries (Stewart, 1973)



# Scaling

## Definition 8.20

- ▶  $Ax = b$
- ▶  $(D, F)$  - ordered pair of non-singular diagonal matrices
- ▶  $A_s = DAF$  - scaled matrix
- ▶ scaled system  $DAFy = Db, y = F^{-1}x$
- ▶  $(D, I)$  defines row scaling
- ▶  $(I, F)$  defines column scaling
- ▶ Why to apply scaling
  - ▶ Make  $\text{cond}(A_s)$  smaller (Golub, VanLoan, 1989)
  - ▶ Get more balanced matrix = eliminate disparity in the size of entries (Stewart, 1973)
  - ▶ Control selection of pivots (Atkinson, 1989)

# Scaling

## Definition 8.20

- ▶  $Ax = b$
- ▶  $(D, F)$  - ordered pair of non-singular diagonal matrices
- ▶  $A_s = DAF$  - scaled matrix
- ▶ scaled system  $DAFy = Db$ ,  $y = F^{-1}x$
- ▶  $(D, I)$  defines row scaling
- ▶  $(I, F)$  defines column scaling
- ▶ Why to apply scaling
  - ▶ Make  $\text{cond}(A_s)$  smaller (Golub, VanLoan, 1989)
  - ▶ Get more balanced matrix = eliminate disparity in the size of entries (Stewart, 1973)
  - ▶ Control selection of pivots (Atkinson, 1989)
  - ▶ Scaling example: textbook, chapter 6, p.135

# Iterative refinement

- ▶ Q: What if computed solution  $\tilde{x}$  is not acceptable?

# Iterative refinement

- ▶ Q: What if computed solution  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement

# Iterative refinement

- ▶ Q: What if computed solution  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$

# Iterative refinement

- ▶ Q: What if computed solution  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$

# Iterative refinement

- ▶ Q: What if computed solution  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$

# Iterative refinement

- ▶ Q: What if computed solution  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$



# Iterative refinement

- ▶ Q: What if computed solution  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:

# Iterative refinement

- ▶ Q: What if computed solution  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:
  - ▶  $x^{(0)} = \tilde{x}, tol = 1, \epsilon = 10^{-l}$

# Iterative refinement

- ▶ Q: What if computed solution  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:
  - ▶  $x^{(0)} = \tilde{x}, tol = 1, \epsilon = 10^{-l}$
  - ▶ Do until  $tol \leq \epsilon$ :

# Iterative refinement

- ▶ Q: What if computed solution  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:
  - ▶  $x^{(0)} = \tilde{x}, tol = 1, \epsilon = 10^{-l}$
  - ▶ Do until  $tol \leq \epsilon$ :
    1.  $r^{(k)} = b - Ax^{(k)}$

# Iterative refinement

- ▶ Q: What if computed solution  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:
  - ▶  $x^{(0)} = \tilde{x}, tol = 1, \epsilon = 10^{-l}$
  - ▶ Do until  $tol \leq \epsilon$ :
    1.  $r^{(k)} = b - Ax^{(k)}$
    2.  $Ac^{(k)} = r^{(k)}$

# Iterative refinement

- ▶ Q: What if computed solution  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:
  - ▶  $x^{(0)} = \tilde{x}, tol = 1, \epsilon = 10^{-l}$
  - ▶ Do until  $tol \leq \epsilon$ :
    1.  $r^{(k)} = b - Ax^{(k)}$
    2.  $Ac^{(k)} = r^{(k)}$
    3.  $x^{(k+1)} = x^{(k)} + c^{(k)}$

# Iterative refinement

- ▶ Q: What if computed solution  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:

- ▶  $x^{(0)} = \tilde{x}, tol = 1, \epsilon = 10^{-l}$
- ▶ Do until  $tol \leq \epsilon$ :
  1.  $r^{(k)} = b - Ax^{(k)}$
  2.  $Ac^{(k)} = r^{(k)}$
  3.  $x^{(k+1)} = x^{(k)} + c^{(k)}$
  4.  $tol = \frac{\|x^{(k+1)} - x^{(k)}\|}{\|x^{(k)}\|}$

# Iterative refinement

- ▶ Q: What if computed solution  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:
  - ▶  $x^{(0)} = \tilde{x}, tol = 1, \epsilon = 10^{-l}$
  - ▶ Do until  $tol \leq \epsilon$ :
    1.  $r^{(k)} = b - Ax^{(k)}$
    2.  $Ac^{(k)} = r^{(k)}$
    3.  $x^{(k+1)} = x^{(k)} + c^{(k)}$
    4.  $tol = \frac{\|x^{(k+1)} - x^{(k)}\|}{\|x^{(k)}\|}$
- ▶ Produces very accurate solution if  $cond(A)$  not too large



# Iterative refinement

- ▶ Q: What if computed solution  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:
  - ▶  $x^{(0)} = \tilde{x}, tol = 1, \epsilon = 10^{-l}$
  - ▶ Do until  $tol \leq \epsilon$ :
    1.  $r^{(k)} = b - Ax^{(k)}$
    2.  $Ac^{(k)} = r^{(k)}$
    3.  $x^{(k+1)} = x^{(k)} + c^{(k)}$
    4.  $tol = \frac{\|x^{(k+1)} - x^{(k)}\|}{\|x^{(k)}\|}$
- ▶ Produces very accurate solution if  $cond(A)$  not too large
- ▶ Convergence depends on  $cond(A)$

Q & A