

Numerical Linear Algebra

Ramaz Botchorishvili

Kutaisi International University

November 30, 2022

Numerical Linear Algebra

Ramaz Botchorishvili

Kutaisi International University

November 30, 2022

Direct and iterative methods for linear systems

- ▶ Recap of Previous Lecture
- ▶ Convergence of Richardson's iterations
- ▶ Optimal parameter parameter of Richardson's iterations
- ▶ Sufficient condition $P > 0.5A$
- ▶ Preconditioning matrices
- ▶ Quadratic functional and linear systems
- ▶ Q & A

Recap of Previous Lecture

- ▶ Sherman-Morrison formula
- ▶ Cholesky factorization of symmetric system
- ▶ Diagonally dominant system
- ▶ Necessary and sufficient conditions for convergence
- ▶ Classic iterative methods
- ▶ Convergence theorems of J,GS

Richardson method 1

► Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

Richardson method 1

- ▶ Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

- ▶ Consistent with $Ax = b$

Richardson method 1

- ▶ Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

- ▶ Consistent with $Ax = b$

- ▶ $\frac{1}{\tau_k} Px^{(k+1)} = (\frac{1}{\tau_k} P - A)x^{(k)} + b, \quad k = 0, 1, 2, \dots$

Richardson method 1

- ▶ Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

- ▶ Consistent with $Ax = b$

- ▶ $\frac{1}{\tau_k} Px^{(k+1)} = (\frac{1}{\tau_k} P - A)x^{(k)} + b, \quad k = 0, 1, 2, \dots$

- ▶ $x^{(k+1)} = B_{R,k}x^{(k)} + f, \quad B_{R,k} = I - \tau_k P^{-1}A$

Richardson method 1

- ▶ Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

- ▶ Consistent with $Ax = b$

- ▶ $\frac{1}{\tau_k} Px^{(k+1)} = (\frac{1}{\tau_k} P - A)x^{(k)} + b, \quad k = 0, 1, 2, \dots$

- ▶ $x^{(k+1)} = B_{R,k}x^{(k)} + f, \quad B_{R,k} = I - \tau_k P^{-1}A$

- ▶ Pre-conditioner - $P \in \mathbb{R}^{n \times n}$

Richardson method 1

- ▶ Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

- ▶ Consistent with $Ax = b$

- ▶ $\frac{1}{\tau_k} Px^{(k+1)} = (\frac{1}{\tau_k} P - A)x^{(k)} + b, \quad k = 0, 1, 2, \dots$

- ▶ $x^{(k+1)} = B_{R,k}x^{(k)} + f, \quad B_{R,k} = I - \tau_k P^{-1}A$

- ▶ Pre-conditioner - $P \in \mathbb{R}^{n \times n}$

- ▶ Numeric parameter - τ_k

Richardson method 1

- ▶ Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

- ▶ Consistent with $Ax = b$
- ▶ $\frac{1}{\tau_k} Px^{(k+1)} = (\frac{1}{\tau_k} P - A)x^{(k)} + b, \quad k = 0, 1, 2, \dots$
- ▶ $x^{(k+1)} = B_{R,k}x^{(k)} + f, \quad B_{R,k} = I - \tau_k P^{-1}A$
- ▶ Pre-conditioner - $P \in \mathbb{R}^{n \times n}$
- ▶ Numeric parameter - τ_k
- ▶ Stationary iteration method if $B_{R,k} \equiv B_R$

Richardson method 1

- ▶ Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

- ▶ Consistent with $Ax = b$
- ▶ $\frac{1}{\tau_k} Px^{(k+1)} = (\frac{1}{\tau_k} P - A)x^{(k)} + b, \quad k = 0, 1, 2, \dots$
- ▶ $x^{(k+1)} = B_{R,k}x^{(k)} + f, \quad B_{R,k} = I - \tau_k P^{-1}A$
- ▶ Pre-conditioner - $P \in \mathbb{R}^{n \times n}$
- ▶ Numeric parameter - τ_k
- ▶ Stationary iteration method if $B_{R,k} \equiv B_R$
- ▶ Converges if $\|B_{R,k}\| < 1$

Richardson method 1

- ▶ Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

- ▶ Consistent with $Ax = b$
- ▶ $\frac{1}{\tau_k} Px^{(k+1)} = (\frac{1}{\tau_k} P - A)x^{(k)} + b, \quad k = 0, 1, 2, \dots$
- ▶ $x^{(k+1)} = B_{R,k}x^{(k)} + f, \quad B_{R,k} = I - \tau_k P^{-1}A$
- ▶ Pre-conditioner - $P \in \mathbb{R}^{n \times n}$
- ▶ Numeric parameter - τ_k
- ▶ Stationary iteration method if $B_{R,k} \equiv B_R$
- ▶ Converges if $\|B_{R,k}\| < 1$
- ▶ Converges iff $\rho(B_{R,k}) < 1$

Richardson method 2

Example 10.1

Jacobi method = preconditioner is diagonal part of A

Richardson method 2

Example 10.1

Jacobi method = preconditioner is diagonal part of A

$$\begin{cases} P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b \\ \tau_k = 1, P = D \end{cases} \Rightarrow \begin{cases} D(x^{(k+1)} - x^{(k)}) + Ax^{(k)} = b \\ x^{(k+1)} = (I - D^{-1}A)x^{(k)} + D^{-1}b \\ B_R = (I - D^{-1}A) = B_J \end{cases}$$

Richardson method 2

Example 10.1

Jacobi method = preconditioner is diagonal part of A

$$\begin{cases} P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b \\ \tau_k = 1, P = D \end{cases} \Rightarrow \begin{cases} D(x^{(k+1)} - x^{(k)}) + Ax^{(k)} = b \\ x^{(k+1)} = (I - D^{-1}A)x^{(k)} + D^{-1}b \\ B_R = (I - D^{-1}A) = B_J \end{cases}$$

Example 10.2

Gaus-Seidel method = preconditioner is lower triangular part of A

Richardson method 2

Example 10.1

Jacobi method = preconditioner is diagonal part of A

$$\begin{cases} P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b \\ \tau_k = 1, P = D \end{cases} \Rightarrow \begin{cases} D(x^{(k+1)} - x^{(k)}) + Ax^{(k)} = b \\ x^{(k+1)} = (I - D^{-1}A)x^{(k)} + D^{-1}b \\ B_R = (I - D^{-1}A) = B_J \end{cases}$$

Example 10.2

Gaus-Seidel method = preconditioner is lower triangular part of A

$$\begin{cases} P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b \\ \tau_k = 1, P = D + L \end{cases} \Rightarrow \begin{cases} A = L + D + U \\ (D + L)(x^{(k+1)} - x^{(k)}) + Ax^{(k)} = b \\ x^{(k+1)} = -(D + L)^{-1}Ux^{(k)} + (D + L)^{-1}b \end{cases}$$

Richardson method 3

Theorem 10.3

The stationary Richardson method converges iff

$$\frac{2\Re(\lambda)}{\tau|\lambda|^2} > 1$$

for all λ - eigenvalues of $P^{-1}A$

Richardson method 3

Theorem 10.3

The stationary Richardson method converges iff

$$\frac{2\Re(\lambda)}{\tau|\lambda|^2} > 1$$

for all λ - eigenvalues of $P^{-1}A$

Proof.

$$\blacktriangleright P \frac{x^{(k+1)} - x^{(k)}}{\tau} + Ax^{(k)} = b$$

Richardson method 3

Theorem 10.3

The stationary Richardson method converges iff

$$\frac{2\Re(\lambda)}{\tau|\lambda|^2} > 1$$

for all λ - eigenvalues of $P^{-1}A$

Proof.

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\tau} + Ax^{(k)} = b$
- ▶ $x^{(k+1)} = B_R x^{(k)} + f, B_R = I - \tau P^{-1}A$

Richardson method 3

Theorem 10.3

The stationary Richardson method converges iff

$$\frac{2\Re(\lambda)}{\tau|\lambda|^2} > 1$$

for all λ - eigenvalues of $P^{-1}A$

Proof.

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\tau} + Ax^{(k)} = b$
- ▶ $x^{(k+1)} = B_R x^{(k)} + f, B_R = I - \tau P^{-1}A$
- ▶ Convergence $\equiv \rho(B_R) < 1$

Richardson method 3

Theorem 10.3

The stationary Richardson method converges iff

$$\frac{2\Re(\lambda)}{\tau|\lambda|^2} > 1$$

for all λ - eigenvalues of $P^{-1}A$

Proof.

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\tau} + Ax^{(k)} = b$
- ▶ $x^{(k+1)} = B_R x^{(k)} + f$, $B_R = I - \tau P^{-1}A$
- ▶ Convergence $\equiv \rho(B_R) < 1$
- ▶ y eigenvector, λ eigenvalue of $P^{-1}A$

Richardson method 3

Theorem 10.3

The stationary Richardson method converges iff

$$\frac{2\Re(\lambda)}{\tau|\lambda|^2} > 1$$

for all λ - eigenvalues of $P^{-1}A$

Proof.

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\tau} + Ax^{(k)} = b$
- ▶ $x^{(k+1)} = B_R x^{(k)} + f$, $B_R = I - \tau P^{-1}A$
- ▶ Convergence $\equiv \rho(B_R) < 1$
- ▶ y eigenvector, λ eigenvalue of $P^{-1}A \Rightarrow (I - \tau P^{-1}A)y = (1 - \tau\lambda)y$

Richardson method 3

Theorem 10.3

The stationary Richardson method converges iff

$$\frac{2\Re(\lambda)}{\tau|\lambda|^2} > 1$$

for all λ - eigenvalues of $P^{-1}A$

Proof.

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\tau} + Ax^{(k)} = b$
- ▶ $x^{(k+1)} = B_R x^{(k)} + f$, $B_R = I - \tau P^{-1}A$
- ▶ Convergence $\equiv \rho(B_R) < 1$
- ▶ y eigenvector, λ eigenvalue of $P^{-1}A \Rightarrow (I - \tau P^{-1}A)y = (1 - \tau\lambda)y$
- ▶ $\rho(B_R) < 1 \Rightarrow |1 - \tau\lambda| < 1$



Richardson method 4

Proof.

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\tau} + Ax^{(k)} = b$
- ▶ $x^{(k+1)} = B_R x^{(k)} + f, B_R = I - \tau P^{-1}A$
- ▶ Convergence $\equiv \rho(B_R) < 1$
- ▶ λ eigenvalue of $P^{-1}A \Rightarrow I - \tau P^{-1}A y = (1 - \tau\lambda)y$
- ▶ $\rho(B_R) < 1 \Rightarrow |1 - \tau\lambda| < 1$

Richardson method 4

Proof.

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\tau} + Ax^{(k)} = b$
- ▶ $x^{(k+1)} = B_R x^{(k)} + f, B_R = I - \tau P^{-1}A$
- ▶ Convergence $\equiv \rho(B_R) < 1$
- ▶ y eigenvalue of $P^{-1}A \Rightarrow I - \tau P^{-1}Ay = (1 - \tau\lambda)y$
- ▶ $\rho(B_R) < 1 \Rightarrow |1 - \tau\lambda| < 1$
- ▶

$$\begin{aligned} |1 - \tau\lambda|^2 &= (1 - \tau\Re(\lambda))^2 + (\tau\Im(\lambda))^2 = \\ &= 1 - 2\tau\Re(\lambda) + (\tau\Re(\lambda))^2 + (\tau\Im(\lambda))^2 = 1 - 2\tau\Re(\lambda) + \tau^2|\lambda|^2 \end{aligned}$$

Richardson method 4

Proof.

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\tau} + Ax^{(k)} = b$
- ▶ $x^{(k+1)} = B_R x^{(k)} + f$, $B_R = I - \tau P^{-1}A$
- ▶ Convergence $\equiv \rho(B_R) < 1$
- ▶ y eigenvalue of $P^{-1}A \Rightarrow I - \tau P^{-1}Ay = (1 - \tau\lambda)y$
- ▶ $\rho(B_R) < 1 \Rightarrow |1 - \tau\lambda| < 1$
- ▶

$$\begin{aligned} |1 - \tau\lambda|^2 &= (1 - \tau\Re(\lambda))^2 + (\tau\Im(\lambda))^2 = \\ &= 1 - 2\tau\Re(\lambda) + (\tau\Re(\lambda))^2 + (\tau\Im(\lambda))^2 = 1 - 2\tau\Re(\lambda) + \tau^2|\lambda|^2 \end{aligned}$$

- ▶ $1 - 2\tau\Re(\lambda) + \tau^2|\lambda|^2 < 1$

Richardson method 4

Proof.

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\tau} + Ax^{(k)} = b$
- ▶ $x^{(k+1)} = B_R x^{(k)} + f, B_R = I - \tau P^{-1}A$
- ▶ Convergence $\equiv \rho(B_R) < 1$
- ▶ λ eigenvalue of $P^{-1}A \Rightarrow I - \tau P^{-1}A y = (1 - \tau\lambda)y$
- ▶ $\rho(B_R) < 1 \Rightarrow |1 - \tau\lambda| < 1$
- ▶

$$\begin{aligned} |1 - \tau\lambda|^2 &= (1 - \tau\Re(\lambda))^2 + (\tau\Im(\lambda))^2 = \\ &= 1 - 2\tau\Re(\lambda) + (\tau\Re(\lambda))^2 + (\tau\Im(\lambda))^2 = 1 - 2\tau\Re(\lambda) + \tau^2|\lambda|^2 \end{aligned}$$

- ▶ $1 - 2\tau\Re(\lambda) + \tau^2|\lambda|^2 < 1 \Rightarrow \frac{2\Re(\lambda)}{\tau|\lambda|^2} > 1$



Optimal Step(Parameter) for Stationary Richardson 1

Theorem 10.4

$$\blacktriangleright P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$$

Optimal Step(Parameter) for Stationary Richardson 1

Theorem 10.4

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$
- ▶ $x^{(k+1)} = B_{R,\alpha}x^{(k)} + f, B_{R,\alpha} = I - \alpha P^{-1}A$

Optimal Step(Parameter) for Stationary Richardson 1

Theorem 10.4

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$
- ▶ $x^{(k+1)} = B_{R,\alpha}x^{(k)} + f, B_{R,\alpha} = I - \alpha P^{-1}A$
- ▶ $\lambda_i(P^{-1}A) \in \mathbb{R}$

Optimal Step(Parameter) for Stationary Richardson 1

Theorem 10.4

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$
- ▶ $x^{(k+1)} = B_{R,\alpha}x^{(k)} + f, B_{R,\alpha} = I - \alpha P^{-1}A$
- ▶ $\lambda_i(P^{-1}A) \in \mathbb{R}$
- ▶ $\lambda_1(P^{-1}A) > \lambda_2(P^{-1}A) > \dots > \lambda_n(P^{-1}A) > 0$

Optimal Step(Parameter) for Stationary Richardson 1

Theorem 10.4

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$
- ▶ $x^{(k+1)} = B_{R,\alpha} x^{(k)} + f, B_{R,\alpha} = I - \alpha P^{-1}A$
- ▶ $\lambda_i(P^{-1}A) \in \mathbb{R}$
- ▶ $\lambda_1(P^{-1}A) > \lambda_2(P^{-1}A) > \dots > \lambda_n(P^{-1}A) > 0$
- ▶ \Downarrow

Optimal Step(Parameter) for Stationary Richardson 1

Theorem 10.4

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$
- ▶ $x^{(k+1)} = B_{R,\alpha} x^{(k)} + f, B_{R,\alpha} = I - \alpha P^{-1}A$
- ▶ $\lambda_i(P^{-1}A) \in \mathbb{R}$
- ▶ $\lambda_1(P^{-1}A) > \lambda_2(P^{-1}A) > \dots > \lambda_n(P^{-1}A) > 0$
- ▶ \Downarrow
- ▶ $\alpha_{opt} = \frac{2}{\lambda_1 + \lambda_n}$

Optimal Step(Parameter) for Stationary Richardson 1

Theorem 10.4

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$
- ▶ $x^{(k+1)} = B_{R,\alpha} x^{(k)} + f, B_{R,\alpha} = I - \alpha P^{-1}A$
- ▶ $\lambda_i(P^{-1}A) \in \mathbb{R}$
- ▶ $\lambda_1(P^{-1}A) > \lambda_2(P^{-1}A) > \dots > \lambda_n(P^{-1}A) > 0$
- ▶ \Downarrow
- ▶ $\alpha_{opt} = \frac{2}{\lambda_1 + \lambda_n}$
- ▶ $\rho_{opt} = \min_{\alpha} \rho(B_{R,\alpha}) = \frac{\lambda_1 - \lambda_n}{\lambda_1 + \lambda_n}$

Optimal Step(Parameter) for Stationary Richardson 1

Theorem 10.4

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$
- ▶ $x^{(k+1)} = B_{R,\alpha} x^{(k)} + f, B_{R,\alpha} = I - \alpha P^{-1}A$
- ▶ $\lambda_i(P^{-1}A) \in \mathbb{R}$
- ▶ $\lambda_1(P^{-1}A) > \lambda_2(P^{-1}A) > \dots > \lambda_n(P^{-1}A) > 0$
- ▶ \Downarrow
- ▶ $\alpha_{opt} = \frac{2}{\lambda_1 + \lambda_n}$
- ▶ $\rho_{opt} = \min_{\alpha} \rho(B_{R,\alpha}) = \frac{\lambda_1 - \lambda_n}{\lambda_1 + \lambda_n}$

Proof.

- ▶ $\lambda_i(B_{R,\alpha}) = 1 - \alpha \lambda_i(P^{-1}A), i = 1, 2, \dots, n$

Optimal Step(Parameter) for Stationary Richardson 1

Theorem 10.4

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$
- ▶ $x^{(k+1)} = B_{R,\alpha} x^{(k)} + f, B_{R,\alpha} = I - \alpha P^{-1}A$
- ▶ $\lambda_i(P^{-1}A) \in \mathbb{R}$
- ▶ $\lambda_1(P^{-1}A) > \lambda_2(P^{-1}A) > \dots > \lambda_n(P^{-1}A) > 0$
- ▶ \Downarrow
- ▶ $\alpha_{opt} = \frac{2}{\lambda_1 + \lambda_n}$
- ▶ $\rho_{opt} = \min_{\alpha} \rho(B_{R,\alpha}) = \frac{\lambda_1 - \lambda_n}{\lambda_1 + \lambda_n}$

Proof.

- ▶ $\lambda_i(B_{R,\alpha}) = 1 - \alpha \lambda_i(P^{-1}A), i = 1, 2, \dots, n$
- ▶ $\rho(B_{R,\alpha}) < 1 \Rightarrow |1 - \alpha \lambda_i(P^{-1}A)| < 1, i = 1, 2, \dots, n$



Optimal Step(Parameter) for Stationary Richardson 2

Proof.

- ▶ $\lambda_i(B_{R,\alpha}) = 1 - \alpha\lambda_i(P^{-1}A), i = 1, 2, \dots, n$
- ▶ $\rho(B_{R,\alpha}) < 1 \Rightarrow |1 - \alpha\lambda_i(P^{-1}A)| < 1, i = 1, 2, ..n$

Optimal Step(Parameter) for Stationary Richardson 2

Proof.

- ▶ $\lambda_i(B_{R,\alpha}) = 1 - \alpha\lambda_i(P^{-1}A), i = 1, 2, \dots, n$
- ▶ $\rho(B_{R,\alpha}) < 1 \Rightarrow |1 - \alpha\lambda_i(P^{-1}A)| < 1, i = 1, 2, \dots, n$




Figure: Eigenvalues of $B_{R,\alpha}$ are functions of α

Optimal Step(Parameter) for Stationary Richardson 3

Proof.

$$\blacktriangleright \lambda_i(B_{R,\alpha}) = 1 - \alpha \lambda_i(P^{-1}A), i = 1, 2, \dots, n$$

Optimal Step(Parameter) for Stationary Richardson 3

Proof.

- ▶ $\lambda_i(B_{R,\alpha}) = 1 - \alpha\lambda_i(P^{-1}A), i = 1, 2, \dots, n$
- ▶ $\rho(B_{R,\alpha}) < 1 \Rightarrow |1 - \alpha\lambda_i(P^{-1}A)| < 1, i = 1, 2, \dots, n$

Optimal Step(Parameter) for Stationary Richardson 3

Proof.

- ▶ $\lambda_i(B_{R,\alpha}) = 1 - \alpha\lambda_i(P^{-1}A), i = 1, 2, \dots, n$
- ▶ $\rho(B_{R,\alpha}) < 1 \Rightarrow |1 - \alpha\lambda_i(P^{-1}A)| < 1, i = 1, 2, \dots, n$
- ▶ $\rho(B_{R,\alpha}) = \max_{1 \leq i \leq n, \alpha} |1 - \alpha\lambda_i(P^{-1}A)|$

Optimal Step(Parameter) for Stationary Richardson 3

Proof.

- ▶ $\lambda_i(B_{R,\alpha}) = 1 - \alpha\lambda_i(P^{-1}A), i = 1, 2, \dots, n$
- ▶ $\rho(B_{R,\alpha}) < 1 \Rightarrow |1 - \alpha\lambda_i(P^{-1}A)| < 1, i = 1, 2, \dots, n$
- ▶ $\rho(B_{R,\alpha}) = \max_{1 \leq i \leq n, \alpha} |1 - \alpha\lambda_i(P^{-1}A)|$




Figure: $\rho(B_{R,\alpha})$ is function of α

Optimal Step(Parameter) for Stationary Richardson 4

Proof.




Figure: ρ_{opt} and $\rho(B_{R,\alpha})$

Optimal Step(Parameter) for Stationary Richardson 4

Proof.




Figure: ρ_{opt} and $\rho(B_{R,\alpha})$

► $\alpha_{opt} = \arg \min \rho(B_{R,\alpha})$

Optimal Step(Parameter) for Stationary Richardson 4

Proof.




Figure: ρ_{opt} and $\rho(B_{R,\alpha})$

- ▶ $\alpha_{opt} = \arg \min \rho(B_{R,\alpha})$
- ▶ $\alpha_{opt}\lambda_1(P^{-1}A) - 1 = 1 - \alpha_{opt}\lambda_n(P^{-1}A) \Rightarrow \alpha_{opt} = \frac{2}{\lambda_1 + \lambda_n}$

Optimal Step(Parameter) for Stationary Richardson 4

Proof.




Figure: ρ_{opt} and $\rho(B_{R,\alpha})$

- ▶ $\alpha_{opt} = \arg \min \rho(B_{R,\alpha})$
- ▶ $\alpha_{opt} \lambda_1 (P^{-1}A) - 1 = 1 - \alpha_{opt} \lambda_n (P^{-1}A) \Rightarrow \alpha_{opt} = \frac{2}{\lambda_1 + \lambda_n}$
- ▶ $\rho_{opt} = \min_{\alpha} \rho(B_{R,\alpha}) = 1 - \alpha_{opt} \lambda_n = 1 - \frac{2}{\lambda_1 + \lambda_n} \lambda_n = \frac{\lambda_1 - \lambda_n}{\lambda_1 + \lambda_n}$

Sufficient condition for convergence $P > 0.5\alpha A, 1$

Theorem 10.5

$$\blacktriangleright P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$$

Sufficient condition for convergence $P > 0.5\alpha A, 1$

Theorem 10.5

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$
- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$

Sufficient condition for convergence $P > 0.5\alpha A, 1$

Theorem 10.5

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$
- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $\alpha > 0$

Sufficient condition for convergence $P > 0.5\alpha A, 1$

Theorem 10.5

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$
- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $\alpha > 0$
- ▶ $P > 0.5\alpha A$


Sufficient condition for convergence $P > 0.5\alpha A, 1$

Theorem 10.5

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$
- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $\alpha > 0$
- ▶ $P > 0.5\alpha A$
- ▶ \Downarrow


Sufficient condition for convergence $P > 0.5\alpha A, 1$

Theorem 10.5

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$
- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $\alpha > 0$
- ▶ $P > 0.5\alpha A$
- ▶ 
- ▶ *Errors decrease monotonically in A-norm*

Sufficient condition for convergence $P > 0.5\alpha A, 1$

Theorem 10.5

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$
- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $\alpha > 0$
- ▶ $P > 0.5\alpha A$
- ▶ 
- ▶ *Errors decrease monotonically in A-norm*
- ▶ *Richardson's method converges*

Sufficient condition for convergence $P > 0.5\alpha A, 1$

Theorem 10.5

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$
- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $\alpha > 0$
- ▶ $P > 0.5\alpha A$
- ▶ \Downarrow
- ▶ *Errors decrease monotonically in A -norm*
- ▶ *Richardson's method converges*

Example 10.6

- ▶ $A > 0 \equiv \exists \beta > 0, (Ay, y) > \beta(y, y), \forall y \neq 0, y \in \mathbb{R}^n, (.,.)$ - inner product

Sufficient condition for convergence $P > 0.5\alpha A, 1$

Theorem 10.5

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b$
- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $\alpha > 0$
- ▶ $P > 0.5\alpha A$
- ▶ \Downarrow
- ▶ *Errors decrease monotonically in A -norm*
- ▶ *Richardson's method converges*

Example 10.6

- ▶ $A > 0 \equiv \exists \beta > 0, (Ay, y) > \beta(y, y), \forall y \neq 0, y \in \mathbb{R}^n, (.,.)$ - inner product
- ▶ $P > 0.5\alpha A \equiv \exists \beta > 0, ((P - 0.5\alpha A)y, y) > \beta(y, y) \forall y \neq 0, y \in \mathbb{R}^n, (.,.)$ - inner product

Sufficient condition for convergence $P > 0.5\alpha A$, 2

Proof.

$$\blacktriangleright \begin{cases} P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b \\ P \frac{x - x}{\alpha} + Ax = b \end{cases} \Rightarrow P \frac{e^{(k+1)} - e^{(k)}}{\alpha} + Ae^{(k)} = 0$$

Sufficient condition for convergence $P > 0.5\alpha A$, 2

Proof.

$$\blacktriangleright \begin{cases} P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b \\ P \frac{x - x}{\alpha} + Ax = b \end{cases} \Rightarrow P \frac{e^{(k+1)} - e^{(k)}}{\alpha} + Ae^{(k)} = 0$$

$$\blacktriangleright e^{(k)} = x^{(k)} - x, \quad e^{(k+1)} = (I - \alpha P^{-1}A)e^{(k)}$$

Sufficient condition for convergence $P > 0.5\alpha A$, 2

Proof.

- ▶
$$\begin{cases} P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b \\ P \frac{x - x}{\alpha} + Ax = b \end{cases} \Rightarrow P \frac{e^{(k+1)} - e^{(k)}}{\alpha} + Ae^{(k)} = 0$$
- ▶ $e^{(k)} = x^{(k)} - x, \quad e^{(k+1)} = (I - \alpha P^{-1}A)e^{(k)}$
- ▶ $s_k = (Ae^{(k)}, e^{(k)}), \quad y_k = P^{-1}Ae^{(k)}, \quad e^{(k)} = A^{-1}Py_k$

Sufficient condition for convergence $P > 0.5\alpha A$, 2

Proof.

- ▶
$$\begin{cases} P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b \\ P \frac{x - x}{\alpha} + Ax = b \end{cases} \Rightarrow P \frac{e^{(k+1)} - e^{(k)}}{\alpha} + Ae^{(k)} = 0$$
- ▶ $e^{(k)} = x^{(k)} - x, \quad e^{(k+1)} = (I - \alpha P^{-1}A)e^{(k)}$
- ▶ $s_k = (Ae^{(k)}, e^{(k)}), \quad y_k = P^{-1}Ae^{(k)}, \quad e^{(k)} = A^{-1}Py_k$
- ▶ $e^{(k+1)} = (I - \alpha P^{-1}A)e^{(k)} = e^{(k)} - \alpha y_k$

Sufficient condition for convergence $P > 0.5\alpha A$, 2

Proof.

- ▶ $\begin{cases} P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b \\ P \frac{x - x}{\alpha} + Ax = b \end{cases} \Rightarrow P \frac{e^{(k+1)} - e^{(k)}}{\alpha} + Ae^{(k)} = 0$
- ▶ $e^{(k)} = x^{(k)} - x, \quad e^{(k+1)} = (I - \alpha P^{-1}A)e^{(k)}$
- ▶ $s_k = (Ae^{(k)}, e^{(k)}), \quad y_k = P^{-1}Ae^{(k)}, \quad e^{(k)} = A^{-1}Py_k$
- ▶ $e^{(k+1)} = (I - \alpha P^{-1}A)e^{(k)} = e^{(k)} - \alpha y_k$
- ▶ $Ae^{(k+1)} = Ae^{(k)} - \alpha Ay_k$

Sufficient condition for convergence $P > 0.5\alpha A$, 2

Proof.

- ▶
$$\begin{cases} P \frac{x^{(k+1)} - x^{(k)}}{\alpha} + Ax^{(k)} = b \\ P \frac{x - x}{\alpha} + Ax = b \end{cases} \Rightarrow P \frac{e^{(k+1)} - e^{(k)}}{\alpha} + Ae^{(k)} = 0$$
- ▶ $e^{(k)} = x^{(k)} - x, \quad e^{(k+1)} = (I - \alpha P^{-1}A)e^{(k)}$
- ▶ $s_k = (Ae^{(k)}, e^{(k)}), \quad y_k = P^{-1}Ae^{(k)}, \quad e^{(k)} = A^{-1}Py_k$
- ▶ $e^{(k+1)} = (I - \alpha P^{-1}A)e^{(k)} = e^{(k)} - \alpha y_k$
- ▶ $Ae^{(k+1)} = Ae^{(k)} - \alpha Ay_k$
- ▶

$$\begin{aligned} s_{k+1} &= (Ae^{(k+1)}, e^{(k+1)}) = (Ae^{(k)} - \alpha Ay_k, e^{(k)} - \alpha y_k) = \\ &= (Ae^{(k)}, e^{(k)}) - \alpha(Ae^{(k)}, y_k) - \alpha(Ay_k, e^{(k)}) + \alpha^2(Ay_k, y_k) = \\ &= (Ae^{(k)}, e^{(k)}) - 2\alpha(Ae^{(k)}, y_k) + \alpha^2(Ay_k, y_k) = \\ &= s_k - 2\alpha(AA^{-1}Py^{(k)}, y_k) + \alpha^2(Ay_k, y_k) = \\ &= s_k - 2\alpha(Py^{(k)}, y_k) + \alpha^2(Ay_k, y_k) = s_k - 2\alpha((P - 0.5\alpha A)y_k, y_k) < s_k \end{aligned}$$

Sufficient condition for convergence $P > 0.5\alpha A$, 3

Proof.

$$\blacktriangleright s_{(k+1)} = s_k - 2\alpha ((P - 0.5\alpha A)y_k, y_k)$$

Sufficient condition for convergence $P > 0.5\alpha A$, 3

Proof.

- ▶ $s_{(k+1)} = s_k - 2\alpha ((P - 0.5\alpha A)y_k, y_k)$
- ▶ $0 < (Ae^{(k+)}, e^{(k+1)}) = s_{k+1} < s_k = (Ae^{(k)}, e^{(k)}) \Rightarrow$ convergence to some limit

Sufficient condition for convergence $P > 0.5\alpha A$, 3

Proof.

- ▶ $s_{(k+1)} = s_k - 2\alpha ((P - 0.5\alpha A)y_k, y_k)$
- ▶ $0 < (Ae^{(k+)}, e^{(k+)}) = s_{k+1} < s_k = (Ae^{(k)}, e^{(k)}) \Rightarrow$ convergence to some limit
- ▶ $s_k - s_{k+1} = 2\alpha ((P - 0.5\alpha A)y_k, y_k) \geq 2\alpha\beta(y_k, y_k), \beta > 0$

Sufficient condition for convergence $P > 0.5\alpha A, 3$

Proof.

- ▶ $s_{(k+1)} = s_k - 2\alpha ((P - 0.5\alpha A)y_k, y_k)$
- ▶ $0 < (Ae^{(k+1)}, e^{(k+1)}) = s_{k+1} < s_k = (Ae^{(k)}, e^{(k)}) \Rightarrow$ convergence to some limit
- ▶ $s_k - s_{k+1} = 2\alpha ((P - 0.5\alpha A)y_k, y_k) \geq 2\alpha\beta(y_k, y_k), \beta > 0$
- ▶ $s_k - s_{k+1} \geq 2\alpha\beta(y_k, y_k) \Rightarrow \lim_{k \rightarrow \infty} \|y_k\|_2 = 0$

Sufficient condition for convergence $P > 0.5\alpha A$, 3

Proof.

- ▶ $s_{(k+1)} = s_k - 2\alpha ((P - 0.5\alpha A)y_k, y_k)$
- ▶ $0 < (Ae^{(k+1)}, e^{(k+1)}) = s_{k+1} < s_k = (Ae^{(k)}, e^{(k)}) \Rightarrow$ convergence to some limit
- ▶ $s_k - s_{k+1} = 2\alpha ((P - 0.5\alpha A)y_k, y_k) \geq 2\alpha\beta(y_k, y_k), \beta > 0$
- ▶ $s_k - s_{k+1} \geq 2\alpha\beta(y_k, y_k) \Rightarrow \lim_{k \rightarrow \infty} \|y_k\|_2 = 0$
- ▶ $\begin{cases} \|e^{(k)}\|_2 = \|A^{-1}Py_k\|_2 \leq \|A^{-1}P\|_2 \cdot \|y_k\|_2 \\ \lim_{k \rightarrow \infty} \|y_k\|_2 = 0 \end{cases} \Rightarrow \lim_{k \rightarrow \infty} \|e^{(k)}\|_2 = 0$



Convergence SOR, 1

Example 10.7

$$\blacktriangleright P \frac{x^{(k+1)} - x^{(k)}}{\omega} + Ax^{(k)} = b$$

Convergence SOR, 1

Example 10.7

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\omega} + Ax^{(k)} = b$
- ▶ $\text{SOR} \equiv P = D + \omega L, A = L + D + U$

Convergence SOR, 1

Example 10.7

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\omega} + Ax^{(k)} = b$
- ▶ $\text{SOR} \equiv P = D + \omega L, A = L + D + U$
- ▶

$$B_{R,\omega} = P^{-1}(P - \omega A) = (D + \omega L)^{-1}(D + \omega L - \omega(L + D + U)) = \\ (D + \omega L)^{-1}((1 - \omega)D - \omega U) \equiv B_{\text{SOR}}$$

Convergence SOR, 1

Example 10.7

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\omega} + Ax^{(k)} = b$
- ▶ $\text{SOR} \equiv P = D + \omega L, A = L + D + U$
- ▶

$$B_{R,\omega} = P^{-1}(P - \omega A) = (D + \omega L)^{-1}(D + \omega L - \omega(L + D + U)) = \\ (D + \omega L)^{-1}((1 - \omega)D - \omega U) \equiv B_{\text{SOR}}$$

Theorem 10.8

- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$

Convergence SOR, 1

Example 10.7

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\omega} + Ax^{(k)} = b$
- ▶ $\text{SOR} \equiv P = D + \omega L, A = L + D + U$
- ▶

$$B_{R,\omega} = P^{-1}(P - \omega A) = (D + \omega L)^{-1}(D + \omega L - \omega(L + D + U)) = \\ (D + \omega L)^{-1}((1 - \omega)D - \omega U) \equiv B_{\text{SOR}}$$

Theorem 10.8

- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $0 < \omega < 2$

Convergence SOR, 1

Example 10.7

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\omega} + Ax^{(k)} = b$
- ▶ $\text{SOR} \equiv P = D + \omega L, A = L + D + U$
- ▶

$$B_{R,\omega} = P^{-1}(P - \omega A) = (D + \omega L)^{-1}(D + \omega L - \omega(L + D + U)) = \\ (D + \omega L)^{-1}((1 - \omega)D - \omega U) \equiv B_{\text{SOR}}$$

Theorem 10.8

- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $0 < \omega < 2$
- ▶ \Downarrow

Convergence SOR, 1

Example 10.7

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\omega} + Ax^{(k)} = b$
- ▶ $\text{SOR} \equiv P = D + \omega L, A = L + D + U$
- ▶

$$B_{R,\omega} = P^{-1}(P - \omega A) = (D + \omega L)^{-1}(D + \omega L - \omega(L + D + U)) = \\ (D + \omega L)^{-1}((1 - \omega)D - \omega U) \equiv B_{\text{SOR}}$$

Theorem 10.8

- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $0 < \omega < 2$
- ▶ \Downarrow
- ▶ *SOR Converges*

Convergence SOR, 2

Theorem 10.9

- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $0 < \omega < 2$
- ▶ \Downarrow
- ▶ *SOR converges*

Convergence SOR, 2

Theorem 10.9

- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $0 < \omega < 2$
- ▶ \Downarrow
- ▶ *SOR converges*

Proof.

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\omega} + Ax^{(k)} = b$

Convergence SOR, 2

Theorem 10.9

- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $0 < \omega < 2$
- ▶ \Downarrow
- ▶ *SOR converges*

Proof.

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\omega} + Ax^{(k)} = b$
- ▶ $\text{SOR} \equiv P = D + \omega L, A = L + D + U$

Convergence SOR, 2

Theorem 10.9

- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $0 < \omega < 2$
- ▶ \Downarrow
- ▶ *SOR converges*

Proof.

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\omega} + Ax^{(k)} = b$
- ▶ $\text{SOR} \equiv P = D + \omega L, A = L + D + U$
- ▶ $P > 0.5\omega A$?

Convergence SOR, 2

Theorem 10.9

- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $0 < \omega < 2$
- ▶ \Downarrow
- ▶ *SOR converges*

Proof.

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\omega} + Ax^{(k)} = b$
- ▶ $\text{SOR} \equiv P = D + \omega L, A = L + D + U$
- ▶ $P > 0.5\omega A$?
- ▶ $A = A^T > 0, y \neq 0 \Rightarrow ((L + D + U)y, y) = ((D + 2L)y, y), (Dy, y) > 0$

Convergence SOR, 2

Theorem 10.9

- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $0 < \omega < 2$
- ▶ \Downarrow
- ▶ *SOR converges*

Proof.

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\omega} + Ax^{(k)} = b$
- ▶ $\text{SOR} \equiv P = D + \omega L, A = L + D + U$
- ▶ $P > 0.5\omega A$?
- ▶ $A = A^T > 0, y \neq 0 \Rightarrow ((L + D + U)y, y) = ((D + 2L)y, y), (Dy, y) > 0$
 $((D + \omega L - 0.5\omega A)y, y) = ((D + \omega L - 0.5\omega(L + D + U))y, y)$

Convergence SOR, 2

Theorem 10.9

- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $0 < \omega < 2$
- ▶ \Downarrow
- ▶ *SOR converges*

Proof.

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\omega} + Ax^{(k)} = b$
- ▶ $\text{SOR} \equiv P = D + \omega L, A = L + D + U$
- ▶ $P > 0.5\omega A$?
- ▶ $A = A^T > 0, y \neq 0 \Rightarrow ((L + D + U)y, y) = ((D + 2L)y, y), (Dy, y) > 0$
 $((D + \omega L - 0.5\omega A)y, y) = ((D + \omega L - 0.5\omega(L + D + U))y, y)$
- ▶ $((D + \omega L - 0.5\omega A)y, y) = ((D + \omega L - 0.5\omega(2L + D))y, y)$

Convergence SOR, 2

Theorem 10.9

- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $0 < \omega < 2$
- ▶ \Downarrow
- ▶ *SOR converges*

Proof.

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\omega} + Ax^{(k)} = b$
- ▶ $\text{SOR} \equiv P = D + \omega L, A = L + D + U$
- ▶ $P > 0.5\omega A$?
- ▶ $A = A^T > 0, y \neq 0 \Rightarrow ((L + D + U)y, y) = ((D + 2L)y, y), (Dy, y) > 0$
 $((D + \omega L - 0.5\omega A)y, y) = ((D + \omega L - 0.5\omega(L + D + U))y, y)$
- ▶ $((D + \omega L - 0.5\omega A)y, y) = ((D + \omega L - 0.5\omega(2L + D))y, y)$
- ▶ $((D + \omega L - 0.5\omega A)y, y) = (1 - 0.5\omega)(Dy, y)$

Convergence SOR, 2

Theorem 10.9

- ▶ $A \in \mathbb{R}^{n \times n}, A = A^T > 0$
- ▶ $0 < \omega < 2$
- ▶ \Downarrow
- ▶ *SOR converges*

Proof.

- ▶ $P \frac{x^{(k+1)} - x^{(k)}}{\omega} + Ax^{(k)} = b$
- ▶ $\text{SOR} \equiv P = D + \omega L, A = L + D + U$
- ▶ $P > 0.5\omega A$?
- ▶ $A = A^T > 0, y \neq 0 \Rightarrow ((L + D + U)y, y) = ((D + 2L)y, y), (Dy, y) > 0$
 $((D + \omega L - 0.5\omega A)y, y) = ((D + \omega L - 0.5\omega(L + D + U))y, y)$
- ▶ $((D + \omega L - 0.5\omega A)y, y) = ((D + \omega L - 0.5\omega(2L + D))y, y)$
- ▶ $((D + \omega L - 0.5\omega A)y, y) = (1 - 0.5\omega)(Dy, y)$
- ▶ \Downarrow
- ▶ $P > 0.5\omega A$ if $1 - 0.5\omega > 0 \Rightarrow$ convergence

Preconditioning matrices

- ▶ Q: How to Build optimal preconditioning matrix?

Preconditioning matrices

- ▶ Q: How to Build optimal preconditioning matrix?
- ▶ A: There is no general method for constructing optimal preconditioning matrix

Preconditioning matrices

- ▶ Q: How to Build optimal preconditioning matrix?
- ▶ A: There is no general method for constructing optimal preconditioning matrix
- ▶ Q: What is optimal preconditioning matrix?

Preconditioning matrices

- ▶ Q: How to Build optimal preconditioning matrix?
 - ▶ A: There is no general method for constructing optimal preconditioning matrix
-
- ▶ Q: What is optimal preconditioning matrix?
 - ▶ A: $P = A$ is optimal preconditioning matrix \Rightarrow **inefficient**

Preconditioning matrices

- ▶ Q: How to Build optimal preconditioning matrix?
 - ▶ A: There is no general method for constructing optimal preconditioning matrix
-
- ▶ Q: What is optimal preconditioning matrix?
 - ▶ A: $P = A$ is optimal preconditioning matrix \Rightarrow **inefficient**
-
- ▶ Richardson's method $P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$

Preconditioning matrices

- ▶ Q: How to Build optimal preconditioning matrix?
 - ▶ A: There is no general method for constructing optimal preconditioning matrix
-
- ▶ Q: What is optimal preconditioning matrix?
 - ▶ A: $P = A$ is optimal preconditioning matrix \Rightarrow **inefficient**
-
- ▶ Richardson's method $P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$
 - ▶ $r^{(k)} = b - Ax^{(k)}, z^{(k)} = x^{(k+1)} - x^{(k)}$

Preconditioning matrices

- ▶ Q: How to Build optimal preconditioning matrix?
 - ▶ A: There is no general method for constructing optimal preconditioning matrix
-
- ▶ Q: What is optimal preconditioning matrix?
 - ▶ A: $P = A$ is optimal preconditioning matrix \Rightarrow **inefficient**
-
- ▶ Richardson's method $P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$
 - ▶ $r^{(k)} = b - Ax^{(k)}, z^{(k)} = x^{(k+1)} - x^{(k)}$
 - ▶ $Pz^{(k)} = \tau_k r^{(k)}$

Preconditioning matrices

- ▶ Q: How to Build optimal preconditioning matrix?
 - ▶ A: There is no general method for constructing optimal preconditioning matrix
-
- ▶ Q: What is optimal preconditioning matrix?
 - ▶ A: $P = A$ is optimal preconditioning matrix \Rightarrow **inefficient**
-
- ▶ Richardson's method $P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$
 - ▶ $r^{(k)} = b - Ax^{(k)}, z^{(k)} = x^{(k+1)} - x^{(k)}$
 - ▶ $Pz^{(k)} = \tau_k r^{(k)}$
 - ▶ $x^{(k+1)} = x^{(k)} + z^{(k)}$

Preconditioning matrices

- ▶ Q: How to Build optimal preconditioning matrix?
 - ▶ A: There is no general method for constructing optimal preconditioning matrix
-
- ▶ Q: What is optimal preconditioning matrix?
 - ▶ A: $P = A$ is optimal preconditioning matrix \Rightarrow **inefficient**
-
- ▶ Richardson's method $P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$
 - ▶ $r^{(k)} = b - Ax^{(k)}, z^{(k)} = x^{(k+1)} - x^{(k)}$
 - ▶ $Pz^{(k)} = \tau_k r^{(k)}$
 - ▶ $x^{(k+1)} = x^{(k)} + z^{(k)}$

What are requirements to preconditioning matrix?

Preconditioning matrices

- ▶ Q: How to Build optimal preconditioning matrix?
- ▶ A: There is no general method for constructing optimal preconditioning matrix

- ▶ Q: What is optimal preconditioning matrix?
- ▶ A: $P = A$ is optimal preconditioning matrix \Rightarrow **inefficient**

- ▶ Richardson's method $P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$
- ▶ $r^{(k)} = b - Ax^{(k)}, z^{(k)} = x^{(k+1)} - x^{(k)}$
- ▶ $Pz^{(k)} = \tau_k r^{(k)}$
- ▶ $x^{(k+1)} = x^{(k)} + z^{(k)}$

What are requirements to preconditioning matrix?

- ▶ Reduce spectral radius of $(I - \tau P^{-1}A)$?

Preconditioning matrices

- ▶ Q: How to Build optimal preconditioning matrix?
- ▶ A: There is no general method for constructing optimal preconditioning matrix

- ▶ Q: What is optimal preconditioning matrix?
- ▶ A: $P = A$ is optimal preconditioning matrix \Rightarrow **inefficient**

- ▶ Richardson's method $P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$
- ▶ $r^{(k)} = b - Ax^{(k)}, z^{(k)} = x^{(k+1)} - x^{(k)}$
- ▶ $Pz^{(k)} = \tau_k r^{(k)}$
- ▶ $x^{(k+1)} = x^{(k)} + z^{(k)}$

What are requirements to preconditioning matrix?

- ▶ Reduce spectral radius of $(I - \tau P^{-1}A)$?
- ▶ $P^{-1}A$ almost normal, eigenvalues clustered in small region

Preconditioning matrices

How to devise preconditioning matrix?

Preconditioning matrices

How to devise preconditioning matrix?

- ▶ Diagonal preconditioners

Preconditioning matrices

How to devise preconditioning matrix?

- ▶ Diagonal preconditioners $p_{ii} = \sum_{j=1}^n |a_{ij}|$

Preconditioning matrices

How to devise preconditioning matrix?

- ▶ Diagonal preconditioners $p_{ii} = \sum_{j=1}^n |a_{ij}|$
- ▶ ILU factorization, for sparse matrices

Preconditioning matrices

How to devise preconditioning matrix?

- ▶ Diagonal preconditioners $p_{ii} = \sum_{j=1}^n |a_{ij}|$
- ▶ ILU factorization, for sparse matrices

```
[n,m]=size(A);  
if n ~= m, error('Only square matrices'); end  
for k=1:n-1  
    for i=k+1:n,  
        if A(i,k) ~= 0  
            if A(k,k) == 0, error('Null pivot element'); end  
            A(i,k)=A(i,k)/A(k,k);  
            for j=k+1:n  
                if A(i,j) ~= 0  
                    A(i,j)=A(i,j)-A(i,k)*A(k,j);  
                end  
            end  
        end  
    end  
end  
end
```

Figure: Quarteroni et al.

Preconditioning matrices

How to devise preconditioning matrix?

- ▶ Diagonal preconditioners $p_{ii} = \sum_{j=1}^n |a_{ij}|$
- ▶ ILU factorization, for sparse matrices

```
[n,m]=size(A);  
if n ~= m, error('Only square matrices'); end  
for k=1:n-1  
    for i=k+1:n,  
        if A(i,k) ~= 0  
            if A(k,k) == 0, error('Null pivot element'); end  
            A(i,k)=A(i,k)/A(k,k);  
            for j=k+1:n  
                if A(i,j) ~= 0  
                    A(i,j)=A(i,j)-A(i,k)*A(k,j);  
                end  
            end  
        end  
    end  
end  
end
```

Figure: Quarteroni et al.

- ▶ Von Neuman method

Preconditioning matrices

How to devise preconditioning matrix?

- ▶ Diagonal preconditioners $p_{ii} = \sum_{j=1}^n |a_{ij}|$
- ▶ ILU factorization, for sparse matrices

```
[n,m]=size(A);  
if n ~= m, error('Only square matrices'); end  
for k=1:n-1  
    for i=k+1:n,  
        if A(i,k) ~= 0  
            if A(k,k) == 0, error('Null pivot element'); end  
            A(i,k)=A(i,k)/A(k,k);  
            for j=k+1:n  
                if A(i,j) ~= 0  
                    A(i,j)=A(i,j)-A(i,k)*A(k,j);  
                end  
            end  
        end  
    end  
end  
end
```

Figure: Quarteroni et al.

- ▶ Von Neuman method $P^{-1} = p(A)$,

Preconditioning matrices

How to devise preconditioning matrix?

- ▶ Diagonal preconditioners $p_{ii} = \sum_{j=1}^n |a_{ij}|$
- ▶ ILU factorization, for sparse matrices

```
[n,m]=size(A);  
if n ~= m, error('Only square matrices'); end  
for k=1:n-1  
    for i=k+1:n,  
        if A(i,k) ~= 0  
            if A(k,k) == 0, error('Null pivot element'); end  
            A(i,k)=A(i,k)/A(k,k);  
            for j=k+1:n  
                if A(i,j) ~= 0  
                    A(i,j)=A(i,j)-A(i,k)*A(k,j);  
                end  
            end  
        end  
    end  
end  
end
```

Figure: Quarteroni et al.

- ▶ Von Neuman method $P^{-1} = p(A)$, $A = (I - CD^{-1})D$

Preconditioning matrices

How to devise preconditioning matrix?

- ▶ Diagonal preconditioners $p_{ii} = \sum_{j=1}^n |a_{ij}|$
- ▶ ILU factorization, for sparse matrices

```
[n,m]=size(A);  
if n ~= m, error('Only square matrices'); end  
for k=1:n-1  
    for i=k+1:n,  
        if A(i,k) ~= 0  
            if A(k,k) == 0, error('Null pivot element'); end  
            A(i,k)=A(i,k)/A(k,k);  
            for j=k+1:n  
                if A(i,j) ~= 0  
                    A(i,j)=A(i,j)-A(i,k)*A(k,j);  
                end  
            end  
        end  
    end  
end  
end
```

Figure: Quarteroni et al.

- ▶ Von Neuman method $P^{-1} = p(A)$, $A = (I - CD^{-1})D$
 $A^{-1} = D^{-1}(I - CD^{-1})^{-1} = D^{-1}(I + CD^{-1} + (CD^{-1})^2 + \dots)$

Q & A