

expected run time of quicksort

run time = number of comparisons

product of probability spaces

$A, B \subseteq S$ independent iff $p(A \cap B) = p(A) \cdot p(B)$

Lemma 1. $W_1 \times W_2$ is a probability space

single coin flip or single throw of dice:
too simple for example

let's consider *two* experiments

$$W_1 = (S_1, p_1), W_2 = (S_2, p_2)$$

set of outcomes for the pair of events

$$S = S_1 \times S_2 = \{(a, b) | a \in S_1, b \in S_2\}$$

$$p(a, b) = p_1(a) \cdot p_2(b)$$

$$W = W_1 \times W_2 = (S, p)$$

random variables from different independent experiments

$W = (S, p)$ probability space

Lemma 9.

$X : S \rightarrow \mathbb{R}$ random variable

$$E(X) = E(X_1) + E(X_2)$$

expected value of random variable X

$$E(X) = \sum_{a \in S} X(a) \cdot p(a)$$

For $i \in \{1, 2\}$ let

$$W_i = (S_i, p_i)$$

be probability spaces and let

$$X_i : S_i \rightarrow \mathbb{R}$$

be random variables in these spaces.

sum if we perform independent experiments

$$X : S_1 \times S_2 \rightarrow \mathbb{R} \quad , \quad X(a, b) = X_1(a) + X_2(b)$$

$W = (S, p)$ probability space

$X : S \rightarrow \mathbb{R}$ random variable

expected value of random variable X

$$E(X) = \sum_{a \in S} X(a) \cdot p(a)$$

probability of B given A

$$A, B \subseteq S, p(A) \neq 0$$

$$p(B|A) = \frac{p(B \cap A)}{p(A)}$$

two experiments:

- throw coin
- if 0 throw coin c, otherwise dice d
- expected total number of points

it might be

$$E(c) + 1/2 \cdot E(c) + 1/2 \cdot E(d) = 1/2 + 1/4 + 7/4$$

sanity check 1

first experiment:

$$W_S = (S, p)$$

second experiments:

$$W_i = (R_i, p_i) \quad i \in S$$

probability space:

$$W_Q = (Q, q)$$

$$Q = \bigcup_{i \in S} \{i\} \times R_i$$

$$a \in R_i \rightarrow q(i, a) = p(i) \cdot p_i(a)$$

Lemma 10. $W_Q = (Q, q)$ is a probability space

first experiment:
 $W_S = (S, p)$

second experiments:
 $W_i = (R_i, p_i) \quad i \in S$

probability space:
 $W_Q = (Q, q)$
 $Q = \bigcup_{i \in S} \{i\} \times R_i$
 $a \in R_i \rightarrow q(i, a) = p(i) \cdot p_i(a)$

random variables on these spaces

$X_0 : S \rightarrow \mathbb{R} \quad , \quad X_i : R_i \rightarrow \mathbb{R}$

$X : Q \rightarrow \mathbb{R}$

$X(i, r) = X_0(i) + X_i(r)$

Lemma 12.

$E(X) = E(X_0) + \sum_{i \in S} p(i) \cdot E(X_i)$

two experiments:

- throw coin
- if 0 throw coin c , otherwise dice d
- expected total number of points

$E(c) + 1/2 \cdot E(c) + 1/2 \cdot E(d) = 1/2 + 1/4 + 7/4$

Quicksort

here: the algorithm does the random experiments

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$ we assume here: $a(i)$ mutually distinct

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$$A_{<} = \{a \in A | a < s\}$$

$$A_{>} = \{a \in A | a > s\}$$

$$\text{sort}(A) = \text{sort}(A_{<}) \circ s \circ \text{sort}(A_{>})$$

$$W_1 = (S_1, p_1), W_2 = (S_2, p_2)$$

$$p(a,b) = p_1(a) \cdot p_2(b)$$

$$W = W_1 \times W_2 = (S, p)$$

Lemma 1. *$W_1 \times W_2$ is a probability space*

For $i \in \{1, 2\}$ let

$$W_i = (S_i, p_i)$$

be probability spaces and let

$$X_i : S_i \rightarrow \mathbb{R}$$

be random variables in these spaces.

$$X : S_1 \times S_2 \rightarrow \mathbb{R} \quad , \quad X(a,b) = X_1(a) + X_2(b)$$

Lemma 9.

$$E(X) = E(X_1) + E(X_2)$$

$$s \in \{a(1), \ldots, a(n)\}$$

$$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$$

first experiment:

$$W_S = (S, p)$$

second experiments:

$$W_i = (R_i, p_i) \qquad i \in S$$

probability space:

$$W_Q = (Q, q)$$

$$Q = \bigcup_{i \in S} \{i\} \times R_i$$

$$a \in R_i \rightarrow q(i, a) = p(i) \cdot p_i(a)$$

Lemma 10. *$W_Q = (Q, q)$ is a probability space*

$$X_0 : S \rightarrow \mathbb{R} \quad , \quad X_i : R_i \rightarrow \mathbb{R}$$

$$X : Q \rightarrow \mathbb{R}$$

$$X(i, r) = X_0(i) + X_i(r)$$

Lemma 12.

$$E(X) = E(X_0) + \Sigma_{i \in S} p(i) \cdot E(X_i)$$

Define by induction on n probability spaces for sorting
 n distinct numbers

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

$$QS_n = (Q_n, q_n)$$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$$A_{<} = \{a \in A \mid a < s\}$$

a(i) mutually distinct

$$A_{>} = \{a \in A \mid a > s\}$$

$$\text{sort}(A) = \text{sort}(A_{<}) \circ s \circ \text{sort}(A_{>})$$

Define by induction on n probability spaces for sorting
 n distinct numbers

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$A_{<} = \{a \in A \mid a < s\}$ **a(i) mutually distinct**

$A_{>} = \{a \in A \mid a > s\}$

$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$

$QS_n = (Q_n, q_n)$ $RS_{n,i} = QS_{i-1} \times QS_{n-i}$

and random variables

$t_n : Q_n \rightarrow \mathbb{R}$

$x \in Q_n \rightarrow t_n(x) = \text{number of comparisons in run } x$

Define by induction on n probability spaces for sorting
 n distinct numbers

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$A_{<} = \{a \in A \mid a < s\}$ **a(i) mutually distinct**

$A_{>} = \{a \in A \mid a > s\}$

$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$

$$QS_n = (Q_n, q_n) \quad RS_{n,i} = QS_{i-1} \times QS_{n-i}$$

and random variables

$$t_n : Q_n \rightarrow \mathbb{R}$$

$x \in Q_n \rightarrow t_n(x) = \text{number of comparisons in run } x$

$$n \in \{0,1\}: \quad Q_n = \{\perp\} \quad q_n(\perp) = 1 \quad t_n(\perp) = 0$$

0 or 1 elements: no randomness
no comparisons

Define by induction on n probability spaces for sorting
 n distinct numbers

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$A_{<} = \{a \in A \mid a < s\}$ a(i) mutually distinct

$A_{>} = \{a \in A \mid a > s\}$

$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$

$$QS_n = (Q_n, q_n) \quad RS_{n,i} = QS_{i-1} \times QS_{n-i}$$

and random variables

$$t_n : Q_n \rightarrow \mathbb{R}$$

$x \in Q_n \rightarrow t_n(x) = \text{number of comparisons in run } x$

$$n \in \{0,1\}: \quad Q_n = \{\perp\} \quad q_n(\perp) = 1 \quad t_n(\perp) = 0$$

0 or 1 elements: no randomness
no comparisons

$$n \geq 2: \quad S_n = \{1, \dots, n\} \quad i \in S_n \quad i = \#A_{<} + 1$$

rank of splitter s : number of $a(i) \leq s$.

Define by induction on n probability spaces for sorting
n distinct numbers

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$A_{<} = \{a \in A | a < s\}$ a(i) mutually distinct

$A_{>} = \{a \in A | a > s\}$

$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$

$$QS_n = (Q_n, q_n) \qquad RS_{n,i} = QS_{i-1} \times QS_{n-i}$$

and random variables

$$t_n : Q_n \rightarrow \mathbb{R}$$

$$x \in Q_n \rightarrow t_n(x) = \text{number of comparisons in run } x$$

$$n \in \{0,1\}: \quad Q_n = \{\perp\} \quad q_n(\perp) = 1 \quad t_n(\perp) = 0$$

0 or 1 elements: no randomness
no comparisons

$$n \geq 2: \quad S_n = \{1, \dots, n\} \quad i \in S_n \quad i = \#A_{<} + 1 \quad p(i) = 1/n$$

$W = (S_n, p)$ is probability space.

rank of splitter s: number of a(i) ≤ s.

Define by induction on n probability spaces for sorting n distinct numbers

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$A_{<} = \{a \in A \mid a < s\}$ **a(i) mutually distinct**

$A_{>} = \{a \in A \mid a > s\}$

$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$

$$QS_n = (Q_n, q_n) \quad RS_{n,i} = QS_{i-1} \times QS_{n-i}$$

and random variables

$$t_n : Q_n \rightarrow \mathbb{R}$$

$x \in Q_n \rightarrow t_n(x) = \text{number of comparisons in run } x$

$$n \in \{0,1\}: \quad Q_n = \{\perp\} \quad q_n(\perp) = 1 \quad t_n(\perp) = 0$$

0 or 1 elements: no randomness
no comparisons

$$n \geq 2: \quad S_n = \{1, \dots, n\} \quad i \in S_n \quad i = \#A_{<} + 1 \quad r_n(i) = 1/n \quad W = (S_n, r_n) \text{ is probability space.}$$

$$R_{n,i} = (Q_{i-1} \times Q_{n-i})$$

$$q_{n,i}(a, b) = q_{i-1}(a) \cdot q_{n-i}(b)$$

$$Q_n = \bigcup_{i \in S_n} \{i\} \times (Q_{i-1} \times Q_{n-i})$$

$$\begin{aligned} Q_2 &= \{1\} \times Q_0 \times Q_1 \cup \{2\} \times Q_1 \times Q_0 = \\ &= \{(1, \perp, \perp), (2, \perp, \perp)\} \end{aligned}$$

$$q_2(1, \perp, \perp) = \frac{1}{2} q_{2,1}(\perp, \perp) = \frac{1}{2} \cdot 1 \cdot 1 = \frac{1}{2} = q_2(2, \perp, \perp)$$

$$q_n(i, (a, b)) = (1/n) \cdot q_{n,i}(a, b)$$

Define by induction on n probability spaces for sorting n distinct numbers

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$A_{<} = \{a \in A \mid a < s\}$ **a(i) mutually distinct**

$A_{>} = \{a \in A \mid a > s\}$

$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$

$$QS_n = (Q_n, q_n) \quad RS_{n,i} = QS_{i-1} \times QS_{n-i}$$

and random variables

$$t_n : Q_n \rightarrow \mathbb{R}$$

$x \in Q_n \rightarrow t_n(x) = \text{number of comparisons in run } x$

$$n \in \{0,1\}: \quad Q_n = \{\perp\} \quad q_n(\perp) = 1 \quad t_n(\perp) = 0$$

0 or 1 elements: no randomness
no comparisons

$$n \geq 2: \quad S_n = \{1, \dots, n\} \quad i \in S_n \quad i = \#A_{<} + 1 \quad r_n(i) = 1/n \quad W = (S_n, r_n) \text{ is probability space.}$$

$$R_{n,i} = (Q_{i-1} \times Q_{n-i})$$

$$q_{n,i}(a, b) = q_{i-1}(a) \cdot q_{n-i}(b)$$

$$Q_n = \bigcup_{i \in S_n} \{i\} \times (Q_{i-1} \times Q_{n-i})$$

$$\begin{aligned} Q_3 &= \{1\} \times Q_0 \times Q_2 \cup \{2\} \times Q_1 \times Q_1 \cup \{3\} \times Q_2 \times Q_0 = \\ &= \left\{ \left(1, (\perp), (1, \perp, \perp) \right), \left(1, (\perp), (2, \perp, \perp) \right), \left(2, \perp, \perp \right), \right. \\ &\quad \left. \left(3, (1, \perp, \perp, \perp) \right), \left(3, (2, \perp, \perp, \perp) \right) \right\} \\ &\quad \frac{1}{3} \cdot 1 \cdot \frac{1}{2} = \frac{1}{6} \quad \frac{1}{6} \quad \frac{1}{3} \end{aligned}$$

$$q_n(i, (a, b)) = (1/n) \cdot q_{n,i}(a, b)$$

Define by induction on n probability spaces for sorting
n distinct numbers

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$A_{<} = \{a \in A | a < s\}$ a(i) mutually distinct

$A_{>} = \{a \in A | a > s\}$

$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$

$$QS_n = (Q_n, q_n) \qquad RS_{n,i} = QS_{i-1} \times QS_{n-i}$$

and random variables

$$t_n : Q_n \rightarrow \mathbb{R}$$

$x \in Q_n \rightarrow t_n(x) = \text{number of comparisons in run } x$

$$n \in \{0,1\}: \quad Q_n = \{\perp\} \quad q_n(\perp) = 1 \quad t_n(\perp) = 0$$

0 or 1 elements: no randomness
no comparisons

$$n \geq 2: \quad S_n = \{1, \dots, n\} \quad i \in S_n \quad i = \#A_{<} + 1 \quad r_n(i) = 1/n \quad W = (S_n, r_n) \text{ is probability space.}$$

$$R_{n,i} = (Q_{i-1} \times Q_{n-i})$$

Lemma: For all i and n: QS_n and $RS_{n,i}$ are probability spaces

$$q_{n,i}(a, b) = q_{i-1}(a) \cdot q_{n-i}(b)$$

$$Q_n = \bigcup_{i \in S_n} \{i\} \times (Q_{i-1} \times Q_{n-i})$$

$$q_n(i, (a, b)) = (1/n) \cdot q_{n,i}(a, b)$$

Define by induction on n probability spaces for sorting
n distinct numbers

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$A_{<} = \{a \in A | a < s\}$ a(i) mutually distinct

$A_{>} = \{a \in A | a > s\}$

$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$

$$QS_n = (Q_n, q_n) \qquad RS_{n,i} = QS_{i-1} \times QS_{n-i}$$

and random variables

$$t_n : Q_n \rightarrow \mathbb{R}$$

$x \in Q_n \rightarrow t_n(x) = \text{number of comparisons in run } x$

$$n \in \{0,1\}: \quad Q_n = \{\perp\} \quad q_n(\perp) = 1 \quad t_n(\perp) = 0$$

0 or 1 elements: no randomness
no comparisons

$$n \geq 2: \quad S_n = \{1, \dots, n\} \quad i \in S_n \quad i = \#A_{<} + 1 \quad r_n(i) = 1/n \quad W = (S_n, r_n) \text{ is probability space.}$$

$$R_{n,i} = (Q_{i-1} \times Q_{n-i})$$

Lemma: For all i and n: QS_n and $RS_{n,i}$ are probability spaces

$$q_{n,i}(a, b) = q_{i-1}(a) \cdot q_{n-i}(b)$$

proof: induction on n.
n = 0 or 1: trivial

$$Q_n = \bigcup_{i \in S_n} \{i\} \times (Q_{i-1} \times Q_{n-i})$$

$$q_n(i, (a, b)) = (1/n) \cdot q_{n,i}(a, b)$$

Define by induction on n probability spaces for sorting
n distinct numbers

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$A_{<} = \{a \in A \mid a < s\}$ a(i) mutually distinct

$A_{>} = \{a \in A \mid a > s\}$

$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$

$$QS_n = (Q_n, q_n) \quad RS_{n,i} = QS_{i-1} \times QS_{n-i}$$

and random variables

$$t_n : Q_n \rightarrow \mathbb{R}$$

$x \in Q_n \rightarrow t_n(x) = \text{number of comparisons in run } x$

$$n \in \{0,1\}: \quad Q_n = \{\perp\} \quad q_n(\perp) = 1 \quad t_n(\perp) = 0$$

0 or 1 elements: no randomness
no comparisons

$$n \geq 2: \quad S_n = \{1, \dots, n\} \quad i \in S_n \quad i = \#A_{<} + 1 \quad r_n(i) = 1/n \quad W = (S_n, r_n) \text{ is probability space.}$$

$$R_{n,i} = (Q_{i-1} \times Q_{n-i})$$

Lemma: For all i and n: QS_n and $RS_{n,i}$ are probability spaces

$$q_{n,i}(a, b) = q_{i-1}(a) \cdot q_{n-i}(b)$$

proof: induction on n.

$$Q_n = \bigcup_{i \in S_n} \{i\} \times (Q_{i-1} \times Q_{n-i})$$

- n = 0 or 1: trivial
- assume true for j < n:

$$q_n(i, (a, b)) = (1/n) \cdot q_{n,i}(a, b)$$

Define by induction on n probability spaces for sorting
n distinct numbers

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$A_{<} = \{a \in A | a < s\}$ a(i) mutually distinct

$A_{>} = \{a \in A | a > s\}$

$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$

$$QS_n = (Q_n, q_n) \quad RS_{n,i} = QS_{i-1} \times QS_{n-i}$$

and random variables

$$t_n : Q_n \rightarrow \mathbb{R}$$

$x \in Q_n \rightarrow t_n(x) = \text{number of comparisons in run } x$

$$n \in \{0,1\}: \quad Q_n = \{\perp\} \quad q_n(\perp) = 1 \quad t_n(\perp) = 0$$

0 or 1 elements: no randomness
no comparisons

$$n \geq 2: \quad S_n = \{1, \dots, n\} \quad i \in S_n \quad i = \#A_{<} + 1 \quad r_n(i) = 1/n \quad W = (S_n, r_n) \text{ is probability space.}$$

$$R_{n,i} = (Q_{i-1} \times Q_{n-i})$$

Lemma: For all i and n: QS_n and $RS_{n,i}$ are probability spaces

$$q_{n,i}(a, b) = q_{i-1}(a) \cdot q_{n-i}(b)$$

$$Q_n = \bigcup_{i \in S_n} \{i\} \times (Q_{i-1} \times Q_{n-i})$$

proof: induction on n.

• n = 0 or 1: trivial

• assume true for j < n:

$$i - 1 < n, n - i < n$$

Lemma 1: $QS_{i-1} \times QS_{n-i}$ is probability space

$$q_n(i, (a, b)) = (1/n) \cdot q_{n,i}(a, b)$$

Define by induction on n probability spaces for sorting
n distinct numbers

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$A_{<} = \{a \in A | a < s\}$ a(i) mutually distinct

$A_{>} = \{a \in A | a > s\}$

$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$

$$QS_n = (Q_n, q_n) \quad RS_{n,i} = QS_{i-1} \times QS_{n-i}$$

and random variables

$$t_n : Q_n \rightarrow \mathbb{R}$$

$x \in Q_n \rightarrow t_n(x) = \text{number of comparisons in run } x$

$$n \in \{0,1\}: \quad Q_n = \{\perp\} \quad q_n(\perp) = 1 \quad t_n(\perp) = 0$$

0 or 1 elements: no randomness
no comparisons

$$n \geq 2: \quad S_n = \{1, \dots, n\} \quad i \in S_n \quad i = \#A_{<} + 1 \quad r_n(i) = 1/n \quad W = (S_n, r_n) \text{ is probability space.}$$

$$R_{n,i} = (Q_{i-1} \times Q_{n-i})$$

Lemma: For all i and n: QS_n and $RS_{n,i}$ are probability spaces

$$q_{n,i}(a, b) = q_{i-1}(a) \cdot q_{n-i}(b)$$

$$Q_n = \bigcup_{i \in S_n} \{i\} \times (Q_{i-1} \times Q_{n-i})$$

proof: induction on n.

• n = 0 or 1: trivial

• assume true for j < n:

$$i - 1 < n, n - i < n$$

Lemma 1: $QS_{i-1} \times QS_{n-i}$ is probability space

Lemma 10: QS_n is probability space

$$q_n(i, (a, b)) = (1/n) \cdot q_{n,i}(a, b)$$

Define by induction on n probability spaces for sorting
n distinct numbers

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$A_{<} = \{a \in A | a < s\}$ a(i) mutually distinct

$A_{>} = \{a \in A | a > s\}$

$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$

$$QS_n = (Q_n, q_n) \quad RS_{n,i} = QS_{i-1} \times QS_{n-i}$$

and random variables

$$t_n : Q_n \rightarrow \mathbb{R}$$

$x \in Q_n \rightarrow t_n(x) = \text{number of comparisons in run } x$

$$n \in \{0,1\}: \quad Q_n = \{\perp\} \quad q_n(\perp) = 1 \quad t_n(\perp) = 0$$

0 or 1 elements: no randomness
no comparisons

$$n \geq 2: \quad S_n = \{1, \dots, n\} \quad i \in S_n \quad i = \#A_{<} + 1 \quad r_n(i) = 1/n \quad W = (S_n, r_n) \text{ is probability space.}$$

$$R_{n,i} = (Q_{i-1} \times Q_{n-i})$$

Lemma: For all i and n: QS_n and $RS_{n,i}$ are probability spaces

$$q_{n,i}(a, b) = q_{i-1}(a) \cdot q_{n-i}(b)$$

$$Q_n = \bigcup_{i \in S_n} \{i\} \times (Q_{i-1} \times Q_{n-i}) \quad t_n(i, (a, b)) = n - 1 + t_{i-1}(a) + t_{n-i}(b)$$

number of comparisons, random variable on Q_n
defined by induction on n

$$q_n(i, (a, b)) = (1/n) \cdot q_{n,i}(a, b)$$

Define by induction on n probability spaces for sorting
n distinct numbers

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$A_{<} = \{a \in A \mid a < s\}$ a(i) mutually distinct

$A_{>} = \{a \in A \mid a > s\}$

$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$

$$QS_n = (Q_n, q_n) \qquad RS_{n,i} = QS_{i-1} \times QS_{n-i}$$

and random variables

$$t_n : Q_n \rightarrow \mathbb{R}$$

$x \in Q_n \rightarrow t_n(x) = \text{number of comparisons in run } x$

$$n \in \{0,1\}: \quad Q_n = \{\perp\} \quad q_n(\perp) = 1 \quad t_n(\perp) = 0$$

0 or 1 elements: no randomness
no comparisons

$$n \geq 2: \quad S_n = \{1, \dots, n\} \quad i \in S_n \quad i = \#A_{<} + 1 \quad r_n(i) = 1/n \quad W = (S_n, r_n) \text{ is probability space.}$$

$$R_{n,i} = (Q_{i-1} \times Q_{n-i})$$

Lemma: For all i and n: QS_n and $RS_{n,i}$ are probability spaces

$$q_{n,i}(a, b) = q_{i-1}(a) \cdot q_{n-i}(b)$$

$$Q_n = \bigcup_{i \in S_n} \{i\} \times (Q_{i-1} \times Q_{n-i}) \qquad t_n(i, (a, b)) = n - 1 + t_{i-1}(a) + t_{n-i}(b) = n - 1 + t'_{n,i}(a, b)$$

t_{i-1} random variable on Q_{i-1}

$$q_n(i, (a, b)) = (1/n) \cdot q_{n,i}(a, b) \qquad t_{n-i} \text{ random variable on } Q_{n-i}$$

Define by induction on n probability spaces for sorting
n distinct numbers

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$A_{<} = \{a \in A | a < s\}$ a(i) mutually distinct

$A_{>} = \{a \in A | a > s\}$

$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$

$QS_n = (Q_n, q_n)$ $RS_{n,i} = QS_{i-1} \times QS_{n-i}$

and random variables

$t_n : Q_n \rightarrow \mathbb{R}$

$x \in Q_n \rightarrow t_n(x) = \text{number of comparisons in run } x$

$n \in \{0,1\}$: $Q_n = \{\perp\}$ $q_n(\perp) = 1$ $t_n(\perp) = 0$

0 or 1 elements: no randomness
no comparisons

$n \geq 2$: $S_n = \{1, \dots, n\}$ $i \in S_n$ $i = \#A_{<} + 1$ $r_n(i) = 1/n$ $W = (S_n, r_n)$ is probability space.

$R_{n,i} = (Q_{i-1} \times Q_{n-i})$

Lemma: For all i and n: QS_n and $RS_{n,i}$ are probability spaces

$q_{n,i}(a, b) = q_{i-1}(a) \cdot q_{n-i}(b)$

$Q_n = \bigcup_{i \in S_n} \{i\} \times (Q_{i-1} \times Q_{n-i})$ $t_n(i, (a, b)) = n - 1 + t_{i-1}(a) + t_{n-i}(b) = n - 1 + t'_{n,i}(a, b)$

t_{i-1} random variable on Q_{i-1} Lemma 9: $E(t'_{n,i}) = E(t_{i-1}) + E(t_{n-i})$

$q_n(i, (a, b)) = (1/n) \cdot q_{n,i}(a, b)$ t_{n-i} random variable on Q_{n-i}

Define by induction on n probability spaces for sorting
n distinct numbers

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$A_{<} = \{a \in A | a < s\}$ a(i) mutually distinct

$A_{>} = \{a \in A | a > s\}$

$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$

$QS_n = (Q_n, q_n)$ $RS_{n,i} = QS_{i-1} \times QS_{n-i}$

and random variables

$t_n : Q_n \rightarrow \mathbb{R}$

$x \in Q_n \rightarrow t_n(x) = \text{number of comparisons in run } x$

$n \in \{0,1\}$: $Q_n = \{\perp\}$ $q_n(\perp) = 1$ $t_n(\perp) = 0$

0 or 1 elements: no randomness
no comparisons

$n \geq 2$: $S_n = \{1, \dots, n\}$ $i \in S_n$ $i = \#A_{<} + 1$ $r_n(i) = 1/n$ $W = (S_n, r_n)$ is probability space.

$R_{n,i} = (Q_{i-1} \times Q_{n-i})$

Lemma: For all i and n: QS_n and $RS_{n,i}$ are probability spaces

$q_{n,i}(a, b) = q_{i-1}(a) \cdot q_{n-i}(b)$

$Q_n = \bigcup_{i \in S_n} \{i\} \times (Q_{i-1} \times Q_{n-i})$ $t_n(i, (a, b)) = n - 1 + t_{i-1}(a) + t_{n-i}(b) = n - 1 + t'_{n,i}(a, b)$

t_{i-1} random variable on Q_{i-1} Lemma 9: $E(t'_{n,i}) = E(t_{i-1}) + E(t_{n-i})$

$q_n(i, (a, b)) = (1/n) \cdot q_{n,i}(a, b)$ t_{n-i} random variable on Q_{n-i} Lemma 12: $E(t_n) = n - 1 + (1/n) \cdot \sum_{i=1}^n (E(t_{i-1}) + E(t_{n-i}))$

Define by induction on n probability spaces for sorting
n distinct numbers

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$A_{<} = \{a \in A | a < s\}$ a(i) mutually distinct

$A_{>} = \{a \in A | a > s\}$

$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$

$QS_n = (Q_n, q_n)$ $RS_{n,i} = QS_{i-1} \times QS_{n-i}$

and random variables

$t_n : Q_n \rightarrow \mathbb{R}$

$x \in Q_n \rightarrow t_n(x) = \text{number of comparisons in run } x$

$n \in \{0,1\}$: $Q_n = \{\perp\}$ $q_n(\perp) = 1$ $t_n(\perp) = 0$

0 or 1 elements: no randomness
no comparisons

$n \geq 2$: $S_n = \{1, \dots, n\}$ $i \in S_n$ $i = \#A_{<} + 1$ $r_n(i) = 1/n$ $W = (S_n, r_n)$ is probability space.

$R_{n,i} = (Q_{i-1} \times Q_{n-i})$

Lemma: For all i and n: QS_n and $RS_{n,i}$ are probability spaces

$q_{n,i}(a, b) = q_{i-1}(a) \cdot q_{n-i}(b)$

$Q_n = \bigcup_{i \in S_n} \{i\} \times (Q_{i-1} \times Q_{n-i})$ $t_n(i, (a, b)) = n - 1 + t_{i-1}(a) + t_{n-i}(b) = n - 1 + t'_{n,i}(a, b)$

t_{i-1} random variable on Q_{i-1}

Lemma 9: $E(t'_{n,i}) = E(t_{i-1}) + E(t_{n-i})$

$q_n(i, (a, b)) = (1/n) \cdot q_{n,i}(a, b)$ t_{n-i} random variable on Q_{n-i}

Lemma 12: $E(t_n) = n - 1 + (1/n) \cdot \sum_{i=1}^n (E(t_{i-1}) + E(t_{n-i}))$

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$$A_{<} = \{a \in A \mid a < s\}$$

a(i) mutually distinct

$$A_{>} = \{a \in A \mid a > s\}$$

$$\text{sort}(A) = \text{sort}(A_{<}) \circ s \circ \text{sort}(A_{>})$$

$$T(n) = E(t_n): \quad T(n) = n - 1 + (1/n) \cdot \sum_{i=1}^n (T(i-1) + T(n-i)) \quad \text{back to the sixties!!!}$$

Lemma: $T(n) \leq 2n \cdot \ln(n)$

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$$A_{<} = \{a \in A \mid a < s\}$$

a(i) mutually distinct

$$A_{>} = \{a \in A \mid a > s\}$$

$$\text{sort}(A) = \text{sort}(A_{<}) \circ s \circ \text{sort}(A_{>})$$

$$T(n) = E(t_n): \quad T(n) = n - 1 + (1/n) \cdot \sum_{i=1}^n (T(i-1) + T(n-i)) \quad \text{back to the sixties!!!}$$

Lemma: $T(n) \leq 2n \cdot \ln(n)$

proof: induction on n

• $n = 1$:

$$T(1) = 0 = 2 \cdot 1 \ln(1)$$

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$A_{<} = \{a \in A \mid a < s\}$ a(i) mutually distinct

$A_{>} = \{a \in A \mid a > s\}$

$sort(A) = sort(A_{<}) \circ s \circ sort(A_{>})$

$T(n) = E(t_n): \quad T(n) = n - 1 + (1/n) \cdot \sum_{i=1}^n (T(i - 1) + T(n - i))$

Lemma: $T(n) \leq 2n \cdot \ln(n)$

proof: induction on n

- $n = 1:$ $T(1) = 0 = 2 \cdot 1 \ln(1)$
- $n > 1:$

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$$A_{<} = \{a \in A \mid a < s\} \quad \text{a(i) mutually distinct}$$

$$A_{>} = \{a \in A \mid a > s\}$$

$$\text{sort}(A) = \text{sort}(A_{<}) \circ s \circ \text{sort}(A_{>})$$

$$\begin{aligned} T(n) &< n + \frac{1}{n} \cdot \sum_{i=1}^n (T(i-1) + T(n-i)) \\ &= n + \frac{1}{n} \cdot \left(\sum_{i=0}^{n-1} T(i) + \sum_{i=0}^{n-1} T(i) \right) \\ &= n + \frac{2}{n} \cdot \left(\sum_{i=0}^{n-1} T(i) \right) \\ &= n + \frac{2}{n} \cdot \left(\sum_{i=1}^{n-1} T(i) \right) \quad (T(0) = T(1) = 0) \\ &\leq n + \frac{2}{n} \cdot \left(\sum_{i=2}^{n-1} 2i \cdot \ln(i) \right) \quad (\text{induction hypothesis}) \\ &\leq n + \frac{2}{n} \cdot \left(\int_2^n 2x \cdot \ln(x) dx \right) \quad (\text{area under the curve}) \end{aligned}$$

$$T(n) = E(t_n): \quad T(n) = n - 1 + (1/n) \cdot \sum_{i=1}^n (T(i-1) + T(n-i))$$

Lemma: $T(n) \leq 2n \cdot \ln(n)$

proof: induction on n

- $n = 1$: $T(1) = 0 = 2 \cdot 1 \ln(1)$
- $n > 1$:

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$$A_{<} = \{a \in A \mid a < s\} \quad \text{a(i) mutually distinct}$$

$$A_{>} = \{a \in A \mid a > s\}$$

$$\text{sort}(A) = \text{sort}(A_{<}) \circ s \circ \text{sort}(A_{>})$$


$$T(n) = E(t_n): \quad T(n) = n - 1 + (1/n) \cdot \sum_{i=1}^n (T(i-1) + T(n-i))$$

$$\text{Lemma: } T(n) \leq 2n \cdot \ln(n)$$

proof: induction on n

- $n = 1$: $T(1) = 0 = 2 \cdot 1 \ln(1)$
- $n > 1$:

$$\begin{aligned} T(n) &< n + \frac{1}{n} \cdot \sum_{i=1}^n (T(i-1) + T(n-i)) \\ &= n + \frac{1}{n} \cdot \left(\sum_{i=0}^{n-1} T(i) + \sum_{i=0}^{n-1} T(i) \right) \\ &= n + \frac{2}{n} \cdot \left(\sum_{i=0}^{n-1} T(i) \right) \\ &= n + \frac{2}{n} \cdot \left(\sum_{i=1}^{n-1} T(i) \right) \quad (T(0) = T(1) = 0) \\ &\leq n + \frac{2}{n} \cdot \left(\sum_{i=2}^{n-1} 2i \cdot \ln(i) \right) \quad (\text{induction hypothesis}) \\ &\leq n + \frac{2}{n} \cdot \left(\int_2^n 2x \cdot \ln(x) dx \right) \quad (\text{area under the curve}) \end{aligned}$$



$$\leq n + \frac{2}{n} \cdot \left(\int_2^n 2x \cdot \ln(x) dx \right) \quad (\text{area under the curve})$$

Product rule of differentiation

$$(uv)' = u'v + uv'$$

Integrate

$$\int u'v = uv - \int uv'$$

$$u' = 2x$$

$$v = \ln(x)$$

$$u = x^2$$

$$v' = 1/x$$

$$\int u'v = x^2 \ln(x) - \int x^2/x$$

$$= x^2 \cdot \ln(x) - \frac{x^2}{2}$$

input: $(a(1), \dots, a(n))$ or set $A = \{a(1), \dots, a(n)\}$

random experiment: choose 'splitter' $s \in \{a(1), \dots, a(n)\}$

all n splitters equally likely

$$A_{<} = \{a \in A \mid a < s\} \quad \text{a(i) mutually distinct}$$

$$A_{>} = \{a \in A \mid a > s\}$$

$$\text{sort}(A) = \text{sort}(A_{<}) \circ s \circ \text{sort}(A_{>})$$


$$T(n) = E(t_n): \quad T(n) = n - 1 + (1/n) \cdot \sum_{i=1}^n (T(i-1) + T(n-i))$$

Lemma: $T(n) \leq 2n \cdot \ln(n)$

proof: induction on n

- $n = 1$: $T(1) = 0 = 2 \cdot 1 \ln(1)$
- $n > 1$:

$$\begin{aligned} T(n) &< n + \frac{1}{n} \cdot \sum_{i=1}^n (T(i-1) + T(n-i)) \\ &= n + \frac{1}{n} \cdot \left(\sum_{i=0}^{n-1} T(i) + \sum_{i=0}^{n-1} T(i) \right) \\ &= n + \frac{2}{n} \cdot \left(\sum_{i=0}^{n-1} T(i) \right) \\ &= n + \frac{2}{n} \cdot \left(\sum_{i=1}^{n-1} T(i) \right) \quad (T(0) = T(1) = 0) \\ &\leq n + \frac{2}{n} \cdot \left(\sum_{i=2}^{n-1} 2i \cdot \ln(i) \right) \quad (\text{induction hypothesis}) \\ &\leq n + \frac{2}{n} \cdot \left(\int_2^n 2x \cdot \ln(x) dx \right) \quad (\text{area under the curve}) \end{aligned}$$



$$\begin{aligned} &\leq n + \frac{2}{n} \cdot \left(\int_2^n 2x \cdot \ln(x) dx \right) \quad (\text{area under the curve}) \\ &= n + \frac{2}{n} \cdot \left[(n^2 \ln(n) - \frac{n^2}{2}) - (2^2 \ln 2 - 2^2/2) \right] \quad (\text{product rule}) \\ &< 2n \cdot \ln(n) \quad (\ln 2 > 0.69 > 1/2) \end{aligned}$$