

# Numerical Linear Algebra

Ramaz Botchorishvili

Kutaisi International University

November 15, 2023

# Numerical Linear Algebra

Ramaz Botchorishvili

Kutaisi International University

November 15, 2023

# Direct and iterative methods for linear systems

- ▶ Recap of Previous Lecture
- ▶ Sherman-Morrison formula
- ▶ Cholesky factorization of symmetric system
- ▶ Diagonally dominant system
- ▶ Necessary and sufficient conditions for convergence
- ▶ Classic iterative methods
- ▶ Convergence theorems of J,GS
- ▶ Q & A

# Recap of Previous Lecture

- ▶ Examples of LU factorization
- ▶ Existence of LU factorization
- ▶ Gaussian elimination(GE) and LU factorization
- ▶ Pivoting
- ▶ Stability of GE

# Shermann-Morrison formula

## Problem 9.1

Suppose  $A$  is square matrix and  $A^{-1}$  is known.

# Shermann-Morrison formula

## Problem 9.1

Suppose  $A$  is square matrix and  $A^{-1}$  is known.

- Compute inverse of  $A + B$  by means of using  $A^{-1}$ ,  $B$  is square matrix.

# Shermann-Morrison formula

## Problem 9.1

Suppose  $A$  is square matrix and  $A^{-1}$  is known.

- ▶ Compute inverse of  $A + B$  by means of using  $A^{-1}$ ,  $B$  is square matrix.
- ▶ For which  $B$  is this possible?

# Shermann-Morrison formula

## Problem 9.1

Suppose  $A$  is square matrix and  $A^{-1}$  is known.

- ▶ Compute inverse of  $A + B$  by means of using  $A^{-1}$ ,  $B$  is square matrix.
- ▶ For which  $B$  is this possible?

## Theorem 9.2

(*Shermann-Morrison formula*)

- ▶  $A \in \mathbb{R}^{n \times n}$ ,  $u, v \in \mathbb{R}^n$



# Shermann-Morrison formula

## Problem 9.1

Suppose  $A$  is square matrix and  $A^{-1}$  is known.

- ▶ Compute inverse of  $A + B$  by means of using  $A^{-1}$ ,  $B$  is square matrix.
- ▶ For which  $B$  is this possible?

## Theorem 9.2

(*Shermann-Morrison formula*)

- ▶  $A \in \mathbb{R}^{n \times n}$ ,  $u, v \in \mathbb{R}^n$
- ▶  $\lambda \equiv v^T A^{-1} u \neq -1$

# Shermann-Morrison formula

## Problem 9.1

Suppose  $A$  is square matrix and  $A^{-1}$  is known.

- ▶ Compute inverse of  $A + B$  by means of using  $A^{-1}$ ,  $B$  is square matrix.
- ▶ For which  $B$  is this possible?

## Theorem 9.2

(*Shermann-Morrison formula*)

- ▶  $A \in \mathbb{R}^{n \times n}$ ,  $u, v \in \mathbb{R}^n$
- ▶  $\lambda \equiv v^T A^{-1} u \neq -1$
- ▶  $\alpha = \frac{1}{1+\lambda}$

# Shermann-Morrison formula

## Problem 9.1

Suppose  $A$  is square matrix and  $A^{-1}$  is known.

- ▶ Compute inverse of  $A + B$  by means of using  $A^{-1}$ ,  $B$  is square matrix.
- ▶ For which  $B$  is this possible?

## Theorem 9.2

(*Shermann-Morrison formula*)

- ▶  $A \in \mathbb{R}^{n \times n}$ ,  $u, v \in \mathbb{R}^n$
- ▶  $\lambda \equiv v^T A^{-1} u \neq -1$
- ▶  $\alpha = \frac{1}{1+\lambda}$
- ▶  $(A + uv^T)^{-1} = A^{-1} - \alpha(A^{-1}uv^T A^{-1})$

# Shermann-Morrison formula

## Problem 9.1

Suppose  $A$  is square matrix and  $A^{-1}$  is known.

- ▶ Compute inverse of  $A + B$  by means of using  $A^{-1}$ ,  $B$  is square matrix.
- ▶ For which  $B$  is this possible?

## Theorem 9.2

(*Shermann-Morrison formula*)

- ▶  $A \in \mathbb{R}^{n \times n}$ ,  $u, v \in \mathbb{R}^n$
- ▶  $\lambda \equiv v^T A^{-1} u \neq -1$
- ▶  $\alpha = \frac{1}{1+\lambda}$
- ▶  $(A + uv^T)^{-1} = A^{-1} - \alpha(A^{-1}uv^T A^{-1})$

Generalization: Sherman-Morrison-Woodbury

# Shermann-Morrison formula

## Theorem 9.3

- ▶  $A \in \mathbb{R}^{n \times n}$ ,  $u, v \in \mathbb{R}^n$
- ▶  $\lambda \equiv v^T A^{-1} u \neq -1$
- ▶  $\alpha = \frac{1}{1+\lambda}$
- ▶  $(A + uv^T)^{-1} = A^{-1} - \alpha(A^{-1}uv^T A^{-1})$

Proof.

# Shermann-Morrison formula

## Theorem 9.3

- ▶  $A \in \mathbb{R}^{n \times n}$ ,  $u, v \in \mathbb{R}^n$
- ▶  $\lambda \equiv v^T A^{-1} u \neq -1$
- ▶  $\alpha = \frac{1}{1+\lambda}$
- ▶  $(A + uv^T)^{-1} = A^{-1} - \alpha(A^{-1}uv^T A^{-1})$

## Proof.

$$(A + uv^T)(A^{-1} - \alpha(A^{-1}uv^T A^{-1})) =$$

# Shermann-Morrison formula

## Theorem 9.3

- ▶  $A \in \mathbb{R}^{n \times n}$ ,  $u, v \in \mathbb{R}^n$
- ▶  $\lambda \equiv v^T A^{-1} u \neq -1$
- ▶  $\alpha = \frac{1}{1+\lambda}$
- ▶  $(A + uv^T)^{-1} = A^{-1} - \alpha(A^{-1}uv^T A^{-1})$

## Proof.

$$\begin{aligned} & (A + uv^T)(A^{-1} - \alpha(A^{-1}uv^T A^{-1})) = \\ & = AA^{-1} + uv^T A^{-1} - \alpha AA^{-1}uv^T A^{-1} - \alpha uv^T A^{-1}uv^T A^{-1} = \end{aligned}$$

# Shermann-Morrison formula

## Theorem 9.3

- ▶  $A \in \mathbb{R}^{n \times n}$ ,  $u, v \in \mathbb{R}^n$
- ▶  $\lambda \equiv v^T A^{-1} u \neq -1$
- ▶  $\alpha = \frac{1}{1+\lambda}$
- ▶  $(A + uv^T)^{-1} = A^{-1} - \alpha(A^{-1}uv^T A^{-1})$

## Proof.

$$\begin{aligned} & (A + uv^T)(A^{-1} - \alpha(A^{-1}uv^T A^{-1})) = \\ &= AA^{-1} + uv^T A^{-1} - \alpha AA^{-1} uv^T A^{-1} - \alpha uv^T A^{-1} uv^T A^{-1} = \\ &= I + uv^T A^{-1} - \alpha uv^T A^{-1} - \alpha u \lambda v^T A^{-1} = \end{aligned}$$



# Shermann-Morrison formula

## Theorem 9.3

- ▶  $A \in \mathbb{R}^{n \times n}$ ,  $u, v \in \mathbb{R}^n$
- ▶  $\lambda \equiv v^T A^{-1} u \neq -1$
- ▶  $\alpha = \frac{1}{1+\lambda}$
- ▶  $(A + uv^T)^{-1} = A^{-1} - \alpha(A^{-1}uv^T A^{-1})$

## Proof.

$$\begin{aligned} & (A + uv^T)(A^{-1} - \alpha(A^{-1}uv^T A^{-1})) = \\ &= AA^{-1} + uv^T A^{-1} - \alpha AA^{-1} uv^T A^{-1} - \alpha uv^T A^{-1} uv^T A^{-1} = \\ &= I + uv^T A^{-1} - \alpha uv^T A^{-1} - \alpha u \lambda v^T A^{-1} = \\ &= I + uv^T A^{-1} - \alpha(1 + \lambda) uv^T A^{-1} = \end{aligned}$$

# Shermann-Morrison formula

## Theorem 9.3

- ▶  $A \in \mathbb{R}^{n \times n}$ ,  $u, v \in \mathbb{R}^n$
- ▶  $\lambda \equiv v^T A^{-1} u \neq -1$
- ▶  $\alpha = \frac{1}{1+\lambda}$
- ▶  $(A + uv^T)^{-1} = A^{-1} - \alpha(A^{-1}uv^T A^{-1})$

## Proof.

$$\begin{aligned} & (A + uv^T)(A^{-1} - \alpha(A^{-1}uv^T A^{-1})) = \\ &= AA^{-1} + uv^T A^{-1} - \alpha AA^{-1} uv^T A^{-1} - \alpha uv^T A^{-1} uv^T A^{-1} = \\ &= I + uv^T A^{-1} - \alpha uv^T A^{-1} - \alpha u \lambda v^T A^{-1} = \\ &= I + uv^T A^{-1} - \alpha(1 + \lambda) uv^T A^{-1} = \\ &= I + uv^T A^{-1} - 1 \cdot uv^T A^{-1} = \end{aligned}$$

# Shermann-Morrison formula

## Theorem 9.3

- ▶  $A \in \mathbb{R}^{n \times n}$ ,  $u, v \in \mathbb{R}^n$
- ▶  $\lambda \equiv v^T A^{-1} u \neq -1$
- ▶  $\alpha = \frac{1}{1+\lambda}$
- ▶  $(A + uv^T)^{-1} = A^{-1} - \alpha(A^{-1}uv^T A^{-1})$

## Proof.

$$\begin{aligned} & (A + uv^T)(A^{-1} - \alpha(A^{-1}uv^T A^{-1})) = \\ &= AA^{-1} + uv^T A^{-1} - \alpha AA^{-1} uv^T A^{-1} - \alpha uv^T A^{-1} uv^T A^{-1} = \\ &= I + uv^T A^{-1} - \alpha uv^T A^{-1} - \alpha u \lambda v^T A^{-1} = \\ &= I + uv^T A^{-1} - \alpha(1 + \lambda) uv^T A^{-1} = \\ &= I + uv^T A^{-1} - 1 \cdot uv^T A^{-1} = \\ &= I \end{aligned}$$

# Sherman-Morrison-Woodbury formula

## Theorem 9.4

*(Sherman-Morrison-Woodbury formula)*

►  $A \in \mathbb{R}^{n \times n}, U \in \mathbb{R}^{n \times k}, C \in \mathbb{R}^{k \times k}, V \in \mathbb{R}^{k \times n},$

# Sherman-Morrison-Woodbury formula

## Theorem 9.4

*(Sherman-Morrison-Woodbury formula)*

- ▶  $A \in \mathbb{R}^{n \times n}$ ,  $U \in \mathbb{R}^{n \times k}$ ,  $C \in \mathbb{R}^{k \times k}$ ,  $V \in \mathbb{R}^{k \times n}$ ,
- ▶  $(A + UCV)^{-1} = A^{-1} - A^{-1}U(C^{-1} + VA^{-1}U)VA^{-1}$

# Sherman-Morrison-Woodbury formula

## Theorem 9.4

(*Sherman-Morrison-Woodbury formula*)

- ▶  $A \in \mathbb{R}^{n \times n}$ ,  $U \in \mathbb{R}^{n \times k}$ ,  $C \in \mathbb{R}^{k \times k}$ ,  $V \in \mathbb{R}^{k \times n}$ ,
- ▶  $(A + UCV)^{-1} = A^{-1} - A^{-1}U(C^{-1} + VA^{-1}U)VA^{-1}$

Proof.

Direct proof similar to Sherman-Morrison formula



# Special matrix: diagonal dominance

## Definition 9.5

- Strictly row diagonally dominant matrix

$$|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}|, i = 1, 2, \dots, n$$

# Special matrix: diagonal dominance

## Definition 9.5

- ▶ Strictly row diagonally dominant matrix

$$|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}|, i = 1, 2, \dots, n$$

- ▶ Strictly column diagonally dominant matrix

$$|a_{jj}| > \sum_{i=1, i \neq j}^n |a_{ij}|, j = 1, 2, \dots, n$$



# Special matrix: diagonal dominance

## Definition 9.5

- ▶ Strictly row diagonally dominant matrix

$$|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}|, i = 1, 2, \dots, n$$

- ▶ Strictly column diagonally dominant matrix

$$|a_{jj}| > \sum_{i=1, i \neq j}^n |a_{ij}|, j = 1, 2, \dots, n$$

- ▶ Row diagonally dominant matrix

$$|a_{ii}| \geq \sum_{j=1, j \neq i}^n |a_{ij}|, i = 1, 2, \dots, n$$

# Special matrix: diagonal dominance

## Definition 9.5

- ▶ Strictly row diagonally dominant matrix

$$|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}|, i = 1, 2, \dots, n$$

- ▶ Strictly column diagonally dominant matrix

$$|a_{jj}| > \sum_{i=1, i \neq j}^n |a_{ij}|, j = 1, 2, \dots, n$$

- ▶ Row diagonally dominant matrix

$$|a_{ii}| \geq \sum_{j=1, j \neq i}^n |a_{ij}|, i = 1, 2, \dots, n$$

- ▶ Column diagonally dominant matrix

$$|a_{jj}| \geq \sum_{i=1, i \neq j}^n |a_{ij}|, j = 1, 2, \dots, n$$

# Special matrix: diagonal dominance

## Definition 9.5

- ▶ Strictly row diagonally dominant matrix

$$|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}|, i = 1, 2, \dots, n$$

- ▶ Strictly column diagonally dominant matrix

$$|a_{jj}| > \sum_{i=1, i \neq j}^n |a_{ij}|, j = 1, 2, \dots, n$$

- ▶ Row diagonally dominant matrix

$$|a_{ii}| \geq \sum_{j=1, j \neq i}^n |a_{ij}|, i = 1, 2, \dots, n$$

- ▶ Column diagonally dominant matrix

$$|a_{jj}| \geq \sum_{i=1, i \neq j}^n |a_{ij}|, j = 1, 2, \dots, n$$

## Proposition 9.6

*Suppose  $A \in \mathbb{R}^{n \times n}$  is column diagonally dominant. Then Gaussian elimination algorithm coincides with Gaussian elimination algorithm with partial pivoting*

# Cholesky decomposition

## Problem 9.7

►  $A \in \mathbb{R}^{n \times n}, A = A^T, A \geq 0$

# Cholesky decomposition

## Problem 9.7

- ▶  $A \in \mathbb{R}^{n \times n}, A = A^T, A \geq 0$
- ▶ Find lower triangular matrix  $H$  such that  $A = HH^T$

# Cholesky decomposition

## Problem 9.7

- ▶  $A \in \mathbb{R}^{n \times n}, A = A^T, A \geq 0$
- ▶ Find lower triangular matrix  $H$  such that  $A = HH^T$

## Theorem 9.8

*Unique Cholesky decomposition exists*

# Cholesky decomposition

## Problem 9.7

- ▶  $A \in \mathbb{R}^{n \times n}, A = A^T, A \geq 0$
- ▶ Find lower triangular matrix  $H$  such that  $A = HH^T$

## Theorem 9.8

*Unique Cholesky decomposition exists*

## Proof.

- ▶ Similar to general  $LU$  decomposition theorem

# Cholesky decomposition

## Problem 9.7

- ▶  $A \in \mathbb{R}^{n \times n}, A = A^T, A \geq 0$
- ▶ Find lower triangular matrix  $H$  such that  $A = HH^T$

## Theorem 9.8

*Unique Cholesky decomposition exists*

## Proof.

- ▶ Similar to general  $LU$  decomposition theorem
- ▶ Apply mathematical induction



# Cholesky decomposition

## Problem 9.7

- ▶  $A \in \mathbb{R}^{n \times n}, A = A^T, A \geq 0$
- ▶ Find lower triangular matrix  $H$  such that  $A = HH^T$

## Theorem 9.8

*Unique Cholesky decomposition exists*

## Proof.

- ▶ Similar to general  $LU$  decomposition theorem
- ▶ Apply mathematical induction
- ▶ Apply decomposition of main sub matrices

# Cholesky decomposition

## Problem 9.7

- ▶  $A \in \mathbb{R}^{n \times n}$ ,  $A = A^T$ ,  $A \geq 0$
- ▶ Find lower triangular matrix  $H$  such that  $A = HH^T$

## Theorem 9.8

*Unique Cholesky decomposition exists*

## Proof.

- ▶ Similar to general  $LU$  decomposition theorem
- ▶ Apply mathematical induction
- ▶ Apply decomposition of main sub matrices

$$A_i = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1i-1} & a_{1i} \\ a_{21} & a_{22} & \cdots & a_{2i-1} & a_{2i} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ a_{i-11} & a_{i-12} & \cdots & a_{i-1i-1} & a_{i-1i} \\ a_{i1} & a_{i2} & \cdots & a_{ii-1} & a_{ii} \end{pmatrix}$$

Diagram illustrating the recursive Cholesky decomposition of a matrix  $A_i$ . The matrix is partitioned into a top-left block  $A_{i-1}$ , a top-right vector  $v$ , a bottom-left vector  $v^T$ , and a bottom-right scalar  $\alpha$ . Green arrows indicate the flow of information from these components into the full matrix  $A_i$ .

# Convergence of iterative methods 1

Iterative method

# Convergence of iterative methods 1

## Iterative method

$$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$$

# Convergence of iterative methods 1

## Iterative method

$$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$$

►  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$

# Convergence of iterative methods 1

## Iterative method

$$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$$

- ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
- ▶  $x^{(0)}$  - initial guess

# Convergence of iterative methods 1

## Iterative method

$$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$$

- ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
- ▶  $x^{(0)}$  - initial guess
- ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence

# Convergence of iterative methods 1

## Iterative method

$$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$$

- ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
- ▶  $x^{(0)}$  - initial guess
- ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence

## Convergence of iterative method



# Convergence of iterative methods 1

## Iterative method

$$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$$

- ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
- ▶  $x^{(0)}$  - initial guess
- ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence

## Convergence of iterative method

Suppose  $\{x_k\}_{k=0}^{\infty}$  converges to  $x$

# Convergence of iterative methods 1

## Iterative method

$$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$$

- ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
- ▶  $x^{(0)}$  - initial guess
- ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence

## Convergence of iterative method

Suppose  $\{x_k\}_{k=0}^{\infty}$  converges to  $x$

- ▶ Does  $x$  satisfy any equation?

# Convergence of iterative methods 1

## Iterative method

$$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$$

- ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
- ▶  $x^{(0)}$  - initial guess
- ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence

## Convergence of iterative method

Suppose  $\{x_k\}_{k=0}^{\infty}$  converges to  $x$

- ▶ Does  $x$  satisfy any equation?

$$x = Bx + f$$

# Convergence of iterative methods 1

## Iterative method

$$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$$

- ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
- ▶  $x^{(0)}$  - initial guess
- ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence

## Convergence of iterative method

Suppose  $\{x_k\}_{k=0}^{\infty}$  converges to  $x$

- ▶ Does  $x$  satisfy any equation?

$$x = Bx + f$$

- ▶ Can  $x$  be solution to  $Ax = b$ ?

# Convergence of iterative methods 1

## Iterative method

$$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$$

- ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
- ▶  $x^{(0)}$  - initial guess
- ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence

## Convergence of iterative method

Suppose  $\{x_k\}_{k=0}^{\infty}$  converges to  $x$

- ▶ Does  $x$  satisfy any equation?

$$x = Bx + f$$

- ▶ Can  $x$  be solution to  $Ax = b$ ?

## Convergence of iterative methods 2

### Convergence of iterative method

Suppose  $\{x_k\}_{k=0}^{\infty}$  converges to  $x$

- ▶ Does  $x$  satisfy any equation?

$$x = Bx + f$$

- ▶ Can  $x$  be solution to  $Ax = b$ ?

# Convergence of iterative methods 2

## Convergence of iterative method

Suppose  $\{x_k\}_{k=0}^{\infty}$  converges to  $x$

- ▶ Does  $x$  satisfy any equation?

$$x = Bx + f$$

- ▶ Can  $x$  be solution to  $Ax = b$ ?

## Definition 9.9

Iterative method  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, \dots$  is compatible with  $Ax = b$  if the latter is equivalent to  $x = Bx + f$

# Convergence of iterative methods 2

## Convergence of iterative method

Suppose  $\{x_k\}_{k=0}^{\infty}$  converges to  $x$

- ▶ Does  $x$  satisfy any equation?

$$x = Bx + f$$

- ▶ Can  $x$  be solution to  $Ax = b$ ?

## Definition 9.9

Iterative method  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, \dots$  is compatible with  $Ax = b$  if the latter is equivalent to  $x = Bx + f$

## Example 9.10

Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b$$



# Convergence of iterative methods 3

- ▶  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$ 
  - ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
  - ▶  $x^{(0)}$  - initial guess
  - ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence

## Convergence of iterative methods 3

- ▶  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$ 
  - ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
  - ▶  $x^{(0)}$  - initial guess
  - ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence
- ▶ Different initial guess defines different sequences  $\{x_k\}_{k=0}^{\infty}$ . Is it possible that for one initial guess method converges and for another does not?

# Convergence of iterative methods 3

- ▶  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$ 
  - ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
  - ▶  $x^{(0)}$  - initial guess
  - ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence
- ▶ Different initial guess defines different sequences  $\{x_k\}_{k=0}^{\infty}$ . Is it possible that for one initial guess method converges and for another does not?

## Definition 9.11

Iterative method  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$  is convergent if it converges to the same  $x$  for any initial guess.

# Convergence of iterative methods 3

- ▶  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$ 
  - ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
  - ▶  $x^{(0)}$  - initial guess
  - ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence
- ▶ Different initial guess defines different sequences  $\{x_k\}_{k=0}^{\infty}$ . Is it possible that for one initial guess method converges and for another does not?

## Definition 9.11

Iterative method  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$  is convergent if it converges to the same  $x$  for any initial guess.

- ▶ Which parameters are important for convergence of ?

# Convergence of iterative methods 3

- ▶  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$ 
  - ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
  - ▶  $x^{(0)}$  - initial guess
  - ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence
- ▶ Different initial guess defines different sequences  $\{x_k\}_{k=0}^{\infty}$ . Is it possible that for one initial guess method converges and for another does not?

## Definition 9.11

Iterative method  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$  is convergent if it converges to the same  $x$  for any initial guess.

- ▶ Which parameters are important for convergence of ?

## Theorem 9.12

*Sufficient condition of convergence:*  $\|B\| < 1$

## Convergence of iterative methods 4

- ▶  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$ 
  - ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
  - ▶  $x^{(0)}$  - initial guess
  - ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence

### Theorem 9.13

*Sufficient condition of convergence:  $\|B\| < 1$*

## Convergence of iterative methods 4

- ▶  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$ 
  - ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
  - ▶  $x^{(0)}$  - initial guess
  - ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence

### Theorem 9.13

*Sufficient condition of convergence:  $\|B\| < 1$*

### Proof.

- ▶ error -  $e^{(k)} = x^{(k)} - x$

## Convergence of iterative methods 4

- ▶  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$ 
  - ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
  - ▶  $x^{(0)}$  - initial guess
  - ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence

### Theorem 9.13

*Sufficient condition of convergence:  $\|B\| < 1$*

### Proof.

- ▶ error -  $e^{(k)} = x^{(k)} - x$
- ▶  $x^{(k+1)} = Bx^{(k)} + f, x = Bx + f \Rightarrow x^{(k+1)} - x = B(x^{(k)} - x)$



## Convergence of iterative methods 4

- ▶  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$ 
  - ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
  - ▶  $x^{(0)}$  - initial guess
  - ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence

### Theorem 9.13

*Sufficient condition of convergence:  $\|B\| < 1$*

### Proof.

- ▶ error -  $e^{(k)} = x^{(k)} - x$
- ▶  $x^{(k+1)} = Bx^{(k)} + f, x = Bx + f \Rightarrow x^{(k+1)} - x = B(x^{(k)} - x)$
- ▶  $e^{(k+1)} = Be^{(k)}$

## Convergence of iterative methods 4

- ▶  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$ 
  - ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
  - ▶  $x^{(0)}$  - initial guess
  - ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence

### Theorem 9.13

*Sufficient condition of convergence:  $\|B\| < 1$*

### Proof.

- ▶ error -  $e^{(k)} = x^{(k)} - x$
- ▶  $x^{(k+1)} = Bx^{(k)} + f, x = Bx + f \Rightarrow x^{(k+1)} - x = B(x^{(k)} - x)$
- ▶  $e^{(k+1)} = Be^{(k)}$
- ▶  $e^{(k)} = Be^{(k-1)}, e^{(k+1)} = Be^{(k)} = BBe^{(k-1)} = B^2e^{(k-1)}$

## Convergence of iterative methods 4

- ▶  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$ 
  - ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
  - ▶  $x^{(0)}$  - initial guess
  - ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence

### Theorem 9.13

*Sufficient condition of convergence:  $\|B\| < 1$*

### Proof.

- ▶ error -  $e^{(k)} = x^{(k)} - x$
- ▶  $x^{(k+1)} = Bx^{(k)} + f, x = Bx + f \Rightarrow x^{(k+1)} - x = B(x^{(k)} - x)$
- ▶  $e^{(k+1)} = Be^{(k)}$
- ▶  $e^{(k)} = Be^{(k-1)}, e^{(k+1)} = Be^{(k)} = BBe^{(k-1)} = B^2e^{(k-1)}$
- ▶ ...

## Convergence of iterative methods 4

- ▶  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$ 
  - ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
  - ▶  $x^{(0)}$  - initial guess
  - ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence

### Theorem 9.13

*Sufficient condition of convergence:  $\|B\| < 1$*

### Proof.

- ▶ error -  $e^{(k)} = x^{(k)} - x$
- ▶  $x^{(k+1)} = Bx^{(k)} + f, x = Bx + f \Rightarrow x^{(k+1)} - x = B(x^{(k)} - x)$
- ▶  $e^{(k+1)} = Be^{(k)}$
- ▶  $e^{(k)} = Be^{(k-1)}, e^{(k+1)} = Be^{(k)} = BB e^{(k-1)} = B^2 e^{(k-1)}$
- ▶ ...
- ▶  $e^{(k+1)} = B^{k+1} e^{(0)}$

## Convergence of iterative methods 4

- ▶  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$ 
  - ▶  $B \in \mathbb{R}^{n \times n}, x^{(k)} \in \mathbb{R}^n$
  - ▶  $x^{(0)}$  - initial guess
  - ▶  $x^{(1)}, x^{(2)}, ..., x^{(k)}, ..$  - infinite sequence

### Theorem 9.13

*Sufficient condition of convergence:  $\|B\| < 1$*

### Proof.

- ▶ error -  $e^{(k)} = x^{(k)} - x$
- ▶  $x^{(k+1)} = Bx^{(k)} + f, x = Bx + f \Rightarrow x^{(k+1)} - x = B(x^{(k)} - x)$
- ▶  $e^{(k+1)} = Be^{(k)}$
- ▶  $e^{(k)} = Be^{(k-1)}, e^{(k+1)} = Be^{(k)} = BB e^{(k-1)} = B^2 e^{(k-1)}$
- ▶ ...
- ▶  $e^{(k+1)} = B^{k+1} e^{(0)}$
- ▶  $\|e^{(k+1)}\| = \|B^{k+1} e^{(0)}\| \leq \|B^{k+1}\| \|e^{(0)}\| \leq \|B\|^{k+1} \|e^{(0)}\| \xrightarrow{k \rightarrow \infty} 0$

# Convergence of iterative methods 5

## Theorem 9.14

$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B) < 1$

## Convergence of iterative methods 5

### Theorem 9.14

$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B) < 1$

### Proof.

Necessary condition: assume convergence and prove  $\rho(B) < 1$

# Convergence of iterative methods 5

## Theorem 9.14

$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B) < 1$

## Proof.

Necessary condition: assume convergence and prove  $\rho(B) < 1$

► Proof by contradiction: assume  $By = \lambda y, |\lambda| > 1$



# Convergence of iterative methods 5

## Theorem 9.14

$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B) < 1$

## Proof.

Necessary condition: assume convergence and prove  $\rho(B) < 1$

- ▶ Proof by contradiction: assume  $By = \lambda y, |\lambda| > 1$
- ▶  $x^{(0)} = x + y, x = Bx + f$

# Convergence of iterative methods 5

## Theorem 9.14

$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B) < 1$

## Proof.

Necessary condition: assume convergence and prove  $\rho(B) < 1$

- ▶ Proof by contradiction: assume  $By = \lambda y, |\lambda| > 1$
- ▶  $x^{(0)} = x + y, x = Bx + f$
- ▶  $e^{(0)} = x^{(0)} - x = y$

# Convergence of iterative methods 5

## Theorem 9.14

$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B) < 1$

## Proof.

Necessary condition: assume convergence and prove  $\rho(B) < 1$

- ▶ Proof by contradiction: assume  $By = \lambda y, |\lambda| > 1$
- ▶  $x^{(0)} = x + y, x = Bx + f$
- ▶  $e^{(0)} = x^{(0)} - x = y$
- ▶  $e^{(1)} = Be^{(0)} = By = \lambda y$

# Convergence of iterative methods 5

## Theorem 9.14

$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B) < 1$

## Proof.

Necessary condition: assume convergence and prove  $\rho(B) < 1$

- ▶ Proof by contradiction: assume  $By = \lambda y, |\lambda| > 1$
- ▶  $x^{(0)} = x + y, x = Bx + f$
- ▶  $e^{(0)} = x^{(0)} - x = y$
- ▶  $e^{(1)} = Be^{(0)} = By = \lambda y$
- ▶  $e^{(2)} = Be^{(1)} = B(\lambda y) = \lambda By = \lambda^2 y$

# Convergence of iterative methods 5

## Theorem 9.14

$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B) < 1$

## Proof.

Necessary condition: assume convergence and prove  $\rho(B) < 1$

- ▶ Proof by contradiction: assume  $By = \lambda y, |\lambda| > 1$
- ▶  $x^{(0)} = x + y, x = Bx + f$
- ▶  $e^{(0)} = x^{(0)} - x = y$
- ▶  $e^{(1)} = Be^{(0)} = By = \lambda y$
- ▶  $e^{(2)} = Be^{(1)} = B(\lambda y) = \lambda By = \lambda^2 y$
- ▶ ..
- ▶  $e^{(k)} = \lambda^k y$

# Convergence of iterative methods 5

## Theorem 9.14

$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B) < 1$

## Proof.

Necessary condition: assume convergence and prove  $\rho(B) < 1$

- ▶ Proof by contradiction: assume  $By = \lambda y, |\lambda| > 1$
- ▶  $x^{(0)} = x + y, x = Bx + f$
- ▶  $e^{(0)} = x^{(0)} - x = y$
- ▶  $e^{(1)} = Be^{(0)} = By = \lambda y$
- ▶  $e^{(2)} = Be^{(1)} = B(\lambda y) = \lambda By = \lambda^2 y$
- ▶ ..
- ▶  $e^{(k)} = \lambda^k y$
- ▶  $\|e^{(k)}\| = \|\lambda^k y\| = |\lambda^k| \|y\| \xrightarrow{k \rightarrow \infty} \infty$



# Convergence of iterative methods 6

## Theorem 9.15

$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B) < 1$

# Convergence of iterative methods 6

## Theorem 9.15

$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, \dots$  converges iff  $\rho(B) < 1$

Proof.

Sufficient condition: assume  $\rho(B) < 1$  and prove convergence



# Convergence of iterative methods 6

## Theorem 9.15

$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, \dots$  converges iff  $\rho(B) < 1$

Proof.

Sufficient condition: assume  $\rho(B) < 1$  and prove convergence

- Proof under additional assumption:  $By_i = \lambda_i y_i$ , eigenvectors are linearly independent

# Convergence of iterative methods 6

## Theorem 9.15

$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, \dots$  converges iff  $\rho(B) < 1$

Proof.

Sufficient condition: assume  $\rho(B) < 1$  and prove convergence

- ▶ Proof under additional assumption:  $By_i = \lambda_i y_i$ , eigenvectors are linearly independent
- ▶  $e^{(0)} = \sum_{i=1}^n c_i y_i$

# Convergence of iterative methods 6

## Theorem 9.15

$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, \dots$  converges iff  $\rho(B) < 1$

Proof.

Sufficient condition: assume  $\rho(B) < 1$  and prove convergence

- ▶ Proof under additional assumption:  $By_i = \lambda_i y_i$ , eigenvectors are linearly independent
- ▶  $e^{(0)} = \sum_{i=1}^n c_i y_i$
- ▶  $e^{(1)} = Be^{(0)} = B \sum_{i=1}^n c_i y_i = \sum_{i=1}^n c_i \lambda_i y_i$

# Convergence of iterative methods 6

## Theorem 9.15

$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, \dots$  converges iff  $\rho(B) < 1$

Proof.

Sufficient condition: assume  $\rho(B) < 1$  and prove convergence

- ▶ Proof under additional assumption:  $By_i = \lambda_i y_i$ , eigenvectors are linearly independent
- ▶  $e^{(0)} = \sum_{i=1}^n c_i y_i$
- ▶  $e^{(1)} = Be^{(0)} = B \sum_{i=1}^n c_i y_i = \sum_{i=1}^n c_i \lambda_i y_i$
- ▶  $e^{(2)} = Be^{(1)} = B \sum_{i=1}^n c_i \lambda_i y_i = \sum_{i=1}^n c_i \lambda_i By_i = \sum_{i=1}^n c_i \lambda_i^2 y_i$

# Convergence of iterative methods 6

## Theorem 9.15

$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, \dots$  converges iff  $\rho(B) < 1$

Proof.

Sufficient condition: assume  $\rho(B) < 1$  and prove convergence

- ▶ Proof under additional assumption:  $By_i = \lambda_i y_i$ , eigenvectors are linearly independent
- ▶  $e^{(0)} = \sum_{i=1}^n c_i y_i$
- ▶  $e^{(1)} = Be^{(0)} = B \sum_{i=1}^n c_i y_i = \sum_{i=1}^n c_i \lambda_i y_i$
- ▶  $e^{(2)} = Be^{(1)} = B \sum_{i=1}^n c_i \lambda_i y_i = \sum_{i=1}^n c_i \lambda_i By_i = \sum_{i=1}^n c_i \lambda_i^2 y_i$
- ▶ ..
- ▶  $e^{(k)} = \sum_{i=1}^n c_i \lambda_i^k y_i$

# Convergence of iterative methods 6

## Theorem 9.15

$x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, \dots$  converges iff  $\rho(B) < 1$

Proof.

Sufficient condition: assume  $\rho(B) < 1$  and prove convergence

► Proof under additional assumption:  $By_i = \lambda_i y_i$ , eigenvectors are linearly independent

►  $e^{(0)} = \sum_{i=1}^n c_i y_i$

►  $e^{(1)} = Be^{(0)} = B \sum_{i=1}^n c_i y_i = \sum_{i=1}^n c_i \lambda_i y_i$

►  $e^{(2)} = Be^{(1)} = B \sum_{i=1}^n c_i \lambda_i y_i = \sum_{i=1}^n c_i \lambda_i By_i = \sum_{i=1}^n c_i \lambda_i^2 y_i$

► ..

►  $e^{(k)} = \sum_{i=1}^n c_i \lambda_i^k y_i$

$$\|e^{(k)}\| = \left\| \sum_{i=1}^n c_i \lambda_i^k y_i \right\| = \sum_{i=1}^n |c_i| |\lambda_i|^k \|y_i\| \leq$$

$$\rho(B)^k \sum_{i=1}^n |c_i| \|y_i\| \xrightarrow{k \rightarrow \infty} 0$$

# Convergence of iterative methods 7

Important theorem on norms and spectral radius

# Convergence of iterative methods 7

Important theorem on norms and spectral radius

Theorem 9.16

*for any  $A \in \mathbb{C}^{n \times n}$  and  $\epsilon > 0$  there exist*



# Convergence of iterative methods 7

## Important theorem on norms and spectral radius

### Theorem 9.16

*for any  $A \in \mathbb{C}^{n \times n}$  and  $\epsilon > 0$  there exist matrix norm  $\|\cdot\|_*$  such that*  
$$\rho(A) < \|A\|_* < \rho(A) + \epsilon$$

# Convergence of iterative methods 7

## Important theorem on norms and spectral radius

### Theorem 9.16

*for any  $A \in \mathbb{C}^{n \times n}$  and  $\epsilon > 0$  there exist matrix norm  $\|\cdot\|_*$  such that  $\rho(A) < \|A\|_* < \rho(A) + \epsilon$*

## Other proof of sufficient condition of convergence

# Convergence of iterative methods 7

## Important theorem on norms and spectral radius

### Theorem 9.16

for any  $A \in \mathbb{C}^{n \times n}$  and  $\epsilon > 0$  there exist matrix norm  $\|\cdot\|_*$  such that  $\rho(A) < \|A\|_* < \rho(A) + \epsilon$

## Other proof of sufficient condition of convergence

►  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$

# Convergence of iterative methods 7

## Important theorem on norms and spectral radius

### Theorem 9.16

for any  $A \in \mathcal{C}^{n \times n}$  and  $\epsilon > 0$  there exist matrix norm  $\|\cdot\|_*$  such that  $\rho(A) < \|A\|_* < \rho(A) + \epsilon$

## Other proof of sufficient condition of convergence

- ▶  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$
- ▶  $\rho(B) < 1 \Rightarrow \|B\|_* < 1$

# Convergence of iterative methods 7

## Important theorem on norms and spectral radius

### Theorem 9.16

for any  $A \in \mathbb{C}^{n \times n}$  and  $\epsilon > 0$  there exist matrix norm  $\|\cdot\|_*$  such that  $\rho(A) < \|A\|_* < \rho(A) + \epsilon$

## Other proof of sufficient condition of convergence

- ▶  $x^{(k+1)} = Bx^{(k)} + f, k = 0, 1, ..$
- ▶  $\rho(B) < 1 \Rightarrow \|B\|_* < 1$
- ▶ Apply sufficient condition:  $\|B\|_* < 1 \Rightarrow$  convergence

## Convergence of iterative methods 8

►  $x^{(k+1)} = B_1 x^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B_1) < 1$

## Convergence of iterative methods 8

- ▶  $x^{(k+1)} = B_1 x^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B_1) < 1$
- ▶  $x^{(k+1)} = B_2 x^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B_2) < 1$

## Convergence of iterative methods 8

- ▶  $x^{(k+1)} = B_1 x^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B_1) < 1$
- ▶  $x^{(k+1)} = B_2 x^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B_2) < 1$
- ▶  $\rho(B_2) < \rho(B_1)$



## Convergence of iterative methods 8

- ▶  $x^{(k+1)} = B_1 x^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B_1) < 1$
- ▶  $x^{(k+1)} = B_2 x^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B_2) < 1$
- ▶  $\rho(B_2) < \rho(B_1)$
- ▶ Which method converges faster?

## Convergence of iterative methods 8

- ▶  $x^{(k+1)} = B_1 x^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B_1) < 1$
- ▶  $x^{(k+1)} = B_2 x^{(k)} + f, k = 0, 1, ..$  converges iff  $\rho(B_2) < 1$
- ▶  $\rho(B_2) < \rho(B_1)$
- ▶ Which method converges faster?

Answer:

$$\begin{aligned}\|e^{(k)}\| &= \left\| \sum_{i=1}^n c_i \lambda_i^k y_i \right\| = \sum_{i=1}^n |c_i| |\lambda_i^k| \|y_i\| \leq \\ &\rho(B)^k \sum_{i=1}^n |c_i| \|y_i\| \xrightarrow{k \rightarrow \infty} 0\end{aligned}$$

# Iterative methods for linear systems of algebraic equations

# Iterative methods for linear systems of algebraic equations: example - iterative refinement

- Problem:  $Ax = b, A \in \mathcal{R}^{n \times n}, x \in \mathcal{R}^n$

# Iterative methods for linear systems of algebraic equations: example - iterative refinement

- ▶ Problem:  $Ax = b, A \in \mathcal{R}^{n \times n}, x \in \mathcal{R}^n$
- ▶ Apply direct method: due to rounding errors computed solution is  $\tilde{x}$

# Iterative methods for linear systems of algebraic equations: example - iterative refinement

- ▶ Problem:  $Ax = b, A \in \mathcal{R}^{n \times n}, x \in \mathcal{R}^n$
- ▶ Apply direct method: due to rounding errors computed solution is  $\tilde{x}$
- ▶ Q: What if computed solution by direct method  $\tilde{x}$  is not acceptable?

# Iterative methods for linear systems of algebraic equations: example - iterative refinement

- ▶ Problem:  $Ax = b, A \in \mathcal{R}^{n \times n}, x \in \mathcal{R}^n$
- ▶ Apply direct method: due to rounding errors computed solution is  $\tilde{x}$
- ▶ Q: What if computed solution by direct method  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement

# Iterative methods for linear systems of algebraic equations: example - iterative refinement

- ▶ Problem:  $Ax = b, A \in \mathcal{R}^{n \times n}, x \in \mathcal{R}^n$
- ▶ Apply direct method: due to rounding errors computed solution is  $\tilde{x}$
- ▶ Q: What if computed solution by direct method  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$



# Iterative methods for linear systems of algebraic equations: example - iterative refinement

- ▶ Problem:  $Ax = b, A \in \mathcal{R}^{n \times n}, x \in \mathcal{R}^n$
- ▶ Apply direct method: due to rounding errors computed solution is  $\tilde{x}$
- ▶ Q: What if computed solution by direct method  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$

# Iterative methods for linear systems of algebraic equations: example - iterative refinement

- ▶ Problem:  $Ax = b, A \in \mathcal{R}^{n \times n}, x \in \mathcal{R}^n$
- ▶ Apply direct method: due to rounding errors computed solution is  $\tilde{x}$
- ▶ Q: What if computed solution by direct method  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$

# Iterative methods for linear systems of algebraic equations: example - iterative refinement

- ▶ Problem:  $Ax = b, A \in \mathcal{R}^{n \times n}, x \in \mathcal{R}^n$
- ▶ Apply direct method: due to rounding errors computed solution is  $\tilde{x}$
- ▶ Q: What if computed solution by direct method  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$

# Iterative methods for linear systems of algebraic equations: example - iterative refinement

- ▶ Problem:  $Ax = b, A \in \mathcal{R}^{n \times n}, x \in \mathcal{R}^n$
- ▶ Apply direct method: due to rounding errors computed solution is  $\tilde{x}$
- ▶ Q: What if computed solution by direct method  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:

# Iterative methods for linear systems of algebraic equations: example - iterative refinement

- ▶ Problem:  $Ax = b, A \in \mathcal{R}^{n \times n}, x \in \mathcal{R}^n$
- ▶ Apply direct method: due to rounding errors computed solution is  $\tilde{x}$
- ▶ Q: What if computed solution by direct method  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:
  - ▶  $x^{(0)} = \tilde{x}, tol = 1, \epsilon = 10^{-l}$

# Iterative methods for linear systems of algebraic equations: example - iterative refinement

- ▶ Problem:  $Ax = b, A \in \mathcal{R}^{n \times n}, x \in \mathcal{R}^n$
- ▶ Apply direct method: due to rounding errors computed solution is  $\tilde{x}$
- ▶ Q: What if computed solution by direct method  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:
  - ▶  $x^{(0)} = \tilde{x}, tol = 1, \epsilon = 10^{-l}$
  - ▶ Do until  $tol \leq \epsilon$ :

# Iterative methods for linear systems of algebraic equations: example - iterative refinement

- ▶ Problem:  $Ax = b, A \in \mathcal{R}^{n \times n}, x \in \mathcal{R}^n$
- ▶ Apply direct method: due to rounding errors computed solution is  $\tilde{x}$
- ▶ Q: What if computed solution by direct method  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:
  - ▶  $x^{(0)} = \tilde{x}, tol = 1, \epsilon = 10^{-l}$
  - ▶ Do until  $tol \leq \epsilon$ :
    1.  $r^{(k)} = b - Ax^{(k)}$

# Iterative methods for linear systems of algebraic equations: example - iterative refinement

- ▶ Problem:  $Ax = b, A \in \mathcal{R}^{n \times n}, x \in \mathcal{R}^n$
- ▶ Apply direct method: due to rounding errors computed solution is  $\tilde{x}$
- ▶ Q: What if computed solution by direct method  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:
  - ▶  $x^{(0)} = \tilde{x}, tol = 1, \epsilon = 10^{-l}$
  - ▶ Do until  $tol \leq \epsilon$ :
    1.  $r^{(k)} = b - Ax^{(k)}$
    2.  $Ac^{(k)} = r^{(k)}$



# Iterative methods for linear systems of algebraic equations: example - iterative refinement

- ▶ Problem:  $Ax = b, A \in \mathcal{R}^{n \times n}, x \in \mathcal{R}^n$
- ▶ Apply direct method: due to rounding errors computed solution is  $\tilde{x}$
- ▶ Q: What if computed solution by direct method  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:
  - ▶  $x^{(0)} = \tilde{x}, tol = 1, \epsilon = 10^{-l}$
  - ▶ Do until  $tol \leq \epsilon$ :
    1.  $r^{(k)} = b - Ax^{(k)}$
    2.  $Ac^{(k)} = r^{(k)}$
    3.  $x^{(k+1)} = x^{(k)} + c^{(k)}$

# Iterative methods for linear systems of algebraic equations: example - iterative refinement

- ▶ Problem:  $Ax = b, A \in \mathcal{R}^{n \times n}, x \in \mathcal{R}^n$
- ▶ Apply direct method: due to rounding errors computed solution is  $\tilde{x}$
- ▶ Q: What if computed solution by direct method  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:
  - ▶  $x^{(0)} = \tilde{x}, tol = 1, \epsilon = 10^{-l}$
  - ▶ Do until  $tol \leq \epsilon$ :
    1.  $r^{(k)} = b - Ax^{(k)}$
    2.  $Ac^{(k)} = r^{(k)}$
    3.  $x^{(k+1)} = x^{(k)} + c^{(k)}$
    4.  $tol = \frac{\|x^{(k+1)} - x^{(k)}\|}{\|x^{(k)}\|}$

# Iterative methods for linear systems of algebraic equations: example - iterative refinement

- ▶ Problem:  $Ax = b, A \in \mathcal{R}^{n \times n}, x \in \mathcal{R}^n$
- ▶ Apply direct method: due to rounding errors computed solution is  $\tilde{x}$
- ▶ Q: What if computed solution by direct method  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:
  - ▶  $x^{(0)} = \tilde{x}, tol = 1, \epsilon = 10^{-l}$
  - ▶ Do until  $tol \leq \epsilon$ :
    1.  $r^{(k)} = b - Ax^{(k)}$
    2.  $Ac^{(k)} = r^{(k)}$
    3.  $x^{(k+1)} = x^{(k)} + c^{(k)}$
    4.  $tol = \frac{\|x^{(k+1)} - x^{(k)}\|}{\|x^{(k)}\|}$
- ▶ Produces very accurate solution if  $cond(A)$  not too large

# Iterative methods for linear systems of algebraic equations: example - iterative refinement

- ▶ Problem:  $Ax = b, A \in \mathcal{R}^{n \times n}, x \in \mathcal{R}^n$
- ▶ Apply direct method: due to rounding errors computed solution is  $\tilde{x}$
- ▶ Q: What if computed solution by direct method  $\tilde{x}$  is not acceptable?
- ▶ A: Apply iterative refinement
  - ▶ Compute residual:  $r = b - A\tilde{x}$
  - ▶ Define correction:  $Ac = r$
  - ▶ Apply correction:  $x = \tilde{x} + c$
  - ▶  $Ax = A\tilde{x} + Ac = (b - r) + r = b \Rightarrow Ax = b$
- ▶ Iterative refinement algorithm:
  - ▶  $x^{(0)} = \tilde{x}, tol = 1, \epsilon = 10^{-l}$
  - ▶ Do until  $tol \leq \epsilon$ :
    1.  $r^{(k)} = b - Ax^{(k)}$
    2.  $Ac^{(k)} = r^{(k)}$
    3.  $x^{(k+1)} = x^{(k)} + c^{(k)}$
    4.  $tol = \frac{\|x^{(k+1)} - x^{(k)}\|}{\|x^{(k)}\|}$
- ▶ Produces very accurate solution if  $cond(A)$  not too large
- ▶ Convergence depends on  $cond(A)$

# Direct and Iterative methods

- ▶ Direct methods:
  - ▶ Finite number of arithmetic operations for finding exact solution

# Direct and Iterative methods

- ▶ Direct methods:
  - ▶ Finite number of arithmetic operations for finding exact solution
- ▶ Iterative methods:
  - ▶ Infinite number of arithmetic operations for finding exact solution

# Direct and Iterative methods

- ▶ Direct methods:
  - ▶ Finite number of arithmetic operations for finding exact solution
- ▶ Iterative methods:
  - ▶ Infinite number of arithmetic operations for finding exact solution
- ▶ Q: Which methods are better, direct or iterative?

# Direct and Iterative methods

- ▶ Direct methods:
  - ▶ Finite number of arithmetic operations for finding exact solution
- ▶ Iterative methods:
  - ▶ Infinite number of arithmetic operations for finding exact solution
- ▶ Q: Which methods are better, direct or iterative?
- ▶ Why do we need iterative refinement?



# Direct and Iterative methods

- ▶ Direct methods:
  - ▶ Finite number of arithmetic operations for finding exact solution
- ▶ Iterative methods:
  - ▶ Infinite number of arithmetic operations for finding exact solution
- ▶ Q: Which methods are better, direct or iterative?
- ▶ Why do we need iterative refinement?
- ▶ Would you like to apply iterative methods for system with  $2 \times 2$  coefficient matrix and two unknowns?

# Direct and Iterative methods

- ▶ Direct methods:
  - ▶ Finite number of arithmetic operations for finding exact solution
- ▶ Iterative methods:
  - ▶ Infinite number of arithmetic operations for finding exact solution
- ▶ Q: Which methods are better, direct or iterative?
- ▶ Why do we need iterative refinement?
- ▶ Would you like to apply iterative methods for system with  $2 \times 2$  coefficient matrix and two unknowns?
- ▶ Would you use Thomas algorithm or iterative method?

# Direct and Iterative methods

- ▶ Direct methods:
  - ▶ Finite number of arithmetic operations for finding exact solution
- ▶ Iterative methods:
  - ▶ Infinite number of arithmetic operations for finding exact solution
- ▶ Q: Which methods are better, direct or iterative?
- ▶ Why do we need iterative refinement?
- ▶ Would you like to apply iterative methods for system with  $2 \times 2$  coefficient matrix and two unknowns?
- ▶ Would you use Thomas algorithm or iterative method?
- ▶ What is term "exact solution" in floating point arithmetic?

# Jacobi Method

1.  $Ax = b, A \in \mathbb{R}^{n \times n}, x, b \in \mathbb{R}^n, x = ?$

# Jacobi Method

1.  $Ax = b, A \in \mathbb{R}^{n \times n}, x, b \in \mathbb{R}^n, x = ?$
2. Componentwise form

$$x_i^{(k+1)} = \frac{1}{a_{ii}}(b_i - \sum_{j=1, j \neq i}^n a_{ij}x_j^{(k)}), i = 1, 2, \dots, n, k = 0, 1, \dots$$

# Jacobi Method

1.  $Ax = b, A \in \mathbb{R}^{n \times n}, x, b \in \mathbb{R}^n, x = ?$
2. Componentwise form

$$x_i^{(k+1)} = \frac{1}{a_{ii}}(b_i - \sum_{j=1, j \neq i}^n a_{ij}x_j^{(k)}), i = 1, 2, \dots, n, k = 0, 1, \dots$$

3. Matrix form

$$\begin{aligned} A &= L + D + U, A, L, D, U \in \mathbb{R}^{n \times n}, a_{ii} \neq 0, \\ l_{ij} &= 0, \text{ if } i < j; d_{ij} = 0, \text{ if } i \neq j; u_{ij} = 0, \text{ if } i > j; \\ x^{(k+1)} &= -D^{-1}(L + U)x^{(k)} + D^{-1}b, k = 0, 1, \dots \\ B_J &= -D^{-1}(L + U), f_J = D^{-1}b, x^{(k+1)} = B_J x^{(k)} + f_J \end{aligned}$$

# Jacobi Method

1.  $Ax = b, A \in \mathbb{R}^{n \times n}, x, b \in \mathbb{R}^n, x = ?$
2. Componentwise form

$$x_i^{(k+1)} = \frac{1}{a_{ii}}(b_i - \sum_{j=1, j \neq i}^n a_{ij}x_j^{(k)}), i = 1, 2, \dots, n, k = 0, 1, \dots$$

3. Matrix form

$$\begin{aligned} A &= L + D + U, A, L, D, U \in \mathbb{R}^{n \times n}, a_{ii} \neq 0, \\ l_{ij} &= 0, \text{ if } i < j; d_{ij} = 0, \text{ if } i \neq j; u_{ij} = 0, \text{ if } i > j; \\ x^{(k+1)} &= -D^{-1}(L + U)x^{(k)} + D^{-1}b, k = 0, 1, \dots \\ B_J &= -D^{-1}(L + U), f_J = D^{-1}b, x^{(k+1)} = B_J x^{(k)} + f_J \end{aligned}$$

4. Convergence:  $\|D^{-1}(L + U)\| < 1, \rho(D^{-1}(L + U)) < 1$

# Gaus-Seidel

1.  $Ax = b, A \in \mathbb{R}^{n \times n}, x, b \in \mathbb{R}^n, x = ?$



# Gaus-Seidel

1.  $Ax = b, A \in \mathbb{R}^{n \times n}, x, b \in \mathbb{R}^n, x = ?$
2. Component-wise form

$$x_i^{(k+1)} = \frac{1}{a_{ii}} \left( b_i - \sum_{j=1}^{i-1} a_{ij} x_j^{(k+1)} - \sum_{j=i+1}^n a_{ij} x_j^{(k)} \right), i = 1, 2, \dots, n, k = 0, 1, \dots$$

# Gaus-Seidel

1.  $Ax = b, A \in \mathbb{R}^{n \times n}, x, b \in \mathbb{R}^n, x = ?$

2. Component-wise form

$$x_i^{(k+1)} = \frac{1}{a_{ii}} \left( b_i - \sum_{j=1}^{i-1} a_{ij} x_j^{(k+1)} - \sum_{j=i+1}^n a_{ij} x_j^{(k)} \right), i = 1, 2, \dots, n, k = 0, 1, \dots$$

3. Matrix form

$$A = L + D + U, A, L, D, U \in \mathbb{R}^{n \times n}, a_{ii} \neq 0,$$

$$l_{ij} = 0, \text{ if } i < j; d_{ij} = 0, \text{ if } i \neq j; u_{ij} = 0, \text{ if } i > j;$$

$$x^{(k+1)} = D^{-1}(b - Lx^{(k+1)} - Ux^{(k)}), k = 0, 1, \dots$$

$$x^{(k+1)} = (D + L)^{-1}(b - Ux^{(k)}), k = 0, 1, \dots$$

$$B_{GS} = -(D + L)^{-1}U, f_{GS} = (D + L)^{-1}f, x^{(k+1)} = B_{GS}x^{(k)} + f_{GS}$$

# Gaus-Seidel

1.  $Ax = b, A \in \mathbb{R}^{n \times n}, x, b \in \mathbb{R}^n, x = ?$
2. Component-wise form

$$x_i^{(k+1)} = \frac{1}{a_{ii}} \left( b_i - \sum_{j=1}^{i-1} a_{ij} x_j^{(k+1)} - \sum_{j=i+1}^n a_{ij} x_j^{(k)} \right), i = 1, 2, \dots, n, k = 0, 1, \dots$$

3. Matrix form

$$\begin{aligned} A &= L + D + U, A, L, D, U \in \mathbb{R}^{n \times n}, a_{ii} \neq 0, \\ l_{ij} &= 0, \text{ if } i < j; d_{ij} = 0, \text{ if } i \neq j; u_{ij} = 0, \text{ if } i > j; \\ x^{(k+1)} &= D^{-1}(b - Lx^{(k+1)} - Ux^{(k)}), k = 0, 1, \dots \\ x^{(k+1)} &= (D + L)^{-1}(b - Ux^{(k)}), k = 0, 1, \dots \end{aligned}$$

$$B_{GS} = -(D + L)^{-1}U, f_{GS} = (D + L)^{-1}f, x^{(k+1)} = B_{GS}x^{(k)} + f_{GS}$$

4. Convergence:  $\|(D + L)^{-1}U\| < 1, \rho((D + L)^{-1}U) < 1$

# JOR - Jacobi over relaxation

1.  $Ax = b, A \in \mathbb{R}^{n \times n}, x, b \in \mathbb{R}^n, x = ?$

# JOR - Jacobi over relaxation

1.  $Ax = b, A \in \mathbb{R}^{n \times n}, x, b \in \mathbb{R}^n, x = ?$
2. Componentwise form

$$x_i^{(k+1)} = \frac{\omega}{a_{ii}} \left( b_i - \sum_{j=1}^{i-1} a_{ij} x_j^{(k)} - \sum_{j=i+1}^n a_{ij} x_j^{(k)} \right) + (1 - \omega) x_i^{(k)}, i = 1, 2, \dots, n$$

# JOR - Jacobi over relaxation

1.  $Ax = b, A \in \mathbb{R}^{n \times n}, x, b \in \mathbb{R}^n, x = ?$
2. Componentwise form

$$x_i^{(k+1)} = \frac{\omega}{a_{ii}} \left( b_i - \sum_{j=1}^{i-1} a_{ij} x_j^{(k)} - \sum_{j=i+1}^n a_{ij} x_j^{(k)} \right) + (1 - \omega) x_i^{(k)}, i = 1, 2, \dots, n$$

3. Matrix form

$$A = L + D + U, A, L, D, U \in \mathbb{R}^{n \times n}, a_{ii} \neq 0,$$

$$l_{ij} = 0, \text{ if } i < j; d_{ij} = 0, \text{ if } i \neq j; u_{ij} = 0, \text{ if } i > j;$$

$$x^{(k+1)} = \omega D^{-1} (b - Lx^{(k)} - Ux^{(k)}) + (1 - \omega)x^{(k)}, k = 0, 1, \dots$$

$$x^{(k+1)} = \left( -\omega D^{-1} (L + U) + (1 - \omega)I \right) x^{(k)} + \omega D^{-1} b, k = 0, 1, \dots$$

$$B_{JOR} = -\omega D^{-1} (L + U) + (1 - \omega)I = \omega B_J + (1 - \omega)I, f_{JOR} = \omega D^{-1} b,$$

$$x^{(k+1)} = B_{JOR} x^{(k)} + f_{JOR}$$

# JOR - Jacobi over relaxation

1.  $Ax = b, A \in \mathbb{R}^{n \times n}, x, b \in \mathbb{R}^n, x = ?$
2. Componentwise form

$$x_i^{(k+1)} = \frac{\omega}{a_{ii}} \left( b_i - \sum_{j=1}^{i-1} a_{ij} x_j^{(k)} - \sum_{j=i+1}^n a_{ij} x_j^{(k)} \right) + (1 - \omega) x_i^{(k)}, i = 1, 2, \dots, n$$

3. Matrix form

$$A = L + D + U, A, L, D, U \in \mathbb{R}^{n \times n}, a_{ii} \neq 0,$$

$$l_{ij} = 0, \text{ if } i < j; d_{ij} = 0, \text{ if } i \neq j; u_{ij} = 0, \text{ if } i > j;$$

$$x^{(k+1)} = \omega D^{-1} (b - Lx^{(k)} - Ux^{(k)}) + (1 - \omega)x^{(k)}, k = 0, 1, \dots$$

$$x^{(k+1)} = \left( -\omega D^{-1} (L + U) + (1 - \omega)I \right) x^{(k)} + \omega D^{-1} b, k = 0, 1, \dots$$

$$B_{JOR} = -\omega D^{-1} (L + U) + (1 - \omega)I = \omega B_J + (1 - \omega)I, f_{JOR} = \omega D^{-1} b,$$

$$x^{(k+1)} = B_{JOR} x^{(k)} + f_{JOR}$$

4. Convergence:  $\|\omega B_J + (1 - \omega)I\| < 1, \rho(\omega B_J + (1 - \omega)I) < 1$

# SOR - Successive over relaxation

1.  $Ax = b, A \in \mathbb{R}^{n \times n}, x, b \in \mathbb{R}^n, x = ?$



## SOR - Successive over relaxation

1.  $Ax = b, A \in \mathbb{R}^{n \times n}, x, b \in \mathbb{R}^n, x = ?$

2. Componentwise form

$$x_i^{(k+1)} = \frac{\omega}{a_{ii}} \left( b_i - \sum_{j=1}^{i-1} a_{ij} x_j^{(k+1)} - \sum_{j=i+1}^n a_{ij} x_j^{(k)} \right) + (1-\omega) x_i^{(k)}, i = 1, 2, \dots, n$$

## SOR - Successive over relaxation

1.  $Ax = b, A \in \mathbb{R}^{n \times n}, x, b \in \mathbb{R}^n, x = ?$

2. Componentwise form

$$x_i^{(k+1)} = \frac{\omega}{a_{ii}} \left( b_i - \sum_{j=1}^{i-1} a_{ij} x_j^{(k+1)} - \sum_{j=i+1}^n a_{ij} x_j^{(k)} \right) + (1-\omega) x_i^{(k)}, i = 1, 2, \dots, n$$

3. Matrix form

$$A = L + D + U, A, L, D, U \in \mathbb{R}^{n \times n}, a_{ii} \neq 0,$$

$$l_{ij} = 0, \text{ if } i < j; d_{ij} = 0, \text{ if } i \neq j; u_{ij} = 0, \text{ if } i > j;$$

$$x^{(k+1)} = \omega D^{-1} (b - Lx^{(k+1)} - Ux^{(k)}) + (1 - \omega)x^{(k)}, k = 0, 1, \dots$$

$$x^{(k+1)} = (D + \omega L)^{-1} (b + (1 - \omega)D - \omega Ux^{(k)}), k = 0, 1, \dots$$

$$B_{SOR} = (D + \omega L)^{-1} ((1 - \omega)D - \omega U), f_{SOR} = (D + \omega L)^{-1} b,$$

$$x^{(k+1)} = B_{SOR}x^{(k)} + f_{SOR}$$

## SOR - Successive over relaxation

1.  $Ax = b, A \in \mathbb{R}^{n \times n}, x, b \in \mathbb{R}^n, x = ?$

2. Componentwise form

$$x_i^{(k+1)} = \frac{\omega}{a_{ii}} \left( b_i - \sum_{j=1}^{i-1} a_{ij} x_j^{(k+1)} - \sum_{j=i+1}^n a_{ij} x_j^{(k)} \right) + (1-\omega) x_i^{(k)}, i = 1, 2, \dots, n$$

3. Matrix form

$$A = L + D + U, A, L, D, U \in \mathbb{R}^{n \times n}, a_{ii} \neq 0,$$

$$l_{ij} = 0, \text{ if } i < j; d_{ij} = 0, \text{ if } i \neq j; u_{ij} = 0, \text{ if } i > j;$$

$$x^{(k+1)} = \omega D^{-1} (b - Lx^{(k+1)} - Ux^{(k)}) + (1-\omega)x^{(k)}, k = 0, 1, \dots$$

$$x^{(k+1)} = (D + \omega L)^{-1} (b + (1-\omega)D - \omega Ux^{(k)}), k = 0, 1, \dots$$

$$B_{SOR} = (D + \omega L)^{-1} ((1-\omega)D - \omega U), f_{SOR} = (D + \omega L)^{-1} b,$$

$$x^{(k+1)} = B_{SOR} x^{(k)} + f_{SOR}$$

4. Convergence:

$$\|(D + \omega L)^{-1} ((1-\omega)D - \omega U)\| < 1, \rho((D + \omega L)^{-1} ((1-\omega)D - \omega U)) < 1$$

# Richardson's method

## ► Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

# Richardson's method

- ▶ Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

- ▶ Consistent with  $Ax = b$

# Richardson's method

- ▶ Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

- ▶ Consistent with  $Ax = b$

- ▶  $\frac{1}{\tau_k} P x^{(k+1)} = (\frac{1}{\tau_k} P - A)x^{(k)} + b, \quad k = 0, 1, 2, \dots$

# Richardson's method

- ▶ Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

- ▶ Consistent with  $Ax = b$

- ▶  $\frac{1}{\tau_k} Px^{(k+1)} = (\frac{1}{\tau_k} P - A)x^{(k)} + b, \quad k = 0, 1, 2, \dots$

- ▶  $x^{(k+1)} = B_{R,k}x^{(k)} + f, \quad B_{R,k} = I - \tau_k P^{-1}A$

# Richardson's method

- ▶ Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

- ▶ Consistent with  $Ax = b$

- ▶  $\frac{1}{\tau_k} Px^{(k+1)} = (\frac{1}{\tau_k} P - A)x^{(k)} + b, \quad k = 0, 1, 2, \dots$

- ▶  $x^{(k+1)} = B_{R,k}x^{(k)} + f, \quad B_{R,k} = I - \tau_k P^{-1}A$

- ▶ Pre-conditioner -  $P \in \mathbb{R}^{n \times n}$



# Richardson's method

- ▶ Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

- ▶ Consistent with  $Ax = b$
- ▶  $\frac{1}{\tau_k} Px^{(k+1)} = (\frac{1}{\tau_k} P - A)x^{(k)} + b, \quad k = 0, 1, 2, \dots$
- ▶  $x^{(k+1)} = B_{R,k}x^{(k)} + f, \quad B_{R,k} = I - \tau_k P^{-1}A$
- ▶ Pre-conditioner -  $P \in \mathbb{R}^{n \times n}$
- ▶ Numeric parameter -  $\tau_k$

# Richardson's method

- ▶ Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

- ▶ Consistent with  $Ax = b$
- ▶  $\frac{1}{\tau_k} Px^{(k+1)} = (\frac{1}{\tau_k} P - A)x^{(k)} + b, \quad k = 0, 1, 2, \dots$
- ▶  $x^{(k+1)} = B_{R,k}x^{(k)} + f, \quad B_{R,k} = I - \tau_k P^{-1}A$
- ▶ Pre-conditioner -  $P \in \mathbb{R}^{n \times n}$
- ▶ Numeric parameter -  $\tau_k$
- ▶ Stationary iteration method if  $B_{R,k} \equiv B_R$

# Richardson's method

- ▶ Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

- ▶ Consistent with  $Ax = b$
- ▶  $\frac{1}{\tau_k} Px^{(k+1)} = (\frac{1}{\tau_k} P - A)x^{(k)} + b, \quad k = 0, 1, 2, \dots$
- ▶  $x^{(k+1)} = B_{R,k}x^{(k)} + f, \quad B_{R,k} = I - \tau_k P^{-1}A$
- ▶ Pre-conditioner -  $P \in \mathbb{R}^{n \times n}$
- ▶ Numeric parameter -  $\tau_k$
- ▶ Stationary iteration method if  $B_{R,k} \equiv B_R$
- ▶ Converges if  $\|B_{R,k}\| < 1$

# Richardson's method

- ▶ Richardson's method

$$P \frac{x^{(k+1)} - x^{(k)}}{\tau_k} + Ax^{(k)} = b, \quad k = 0, 1, 2, \dots$$

- ▶ Consistent with  $Ax = b$
- ▶  $\frac{1}{\tau_k} Px^{(k+1)} = (\frac{1}{\tau_k} P - A)x^{(k)} + b, \quad k = 0, 1, 2, \dots$
- ▶  $x^{(k+1)} = B_{R,k}x^{(k)} + f, \quad B_{R,k} = I - \tau_k P^{-1}A$
- ▶ Pre-conditioner -  $P \in \mathbb{R}^{n \times n}$
- ▶ Numeric parameter -  $\tau_k$
- ▶ Stationary iteration method if  $B_{R,k} \equiv B_R$
- ▶ Converges if  $\|B_{R,k}\| < 1$
- ▶ Converges iff  $\rho(B_{R,k}) < 1$

# Convergence Jacobi, GS, JOR 1

## Theorem 9.17

*A strictly row diagonally dominant  $\Rightarrow$  Jacobi method converges*

# Convergence Jacobi, GS, JOR 1

## Theorem 9.17

*A strictly row diagonally dominant  $\Rightarrow$  Jacobi method converges*

Proof.

- ▶ Strictly row diagonally dominance  $|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}|, i = 1, 2, \dots, n$

# Convergence Jacobi, GS, JOR 1

## Theorem 9.17

*A strictly row diagonally dominant  $\Rightarrow$  Jacobi method converges*

Proof.

- ▶ Strictly row diagonally dominance  $|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}|, i = 1, 2, \dots, n$
- ▶  $|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}| \Rightarrow \frac{1}{|a_{ii}|} \sum_{j=1, j \neq i}^n |a_{ij}| < 1, i = 1, 2, \dots, n$

# Convergence Jacobi, GS, JOR 1

## Theorem 9.17

*A strictly row diagonally dominant  $\Rightarrow$  Jacobi method converges*

Proof.

- ▶ Strictly row diagonally dominance  $|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}|, i = 1, 2, \dots, n$
- ▶  $|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}| \Rightarrow \frac{1}{|a_{ii}|} \sum_{j=1, j \neq i}^n |a_{ij}| < 1, i = 1, 2, \dots, n$
- ▶  $\sum_{j=1, j \neq i}^n \frac{|a_{ij}|}{|a_{ii}|} < 1, i = 1, 2, \dots, n \Rightarrow \max_{1 \leq i \leq n} \frac{1}{|a_{ii}|} \sum_{j=1, j \neq i}^n |a_{ij}| < 1$



# Convergence Jacobi, GS, JOR 1

## Theorem 9.17

*A strictly row diagonally dominant  $\Rightarrow$  Jacobi method converges*

### Proof.

- ▶ Strictly row diagonally dominance  $|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}|, i = 1, 2, \dots, n$
- ▶  $|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}| \Rightarrow \frac{1}{|a_{ii}|} \sum_{j=1, j \neq i}^n |a_{ij}| < 1, i = 1, 2, \dots, n$
- ▶  $\sum_{j=1, j \neq i}^n \frac{|a_{ij}|}{|a_{ii}|} < 1, i = 1, 2, \dots, n \Rightarrow \max_{1 \leq i \leq n} \frac{1}{|a_{ii}|} \sum_{j=1, j \neq i}^n |a_{ij}| < 1$
- ▶  $\frac{1}{|a_{ii}|} \sum_{j=1, j \neq i}^n |a_{ij}| = \frac{1}{|a_{ii}|} (\sum_{j=1}^{i-1} |a_{ij}| + \sum_{j=i+1}^n |a_{ij}|)$

# Convergence Jacobi, GS, JOR 1

## Theorem 9.17

*A strictly row diagonally dominant  $\Rightarrow$  Jacobi method converges*

### Proof.

- ▶ Strictly row diagonally dominance  $|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}|, i = 1, 2, \dots, n$
- ▶  $|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}| \Rightarrow \frac{1}{|a_{ii}|} \sum_{j=1, j \neq i}^n |a_{ij}| < 1, i = 1, 2, \dots, n$
- ▶  $\sum_{j=1, j \neq i}^n \frac{|a_{ij}|}{|a_{ii}|} < 1, i = 1, 2, \dots, n \Rightarrow \max_{1 \leq i \leq n} \frac{1}{|a_{ii}|} \sum_{j=1, j \neq i}^n |a_{ij}| < 1$
- ▶  $\frac{1}{|a_{ii}|} \sum_{j=1, j \neq i}^n |a_{ij}| = \frac{1}{|a_{ii}|} (\sum_{j=1}^{i-1} |a_{ij}| + \sum_{j=i+1}^n |a_{ij}|)$
- ▶  $1 > \max_{1 \leq i \leq n} \frac{1}{|a_{ii}|} (\sum_{j=1}^{i-1} |a_{ij}| + \sum_{j=i+1}^n |a_{ij}|) = \|D^{-1}(L + U)\|_{\infty}$

# Convergence Jacobi, GS, JOR 1

## Theorem 9.17

*A strictly row diagonally dominant  $\Rightarrow$  Jacobi method converges*

### Proof.

- ▶ Strictly row diagonally dominance  $|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}|, i = 1, 2, \dots, n$
- ▶  $|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}| \Rightarrow \frac{1}{|a_{ii}|} \sum_{j=1, j \neq i}^n |a_{ij}| < 1, i = 1, 2, \dots, n$
- ▶  $\sum_{j=1, j \neq i}^n \frac{|a_{ij}|}{|a_{ii}|} < 1, i = 1, 2, \dots, n \Rightarrow \max_{1 \leq i \leq n} \frac{1}{|a_{ii}|} \sum_{j=1, j \neq i}^n |a_{ij}| < 1$
- ▶  $\frac{1}{|a_{ii}|} \sum_{j=1, j \neq i}^n |a_{ij}| = \frac{1}{|a_{ii}|} (\sum_{j=1}^{i-1} |a_{ij}| + \sum_{j=i+1}^n |a_{ij}|)$
- ▶  $1 > \max_{1 \leq i \leq n} \frac{1}{|a_{ii}|} (\sum_{j=1}^{i-1} |a_{ij}| + \sum_{j=i+1}^n |a_{ij}|) = \|D^{-1}(L + U)\|_{\infty}$
- ▶  $\|D^{-1}(L + U)\|_{\infty} = \| - D^{-1}(L + U) \|_{\infty} = \|B_J\|_{\infty} < 1$



# Convergence Jacobi, GS, JOR 2

## Theorem 9.18

*A strictly row diagonally dominant  $\Rightarrow$  GS method converges*

# Convergence Jacobi, GS, JOR 2

## Theorem 9.18

*A strictly row diagonally dominant  $\Rightarrow$  GS method converges*

Proof.

$$\blacktriangleright |a_{ii}| - \sum_{j=1}^{i-1} |a_{ij}| > \sum_{j=i+1}^n |a_{ij}|, , i = 1, 2, \dots, n$$

# Convergence Jacobi, GS, JOR 2

## Theorem 9.18

*A strictly row diagonally dominant  $\Rightarrow$  GS method converges*

Proof.

- ▶  $|a_{ii}| - \sum_{j=1}^{i-1} |a_{ij}| > \sum_{j=i+1}^n |a_{ij}|, , i = 1, 2, \dots, n$
- ▶  $B_{GS} = -(D + L)^{-1}U$ , if  $\rho(B_{GS}) < 1 \Rightarrow$  convergence

# Convergence Jacobi, GS, JOR 2

## Theorem 9.18

*A strictly row diagonally dominant  $\Rightarrow$  GS method converges*

**Proof.**

- ▶  $|a_{ii}| - \sum_{j=1}^{i-1} |a_{ij}| > \sum_{j=i+1}^n |a_{ij}|, , i = 1, 2, \dots, n$
- ▶  $B_{GS} = -(D + L)^{-1}U$ , if  $\rho(B_{GS}) < 1 \Rightarrow$  convergence
- ▶  $B_{GS}y = \lambda y \Rightarrow -(D + L)^{-1}Uy = \lambda y$

# Convergence Jacobi, GS, JOR 2

## Theorem 9.18

*A strictly row diagonally dominant  $\Rightarrow$  GS method converges*

**Proof.**

- ▶  $|a_{ii}| - \sum_{j=1}^{i-1} |a_{ij}| > \sum_{j=i+1}^n |a_{ij}|, i = 1, 2, \dots, n$
- ▶  $B_{GS} = -(D + L)^{-1}U$ , if  $\rho(B_{GS}) < 1 \Rightarrow$  convergence
- ▶  $B_{GS}y = \lambda y \Rightarrow -(D + L)^{-1}Uy = \lambda y$
- ▶  $-(D + L)^{-1}Uy = \lambda y \Rightarrow Uy = -\lambda(D + L)y$



# Convergence Jacobi, GS, JOR 2

## Theorem 9.18

*A strictly row diagonally dominant  $\Rightarrow$  GS method converges*

**Proof.**

- ▶  $|a_{ii}| - \sum_{j=1}^{i-1} |a_{ij}| > \sum_{j=i+1}^n |a_{ij}|, i = 1, 2, \dots, n$
- ▶  $B_{GS} = -(D + L)^{-1}U$ , if  $\rho(B_{GS}) < 1 \Rightarrow$  convergence
- ▶  $B_{GS}y = \lambda y \Rightarrow -(D + L)^{-1}Uy = \lambda y$
- ▶  $-(D + L)^{-1}Uy = \lambda y \Rightarrow Uy = -\lambda(D + L)y$
- ▶  $Uy = -\lambda(D + L)y \Rightarrow -\lambda Dy = (\lambda L + U)y$

# Convergence Jacobi, GS, JOR 2

## Theorem 9.18

*A strictly row diagonally dominant  $\Rightarrow$  GS method converges*

**Proof.**

- ▶  $|a_{ii}| - \sum_{j=1}^{i-1} |a_{ij}| > \sum_{j=i+1}^n |a_{ij}|, i = 1, 2, \dots, n$
- ▶  $B_{GS} = -(D + L)^{-1}U$ , if  $\rho(B_{GS}) < 1 \Rightarrow$  convergence
- ▶  $B_{GS}y = \lambda y \Rightarrow -(D + L)^{-1}Uy = \lambda y$
- ▶  $-(D + L)^{-1}Uy = \lambda y \Rightarrow Uy = -\lambda(D + L)y$
- ▶  $Uy = -\lambda(D + L)y \Rightarrow -\lambda Dy = (\lambda L + U)y$
- ▶ Component-wise:  $-\lambda a_{ii}y_i = \lambda \sum_{j=1}^{i-1} a_{ij}y_j + \sum_{j=i+1}^n a_{ij}y_j$

# Convergence Jacobi, GS, JOR 2

## Theorem 9.18

*A strictly row diagonally dominant  $\Rightarrow$  GS method converges*

**Proof.**

- ▶  $|a_{ii}| - \sum_{j=1}^{i-1} |a_{ij}| > \sum_{j=i+1}^n |a_{ij}|, i = 1, 2, \dots, n$
- ▶  $B_{GS} = -(D + L)^{-1}U$ , if  $\rho(B_{GS}) < 1 \Rightarrow$  convergence
- ▶  $B_{GS}y = \lambda y \Rightarrow -(D + L)^{-1}Uy = \lambda y$
- ▶  $-(D + L)^{-1}Uy = \lambda y \Rightarrow Uy = -\lambda(D + L)y$
- ▶  $Uy = -\lambda(D + L)y \Rightarrow -\lambda Dy = (\lambda L + U)y$
- ▶ Component-wise:  $-\lambda a_{ii}y_i = \lambda \sum_{j=1}^{i-1} a_{ij}y_j + \sum_{j=i+1}^n a_{ij}y_j$
- ▶  $|\lambda| \cdot |a_{ii}| \cdot |y_i| \leq |\lambda| \sum_{j=1}^{i-1} |a_{ij}| \cdot |y_j| + \sum_{j=i+1}^n |a_{ij}| \cdot |y_j|$

# Convergence Jacobi, GS, JOR 2

## Theorem 9.18

*A strictly row diagonally dominant  $\Rightarrow$  GS method converges*

### Proof.

- ▶  $|a_{ii}| - \sum_{j=1}^{i-1} |a_{ij}| > \sum_{j=i+1}^n |a_{ij}|, i = 1, 2, \dots, n$
- ▶  $B_{GS} = -(D + L)^{-1}U$ , if  $\rho(B_{GS}) < 1 \Rightarrow$  convergence
- ▶  $B_{GS}y = \lambda y \Rightarrow -(D + L)^{-1}Uy = \lambda y$
- ▶  $-(D + L)^{-1}Uy = \lambda y \Rightarrow Uy = -\lambda(D + L)y$
- ▶  $Uy = -\lambda(D + L)y \Rightarrow -\lambda Dy = (\lambda L + U)y$
- ▶ Component-wise:  $-\lambda a_{ii}y_i = \lambda \sum_{j=1}^{i-1} a_{ij}y_j + \sum_{j=i+1}^n a_{ij}y_j$
- ▶  $|\lambda| \cdot |a_{ii}| \cdot |y_i| \leq |\lambda| \sum_{j=1}^{i-1} |a_{ij}| \cdot |y_j| + \sum_{j=i+1}^n |a_{ij}| \cdot |y_j|$
- ▶  $|y_k| = \|y\|_{\infty} \Rightarrow |\lambda| \cdot |a_{kk}| \leq |\lambda| \sum_{j=1}^{i-1} |a_{ij}| \frac{|y_j|}{|y_k|} + \sum_{j=i+1}^n |a_{ij}| \frac{|y_j|}{|y_k|}$

# Convergence Jacobi, GS, JOR 2

## Theorem 9.18

*A strictly row diagonally dominant  $\Rightarrow$  GS method converges*

**Proof.**

- ▶  $|a_{ii}| - \sum_{j=1}^{i-1} |a_{ij}| > \sum_{j=i+1}^n |a_{ij}|, i = 1, 2, \dots, n$
- ▶  $B_{GS} = -(D + L)^{-1}U$ , if  $\rho(B_{GS}) < 1 \Rightarrow$  convergence
- ▶  $B_{GS}y = \lambda y \Rightarrow -(D + L)^{-1}Uy = \lambda y$
- ▶  $-(D + L)^{-1}Uy = \lambda y \Rightarrow Uy = -\lambda(D + L)y$
- ▶  $Uy = -\lambda(D + L)y \Rightarrow -\lambda Dy = (\lambda L + U)y$
- ▶ Component-wise:  $-\lambda a_{ii}y_i = \lambda \sum_{j=1}^{i-1} a_{ij}y_j + \sum_{j=i+1}^n a_{ij}y_j$
- ▶  $|\lambda| \cdot |a_{ii}| \cdot |y_i| \leq |\lambda| \sum_{j=1}^{i-1} |a_{ij}| \cdot |y_j| + \sum_{j=i+1}^n |a_{ij}| \cdot |y_j|$
- ▶  $|y_k| = \|y\|_{\infty} \Rightarrow |\lambda| \cdot |a_{kk}| \leq |\lambda| \sum_{j=1}^{i-1} |a_{ij}| \frac{|y_j|}{|y_k|} + \sum_{j=i+1}^n |a_{ij}| \frac{|y_j|}{|y_k|}$
- ▶  $|\lambda| \cdot |a_{kk}| \leq |\lambda| \sum_{j=1}^{i-1} |a_{ij}| + \sum_{j=i+1}^n |a_{ij}|$

# Convergence Jacobi, GS, JOR 2

## Theorem 9.18

A strictly row diagonally dominant  $\Rightarrow$  GS method converges

Proof.

- ▶  $|a_{ii}| - \sum_{j=1}^{i-1} |a_{ij}| > \sum_{j=i+1}^n |a_{ij}|, i = 1, 2, \dots, n$
- ▶  $B_{GS} = -(D + L)^{-1}U$ , if  $\rho(B_{GS}) < 1 \Rightarrow$  convergence
- ▶  $B_{GS}y = \lambda y \Rightarrow -(D + L)^{-1}Uy = \lambda y$
- ▶  $-(D + L)^{-1}Uy = \lambda y \Rightarrow Uy = -\lambda(D + L)y$
- ▶  $Uy = -\lambda(D + L)y \Rightarrow -\lambda Dy = (\lambda L + U)y$
- ▶ Component-wise:  $-\lambda a_{ii}y_i = \lambda \sum_{j=1}^{i-1} a_{ij}y_j + \sum_{j=i+1}^n a_{ij}y_j$
- ▶  $|\lambda| \cdot |a_{ii}| \cdot |y_i| \leq |\lambda| \sum_{j=1}^{i-1} |a_{ij}| \cdot |y_j| + \sum_{j=i+1}^n |a_{ij}| \cdot |y_j|$
- ▶  $|y_k| = \|y\|_{\infty} \Rightarrow |\lambda| \cdot |a_{kk}| \leq |\lambda| \sum_{j=1}^{i-1} |a_{ij}| \frac{|y_j|}{|y_k|} + \sum_{j=i+1}^n |a_{ij}| \frac{|y_j|}{|y_k|}$
- ▶  $|\lambda| \cdot |a_{kk}| \leq |\lambda| \sum_{j=1}^{i-1} |a_{ij}| + \sum_{j=i+1}^n |a_{ij}|$
- ▶  $|\lambda|(|a_{kk}| - \sum_{j=1}^{i-1} |a_{ij}|) \leq \sum_{j=i+1}^n |a_{ij}| \Rightarrow |\lambda| \leq \frac{\sum_{j=i+1}^n |a_{ij}|}{|a_{kk}| - \sum_{j=1}^{i-1} |a_{ij}|} < 1$

# Convergence Jacobi, GS, JOR 3

## Theorem 9.19

*Jacobi method converges  $\Rightarrow$  JOR method converges for  $0 < \omega \leq 1$*

# Convergence Jacobi, GS, JOR 3

## Theorem 9.19

*Jacobi method converges  $\Rightarrow$  JOR method converges for  $0 < \omega \leq 1$*

## Proof.

► Jacobi method converges  $\Rightarrow \rho(B_J) < 1 \Rightarrow B_J y = \lambda y, |\lambda| < 1$



# Convergence Jacobi, GS, JOR 3

## Theorem 9.19

*Jacobi method converges  $\Rightarrow$  JOR method converges for  $0 < \omega \leq 1$*

## Proof.

- ▶ Jacobi method converges  $\Rightarrow \rho(B_J) < 1 \Rightarrow B_J y = \lambda y, |\lambda| < 1$
- ▶  $B_{JOR} = \omega B_J + (1 - \omega)I \Rightarrow B_{JOR} y = \omega B_J y + (1 - \omega)y = \omega \lambda y + (1 - \omega)y$

# Convergence Jacobi, GS, JOR 3

## Theorem 9.19

*Jacobi method converges  $\Rightarrow$  JOR method converges for  $0 < \omega \leq 1$*

## Proof.

- ▶ Jacobi method converges  $\Rightarrow \rho(B_J) < 1 \Rightarrow B_J y = \lambda y, |\lambda| < 1$
- ▶  $B_{JOR} = \omega B_J + (1 - \omega)I \Rightarrow B_{JOR} y = \omega B_J y + (1 - \omega)y = \omega \lambda y + (1 - \omega)y$
- ▶  $B_{JOR} y = \mu y \Rightarrow \mu = \omega \lambda + (1 - \omega)$

# Convergence Jacobi, GS, JOR 3

## Theorem 9.19

*Jacobi method converges  $\Rightarrow$  JOR method converges for  $0 < \omega \leq 1$*

### Proof.

- ▶ Jacobi method converges  $\Rightarrow \rho(B_J) < 1 \Rightarrow B_J y = \lambda y, |\lambda| < 1$
- ▶  $B_{JOR} = \omega B_J + (1 - \omega)I \Rightarrow B_{JOR} y = \omega B_J y + (1 - \omega)y = \omega \lambda y + (1 - \omega)y$
- ▶  $B_{JOR} y = \mu y \Rightarrow \mu = \omega \lambda + (1 - \omega)$
- ▶  $\lambda = r e^{i\theta}, e^{i\theta} = \cos(\theta) + i \sin(\theta), r < 1$

# Convergence Jacobi, GS, JOR 3

## Theorem 9.19

*Jacobi method converges  $\Rightarrow$  JOR method converges for  $0 < \omega \leq 1$*

### Proof.

- ▶ Jacobi method converges  $\Rightarrow \rho(B_J) < 1 \Rightarrow B_J y = \lambda y, |\lambda| < 1$
- ▶  $B_{JOR} = \omega B_J + (1 - \omega)I \Rightarrow B_{JOR} y = \omega B_J y + (1 - \omega)y = \omega \lambda y + (1 - \omega)y$
- ▶  $B_{JOR} y = \mu y \Rightarrow \mu = \omega \lambda + (1 - \omega)$
- ▶  $\lambda = re^{i\theta}, e^{i\theta} = \cos(\theta) + i \sin(\theta), r < 1$
- ▶  $\mu = \omega r(\cos(\theta) + i \sin(\theta)) + 1 - \omega = \omega r \cos(\theta) + 1 - \omega + i \omega r \sin(\theta)$

# Convergence Jacobi, GS, JOR 3

## Theorem 9.19

*Jacobi method converges  $\Rightarrow$  JOR method converges for  $0 < \omega \leq 1$*

### Proof.

- ▶ Jacobi method converges  $\Rightarrow \rho(B_J) < 1 \Rightarrow B_J y = \lambda y, |\lambda| < 1$
- ▶  $B_{JOR} = \omega B_J + (1 - \omega)I \Rightarrow B_{JOR} y = \omega B_J y + (1 - \omega)y = \omega \lambda y + (1 - \omega)y$
- ▶  $B_{JOR} y = \mu y \Rightarrow \mu = \omega \lambda + (1 - \omega)$
- ▶  $\lambda = re^{i\theta}, e^{i\theta} = \cos(\theta) + i \sin(\theta), r < 1$
- ▶  $\mu = \omega r(\cos(\theta) + i \sin(\theta)) + 1 - \omega = \omega r \cos(\theta) + 1 - \omega + i \omega r \sin(\theta)$

$$\begin{aligned} |\mu|^2 &= (\omega r \cos(\theta) + 1 - \omega)^2 + \omega^2 r^2 \sin^2(\theta) = \\ &= \omega^2 r^2 \cos^2(\theta) + 2\omega r(1 - \omega) \cos(\theta) + (1 - \omega)^2 + \omega^2 r^2 \sin^2(\theta) \\ &= \omega^2 r^2 + 2\omega r(1 - \omega) \cos(\theta) + (1 - \omega)^2 \leq (\omega \cdot r + (1 - \omega) \cdot 1)^2 < 1 \end{aligned}$$

# Convergence Jacobi, GS, JOR 3

## Theorem 9.19

*Jacobi method converges  $\Rightarrow$  JOR method converges for  $0 < \omega \leq 1$*

### Proof.

- ▶ Jacobi method converges  $\Rightarrow \rho(B_J) < 1 \Rightarrow B_J y = \lambda y, |\lambda| < 1$
- ▶  $B_{JOR} = \omega B_J + (1 - \omega)I \Rightarrow B_{JOR} y = \omega B_J y + (1 - \omega)y = \omega \lambda y + (1 - \omega)y$
- ▶  $B_{JOR} y = \mu y \Rightarrow \mu = \omega \lambda + (1 - \omega)$
- ▶  $\lambda = re^{i\theta}, e^{i\theta} = \cos(\theta) + i \sin(\theta), r < 1$
- ▶  $\mu = \omega r(\cos(\theta) + i \sin(\theta)) + 1 - \omega = \omega r \cos(\theta) + 1 - \omega + i \omega r \sin(\theta)$

$$\begin{aligned} |\mu|^2 &= (\omega r \cos(\theta) + 1 - \omega)^2 + \omega^2 r^2 \sin^2(\theta) = \\ &= \omega^2 r^2 \cos^2(\theta) + 2\omega r(1 - \omega) \cos(\theta) + (1 - \omega)^2 + \omega^2 r^2 \sin^2(\theta) \\ &= \omega^2 r^2 + 2\omega r(1 - \omega) \cos(\theta) + (1 - \omega)^2 \leq (\omega \cdot r + (1 - \omega) \cdot 1)^2 < 1 \end{aligned}$$

# Convergence Jacobi, GS, JOR 4

## Comparing Jacobi and Gaus-Seidel: Stein-Rosenberg Theorem

### Theorem 9.20

►  $a_{ij} < 0, i \neq j, \quad i, j = 1, 2, \dots, n$

# Convergence Jacobi, GS, JOR 4

## Comparing Jacobi and Gaus-Seidel: Stein-Rosenberg Theorem

### Theorem 9.20

- ▶  $a_{ij} < 0, i \neq j, \quad i, j = 1, 2, \dots, n$
- ▶  $a_{ii} > 0, i = 1, 2, \dots, n$



# Convergence Jacobi, GS, JOR 4

## Comparing Jacobi and Gaus-Seidel: Stein-Rosenberg Theorem

### Theorem 9.20

- ▶  $a_{ij} < 0, i \neq j, \quad i, j = 1, 2, \dots, n$
- ▶  $a_{ii} > 0, i = 1, 2, \dots, n$

*Only one of the following statements holds true*

# Convergence Jacobi, GS, JOR 4

## Comparing Jacobi and Gaus-Seidel: Stein-Rosenberg Theorem

### Theorem 9.20

- ▶  $a_{ij} < 0, i \neq j, \quad i, j = 1, 2, \dots, n$
- ▶  $a_{ii} > 0, i = 1, 2, \dots, n$

*Only one of the following statements holds true*

1.  $0 \leq \rho(G_{GS}) < \rho(B_J) < 1$

# Convergence Jacobi, GS, JOR 4

## Comparing Jacobi and Gaus-Seidel: Stein-Rosenberg Theorem

### Theorem 9.20

- ▶  $a_{ij} < 0, i \neq j, \quad i, j = 1, 2, \dots, n$
- ▶  $a_{ii} > 0, i = 1, 2, \dots, n$

*Only one of the following statements holds true*

1.  $0 \leq \rho(G_{GS}) < \rho(B_J) < 1$
2.  $1 < \rho(B_J) < \rho(G_{GS})$

# Convergence Jacobi, GS, JOR 4

## Comparing Jacobi and Gaus-Seidel: Stein-Rosenberg Theorem

### Theorem 9.20

- ▶  $a_{ij} < 0, i \neq j, \quad i, j = 1, 2, \dots, n$
- ▶  $a_{ii} > 0, i = 1, 2, \dots, n$

*Only one of the following statements holds true*

1.  $0 \leq \rho(G_{GS}) < \rho(B_J) < 1$
2.  $1 < \rho(B_J) < \rho(G_{GS})$
3.  $\rho(B_J) = \rho(B_{GSJ}) = 0$

# Convergence Jacobi, GS, JOR 4

## Comparing Jacobi and Gaus-Seidel: Stein-Rosenberg Theorem

### Theorem 9.20

- ▶  $a_{ij} < 0, i \neq j, \quad i, j = 1, 2, \dots, n$
- ▶  $a_{ii} > 0, i = 1, 2, \dots, n$

*Only one of the following statements holds true*

1.  $0 \leq \rho(G_{GS}) < \rho(B_J) < 1$
2.  $1 < \rho(B_J) < \rho(G_{GS})$
3.  $\rho(B_J) = \rho(B_{GSJ}) = 0$
4.  $\rho(B_J) = \rho(B_{GSJ}) = 1$

# Convergence Jacobi, GS, JOR 4

## Comparing Jacobi and Gaus-Seidel: Stein-Rosenberg Theorem

### Theorem 9.20

- ▶  $a_{ij} < 0, i \neq j, \quad i, j = 1, 2, \dots, n$
- ▶  $a_{ii} > 0, i = 1, 2, \dots, n$

*Only one of the following statements holds true*

1.  $0 \leq \rho(G_{GS}) < \rho(B_J) < 1$
2.  $1 < \rho(B_J) < \rho(G_{GS})$
3.  $\rho(B_J) = \rho(B_{GSJ}) = 0$
4.  $\rho(B_J) = \rho(B_{GSJ}) = 1$

### Theorem 9.21

- ▶ *Tridiagonal system*
- ▶ *Positive definite*

# Convergence Jacobi, GS, JOR 4

## Comparing Jacobi and Gaus-Seidel: Stein-Rosenberg Theorem

### Theorem 9.20

- ▶  $a_{ij} < 0, i \neq j, \quad i, j = 1, 2, \dots, n$
- ▶  $a_{ii} > 0, i = 1, 2, \dots, n$

*Only one of the following statements holds true*

1.  $0 \leq \rho(G_{GS}) < \rho(B_J) < 1$
2.  $1 < \rho(B_J) < \rho(G_{GS})$
3.  $\rho(B_J) = \rho(B_{GSJ}) = 0$
4.  $\rho(B_J) = \rho(B_{GSJ}) = 1$

### Theorem 9.21

- ▶ *Tridiagonal system*
- ▶ *Positive definite*

$$\Rightarrow \rho(B_{GS}) = \rho^2(B_J)$$

Q & A