

Numerical Linear Algebra

Ramaz Botchorishvili

Kutaisi International University

December 21, 2022

Numerical Linear Algebra

Ramaz Botchorishvili

Kutaisi International University

December 21, 2022

QR factorization, Least squares

- ▶ Invertibility of KKT matrix
- ▶ QR factorization for constrained least squares problem
- ▶ Householder transformations and QR factorization
- ▶ Numerical eigenvalue problem
- ▶ Q & A

Recap of Previous Lecture

- ▶ Recap of Previous Lecture
- ▶ QR and reduced QR factorization
- ▶ Linear least squares
- ▶ Constrained least squares

Constrained Least Squares

Problem 14.1

Constrained Least Squares Problem

Constrained Least Squares

Problem 14.1

Constrained Least Squares Problem

► $A \in \mathbb{R}^{n \times m}, n > m$

Constrained Least Squares

Problem 14.1

Constrained Least Squares Problem

- ▶ $A \in \mathbb{R}^{n \times m}, n > m$
- ▶ $C \in \mathbb{R}^{p \times m}$

Constrained Least Squares

Problem 14.1

Constrained Least Squares Problem

- ▶ $A \in \mathbb{R}^{n \times m}, n > m$
- ▶ $C \in \mathbb{R}^{p \times m}$
- ▶ $x \in \mathbb{R}^m, b \in \mathbb{R}^n, d \in \mathbb{R}^p$

Constrained Least Squares

Problem 14.1

Constrained Least Squares Problem

- ▶ $A \in \mathbb{R}^{n \times m}, n > m$
- ▶ $C \in \mathbb{R}^{p \times m}$
- ▶ $x \in \mathbb{R}^m, b \in \mathbb{R}^n, d \in \mathbb{R}^p$
- ▶ $Ax = b$ - over determined system

Constrained Least Squares

Problem 14.1

Constrained Least Squares Problem

- ▶ $A \in \mathbb{R}^{n \times m}, n > m$
- ▶ $C \in \mathbb{R}^{p \times m}$
- ▶ $x \in \mathbb{R}^m, b \in \mathbb{R}^n, d \in \mathbb{R}^p$
- ▶ $Ax = b$ - over determined system
- ▶ $f(x) = \|Ax - b\|_2$

Constrained Least Squares

Problem 14.1

Constrained Least Squares Problem

- ▶ $A \in \mathbb{R}^{n \times m}, n > m$
- ▶ $C \in \mathbb{R}^{p \times m}$
- ▶ $x \in \mathbb{R}^m, b \in \mathbb{R}^n, d \in \mathbb{R}^p$
- ▶ $Ax = b$ - over determined system
- ▶ $f(x) = \|Ax - b\|_2$
- ▶ $x_* = \arg \min_{Cx=d} f(x)$

Constrained Least Squares

Problem 14.1

Constrained Least Squares Problem

- ▶ $A \in \mathbb{R}^{n \times m}, n > m$
- ▶ $C \in \mathbb{R}^{p \times m}$
- ▶ $x \in \mathbb{R}^m, b \in \mathbb{R}^n, d \in \mathbb{R}^p$
- ▶ $Ax = b$ - over determined system
- ▶ $f(x) = \|Ax - b\|_2$
- ▶ $x_* = \arg \min_{Cx=d} f(x)$

KKT equation

$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} = \begin{pmatrix} 2A^T b \\ d \end{pmatrix}$$

Constrained Least Squares

Problem 14.1

Constrained Least Squares Problem

- ▶ $A \in \mathbb{R}^{n \times m}, n > m$
- ▶ $C \in \mathbb{R}^{p \times m}$
- ▶ $x \in \mathbb{R}^m, b \in \mathbb{R}^n, d \in \mathbb{R}^p$
- ▶ $Ax = b$ - over determined system
- ▶ $f(x) = \|Ax - b\|_2$
- ▶ $x_* = \arg \min_{Cx=d} f(x)$

KKT equation

$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} = \begin{pmatrix} 2A^T b \\ d \end{pmatrix}$$

KKT equation (William Karush, Harold Kuhn and Albert Tucker)

Constrained Least Squares, KKT matrix

KKT matrix

$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix}$$

Constrained Least Squares, KKT matrix

KKT matrix

$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix}$$

Theorem 14.2

KKT matrix is invertible iff

Constrained Least Squares, KKT matrix

KKT matrix

$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix}$$

Theorem 14.2

KKT matrix is invertible iff

- a. *C has linearly independent rows*

Constrained Least Squares, KKT matrix

KKT matrix

$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix}$$

Theorem 14.2

KKT matrix is invertible iff

- a. *C has linearly independent rows*
- b. *$\begin{pmatrix} A \\ C \end{pmatrix}$ has linearly independent columns*

Constrained Least Squares, KKT matrix

KKT matrix

$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix}$$

Theorem 14.2

KKT matrix is invertible iff

- a. *C has linearly independent rows*
- b. $\begin{pmatrix} A \\ C \end{pmatrix}$ *has linearly independent columns*

► Necessary condition:

Constrained Least Squares, KKT matrix

KKT matrix

$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix}$$

Theorem 14.2

KKT matrix is invertible iff

- a. *C has linearly independent rows*
- b. $\begin{pmatrix} A \\ C \end{pmatrix}$ *has linearly independent columns*

► Necessary condition:

KKT matrix is invertible \Rightarrow a. & b. of the theorem 14.2 are satisfied

Constrained Least Squares, KKT matrix

KKT matrix

$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix}$$

Theorem 14.2

KKT matrix is invertible iff

- C has linearly independent rows*
- $\begin{pmatrix} A \\ C \end{pmatrix}$ has linearly independent columns*

► Necessary condition:

KKT matrix is invertible \Rightarrow a. & b. of the theorem 14.2 are satisfied

► Sufficient condition:

Constrained Least Squares, KKT matrix

KKT matrix

$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix}$$

Theorem 14.2

KKT matrix is invertible iff

- C has linearly independent rows*
- $\begin{pmatrix} A \\ C \end{pmatrix}$ has linearly independent columns*

► Necessary condition:

KKT matrix is invertible \Rightarrow a. & b. of the theorem 14.2 are satisfied

► Sufficient condition:

a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible

Constrained Least Squares, KKT matrix

Proof of necessary condition, part a

- Necessary condition:

KKT matrix is invertible \Rightarrow a. & b. of the theorem 14.2 are satisfied

Constrained Least Squares, KKT matrix

Proof of necessary condition, part a

- ▶ Necessary condition:
KKT matrix is invertible \Rightarrow a. & b. of the theorem 14.2 are satisfied
- ▶ **ASSUME**: KKT matrix is invertible **AND** a. of the theorem 14.2 are **NOT** satisfied

Constrained Least Squares, KKT matrix

Proof of necessary condition, part a

- ▶ Necessary condition:
KKT matrix is invertible \Rightarrow a. & b. of the theorem 14.2 are satisfied
- ▶ **ASSUME**: KKT matrix is invertible **AND** a. of the theorem 14.2 are **NOT** satisfied



- ▶ Rows of C are linearly dependent

Constrained Least Squares, KKT matrix

Proof of necessary condition, part a

- ▶ Necessary condition:
KKT matrix is invertible \Rightarrow a. & b. of the theorem 14.2 are satisfied
- ▶ **ASSUME**: KKT matrix is invertible **AND** a. of the theorem 14.2 are **NOT** satisfied



- ▶ Rows of C are linearly dependent $\Rightarrow \exists \tilde{\lambda} \neq 0, C^T \tilde{\lambda} = 0$

Constrained Least Squares, KKT matrix

Proof of necessary condition, part a

- Necessary condition:

KKT matrix is invertible \Rightarrow a. & b. of the theorem 14.2 are satisfied

- **ASSUME**: KKT matrix is invertible **AND** a. of the theorem 14.2 are **NOT** satisfied

\Downarrow

- Rows of C are linearly dependent $\Rightarrow \exists \tilde{\lambda} \neq 0, C^T \tilde{\lambda} = 0$

\Downarrow



$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \tilde{x} \\ \tilde{\lambda} \end{pmatrix} = 0$$

Constrained Least Squares, KKT matrix

Proof of necessary condition, part a

- Necessary condition:

KKT matrix is invertible \Rightarrow a. & b. of the theorem 14.2 are satisfied

- **ASSUME**: KKT matrix is invertible **AND** a. of the theorem 14.2 are **NOT** satisfied

\Downarrow

- Rows of C are linearly dependent $\Rightarrow \exists \tilde{\lambda} \neq 0, C^T \tilde{\lambda} = 0$

\Downarrow



$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \tilde{x} \\ \tilde{\lambda} \end{pmatrix} = 0$$

\Downarrow

- **Contradiction**: KKT matrix is not invertible

Constrained Least Squares, KKT matrix

Proof of necessary condition, part b

- Necessary condition:

KKT matrix is invertible \Rightarrow a. & b. of the theorem 14.2 are satisfied

Constrained Least Squares, KKT matrix

Proof of necessary condition, part b

- ▶ Necessary condition:
KKT matrix is invertible \Rightarrow a. & b. of the theorem 14.2 are satisfied
- ▶ **ASSUME**: KKT matrix is invertible **AND** a. of the theorem 14.2 is **NOT** satisfied

Constrained Least Squares, KKT matrix

Proof of necessary condition, part b

- ▶ Necessary condition:
KKT matrix is invertible \Rightarrow a. & b. of the theorem 14.2 are satisfied
- ▶ **ASSUME**: KKT matrix is invertible **AND** a. of the theorem 14.2 is **NOT** satisfied



- ▶ Rows of C are linearly dependent

Constrained Least Squares, KKT matrix

Proof of necessary condition, part b

- ▶ Necessary condition:
KKT matrix is invertible \Rightarrow a. & b. of the theorem 14.2 are satisfied
- ▶ **ASSUME**: KKT matrix is invertible **AND** a. of the theorem 14.2 is **NOT** satisfied



- ▶ Rows of C are linearly dependent $\Rightarrow \exists \tilde{\lambda} \neq 0, C^T \tilde{\lambda} = 0$

Constrained Least Squares, KKT matrix

Proof of necessary condition, part b

- Necessary condition:

KKT matrix is invertible \Rightarrow a. & b. of the theorem 14.2 are satisfied

- **ASSUME**: KKT matrix is invertible **AND** a. of the theorem 14.2 is **NOT** satisfied

\Downarrow

- Rows of C are linearly dependent $\Rightarrow \exists \tilde{\lambda} \neq 0, C^T \tilde{\lambda} = 0$

\Downarrow



$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} 0 \\ \tilde{\lambda} \end{pmatrix} = 0$$

Constrained Least Squares, KKT matrix

Proof of necessary condition, part b

- ▶ Necessary condition:

KKT matrix is invertible \Rightarrow a. & b. of the theorem 14.2 are satisfied

- ▶ **ASSUME**: KKT matrix is invertible **AND** a. of the theorem 14.2 is **NOT** satisfied

\Downarrow

- ▶ Rows of C are linearly dependent $\Rightarrow \exists \tilde{\lambda} \neq 0, C^T \tilde{\lambda} = 0$

\Downarrow



$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} 0 \\ \tilde{\lambda} \end{pmatrix} = 0$$

\Downarrow

- ▶ **Contradiction**: KKT matrix is not invertible

Constrained Least Squares, KKT matrix

Proof of sufficient condition

- ▶ Sufficient condition:
 - a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible

Constrained Least Squares, KKT matrix

Proof of sufficient condition

- ▶ Sufficient condition:
 - a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶ **ASSUME**: a.& b. of the theorem 14.2 is satisfied **AND** KKT matrix is **NOT** invertible

Constrained Least Squares, KKT matrix

Proof of sufficient condition

- ▶ Sufficient condition:
 - a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶ **ASSUME**: a.& b. of the theorem 14.2 is satisfied **AND** KKT matrix is **NOT** invertible \Downarrow
- ▶ $\exists \tilde{x} \neq 0, \exists \tilde{\lambda} \neq 0, \begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \tilde{x} \\ \tilde{\lambda} \end{pmatrix} = 0$

Constrained Least Squares, KKT matrix

Proof of sufficient condition

- ▶ Sufficient condition:
 - a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶ **ASSUME**: a.& b. of the theorem 14.2 is satisfied **AND** KKT matrix is **NOT** invertible \Downarrow
- ▶ $\exists \tilde{x} \neq 0, \exists \tilde{\lambda} \neq 0, \begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \tilde{x} \\ \tilde{\lambda} \end{pmatrix} = 0 \Rightarrow 2A^T A \tilde{x} + C^T \tilde{\lambda} = 0$

Constrained Least Squares, KKT matrix

Proof of sufficient condition

- ▶ Sufficient condition:
 - a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶ **ASSUME**: a.& b. of the theorem 14.2 is satisfied **AND** KKT matrix is **NOT** invertible \Downarrow
- ▶ $\exists \tilde{x} \neq 0, \exists \tilde{\lambda} \neq 0, \begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \tilde{x} \\ \tilde{\lambda} \end{pmatrix} = 0 \Rightarrow 2A^T A \tilde{x} + C^T \tilde{\lambda} = 0 \Downarrow$
- ▶ $x^T (2A^T A \tilde{x} + C^T \tilde{\lambda}) = 0$

Constrained Least Squares, KKT matrix

Proof of sufficient condition

- ▶ Sufficient condition:
 - a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶ **ASSUME**: a.& b. of the theorem 14.2 is satisfied **AND** KKT matrix is **NOT** invertible \Downarrow
- ▶ $\exists \tilde{x} \neq 0, \exists \tilde{\lambda} \neq 0, \begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \tilde{x} \\ \tilde{\lambda} \end{pmatrix} = 0 \Rightarrow 2A^T A \tilde{x} + C^T \tilde{\lambda} = 0 \Downarrow$
- ▶ $x^T (2A^T A \tilde{x} + C^T \tilde{\lambda}) = 0 \Rightarrow 2\|A\tilde{x}\|_2^2 + \tilde{x}^T C^T \tilde{\lambda} = 0$

Constrained Least Squares, KKT matrix

Proof of sufficient condition

- ▶ Sufficient condition:
 - a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶ **ASSUME**: a.& b. of the theorem 14.2 is satisfied **AND** KKT matrix is **NOT** invertible \Downarrow
- ▶ $\exists \tilde{x} \neq 0, \exists \tilde{\lambda} \neq 0, \begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \tilde{x} \\ \tilde{\lambda} \end{pmatrix} = 0 \Rightarrow 2A^T A \tilde{x} + C^T \tilde{\lambda} = 0 \Downarrow$
- ▶ $x^T (2A^T A \tilde{x} + C^T \tilde{\lambda}) = 0 \Rightarrow 2\|A\tilde{x}\|_2^2 + \tilde{x}^T C^T \tilde{\lambda} = 0$
- ▶ $C\tilde{x} = 0 \equiv \tilde{x}^T C^T = 0$

Constrained Least Squares, KKT matrix

Proof of sufficient condition

- ▶ Sufficient condition:
 - a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶ **ASSUME**: a.& b. of the theorem 14.2 is satisfied **AND** KKT matrix is **NOT** invertible \Downarrow
- ▶ $\exists \tilde{x} \neq 0, \exists \tilde{\lambda} \neq 0, \begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \tilde{x} \\ \tilde{\lambda} \end{pmatrix} = 0 \Rightarrow 2A^T A \tilde{x} + C^T \tilde{\lambda} = 0 \Downarrow$
- ▶ $x^T (2A^T A \tilde{x} + C^T \tilde{\lambda}) = 0 \Rightarrow 2\|A\tilde{x}\|_2^2 + \tilde{x}^T C^T \tilde{\lambda} = 0$
- ▶ $C\tilde{x} = 0 \equiv \tilde{x}^T C^T = 0 \Rightarrow \tilde{x}^T C^T \tilde{\lambda} = 0$

Constrained Least Squares, KKT matrix

Proof of sufficient condition

- ▶ Sufficient condition:
 - a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶ **ASSUME**: a.& b. of the theorem 14.2 is satisfied **AND** KKT matrix is **NOT** invertible \Downarrow
- ▶ $\exists \tilde{x} \neq 0, \exists \tilde{\lambda} \neq 0, \begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \tilde{x} \\ \tilde{\lambda} \end{pmatrix} = 0 \Rightarrow 2A^T A \tilde{x} + C^T \tilde{\lambda} = 0 \Downarrow$
- ▶ $x^T (2A^T A \tilde{x} + C^T \tilde{\lambda}) = 0 \Rightarrow 2\|A\tilde{x}\|_2^2 + \tilde{x}^T C^T \tilde{\lambda} = 0$
- ▶ $C\tilde{x} = 0 \equiv \tilde{x}^T C^T = 0 \Rightarrow \tilde{x}^T C^T \tilde{\lambda} = 0 \Rightarrow \|A\tilde{x}\|_2^2 = 0$

Constrained Least Squares, KKT matrix

Proof of sufficient condition

- ▶ Sufficient condition:
 - a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶ **ASSUME**: a.& b. of the theorem 14.2 is satisfied **AND** KKT matrix is **NOT** invertible \Downarrow
- ▶ $\exists \tilde{x} \neq 0, \exists \tilde{\lambda} \neq 0, \begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \tilde{x} \\ \tilde{\lambda} \end{pmatrix} = 0 \Rightarrow 2A^T A \tilde{x} + C^T \tilde{\lambda} = 0 \Downarrow$
- ▶ $\tilde{x}^T (2A^T A \tilde{x} + C^T \tilde{\lambda}) = 0 \Rightarrow 2\|A\tilde{x}\|_2^2 + \tilde{x}^T C^T \tilde{\lambda} = 0$
- ▶ $C\tilde{x} = 0 \equiv \tilde{x}^T C^T = 0 \Rightarrow \tilde{x}^T C^T \tilde{\lambda} = 0 \Rightarrow \|A\tilde{x}\|_2^2 = 0 \Rightarrow A\tilde{x} = 0 \Downarrow$
- ▶ $\begin{pmatrix} A \\ C \end{pmatrix} \tilde{x} = 0$

Constrained Least Squares, KKT matrix

Proof of sufficient condition

- ▶ Sufficient condition:
 - a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶ **ASSUME**: a.& b. of the theorem 14.2 is satisfied **AND** KKT matrix is **NOT** invertible \Downarrow
- ▶ $\exists \tilde{x} \neq 0, \exists \tilde{\lambda} \neq 0, \begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \tilde{x} \\ \tilde{\lambda} \end{pmatrix} = 0 \Rightarrow 2A^T A \tilde{x} + C^T \tilde{\lambda} = 0 \Downarrow$
- ▶ $\tilde{x}^T (2A^T A \tilde{x} + C^T \tilde{\lambda}) = 0 \Rightarrow 2\|A\tilde{x}\|_2^2 + \tilde{x}^T C^T \tilde{\lambda} = 0$
- ▶ $C\tilde{x} = 0 \equiv \tilde{x}^T C^T = 0 \Rightarrow \tilde{x}^T C^T \tilde{\lambda} = 0 \Rightarrow \|A\tilde{x}\|_2^2 = 0 \Rightarrow A\tilde{x} = 0 \Downarrow$
- ▶ $\begin{pmatrix} A \\ C \end{pmatrix} \tilde{x} = 0$ & + condition b. (linearly independent columns of $\begin{pmatrix} A \\ C \end{pmatrix}$)

Constrained Least Squares, KKT matrix

Proof of sufficient condition

- ▶ Sufficient condition:
 - a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶ **ASSUME:** a.& b. of the theorem 14.2 is satisfied **AND** KKT matrix is **NOT** invertible \Downarrow
- ▶ $\exists \tilde{x} \neq 0, \exists \tilde{\lambda} \neq 0, \begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \tilde{x} \\ \tilde{\lambda} \end{pmatrix} = 0 \Rightarrow 2A^T A \tilde{x} + C^T \tilde{\lambda} = 0 \Downarrow$
- ▶ $\tilde{x}^T (2A^T A \tilde{x} + C^T \tilde{\lambda}) = 0 \Rightarrow 2\|A\tilde{x}\|_2^2 + \tilde{x}^T C^T \tilde{\lambda} = 0$
- ▶ $C\tilde{x} = 0 \equiv \tilde{x}^T C^T = 0 \Rightarrow \tilde{x}^T C^T \tilde{\lambda} = 0 \Rightarrow \|A\tilde{x}\|_2^2 = 0 \Rightarrow A\tilde{x} = 0 \Downarrow$
- ▶ $\begin{pmatrix} A \\ C \end{pmatrix} \tilde{x} = 0$ &+ condition b. (linearly independent columns of $\begin{pmatrix} A \\ C \end{pmatrix}$) \Downarrow
- ▶ **Contradiction:** $\tilde{x} = 0$
- ▶ $\tilde{x} = 0, 2A^T A \tilde{x} + C^T \tilde{\lambda} = 0$

Constrained Least Squares, KKT matrix

Proof of sufficient condition

- ▶ Sufficient condition:
 - a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶ **ASSUME:** a.& b. of the theorem 14.2 is satisfied **AND** KKT matrix is **NOT** invertible \Downarrow
- ▶ $\exists \tilde{x} \neq 0, \exists \tilde{\lambda} \neq 0, \begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \tilde{x} \\ \tilde{\lambda} \end{pmatrix} = 0 \Rightarrow 2A^T A \tilde{x} + C^T \tilde{\lambda} = 0 \Downarrow$
- ▶ $\tilde{x}^T (2A^T A \tilde{x} + C^T \tilde{\lambda}) = 0 \Rightarrow 2\|A\tilde{x}\|_2^2 + \tilde{x}^T C^T \tilde{\lambda} = 0$
- ▶ $C\tilde{x} = 0 \equiv \tilde{x}^T C^T = 0 \Rightarrow \tilde{x}^T C^T \tilde{\lambda} = 0 \Rightarrow \|A\tilde{x}\|_2^2 = 0 \Rightarrow A\tilde{x} = 0 \Downarrow$
- ▶ $\begin{pmatrix} A \\ C \end{pmatrix} \tilde{x} = 0$ &+ condition b. (linearly independent columns of $\begin{pmatrix} A \\ C \end{pmatrix}$) \Downarrow
- ▶ **Contradiction:** $\tilde{x} = 0$
- ▶ $\tilde{x} = 0, 2A^T A \tilde{x} + C^T \tilde{\lambda} = 0 \Rightarrow C^T \tilde{\lambda} = 0$

Constrained Least Squares, KKT matrix

Proof of sufficient condition

- ▶ Sufficient condition:
 - a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶ **ASSUME:** a.& b. of the theorem 14.2 is satisfied **AND** KKT matrix is **NOT** invertible \Downarrow
- ▶ $\exists \tilde{x} \neq 0, \exists \tilde{\lambda} \neq 0, \begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \tilde{x} \\ \tilde{\lambda} \end{pmatrix} = 0 \Rightarrow 2A^T A \tilde{x} + C^T \tilde{\lambda} = 0 \Downarrow$
- ▶ $\tilde{x}^T (2A^T A \tilde{x} + C^T \tilde{\lambda}) = 0 \Rightarrow 2\|A\tilde{x}\|_2^2 + \tilde{x}^T C^T \tilde{\lambda} = 0$
- ▶ $C\tilde{x} = 0 \equiv \tilde{x}^T C^T = 0 \Rightarrow \tilde{x}^T C^T \tilde{\lambda} = 0 \Rightarrow \|A\tilde{x}\|_2^2 = 0 \Rightarrow A\tilde{x} = 0 \Downarrow$
- ▶ $\begin{pmatrix} A \\ C \end{pmatrix} \tilde{x} = 0$ &+ condition b. (linearly independent columns of $\begin{pmatrix} A \\ C \end{pmatrix}$) \Downarrow
- ▶ **Contradiction:** $\tilde{x} = 0$
- ▶ $\tilde{x} = 0, 2A^T A \tilde{x} + C^T \tilde{\lambda} = 0 \Rightarrow C^T \tilde{\lambda} = 0$ &+ condition a. (linearly independent rows of C)

Constrained Least Squares, KKT matrix

Proof of sufficient condition

- ▶ Sufficient condition:
 - a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶ **ASSUME:** a.& b. of the theorem 14.2 is satisfied **AND** KKT matrix is **NOT** invertible \Downarrow
- ▶ $\exists \tilde{x} \neq 0, \exists \tilde{\lambda} \neq 0, \begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \tilde{x} \\ \tilde{\lambda} \end{pmatrix} = 0 \Rightarrow 2A^T A \tilde{x} + C^T \tilde{\lambda} = 0 \Downarrow$
- ▶ $\tilde{x}^T (2A^T A \tilde{x} + C^T \tilde{\lambda}) = 0 \Rightarrow 2\|A\tilde{x}\|_2^2 + \tilde{x}^T C^T \tilde{\lambda} = 0$
- ▶ $C\tilde{x} = 0 \equiv \tilde{x}^T C^T = 0 \Rightarrow \tilde{x}^T C^T \tilde{\lambda} = 0 \Rightarrow \|A\tilde{x}\|_2^2 = 0 \Rightarrow A\tilde{x} = 0 \Downarrow$
- ▶ $\begin{pmatrix} A \\ C \end{pmatrix} \tilde{x} = 0$ &+ condition b. (linearly independent columns of $\begin{pmatrix} A \\ C \end{pmatrix}$) \Downarrow
- ▶ **Contradiction:** $\tilde{x} = 0$
- ▶ $\tilde{x} = 0, 2A^T A \tilde{x} + C^T \tilde{\lambda} = 0 \Rightarrow C^T \tilde{\lambda} = 0$ &+ condition a. (linearly independent rows of C) \Downarrow
- ▶ **Contradiction:** $\tilde{\lambda} = 0$

Constrained Least Squares, KKT matrix

Solution of KKT equation

► KKT equation:
$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} = \begin{pmatrix} 2A^T b \\ d \end{pmatrix}$$

Constrained Least Squares, KKT matrix

Solution of KKT equation

- ▶ KKT equation:
$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} = \begin{pmatrix} 2A^T b \\ d \end{pmatrix}$$
- ▶ a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible

Constrained Least Squares, KKT matrix

Solution of KKT equation

- ▶ KKT equation:
$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} = \begin{pmatrix} 2A^T b \\ d \end{pmatrix}$$
- ▶ a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶
$$\begin{pmatrix} \hat{x} \\ \hat{\lambda} \end{pmatrix} = \begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix}^{-1} \begin{pmatrix} 2A^T b \\ d \end{pmatrix}$$

Constrained Least Squares, KKT matrix

Solution of KKT equation

- ▶ KKT equation:
$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} = \begin{pmatrix} 2A^T b \\ d \end{pmatrix}$$
- ▶ a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶
$$\begin{pmatrix} \hat{x} \\ \hat{\lambda} \end{pmatrix} = \begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix}^{-1} \begin{pmatrix} 2A^T b \\ d \end{pmatrix}$$

Algorithm 14.3

- ▶ Compute $H = 2A^T A$ and $f = 2A^T b$

Constrained Least Squares, KKT matrix

Solution of KKT equation

- ▶ KKT equation:
$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} = \begin{pmatrix} 2A^T b \\ d \end{pmatrix}$$
- ▶ a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶
$$\begin{pmatrix} \hat{x} \\ \hat{\lambda} \end{pmatrix} = \begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix}^{-1} \begin{pmatrix} 2A^T b \\ d \end{pmatrix}$$

Algorithm 14.3

- ▶ Compute $H = 2A^T A$ and $f = 2A^T b$
- ▶ Solve
$$\begin{pmatrix} H & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} = \begin{pmatrix} f \\ d \end{pmatrix}$$

Constrained Least Squares, KKT matrix

Solution of KKT equation

- ▶ KKT equation:
$$\begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} = \begin{pmatrix} 2A^T b \\ d \end{pmatrix}$$
- ▶ a. & b. of the theorem 14.2 are satisfied \Rightarrow KKT matrix is invertible
- ▶
$$\begin{pmatrix} \hat{x} \\ \hat{\lambda} \end{pmatrix} = \begin{pmatrix} 2A^T A & C^T \\ C & 0 \end{pmatrix}^{-1} \begin{pmatrix} 2A^T b \\ d \end{pmatrix}$$

Algorithm 14.3

- ▶ Compute $H = 2A^T A$ and $f = 2A^T b$
- ▶ Solve
$$\begin{pmatrix} H & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} = \begin{pmatrix} f \\ d \end{pmatrix}$$

Remark 14.4

Computing $A^T A$ is not practical: expensive, may introduce errors

QR factorization for constrained least squares

Different Lagrangian \Rightarrow different KKT equation

$$L(x, \lambda) = (Ax - b, Ax - b) + 2(Cx - d, \lambda)$$

QR factorization for constrained least squares

Different Lagrangian \Rightarrow different KKT equation

$$L(x, \lambda) = (Ax - b, Ax - b) + 2(Cx - d, \lambda)$$

$$(A^T Ax - A^T b) + C^T \lambda = 0$$

$$Cx = d$$

QR factorization for constrained least squares

Different Lagrangian \Rightarrow different KKT equation

$$L(x, \lambda) = (Ax - b, Ax - b) + 2(Cx - d, \lambda)$$

$$(A^T Ax - A^T b) + C^T \lambda = 0$$

$$Cx = d$$

$$\begin{pmatrix} A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$

QR factorization for constrained least squares

Different Lagrangian \Rightarrow different KKT equation

$$L(x, \lambda) = (Ax - b, Ax - b) + 2(Cx - d, \lambda)$$

$$(A^T Ax - A^T b) + C^T \lambda = 0$$

$$Cx = d$$

$$\begin{pmatrix} A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$

$$L(x, \lambda) = \frac{1}{2}(Ax - b, Ax - b) + (Cx - d, \lambda), KKT = ?$$

QR factorization for constrained least squares

Different Lagrangian \Rightarrow different KKT equation

$$L(x, \lambda) = (Ax - b, Ax - b) + 2(Cx - d, \lambda)$$

$$(A^T Ax - A^T b) + C^T \lambda = 0$$

$$Cx = d$$

$$\begin{pmatrix} A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \lambda \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$

$$L(x, \lambda) = \frac{1}{2}(Ax - b, Ax - b) + (Cx - d, \lambda), KKT = ?$$

$$L(x, \lambda) = \alpha(Ax - b, Ax - b) + \beta(Cx - d, \lambda), KKT = ?$$

QR factorization for constrained least squares

Change of variables, equivalent KKT equation

$$\begin{aligned}(A^T A x - A^T b) + C^T \lambda &= 0 \\ Cx &= d\end{aligned}$$

QR factorization for constrained least squares

Change of variables, equivalent KKT equation

$$\begin{aligned}(A^T A x - A^T b) + C^T \lambda &= 0 \\ Cx &= d\end{aligned}$$

► $\lambda = \xi + d,$

QR factorization for constrained least squares

Change of variables, equivalent KKT equation

$$\begin{aligned}(A^T A x - A^T b) + C^T \lambda &= 0 \\ Cx &= d\end{aligned}$$

► $\lambda = \xi + d, Cx = d$

QR factorization for constrained least squares

Change of variables, equivalent KKT equation

$$\begin{aligned}(A^T A x - A^T b) + C^T \lambda &= 0 \\ Cx &= d\end{aligned}$$

► $\lambda = \xi + d$, $Cx = d \Rightarrow \lambda = \xi + Cx$

QR factorization for constrained least squares

Change of variables, equivalent KKT equation

$$\begin{aligned}(A^T A x - A^T b) + C^T \lambda &= 0 \\ Cx &= d\end{aligned}$$

- ▶ $\lambda = \xi + d$, $Cx = d \Rightarrow \lambda = \xi + Cx$
- ▶ $(A^T A x - A^T b) + C^T (\xi + d) = (A^T A x - A^T b) + C^T (\xi + Cx)$

QR factorization for constrained least squares

Change of variables, equivalent KKT equation

$$\begin{aligned}(A^T A x - A^T b) + C^T \lambda &= 0 \\ Cx &= d\end{aligned}$$

- ▶ $\lambda = \xi + d$, $Cx = d \Rightarrow \lambda = \xi + Cx$
- ▶ $(A^T A x - A^T b) + C^T (\xi + d) = (A^T A x - A^T b) + C^T (\xi + Cx)$
- ▶ $(A^T A x - A^T b) + C^T (\xi + Cx) = (A^T A + C^T C)x - A^T b + C^T \xi$

QR factorization for constrained least squares

Change of variables, equivalent KKT equation

$$\begin{aligned}(A^T A x - A^T b) + C^T \lambda &= 0 \\ Cx &= d\end{aligned}$$

- ▶ $\lambda = \xi + d$, $Cx = d \Rightarrow \lambda = \xi + Cx$
- ▶ $(A^T A x - A^T b) + C^T (\xi + d) = (A^T A x - A^T b) + C^T (\xi + Cx)$
- ▶ $(A^T A x - A^T b) + C^T (\xi + Cx) = (A^T A + C^T C)x - A^T b + C^T \xi$
- ▶

$$\begin{pmatrix} A^T A + C^T C & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \xi \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$

QR factorization for constrained least squares

QR factorization of $\begin{pmatrix} A \\ C \end{pmatrix}$, equivalent KKT equation

- Assumption b.

QR factorization for constrained least squares

QR factorization of $\begin{pmatrix} A \\ C \end{pmatrix}$, equivalent KKT equation

► Assumption b. $\Rightarrow \begin{pmatrix} A \\ C \end{pmatrix} = QR,$

QR factorization for constrained least squares

QR factorization of $\begin{pmatrix} A \\ C \end{pmatrix}$, equivalent KKT equation

► Assumption b. $\Rightarrow \begin{pmatrix} A \\ C \end{pmatrix} = QR, \Rightarrow \begin{pmatrix} A \\ C \end{pmatrix} = \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix} R$

QR factorization for constrained least squares

QR factorization of $\begin{pmatrix} A \\ C \end{pmatrix}$, equivalent KKT equation

- ▶ Assumption b. $\Rightarrow \begin{pmatrix} A \\ C \end{pmatrix} = QR, \Rightarrow \begin{pmatrix} A \\ C \end{pmatrix} = \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix} R$
- ▶ $A^T = R^T Q_1^T, C^T = R^T Q_2^T, A^T A = R^T R, C^T C = R^T R$

QR factorization for constrained least squares

QR factorization of $\begin{pmatrix} A \\ C \end{pmatrix}$, equivalent KKT equation

► Assumption b. $\Rightarrow \begin{pmatrix} A \\ C \end{pmatrix} = QR, \Rightarrow \begin{pmatrix} A \\ C \end{pmatrix} = \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix} R$

► $A^T = R^T Q_1^T, C^T = R^T Q_2^T, A^T A = R^T R, C^T C = R^T R$

►
$$\begin{pmatrix} A^T A + C^T C & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \xi \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$

QR factorization for constrained least squares

QR factorization of $\begin{pmatrix} A \\ C \end{pmatrix}$, equivalent KKT equation

► Assumption b. $\Rightarrow \begin{pmatrix} A \\ C \end{pmatrix} = QR, \Rightarrow \begin{pmatrix} A \\ C \end{pmatrix} = \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix} R$

► $A^T = R^T Q_1^T, C^T = R^T Q_2^T, A^T A = R^T R, C^T C = R^T R$

►
$$\begin{pmatrix} A^T A + C^T C & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \xi \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$

►
$$\begin{pmatrix} 2R^T R & R^T Q_2^T \\ Q_2 R & 0 \end{pmatrix} \begin{pmatrix} x \\ \xi \end{pmatrix} = \begin{pmatrix} R^T Q_1^T b \\ d \end{pmatrix}$$

QR factorization for constrained least squares

QR factorization of $\begin{pmatrix} A \\ C \end{pmatrix}$, equivalent KKT equation

► Assumption b. $\Rightarrow \begin{pmatrix} A \\ C \end{pmatrix} = QR, \Rightarrow \begin{pmatrix} A \\ C \end{pmatrix} = \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix} R$

► $A^T = R^T Q_1^T, C^T = R^T Q_2^T, A^T A = R^T R, C^T C = R^T R$



$$\begin{pmatrix} A^T A + C^T C & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \xi \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$



$$\begin{pmatrix} 2R^T R & R^T Q_2^T \\ Q_2 R & 0 \end{pmatrix} \begin{pmatrix} x \\ \xi \end{pmatrix} = \begin{pmatrix} R^T Q_1^T b \\ d \end{pmatrix}$$



$$\begin{pmatrix} 2R & Q_2^T \\ Q_2 R & 0 \end{pmatrix} \begin{pmatrix} x \\ \xi \end{pmatrix} = \begin{pmatrix} Q_1^T b \\ d \end{pmatrix}$$

QR factorization for constrained least squares

QR factorization of $\begin{pmatrix} A \\ C \end{pmatrix}$, equivalent KKT equation

► Assumption b. $\Rightarrow \begin{pmatrix} A \\ C \end{pmatrix} = QR, \Rightarrow \begin{pmatrix} A \\ C \end{pmatrix} = \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix} R$

► $A^T = R^T Q_1^T, C^T = R^T Q_2^T, A^T A = R^T R, C^T C = R^T R$

►
$$\begin{pmatrix} A^T A + C^T C & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \xi \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$

►
$$\begin{pmatrix} 2R^T R & R^T Q_2^T \\ Q_2 R & 0 \end{pmatrix} \begin{pmatrix} x \\ \xi \end{pmatrix} = \begin{pmatrix} R^T Q_1^T b \\ d \end{pmatrix}$$

►
$$\begin{pmatrix} 2R & Q_2^T \\ Q_2 R & 0 \end{pmatrix} \begin{pmatrix} x \\ \xi \end{pmatrix} = \begin{pmatrix} Q_1^T b \\ d \end{pmatrix}$$

► New variable: $y = Rx \Rightarrow$

QR factorization for constrained least squares

QR factorization of $\begin{pmatrix} A \\ C \end{pmatrix}$, equivalent KKT equation

► Assumption b. $\Rightarrow \begin{pmatrix} A \\ C \end{pmatrix} = QR, \Rightarrow \begin{pmatrix} A \\ C \end{pmatrix} = \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix} R$

► $A^T = R^T Q_1^T, C^T = R^T Q_2^T, A^T A = R^T R, C^T C = R^T R$

►
$$\begin{pmatrix} A^T A + C^T C & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} x \\ \xi \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$

►
$$\begin{pmatrix} 2R^T R & R^T Q_2^T \\ Q_2 R & 0 \end{pmatrix} \begin{pmatrix} x \\ \xi \end{pmatrix} = \begin{pmatrix} R^T Q_1^T b \\ d \end{pmatrix}$$

►
$$\begin{pmatrix} 2R & Q_2^T \\ Q_2 R & 0 \end{pmatrix} \begin{pmatrix} x \\ \xi \end{pmatrix} = \begin{pmatrix} Q_1^T b \\ d \end{pmatrix}$$

► New variable: $y = Rx \Rightarrow \begin{pmatrix} 2I & Q_2^T \\ Q_2 & 0 \end{pmatrix} \begin{pmatrix} y \\ \xi \end{pmatrix} = \begin{pmatrix} Q_1^T b \\ d \end{pmatrix}$

QR factorization for constrained least squares

Proposition 14.5

Assumption a. is valid and Q_2 has linearly independent rows

Proof.

- ▶ Assume: Q_2 has linearly dependent rows

QR factorization for constrained least squares

Proposition 14.5

Assumption a. is valid and Q_2 has linearly independent rows

Proof.

- ▶ Assume: Q_2 has linearly dependent rows
- ▶ \Downarrow
- ▶ $\exists \tilde{\xi} \neq 0, Q_2^T \tilde{\xi} = 0$

QR factorization for constrained least squares

Proposition 14.5

Assumption a. is valid and Q_2 has linearly independent rows

Proof.

- ▶ Assume: Q_2 has linearly dependent rows
- ▶ \Downarrow
- ▶ $\exists \tilde{\xi} \neq 0, Q_2^T \tilde{\xi} = 0$
- ▶ $C = Q_2 R$

QR factorization for constrained least squares

Proposition 14.5

Assumption a. is valid and Q_2 has linearly independent rows

Proof.

- ▶ Assume: Q_2 has linearly dependent rows
- ▶ \Downarrow
- ▶ $\exists \tilde{\xi} \neq 0, Q_2^T \tilde{\xi} = 0$
- ▶ $C = Q_2 R \Rightarrow Q_2 = CR^{-1}$

QR factorization for constrained least squares

Proposition 14.5

Assumption a. is valid and Q_2 has linearly independent rows

Proof.

- ▶ Assume: Q_2 has linearly dependent rows
- ▶ \Downarrow
- ▶ $\exists \tilde{\xi} \neq 0, Q_2^T \tilde{\xi} = 0$
- ▶ $C = Q_2 R \Rightarrow Q_2 = CR^{-1}$
- ▶ $Q_2^T \tilde{\xi} = (CR^{-1})^T \tilde{\xi} = R^{-T} C^T \tilde{\xi}$

QR factorization for constrained least squares

Proposition 14.5

Assumption a. is valid and Q_2 has linearly independent rows

Proof.

- ▶ Assume: Q_2 has linearly dependent rows
- ▶ \Downarrow
- ▶ $\exists \tilde{\xi} \neq 0, Q_2^T \tilde{\xi} = 0$
- ▶ $C = Q_2 R \Rightarrow Q_2 = CR^{-1}$
- ▶ $Q_2^T \tilde{\xi} = (CR^{-1})^T \tilde{\xi} = R^{-T} C^T \tilde{\xi}$
- ▶ \Downarrow
- ▶ $C^T \tilde{\xi} = 0$

QR factorization for constrained least squares

Proposition 14.5

Assumption a. is valid and Q_2 has linearly independent rows

Proof.

- ▶ Assume: Q_2 has linearly dependent rows
- ▶ \Downarrow
- ▶ $\exists \tilde{\xi} \neq 0, Q_2^T \tilde{\xi} = 0$
- ▶ $C = Q_2 R \Rightarrow Q_2 = CR^{-1}$
- ▶ $Q_2^T \tilde{\xi} = (CR^{-1})^T \tilde{\xi} = R^{-T} C^T \tilde{\xi}$
- ▶ \Downarrow
- ▶ $C^T \tilde{\xi} = 0$
- ▶ C^T has linearly independent columns

QR factorization for constrained least squares

Proposition 14.5

Assumption a. is valid and Q_2 has linearly independent rows

Proof.

- ▶ Assume: Q_2 has linearly dependent rows
- ▶ \Downarrow
- ▶ $\exists \tilde{\xi} \neq 0, Q_2^T \tilde{\xi} = 0$
- ▶ $C = Q_2 R \Rightarrow Q_2 = CR^{-1}$
- ▶ $Q_2^T \tilde{\xi} = (CR^{-1})^T \tilde{\xi} = R^{-T} C^T \tilde{\xi}$
- ▶ \Downarrow
- ▶ $C^T \tilde{\xi} = 0$
- ▶ C^T has linearly independent columns
- ▶ \Downarrow
- ▶ Contradiction: $\xi = 0$



QR factorization for constrained least squares

QR factorization of Q_2 , equivalent KKT equation

QR factorization for constrained least squares

QR factorization of Q_2 , equivalent KKT equation

- ▶ Q_2 has linearly independent rows $\Rightarrow Q_2^T = \tilde{Q}\tilde{R}$

QR factorization for constrained least squares

QR factorization of Q_2 , equivalent KKT equation

- ▶ Q_2 has linearly independent rows $\Rightarrow Q_2^T = \tilde{Q}\tilde{R}$
- ▶ KKT equation
$$\begin{pmatrix} 2I & Q_2^T \\ Q_2 & 0 \end{pmatrix} \begin{pmatrix} y \\ \xi \end{pmatrix} = \begin{pmatrix} Q_1^T b \\ d \end{pmatrix}$$

QR factorization for constrained least squares

QR factorization of Q_2 , equivalent KKT equation

- ▶ Q_2 has linearly independent rows $\Rightarrow Q_2^T = \tilde{Q}\tilde{R}$
- ▶ KKT equation $\begin{pmatrix} 2I & Q_2^T \\ Q_2 & 0 \end{pmatrix} \begin{pmatrix} y \\ \xi \end{pmatrix} = \begin{pmatrix} Q_1^T b \\ d \end{pmatrix}$
- ▶ $Q_2 y = d, 2y + Q_2^T \xi = Q_1^T b$

QR factorization for constrained least squares

QR factorization of Q_2 , equivalent KKT equation

- ▶ Q_2 has linearly independent rows $\Rightarrow Q_2^T = \tilde{Q}\tilde{R}$
- ▶ KKT equation $\begin{pmatrix} 2I & Q_2^T \\ Q_2 & 0 \end{pmatrix} \begin{pmatrix} y \\ \xi \end{pmatrix} = \begin{pmatrix} Q_1^T b \\ d \end{pmatrix}$
- ▶ $Q_2 y = d, 2y + Q_2^T \xi = Q_1^T b$
- ▶ $y = 0.5(Q_1^T b - Q_2^T \xi)$

QR factorization for constrained least squares

QR factorization of Q_2 , equivalent KKT equation

- ▶ Q_2 has linearly independent rows $\Rightarrow Q_2^T = \tilde{Q}\tilde{R}$
- ▶ KKT equation $\begin{pmatrix} 2I & Q_2^T \\ Q_2 & 0 \end{pmatrix} \begin{pmatrix} y \\ \xi \end{pmatrix} = \begin{pmatrix} Q_1^T b \\ d \end{pmatrix}$
- ▶ $Q_2 y = d, 2y + Q_2^T \xi = Q_1^T b$
- ▶ $y = 0.5(Q_1^T b - Q_2^T \xi)$
- ▶ $Q_2 y = 0.5(Q_2 Q_1^T b - Q_2 Q_2^T \xi)$

QR factorization for constrained least squares

QR factorization of Q_2 , equivalent KKT equation

- ▶ Q_2 has linearly independent rows $\Rightarrow Q_2^T = \tilde{Q}\tilde{R}$
- ▶ KKT equation $\begin{pmatrix} 2I & Q_2^T \\ Q_2 & 0 \end{pmatrix} \begin{pmatrix} y \\ \xi \end{pmatrix} = \begin{pmatrix} Q_1^T b \\ d \end{pmatrix}$
- ▶ $Q_2 y = d, 2y + Q_2^T \xi = Q_1^T b$
- ▶ $y = 0.5(Q_1^T b - Q_2^T \xi)$
- ▶ $Q_2 y = 0.5(Q_2 Q_1^T b - Q_2 Q_2^T \xi)$
- ▶ $0.5(Q_2 Q_1^T b - Q_2 Q_2^T \xi) = d$

QR factorization for constrained least squares

QR factorization of Q_2 , equivalent KKT equation

- ▶ Q_2 has linearly independent rows $\Rightarrow Q_2^T = \tilde{Q}\tilde{R}$
- ▶ KKT equation
$$\begin{pmatrix} 2I & Q_2^T \\ Q_2 & 0 \end{pmatrix} \begin{pmatrix} y \\ \xi \end{pmatrix} = \begin{pmatrix} Q_1^T b \\ d \end{pmatrix}$$
- ▶ $Q_2 y = d, 2y + Q_2^T \xi = Q_1^T b$
- ▶ $y = 0.5(Q_1^T b - Q_2^T \xi)$
- ▶ $Q_2 y = 0.5(Q_2 Q_1^T b - Q_2 Q_2^T \xi)$
- ▶ $0.5(Q_2 Q_1^T b - Q_2 Q_2^T \xi) = d$
- ▶ $Q_2 Q_2^T \xi = Q_2 Q_1^T b - 2d$

QR factorization for constrained least squares

QR factorization of Q_2 , equivalent KKT equation

- ▶ Q_2 has linearly independent rows $\Rightarrow Q_2^T = \tilde{Q}\tilde{R}$
- ▶ KKT equation $\begin{pmatrix} 2I & Q_2^T \\ Q_2 & 0 \end{pmatrix} \begin{pmatrix} y \\ \xi \end{pmatrix} = \begin{pmatrix} Q_1^T b \\ d \end{pmatrix}$
- ▶ $Q_2 y = d, 2y + Q_2^T \xi = Q_1^T b$
- ▶ $y = 0.5(Q_1^T b - Q_2^T \xi)$
- ▶ $Q_2 y = 0.5(Q_2 Q_1^T b - Q_2 Q_2^T \xi)$
- ▶ $0.5(Q_2 Q_1^T b - Q_2 Q_2^T \xi) = d$
- ▶ $Q_2 Q_2^T \xi = Q_2 Q_1^T b - 2d$
- ▶ $Q_2 Q_2^T = \tilde{R}^T \tilde{R}$

QR factorization for constrained least squares

QR factorization of Q_2 , equivalent KKT equation

- ▶ Q_2 has linearly independent rows $\Rightarrow Q_2^T = \tilde{Q}\tilde{R}$
- ▶ KKT equation $\begin{pmatrix} 2I & Q_2^T \\ Q_2 & 0 \end{pmatrix} \begin{pmatrix} y \\ \xi \end{pmatrix} = \begin{pmatrix} Q_1^T b \\ d \end{pmatrix}$
- ▶ $Q_2 y = d, 2y + Q_2^T \xi = Q_1^T b$
- ▶ $y = 0.5(Q_1^T b - Q_2^T \xi)$
- ▶ $Q_2 y = 0.5(Q_2 Q_1^T b - Q_2 Q_2^T \xi)$
- ▶ $0.5(Q_2 Q_1^T b - Q_2 Q_2^T \xi) = d$
- ▶ $Q_2 Q_2^T \xi = Q_2 Q_1^T b - 2d$
- ▶ $Q_2 Q_2^T = \tilde{R}^T \tilde{R}$
- ▶ $\tilde{R}^T \tilde{R} \xi = \tilde{R}^T \tilde{Q}^T Q_1^T b - 2d$

QR factorization for constrained least squares

QR factorization of Q_2 , equivalent KKT equation

► Q_2 has linearly independent rows $\Rightarrow Q_2^T = \tilde{Q}\tilde{R}$

► KKT equation $\begin{pmatrix} 2I & Q_2^T \\ Q_2 & 0 \end{pmatrix} \begin{pmatrix} y \\ \xi \end{pmatrix} = \begin{pmatrix} Q_1^T b \\ d \end{pmatrix}$

► $Q_2 y = d, 2y + Q_2^T \xi = Q_1^T b$

► $y = 0.5(Q_1^T b - Q_2^T \xi)$

► $Q_2 y = 0.5(Q_2 Q_1^T b - Q_2 Q_2^T \xi)$

► $0.5(Q_2 Q_1^T b - Q_2 Q_2^T \xi) = d$

► $Q_2 Q_2^T \xi = Q_2 Q_1^T b - 2d$

► $Q_2 Q_2^T = \tilde{R}^T \tilde{R}$

► $\tilde{R}^T \tilde{R} \xi = \tilde{R}^T \tilde{Q}^T Q_1^T b - 2d$

►

$$\tilde{R} \xi = \tilde{Q}^T Q_1^T b - 2\tilde{R}^{-T} d$$

QR factorization for constrained least squares

QR factorization of Q_2 , equivalent KKT equation

► Q_2 has linearly independent rows $\Rightarrow Q_2^T = \tilde{Q}\tilde{R}$

► KKT equation $\begin{pmatrix} 2I & Q_2^T \\ Q_2 & 0 \end{pmatrix} \begin{pmatrix} y \\ \xi \end{pmatrix} = \begin{pmatrix} Q_1^T b \\ d \end{pmatrix}$

► $Q_2 y = d, 2y + Q_2^T \xi = Q_1^T b$

► $y = 0.5(Q_1^T b - Q_2^T \xi)$

► $Q_2 y = 0.5(Q_2 Q_1^T b - Q_2 Q_2^T \xi)$

► $0.5(Q_2 Q_1^T b - Q_2 Q_2^T \xi) = d$

► $Q_2 Q_2^T \xi = Q_2 Q_1^T b - 2d$

► $Q_2 Q_2^T = \tilde{R}^T \tilde{R}$

► $\tilde{R}^T \tilde{R} \xi = \tilde{R}^T \tilde{Q}^T Q_1^T b - 2d$

►

$$\tilde{R} \xi = \tilde{Q}^T Q_1^T b - 2\tilde{R}^{-T} d$$

► Equation for x : $Rx = y$

QR factorization for constrained least squares

Equivalent KKT equation

QR factorization for constrained least squares

Equivalent KKT equation

► KKT equation
$$\begin{pmatrix} A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \hat{x} \\ \lambda \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$

QR factorization for constrained least squares

Equivalent KKT equation

► KKT equation
$$\begin{pmatrix} A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \hat{x} \\ \lambda \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$

► Equivalent equation

$$\tilde{R}\xi = \tilde{Q}^T Q_1^T b - 2\tilde{R}^{-T} d,$$

$$y = 0.5(Q_1^T b - Q_2^T \xi),$$

$$R\hat{x} = y$$

Algorithm 14.6

QR factorization for constrained least squares

Equivalent KKT equation

► KKT equation
$$\begin{pmatrix} A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \hat{x} \\ \lambda \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$

► Equivalent equation

$$\tilde{R}\xi = \tilde{Q}^T Q_1^T b - 2\tilde{R}^{-T} d,$$

$$y = 0.5(Q_1^T b - Q_2^T \xi),$$

$$R\hat{x} = y$$

Algorithm 14.6

1. Compute QR factorization:

$$\begin{pmatrix} A \\ C \end{pmatrix} = \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix} R$$

QR factorization for constrained least squares

Equivalent KKT equation

► KKT equation
$$\begin{pmatrix} A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \hat{x} \\ \lambda \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$

► Equivalent equation

$$\tilde{R}\xi = \tilde{Q}^T Q_1^T b - 2\tilde{R}^{-T} d,$$

$$y = 0.5(Q_1^T b - Q_2^T \xi),$$

$$R\hat{x} = y$$

Algorithm 14.6

1. Compute QR factorization:

2. Compute QR factorization: $Q_2^T = \tilde{Q}\tilde{R}$

$$\begin{pmatrix} A \\ C \end{pmatrix} = \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix} R$$

QR factorization for constrained least squares

Equivalent KKT equation

► KKT equation
$$\begin{pmatrix} A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \hat{x} \\ \lambda \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$

► Equivalent equation

$$\tilde{R}\xi = \tilde{Q}^T Q_1^T b - 2\tilde{R}^{-T} d,$$

$$y = 0.5(Q_1^T b - Q_2^T \xi),$$

$$R\hat{x} = y$$

Algorithm 14.6

1. Compute QR factorization:
2. Compute QR factorization: $Q_2^T = \tilde{Q}\tilde{R}$
3. Solve $\tilde{R}^T \eta = d$

$$\begin{pmatrix} A \\ C \end{pmatrix} = \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix} R$$

QR factorization for constrained least squares

Equivalent KKT equation

► KKT equation
$$\begin{pmatrix} A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \hat{x} \\ \lambda \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$

► Equivalent equation

$$\tilde{R}\xi = \tilde{Q}^T Q_1^T b - 2\tilde{R}^{-T} d,$$

$$y = 0.5(Q_1^T b - Q_2^T \xi),$$

$$R\hat{x} = y$$

Algorithm 14.6

1. Compute QR factorization:
2. Compute QR factorization: $Q_2^T = \tilde{Q}\tilde{R}$
3. Solve $\tilde{R}^T \eta = d$
4. Compute $\tilde{\eta} = \tilde{Q}^T Q_1^T b - 2\eta$

$$\begin{pmatrix} A \\ C \end{pmatrix} = \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix} R$$

QR factorization for constrained least squares

Equivalent KKT equation

► KKT equation
$$\begin{pmatrix} A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \hat{x} \\ \lambda \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$

► Equivalent equation

$$\tilde{R}\xi = \tilde{Q}^T Q_1^T b - 2\tilde{R}^{-T} d,$$

$$y = 0.5(Q_1^T b - Q_2^T \xi),$$

$$R\hat{x} = y$$

Algorithm 14.6

1. Compute QR factorization:
2. Compute QR factorization: $Q_2^T = \tilde{Q}\tilde{R}$
3. Solve $\tilde{R}^T \eta = d$
4. Compute $\tilde{\eta} = \tilde{Q}^T Q_1^T b - 2\eta$
5. Solve $\tilde{R}\xi = \tilde{\eta}$

$$\begin{pmatrix} A \\ C \end{pmatrix} = \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix} R$$

QR factorization for constrained least squares

Equivalent KKT equation

► KKT equation
$$\begin{pmatrix} A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \hat{x} \\ \lambda \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$

► Equivalent equation

$$\tilde{R}\xi = \tilde{Q}^T Q_1^T b - 2\tilde{R}^{-T} d,$$

$$y = 0.5(Q_1^T b - Q_2^T \xi),$$

$$R\hat{x} = y$$

Algorithm 14.6

1. Compute QR factorization:

$$\begin{pmatrix} A \\ C \end{pmatrix} = \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix} R$$

2. Compute QR factorization: $Q_2^T = \tilde{Q}\tilde{R}$

3. Solve $\tilde{R}^T \eta = d$

4. Compute $\tilde{\eta} = \tilde{Q}^T Q_1^T b - 2\eta$

5. Solve $\tilde{R}\xi = \tilde{\eta}$

6. Compute $y = 0.5(Q_1^T b - Q_2^T \xi)$

QR factorization for constrained least squares

Equivalent KKT equation

► KKT equation
$$\begin{pmatrix} A^T A & C^T \\ C & 0 \end{pmatrix} \begin{pmatrix} \hat{x} \\ \lambda \end{pmatrix} = \begin{pmatrix} A^T b \\ d \end{pmatrix}$$

► Equivalent equation

$$\tilde{R}\xi = \tilde{Q}^T Q_1^T b - 2\tilde{R}^{-T} d,$$

$$y = 0.5(Q_1^T b - Q_2^T \xi),$$

$$R\hat{x} = y$$

Algorithm 14.6

1. Compute QR factorization:

$$\begin{pmatrix} A \\ C \end{pmatrix} = \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix} R$$

2. Compute QR factorization: $Q_2^T = \tilde{Q}\tilde{R}$

3. Solve $\tilde{R}^T \eta = d$

4. Compute $\tilde{\eta} = \tilde{Q}^T Q_1^T b - 2\eta$

5. Solve $\tilde{R}\xi = \tilde{\eta}$

6. Compute $y = 0.5(Q_1^T b - Q_2^T \xi)$

7. Find solution of constrained least squares problem: solve $R\hat{x} = y$

Hausholder transformation, full QR factorization

Definition 14.7

$$H = I - \frac{2vv^T}{v^Tv}$$

Hausholder transformation, full QR factorization

Definition 14.7

$$H = I - \frac{2vv^T}{v^Tv}$$

H - Hausholder matrix

Hausholder transformation, full QR factorization

Definition 14.7

$$H = I - \frac{2vv^T}{v^Tv}$$

H - Hausholder matrix

v - Hausholder vector

Hausholder transformation, full QR factorization

Definition 14.7

$$H = I - \frac{2vv^T}{v^Tv}$$

H - Hausholder matrix

v - Hausholder vector

Alston Hausholder (1904-1993)

American mathematician, celebrated numerical analyst, former director of Mathematics and Computer Science Division of Oak Ridge National Laboratory.

Hausholder transformation, full QR factorization

Hausholder transformation

Hausholder transformation, full QR factorization

Hausholder transformation

- ▶ v - unit vector, $\|v\|_2 = 1$, normal to hyperplane

Hausholder transformation, full QR factorization

Hausholder transformation

- ▶ v - unit vector, $\|v\|_2 = 1$, normal to hyperplane
- ▶ $v^T x = \|v\| \|x\| \cos(\hat{v}\hat{x})$

Hausholder transformation, full QR factorization

Hausholder transformation

- ▶ v - unit vector, $\|v\|_2 = 1$, normal to hyperplane
- ▶ $v^T x = \|v\| \|x\| \cos(\hat{v}x)$
- ▶ $\hat{v}x$ - angle between vectors v and x

Hausholder transformation, full QR factorization

Hausholder transformation

- ▶ v - unit vector, $\|v\|_2 = 1$, normal to hyperplane
- ▶ $v^T x = \|v\| \|x\| \cos(\hat{v}\hat{x})$
- ▶ $\hat{v}\hat{x}$ - angle between vectors v and x
- ▶ Geometric interpretation




Figure: Source: Biswa Nath Datta

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

► $\|Hx\|_2 = \|x\|_2$

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

► $\|Hx\|_2 = \|x\|_2$ ↓↓

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

- ▶ $\|Hx\|_2 = \|x\|_2$ ↓
- ▶ H is an orthogonal matrix:

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

- ▶ $\|Hx\|_2 = \|x\|_2$ \Downarrow
- ▶ H is an orthogonal matrix: $\|Hx\|_2^2 = \|x\|_2^2, (Hx, Hx) = (H^T Hx, x)$

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

- ▶ $\|Hx\|_2 = \|x\|_2$ \Downarrow
- ▶ H is an orthogonal matrix: $\|Hx\|_2^2 = \|x\|_2^2, (Hx, Hx) = (H^T Hx, x) \Rightarrow (H^T Hx, x) = (x, x)$

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

- ▶ $\|Hx\|_2 = \|x\|_2$ \Downarrow
- ▶ H is an orthogonal matrix: $\|Hx\|_2^2 = \|x\|_2^2, (Hx, Hx) = (H^T Hx, x) \Rightarrow (H^T Hx, x) = (x, x) \Rightarrow ((H^T H - I)x, x) = 0 \forall x$

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

- ▶ $\|Hx\|_2 = \|x\|_2$ ↓
- ▶ H is an orthogonal matrix: $\|Hx\|_2^2 = \|x\|_2^2, (Hx, Hx) = (H^T Hx, x) \Rightarrow (H^T Hx, x) = (x, x) \Rightarrow ((H^T H - I)x, x) = 0 \forall x \Rightarrow H^T H = I$

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

- ▶ $\|Hx\|_2 = \|x\|_2$ ↓
- ▶ H is an orthogonal matrix: $\|Hx\|_2^2 = \|x\|_2^2, (Hx, Hx) = (H^T Hx, x) \Rightarrow (H^T Hx, x) = (x, x) \Rightarrow ((H^T H - I)x, x) = 0 \forall x \Rightarrow H^T H = I$
- ▶ $H^2 = I$: reflect on other side of hyperplane and reflect back, $H^2 x = H(Hx)$

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

- ▶ $\|Hx\|_2 = \|x\|_2$ \Downarrow
- ▶ H is an orthogonal matrix: $\|Hx\|_2^2 = \|x\|_2^2, (Hx, Hx) = (H^T Hx, x) \Rightarrow (H^T Hx, x) = (x, x) \Rightarrow ((H^T H - I)x, x) = 0 \forall x \Rightarrow H^T H = I$
- ▶ $H^2 = I$: reflect on other side of hyperplane and reflect back, $H^2 x = H(Hx)$
- ▶ $y^T v = 0 \Rightarrow Hy = y$: y remain in the hyperplane after reflection.

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

► $\|Hx\|_2 = \|x\|_2$

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

► $\|Hx\|_2 = \|x\|_2$ ↓↓

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

- ▶ $\|Hx\|_2 = \|x\|_2$ \Downarrow
- ▶ H is an orthogonal matrix:

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

- ▶ $\|Hx\|_2 = \|x\|_2$ \Downarrow
- ▶ H is an orthogonal matrix: $\|Hx\|_2^2 = \|x\|_2^2, (Hx, Hx) = (H^T Hx, x)$

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

- ▶ $\|Hx\|_2 = \|x\|_2$ ↓
- ▶ H is an orthogonal matrix: $\|Hx\|_2^2 = \|x\|_2^2, (Hx, Hx) = (H^T Hx, x) \Rightarrow (H^T Hx, x) = (x, x)$

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

- ▶ $\|Hx\|_2 = \|x\|_2$ ↓
- ▶ H is an orthogonal matrix: $\|Hx\|_2^2 = \|x\|_2^2, (Hx, Hx) = (H^T Hx, x) \Rightarrow (H^T Hx, x) = (x, x) \Rightarrow ((H^T H - I)x, x) = 0 \forall x \Rightarrow H^T H = I$

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

- ▶ $\|Hx\|_2 = \|x\|_2$ ↓
- ▶ H is an orthogonal matrix: $\|Hx\|_2^2 = \|x\|_2^2, (Hx, Hx) = (H^T Hx, x) \Rightarrow (H^T Hx, x) = (x, x) \Rightarrow ((H^T H - I)x, x) = 0 \forall x \Rightarrow H^T H = I$
- ▶ $H^2 = I$:

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

- ▶ $\|Hx\|_2 = \|x\|_2$ ↓
- ▶ H is an orthogonal matrix: $\|Hx\|_2^2 = \|x\|_2^2, (Hx, Hx) = (H^T Hx, x) \Rightarrow (H^T Hx, x) = (x, x) \Rightarrow ((H^T H - I)x, x) = 0 \forall x \Rightarrow H^T H = I$
- ▶ $H^2 = I$: reflect on other side of hyperplane and reflect back, $H^2 x = H(Hx)$

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

- ▶ $\|Hx\|_2 = \|x\|_2$ \Downarrow
- ▶ H is an orthogonal matrix: $\|Hx\|_2^2 = \|x\|_2^2, (Hx, Hx) = (H^T Hx, x) \Rightarrow (H^T Hx, x) = (x, x) \Rightarrow ((H^T H - I)x, x) = 0 \forall x \Rightarrow H^T H = I$
- ▶ $H^2 = I$: reflect on other side of hyperplane and reflect back,
 $H^2 x = H(Hx)$
- ▶ $y^T v = 0 \Rightarrow Hy = y$:

Hausholder transformation, full QR factorization

Properties of Hausholder transformation, reflection




Figure: Source: Biswa Nath Datta

- ▶ $\|Hx\|_2 = \|x\|_2$ \Downarrow
- ▶ H is an orthogonal matrix: $\|Hx\|_2^2 = \|x\|_2^2, (Hx, Hx) = (H^T Hx, x) \Rightarrow (H^T Hx, x) = (x, x) \Rightarrow ((H^T H - I)x, x) = 0 \forall x \Rightarrow H^T H = I$
- ▶ $H^2 = I$: reflect on other side of hyperplane and reflect back, $H^2 x = H(Hx)$
- ▶ $y^T v = 0 \Rightarrow Hy = y$: y remain in the hyperplane after reflection.

Hausholder transformation, full QR factorization

Theorem 14.8

$$v \in \mathcal{R}^n, H = I - \frac{2vv^T}{v^T v}$$

Hausholder transformation, full QR factorization

Theorem 14.8

$$v \in \mathcal{R}^n, H = I - \frac{2vv^T}{v^T v}$$

► *H is symmetric*

Hausholder transformation, full QR factorization

Theorem 14.8

$$v \in \mathcal{R}^n, H = I - \frac{2vv^T}{v^T v}$$

- ▶ H is symmetric
- ▶ H is an orthogonal matrix

Hausholder transformation, full QR factorization

Theorem 14.8

$$v \in \mathcal{R}^n, H = I - \frac{2vv^T}{v^T v}$$

- ▶ H is symmetric
- ▶ H is an orthogonal matrix
- ▶ $H^2 = I$

Hausholder transformation, full QR factorization

Theorem 14.8

$$v \in \mathcal{R}^n, H = I - \frac{2vv^T}{v^Tv}$$

- ▶ H is symmetric
- ▶ H is an orthogonal matrix
- ▶ $H^2 = I$
- ▶ $Hv = -v$

Hausholder transformation, full QR factorization

Theorem 14.8

$$v \in \mathcal{R}^n, H = I - \frac{2vv^T}{v^T v}$$

- ▶ H is symmetric
- ▶ H is an orthogonal matrix
- ▶ $H^2 = I$
- ▶ $Hv = -v$
- ▶ $Hy = y$ **if** $y^T v = 0$

Hausholder transformation, full QR factorization

Theorem 14.8

$$v \in \mathcal{R}^n, H = I - \frac{2vv^T}{v^T v}$$

- ▶ H is symmetric
- ▶ H is an orthogonal matrix
- ▶ $H^2 = I$
- ▶ $Hv = -v$
- ▶ $Hy = y$ **if** $y^T v = 0$

Proof by direct verification (Exercise)

Hausholder transformation, full QR factorization

Theorem 14.9

$$v \in \mathcal{R}^n, H = I - \frac{2vv^T}{v^Tv} \text{ (Creating zeros in a vector)}$$

Hausholder transformation, full QR factorization

Theorem 14.9

$v \in \mathcal{R}^n, H = I - \frac{2vv^T}{v^Tv}$ (Creating zeros in a vector)

► $e_1, x \in \mathcal{R}^n, e_1 = (1, 0, \dots, 0)^T$

Hausholder transformation, full QR factorization

Theorem 14.9

$v \in \mathcal{R}^n, H = I - \frac{2vv^T}{v^Tv}$ (Creating zeros in a vector)

- ▶ $e_1, x \in \mathcal{R}^n, e_1 = (1, 0, \dots, 0)^T$
- ▶ $v = x + s_1 \|x\|_2 e_1, s_1 = \text{sign}(x_1)$

Hausholder transformation, full QR factorization

Theorem 14.9

$v \in \mathcal{R}^n, H = I - \frac{2vv^T}{v^Tv}$ (Creating zeros in a vector)

- ▶ $e_1, x \in \mathcal{R}^n, e_1 = (1, 0, \dots, 0)^T$
- ▶ $v = x + s_1 \|x\|_2 e_1, s_1 = \text{sign}(x_1)$
- ▶ $H = I - \frac{2vv^T}{v^Tv}$

Hausholder transformation, full QR factorization

Theorem 14.9

$v \in \mathcal{R}^n, H = I - \frac{2vv^T}{v^Tv}$ (Creating zeros in a vector)

- ▶ $e_1, x \in \mathcal{R}^n, e_1 = (1, 0, \dots, 0)^T$
- ▶ $v = x + s_1 \|x\|_2 e_1, s_1 = \text{sign}(x_1)$
- ▶ $H = I - \frac{2vv^T}{v^Tv}$
- ▶ \Downarrow

Hausholder transformation, full QR factorization

Theorem 14.9

$v \in \mathcal{R}^n, H = I - \frac{2vv^T}{v^T v}$ (Creating zeros in a vector)

- ▶ $e_1, x \in \mathcal{R}^n, e_1 = (1, 0, \dots, 0)^T$
- ▶ $v = x + s_1 \|x\|_2 e_1, s_1 = \text{sign}(x_1)$
- ▶ $H = I - \frac{2vv^T}{v^T v}$
- ▶ \Downarrow
- ▶ $Hx = s_1 \|x\|_2 e_1$

Hausholder transformation, full QR factorization

Theorem 14.9

$v \in \mathcal{R}^n, H = I - \frac{2vv^T}{v^T v}$ (Creating zeros in a vector)

- ▶ $e_1, x \in \mathcal{R}^n, e_1 = (1, 0, \dots, 0)^T$
- ▶ $v = x + s_1 \|x\|_2 e_1, s_1 = \text{sign}(x_1)$
- ▶ $H = I - \frac{2vv^T}{v^T v}$
- ▶ \Downarrow
- ▶ $Hx = s_1 \|x\|_2 e_1$

Proof by direct verification (Exercise)

Hausholder transformation, full QR factorization

Hausholder's method of QR factorization

Hausholder transformation, full QR factorization

Hausholder's method of QR factorization

- ▶ $A \in \mathcal{R}^{n \times m}$, tall matrix

Hausholder transformation, full QR factorization

Hausholder's method of QR factorization

- ▶ $A \in \mathcal{R}^{n \times m}$, tall matrix
- ▶ $H_n H_{n-1} \cdots H_1 A = R, A \in \mathcal{R}^{n \times m}$

Hausholder transformation, full QR factorization

Hausholder's method of QR factorization

- ▶ $A \in \mathcal{R}^{n \times m}$, tall matrix
- ▶ $H_n H_{n-1} \cdots H_1 A = R, A \in \mathcal{R}^{n \times m}$
- ▶ H_i -orthogonal

Hausholder transformation, full QR factorization

Hausholder's method of QR factorization

- ▶ $A \in \mathcal{R}^{n \times m}$, tall matrix
- ▶ $H_n H_{n-1} \cdots H_1 A = R, A \in \mathcal{R}^{n \times m}$
- ▶ H_i -orthogonal
- ▶ $Q = H_1 H_2 \cdots H_n$

Hausholder transformation, full QR factorization

Hausholder's method of QR factorization

- ▶ $A \in \mathcal{R}^{n \times m}$, tall matrix
- ▶ $H_n H_{n-1} \cdots H_1 A = R, A \in \mathcal{R}^{n \times m}$
- ▶ H_i -orthogonal
- ▶ $Q = H_1 H_2 \cdots H_n$
- ▶ $A = QR$

Q & A