

1 True or False

- (a) True
- (b) True
- (c) True
- (d) False
- (e) False
- (f) True
- (g) False
- (h) True
- (i) False
- (j) True
- (k) False

2 Condition Numbers

1. $\text{Cond}_2(A) = \frac{1}{\text{Cond}_2(A^{-1})}$
2. $\text{Cond}_2(cA) = \|cA\| \cdot \|A^{-1}c^{-1}\| = c\|A\| \cdot c^{-1}\|A^{-1}\| = \|A\| \cdot \|A^{-1}\| = \text{Cond}_2(A)$

3 Othogonal Matrices

since $\text{Cond}_2(cA) = \|cA\| \cdot \|A^{-1}c^{-1}\| = c\|A\| \cdot c^{-1}\|A^{-1}\| = \|A\| \cdot \|A^{-1}\| = \text{Cond}_2(A)$ and $\text{Cond}_2(Q) = 1$ where Q is an orthogonal matrix, we can say that $\text{Cond}_2(A) = \text{Cond}_2(cQ) = \text{Cond}_2(Q)$

4 Properties of Matrix Norms

By the definition, the norm $\|\cdot\|$ is an induced norm.

(a)

$$\|Ay\| \leq \|A\| \cdot \|y\| \implies \|Ay\| \leq \max_{\|x\|=1} \|Ax\| \cdot \|y\|$$

5 Matrix Products

$$\text{Cond}(AB) = \underbrace{\|AB\| \cdot \|B^{-1}A^{-1}\| \leq \|A\| \cdot \|A^{-1}\| \cdot \|B\| \cdot \|B^{-1}\|}_{\text{by sub-multiplicativity}} = \text{Cond}(A) \cdot \text{Cond}(B)$$

Although, for this, the norm must exhibit sub-multiplicativity.