

Table S2. Constant predictor against calibrated free-space loss by cell

Draws: split draws at $g = 10$ km, $b = 2$ km with at least one valid test node, by cell. Free-space loss (FSPL) is calibrated on the training median of received power plus modelled path loss, over all training nodes (first calibration) or over the valid training nodes only (second calibration). Counts are draws in which FSPL errs less than the constant predictor on the valid test nodes, and draws in which the constant errs less over all test nodes; the last two rows total the sixteen cells, the second one with no buffer ($b = 0$). Medians are mean absolute errors in dB over the valid test nodes.

Cell	Draws	Valid nodes: FSPL lower (first calibration)	Valid nodes: FSPL lower (second calibration)	All nodes: constant lower (first calibration)	All nodes: constant lower (second calibration)	Constant, valid (median, dB)	FSPL (first calibration), valid (median, dB)
Bauru Q1	59	55/59	55/59	37/59	49/59	18.101	6.210
Bauru Q2	56	45/56	45/56	42/56	45/56	15.080	4.906
Bauru Q3	56	53/56	52/56	43/56	52/56	17.826	5.615
Bauru Q4	56	50/56	50/56	37/56	44/56	15.492	5.406
Campinas Q1	59	51/59	50/59	40/59	39/59	14.114	5.067
Campinas Q2	59	53/59	52/59	34/59	33/59	16.940	6.258
Campinas Q3	57	42/57	43/57	42/57	54/57	15.767	9.719
Campinas Q4	58	51/58	51/58	37/58	40/58	15.512	6.661
Lins Q1	56	56/56	56/56	44/56	55/56	18.888	6.392
Lins Q2	59	59/59	59/59	26/59	48/59	21.220	6.913
Lins Q3	48	48/48	47/48	41/48	48/48	19.481	6.912
Lins Q4	53	53/53	52/53	39/53	50/53	19.775	5.799
Sorocaba Q1	58	38/58	39/58	44/58	48/58	11.665	7.231
Sorocaba Q2	59	52/59	52/59	40/59	39/59	14.749	6.283
Sorocaba Q3	55	43/55	43/55	43/55	44/55	20.172	8.030
Sorocaba Q4	56	50/56	50/56	40/56	41/56	14.598	4.815
All cells	904	799/904 (88.4%)	796/904 (88.1%)	629/904 (69.6%)	729/904 (80.6%)		
All cells, no buffer	938	844/938 (90.0%)	847/938 (90.3%)	691/938 (73.7%)	801/938 (85.4%)		