

Creating sample tables and inserting values.

Create the following tables with the given structures and insert sample data as specified: -

A) SALESPeOPLE Snum int(4) Sname varchar(10) City varchar(10) Comm float(3,2)

SALES PEOPLE

SNUM SNAME CITY COMM

1001 Peel London .12

1002 Serres San Jose .13

1004 Motika London .11

1007 Rifkin Barcelona .15

1003 Axelrod New York 10

```
D1_86939_Prajwal>select * from salespeople;
```

Snum	Sname	City	Comm
1001	Peel	London	0.12
1002	Serres	San Jose	0.13
1004	Motika	London	0.11
1007	Rifkin	Barcelona	0.15
1003	Axelrod	New York	0.10

5 rows in set (0.00 sec)

B) CUSTOMERS Cnum int(4) Cname varchar(10) City varchar(10) Rating int(4) Snum int(4)

CUSTOMERS CNUM CNAME CITY RATING SNUM

2001 Hoffman London 100 1001

2002 Giovanni Rome 200 1003

2003 Liu San Jose 200 1002

2004 Grass Berlin 300 1002

2006 Clemens London 100 1001

2008 Cisneros San Jose 300 1007

2007 Pereira Rome 100 1004

```
D1_86939_Prajwal>select * from customers;
```

Cnum	Cname	City	Rating	Snum
2001	Hoffman	London	100	1001
2002	Giovanni	Rome	200	1003
2003	Liu	San Jose	200	1002
2004	Grass	Berlin	300	1007
2006	Clemens	London	100	1001
2008	Cisneros	San Jose	300	1007
2007	Pereira	Rome	100	1004

7 rows in set (0.00 sec)

C) ORDERS Onum int(4) Amt float(7,2) Odate date Cnum int(4) Snum int(4)

ORDERS ONUM AMT ODATE CNUM SNUM

3001 18.69 03-OCT-1990 2008 1007

3003 767.19 03- OCT -1990 2001 1001

3002 1900.10 03- OCT -1990 2007 1004

3005 5160.45 03- OCT -1990 2003 1002

3006 1098.16 03- OCT -1990 2008 1007

3009 1713.23 04- OCT -1990 2002 1003

3007 75.75 04- OCT -1990 2004 1002

3008 4723.00 05- OCT -1990 2006 1001

3010 1309.95 06- OCT -1990 2004 1002

3011 9891.88 06- OCT -1990 2006 1001

```
D1_86939_Prajwal>select * from orders;
```

Onum	Amt	Odate	Cnum	Snum
3001	18.69	1990-10-03	2008	1007
3003	767.19	1990-10-03	2001	1001
3002	1900.10	1990-10-03	2007	1004
3005	5160.45	1990-10-03	2003	1002
3006	1098.16	1990-10-03	2008	1007
3009	1713.23	1990-10-04	2002	1003
3007	75.72	1990-10-05	2004	1002
3008	4723.00	1990-10-06	2006	1001
3010	1309.95	1990-10-06	2004	1002
3011	9891.88	1990-10-06	2006	1001

```
10 rows in set (0.00 sec)
```