

University of Warsaw
Faculty of Mathematics, Informatics and Mechanics

Joanna Dąbrowska

Student no. 438446

Modeling Genotype to Transcription Interactions in Follicular Lymphoma Using Variational Autoencoder

Master's thesis
in BIOINFORMATICS AND SYSTEMS BIOLOGY

Supervisor:
dr. Małgorzata Łazęcka
Institute of Informatics

Warsaw, September 2024

Abstract

Tumors consist of diverse cell populations exhibiting both genetic and functional heterogeneity, which together influence disease progression and treatment response. In cancer research, genotype-to-phenotype mapping aims to associate specific genetic alterations in individual cells or clones with functional traits such as transcriptional programs. Follicular lymphoma, a common non-Hodgkin lymphoma, displays pronounced clonal diversity, yet its genetic and phenotypic heterogeneity have often been studied separately. A previously developed probabilistic graphical model CaClust integrated multimodal data, including whole-exome, single-cell RNA, and B-cell receptor sequencing, to jointly infer clone genotypes and assign cells to clones, enabling single-cell genotyping. Building on this foundation, this study proposes a deep learning framework based on variational autoencoders (VAEs) to capture more complex genotype-to-phenotype relationships. This approach aims to enhance the resolution and interpretability of these links and reveal the functional consequences of somatic evolution in follicular lymphoma.

Keywords

neural networks, oncology, single cell sequencing, variation autoencoders, VAE

Thesis domain (Socrates-Erasmus subject area codes)

11.3 Informatics, Computer Science

Subject classification

Applied computing
Life and medical sciences
Bioinformatics

Tytuł pracy w języku polskim

Modelowanie zależności między genotypem a transkrypcją w chłoniaku grudekowym z wykorzystaniem wariacyjnego autoenkodera

Contents

1. Introduction	7
1.1. General introduction to the topic	7
1.2. Biological background	7
1.2.1. B cell lifecycle	7
1.2.2. Follicular lymphoma	8
1.2.3. Current diagnosis and prognostic treatments	9
1.3. the goal of the study	10
1.4. Approach we will use	10
1.4.1. previous probabilistic approach - graph methods	10
1.4.2. New deep learning approach	10
2. Materials and methods	11
2.1. Data	11
2.1.1. Sample collection and sequencing	11
2.1.2. Data processing	11
2.1.3. Generating sample data	12
2.2. Quick introduction to variation auto encoder	12
2.3. Loss function approach	12
2.4. Model 1 description	12
2.5. Model 2 description	12
3. Results	13
3.1. Metrics	13
3.2. Model comparison	13
4. Discussion	15
Bibliography	17
List of figures	19
List of tables	21

Chapter 1

Introduction

1.1. General introduction to the topic

TODO

1.2. Biological background

1.2.1. B cell lifecycle

The B cells are an essential component of adaptive immunity, responsible for producing antibodies. Their development is tightly regulated to generate a highly diverse range of antibodies (repertoire diversity) while ensuring the cells do not attack the body's own tissues (immune tolerance). This developmental progression begins in the bone marrow, where hematopoietic stem cells differentiate into common lymphoid progenitors to seed the B-cell lineage.

Lineage commitment is driven by a network of transcription factors and microenvironmental cues that guide the cells through distinct developmental phases. During the pro-B stage, cells initiate V(D)J recombination to rearrange their immunoglobulin heavy chains. Upon successful heavy-chain rearrangement, they transition to the pre-B stage to rearrange their light chains. Once a complete, functional receptor is formed, these newly generated immature B cells leave the bone marrow and migrate to peripheral lymphoid organs, such as the spleen, to complete their maturation.

When these mature B cells encounter an antigen in secondary lymphoid organs, they are triggered to form germinal centers (GCs). Within these specialized microenvironments, B cells undergo rapid proliferation and activation-induced cytidine deaminase (AID)-dependent modifications, specifically somatic hypermutation (SHM) and class-switch recombination. Following these genetic modifications, the cells undergo stringent selection; only the clones with the highest affinity for the antigen survive to exit the GC as memory B cells or plasma cells.

Given that GC reactions deliberately introduce DNA breaks and point mutations into immunoglobulin genes to optimize antigen affinity, off-target mutations are an unavoidable byproduct. Consequently, this AID-mediated mutagenesis is the primary mechanism by which B cells acquire secondary, oncogenic lesions that can initiate or promote lymphoma. In summary, while the bone marrow establishes the foundational B-cell repertoire, the germinal centers act as both the crucible for functional optimization and the primary site for oncogenic mutagenesis. [1]

1.2.2. Follicular lymphoma

Follicular lymphoma (FL) is an indolent neoplasm derived from germinal center (GC) B cells, defined by its characteristic nodal architecture of abnormal follicles and by the persistence of a GC-like transcriptional program that circumvents normal selection and exit mechanisms [2]. The hallmark initiating lesion in the majority of cases is the t(14;18)(q32;q21) translocation. This rearrangement places the anti-apoptotic gene BCL2 under the control of the Immunoglobulin Heavy Chain (IGH) locus enhancers. Since the IGH locus is constitutively and highly active in B cells to support antibody production, its enhancers drive the pathological overexpression of BCL2, providing a profound survival advantage by inhibiting apoptosis.

The presence of t(14;18) alone, however, is not sufficient to establish a precursor state. Small numbers of B cells carrying this rearrangement can be detected in the peripheral blood of a large proportion of healthy adults, reflecting the fact that this event often arises during normal GC reactions. These cells, although genetically altered, do not on their own represent a premalignant stage, and in most individuals they never progress.

A subset of these t(14;18)-bearing cells can evolve into follicular lymphoma-like cells (FLLCs). These are atypical memory B cells that have transited through the GC and acquired characteristics reminiscent of centrocytes, including CD10+/-, CXCR4^{low}, CD27+, Ki67^{low}, and BCL6 expression, along with molecular footprints of AID activity. FLLCs display the unusual combination of genomic instability and broad dissemination to lymphoid sites such as lymph nodes and bone marrow, yet they remain biologically inert with respect to malignant progression. One of their distinguishing features is the preferential retention of surface IgM expression, even when the nonproductive allele has undergone class switching, suggesting selective pressure to maintain IgM-mediated signaling. [3]

By contrast, cancer precursor cells (CPCs) represent a further evolutionary step and are considered true premalignant entities. CPCs accumulate cooperating mutations in epigenetic regulators, as well as transcriptional factors and immune modulators. These alterations lock the cells in a GC-like program while progressively eroding normal checkpoints of differentiation and selection. Unlike FLLCs, CPCs are committed to eventual lymphomagenesis and provide the pool from which clinically overt FL arises. Importantly, CPCs also function as the genetic reservoir for relapse and transformation. While early relapses may emerge through linear evolution of the dominant FL clone, late relapses and transformation events typically reflect divergent evolution from a shared CPC ancestor that has branched into genetically distinct sublineages [3].

Once FL is clinically established, the malignant cells remain locked in a GC-like state and persist through iterative GC re-entry, ongoing AID-driven mutagenesis, and selection within the lymphoid niche. FL primarily involves lymph nodes but frequently extends to the bone marrow and, in some cases, to the peripheral blood.

The tumor microenvironment (TME) plays a decisive role in sustaining the disease. T_{FH} cells provide CD40L, IL-4, and IL-21, which contribute to NF- κ B and JAK-STAT pathway activation in tumor B cells [2][3]. Follicular dendritic and stromal cells produce CXCL12 and CXCL13 that support migration and survival, while macrophages together with regulatory T cells contribute to local immune suppression[3].

Tumor-intrinsic mechanisms take advantage of these interactions. N-glycosylation of the BCR enables antigen-independent signaling, which drives clonal expansion without the need for antigen recognition and increases the probability of acquiring somatic mutations. Among the most crucial of these are alterations in chromatin regulators such as CREBBP and EZH2, which promote immune evasion by mechanisms including downregulation of MHC expression. Loss of TNFRSF14 further strengthens T_{FH}-stromal crosstalk, and stromal polarization by




Figure 1.1: The Stepwise Evolutionary Landscape of Follicular Lymphoma.

The development of FL is a long-term process that spans years of subclinical disease. (Left) The process starts with $t(14;18)$ translocations occurring during early B-cell development in the bone marrow or during primary germinal center (GC) reactions. This rearrangement leads to BCL2 overexpression, which prevents apoptosis in cells that would otherwise be eliminated during the normal biological selection process. (Center) These BCL2-expressing cells give rise to follicular lymphoma-like cells (FLLCs) that undergo repeated GC re-entry and spread throughout the lymphoid system. The acquisition of additional mutations, particularly in epigenetic regulators, leads to the emergence of committed cancer precursor cells (CPCs). (Right) CPCs act as a shared ancestral reservoir. While some sublineages evolve directly into the primary tumor found at diagnosis, others remain as a separate pool. This reservoir drives clinical relapse and histological transformation (t-FL) through branched evolution, explaining why relapsed disease is often genetically different from the primary tumor. Figure from [3]

neutrophils and macrophages through NF- κ B pathways reinforces the supportive niche. [3]

Although usually indolent, FL carries a persistent risk of histologic transformation to an aggressive form, typically diffuse large B-cell lymphoma (DLBCL). Transformation involves the acquisition of aggressive lesions such as TP53 loss, CDKN2A/B deletion, BCL6 rearrangements, or MYC activation, which drive genomic instability, impair cell-cycle checkpoints, and reduce reliance on microenvironmental cues [4]. These genetic and phenotypic changes correspond to the disappearance of follicular architecture, the emergence of diffuse growth, and clinical acceleration with extranodal spread. Whether transformation occurs through a linear accumulation of lesions in the dominant FL clone or through divergent evolution from common precursor cells, both models have been observed, and in either case the outcome is an abrupt shift from indolent to aggressive biology.

1.2.3. Current diagnosis and prognostic treatments

TODO

1.3. the goal of the study

TODO - current problems - molecular diagnosis - what does it means for other/how can it help

1.4. Approach we will use

TODO

1.4.1. previous probabilistic approach - graph methods

1.4.2. New deep learning approach

Chapter 2

Materials and methods

2.1. Data

2.1.1. Sample collection and sequencing

TO BE PARAPHRASED

Samples with histologically confirmed infiltration of follicular lymphoma (FL) grade 1-2 were collected under authorization of applicable biobank regulations of Leiden University Medical Center and the Ethical Committee of Leiden University Medical Center (reference HEM 008/SH/sh). Excisional lymph node biopsies were processed immediately by gentle mechanical disruption and mesh filtration. Single cell suspensions were frozen in 10% DMSO and remaining tissue fractions were cultured in low-glucose (1g/L) DMEM with 8% fetal bovine serum to obtain adherent cell cultures for isolation of DNA of cells representing normal counterpart. Cell processing, library preparation and sequencing FL cells were thawed and purified by flowcytometry using anti-CD19-APC (Becton Dickinson, Franklin Lakes, NJ) and anti-CD10-PECy7 (Becton Dickinson) followed by removal of dead cells (MACS Dead Cell Removal Kit, Miltenyi Biotech, Bergisch Gladbach, Germany). For whole exome sequencing, DNA was isolated from $1 \cdot 10^6$ purified FL cells and from $0.5 \cdot 10^6$ cultured adherent cells (Allprep DNA/RNA Mini Kit, Qiagen, Hilden, Germany). Whole exomes were sequenced using SureSelect Human All Exon V7 baits (Agilent, Santa Clara, CA). Adherent cells representing normal cells were sequenced at 50x coverage. To discriminate between early clonal variants and more recently acquired subclonal variants as putative drivers of distinct clones, FL bulk DNA was sequenced at 1500x coverage to allow reliable calling of rare variants down to a variant allele frequency (VAF) of 0.02. For 5' based single cell transcriptome sequencing, $1 \cdot 10^5$ similarly purified viable cells were loaded on a Chromium X single cell device to generate cDNA libraries for an expected $6 \cdot 10^3$ - $8 \cdot 10^3$ cells per sample. The amplified library is split in 3 fractions for 1) full transcriptome sequencing, 2) enrichment of BCR transcripts followed by full length sequencing, 3) targeted resequencing of subclonal somatic variants. Sequencing for WES and single cells was performed on HiSeq2500 or HiSeq4000 devices.

2.1.2. Data processing

TO BE PARAPHRASED

FASTQ files from whole exome sequencing (WES) were processed using the Sarek workflow v2.7 and aligned to the human reference genome GRCh38 using Burrows Wheeler Algorithm (BWA) v0.7.17. [33, 34] Duplicated mapped reads were marked, local realignment of regions flanking indels and recalibration of base quality scores were performed to obtain more accurate

bases according to the Genome Analysis ToolKit (GATK) best practices version v4.1.7.0. [35] Single nucleotide variants (SNV) and short insertions and deletions (INDELS) were called using Strelka2 v2.9.10. [36] Only high confidence variants defined by quality scores (GQX) of at least 15 for SNV and 30 for INDELS were kept. Pathogenicity of variants was determined using the Geneticist Assistant NGS Interpretive Workbench (SoftGenetics) based on publicly available variant-databases (dbSNP, ClinVar, and COSMIC) and literature, into class 1 (benign), class 2 (likely benign), class 3 (unknown significance), class 4 (likely pathogenic), or class 5 (pathogenic) [37–39].

20 Variant selection for CaClust modeling We chose only somatic single nucleotide variants called from Strelka that could also be observed in the scRNA data of the cells with complete BCR heavy and light chain sequences, and that showed at least one alternate read across the cells. Some germline variants were present in the Strelka output, as they had an increased frequency of the alternate nucleotide over the normal sample. We chose to include only those variants that showed over 3-fold increase in the frequency of the alternate nucleotide between the normal and tumour sample. The full table of included variants per sample can be seen in Additional File 3: Output profiles

2.1.3. Generating sample data

- hyperclusters -> Chinese restaurant process - hc to clone assignment -> categorical distribution - bcr sequence -> generate bcr profile using dirichlet distribution for each hypercluster, and then sample sequence from categorical distribution across ATGC with probabilities from the profile - generate raw counts of genes using poisson distribution -> should be NB in fact ;) - generate clone profiles using binomial distribution - generate mutation counts matrix

2.2. Quick introduction to variation auto encoder

2.3. Loss function approach

2.4. Model 1 description

2.5. Model 2 description

Chapter 3

Results

3.1. Metrics

3.2. Model comparison

Chapter 4

Discussion

Bibliography

1. Murphy, K. & Weaver, C. *Janeway's immunobiology* 9th ed. (New York : Garland Science, 2017).
2. Krull, J. E. *et al.* Follicular lymphoma B cells exhibit heterogeneous transcriptional states with associated somatic alterations and tumor microenvironments. *Cell Reports Medicine* **5**, 101443. ISSN: 2666-3791. <https://www.sciencedirect.com/science/article/pii/S2666379124000661> (2025) (Mar. 2024).
3. Carbone, A. *et al.* Follicular lymphoma. en. *Nature Reviews Disease Primers* **5**. Publisher: Nature Publishing Group, 1–20. ISSN: 2056-676X. <https://www.nature.com/articles/s41572-019-0132-x> (2025) (Dec. 2019).
4. Pasqualucci, L. *et al.* Genetics of follicular lymphoma transformation. *Cell reports* **6**, 130–140. ISSN: 2211-1247. <https://pmc.ncbi.nlm.nih.gov/articles/PMC4100800/> (2025) (Jan. 2014).

List of Figures

1.1. The Stepwise Evolutionary Landscape of Follicular Lymphoma.	9
--	---

List of Tables