

Exercise 7 Answer Sheet — Axiomatic Set Theory, 80650

January 15, 2025



Question 1

We will assume ZFC unless stated otherwise.

Let $j : V \rightarrow M$ be a non-trivial elementary embedding such that $\text{crit } j = \kappa$, where M is transitive. Let $\mathcal{U} = \mathcal{U}_j = \{X \subseteq \kappa \mid \kappa \in j(X)\}$, we know that \mathcal{U} is a κ -complete ultrafilter on κ .

a

We will show that for every $A \in \mathcal{U}$ and regressive function $f : A \rightarrow \kappa$ there is $B \subseteq A$, $B \in \mathcal{U}$, such that $\text{rng}(f \restriction B) = \{\gamma\}$ for some $\gamma < \kappa$.

Proof. Assumption that $A = \kappa$ will result in contradiction to f being regressive, then $|\text{dom } f| < \kappa$. $\forall x \in A, f(x) < x$ then $\sup x \in A, f(x) < \kappa$. Then we will notate $\mu = |\text{Im } f|$, we can assume $\mu < \kappa$.

We will imitate Fodor's lemmas proof.

If there is $\gamma < \kappa$ such that the conditions are met then the proof is done, then we will assume there is no such set B .

For every $\gamma < \kappa$ we define $B_\gamma = f^{-1}(\{\gamma\})$, $B_\gamma \notin \mathcal{U}$ then $U_\gamma = \kappa \setminus B_\gamma \in \mathcal{U}$ and disjoint from B_γ . We then use \mathcal{U} 's κ -completeness to deduce that for $U = \bigcap_{\gamma < \mu} U_\gamma$, $U \in \mathcal{U}$. Then $A \cap U \in \mathcal{U}$ as well, and in particular $A \cap U \neq \emptyset$, let $\alpha \in A \cap U$. $f(\alpha) = \gamma < \alpha$ for certain γ , but then $\alpha \in B_\alpha$, meaning $\alpha \notin U$, a contradiction. \square

b

We notate $\text{Ma}(x) \iff x$ is Mahlo cardinal.

We will show that $\{\alpha \mid \text{Ma}(\alpha)\} \in \mathcal{U}$.

Proof. Let $A = \{\alpha \leq \kappa \mid \text{Ma}(\alpha)\}$, from a theorem $\kappa \in A$.

It follows that $j(A) = \{\alpha \leq j(\kappa) \mid M \models \text{Ma}(\alpha)\}$, but $\kappa < j(\kappa)$ and κ is definable and therefore $\kappa \in M$ and $M \models \text{Ma}(\kappa)$.

We can assume then $\kappa \in j(A)$, this is of course implies that $A \in \mathcal{U}$ as we wanted to show. \square

c

Let $S \subseteq \kappa$ be stationary set. We define $\varphi_s(x, y) \iff x$ is stationary in y .

We will prove that $A = \{\alpha \subseteq \kappa \mid \varphi_s(S \cap \alpha, \alpha)\} \in \mathcal{U}$.

Proof. $V \models \varphi_s(S, \kappa) \iff M \models \varphi_s(j(S), j(\kappa))$ as j is elementary embedding and φ_s is first order (I carefully hope). Also $j(A) = \{\alpha \leq j(\kappa) \mid M \models \varphi(j(S \cap \alpha), j(\alpha))\}$, but $\kappa \leq j(\kappa)$, and $j(S \cap \kappa) = j(S)$ then by the last statement $\kappa \in j(A)$ as intended. \square

Question 2

Let κ be a measurable cardinal and let \mathcal{U} be a non-principle κ -complete normal ultrafilter on κ .

Let j be the elementary embedding derived from the ultrapower by \mathcal{U} and let $\mathcal{V} = \{X \subseteq \kappa \mid \kappa \in j(X)\}$.

We will show that $\mathcal{V} = \mathcal{U}$.

Proof. $\{i \mid id(i) > c_\alpha(i)\} = \kappa \setminus \alpha \in \mathcal{U}$ then by Łoś theorem $[id] \geq \kappa$. In similar way $\{i \mid id(i) < c_\alpha(i)\} = \kappa \in \mathcal{U}$ then $[id] \leq \kappa$, in particular $[id] = \kappa$. For $g : \kappa \rightarrow \kappa$ regressive it follows that $g(x) < x = id(x)$ then $\{i \mid g(i) < id(i)\} = \kappa \in \mathcal{U}$ implies $[g] < [id] = \kappa$. From normality $\{i \mid g(i) = c_\gamma(i)\} \in \mathcal{U}$, then $[g] = \gamma$. The idea is to assume there is $x \in \mathcal{V}$, meaning $\kappa \in j(x)$, this is equivalent to $\{i \mid id(i) < g(i)\} \in \mathcal{U}$. We know that $\kappa \leq [id]$ then $\{i \mid g(i) < id(i)\}$ and this is normal then there is $\gamma < \kappa$ such that $\{i \mid g(i) = \gamma\} \in \mathcal{U}$ then $[g] = \gamma$ but this does not make sense. It gives us $\kappa = [id]$ then $\kappa \in j(x) \iff \{i \mid id(i) < f(i)\} \in \mathcal{U}$ □