Space of analytic functions over unit disk — Notes

October 25, 2026



Contents

1 Lecture 1 — 18.10.2026 **3**

1.1 Harmonic functions 3

1.2 Poisson integral formula 4

2 Lecture 2 — 25.10.2026 **6**

2.1 Measures and total variation 6

2.2 Harmonic functions in R-disk 6

2.3 Harmonic functions with bounded Lp means 7

2.4 Elementary study of boundary behavior 8

1 Lecture 1 — 18.10.2026

The main object of this course is a Hardy Space, for $0 < p < \infty$, the Hardy space $\mathbb{H}^p$ consists of all holomorphic functions $f : \mathbb{D} \to \mathbb{C}$ such that,

$$\sup_{0<r<1} \int_{-\pi}^{\pi} |f(re^{it})|^p \, dt < \infty$$

and,

$$\mathbb{H}^\infty = \{\text{bounded analytic functions on } \mathbb{D}\}$$

for $p < q$ it is clear that $\mathbb{H}^p \supseteq \mathbb{H}^q \supseteq \ldots \supseteq BMO \supseteq \mathbb{H}^\infty$. We study boundary behavior, in particular Fatou's theory; for a.e. $\theta \in [0, 2\pi)$, a function $f \in \mathbb{H}^\infty$ has a non-tangential limit at $e^{i\theta}$,

$$z_n to e^{i\theta} \implies \lim_{n\to\infty} f(z_n) \text{ exists}$$

in particular $\lim_{r\to 1^-} f(re^{i\theta})$ exists.

Another question we study is Zero sets, a set of points $\{a_n\} \subseteq \mathbb{D}$ is the Zero set of an $\mathbb{H}^\infty$ function if and only if $\sum(1 - |a_n|) < \infty$, in other words it converges to the boundary quickly (Blaschke condition).

We also study Invariant subspaces $zX \subseteq X$, if $z(z) \in X$ then $z \cdot f(z) \in X$ (Beurling, 1949).

Interpolating sequences. A sequence $\{a_n\} \subseteq \mathbb{D}$ is interpolating if for any $b_n \in l^\infty$ there exists $f \in \mathbb{H}^\infty$ such that $f(a_n) = b_n$. The answer was given by Carleson: a sequence is interpolating if and only if it is separated and has Carleson condition.

Finally, Corona problem: given $f_1, f_2 \in \mathbb{H}^\infty$ such that $|f_1(z)| + |f_2(z)| > \delta$ for all $z \in \mathbb{D}$, then there are $g_1, g_2 \in \mathbb{H}^\infty$ such that $f_1 g_1 + f_2 g_2 = 1$. The Space BMO is going to be involved with the solution to this problem. There are also Bergman spaces, $A^p = \{f : \mathbb{D} \xrightarrow{\text{hol}} \mathbb{C} \mid \int_{\mathbb{D}} |f(z)|^p \, dA < \infty\}$ and Dirichlet space,

$$\mathcal{D} = \{f : \mathbb{D} \xrightarrow{\text{hol}} \mathbb{C} \mid \int_{\mathbb{D}} |f'(z)|^2 \, dA < \infty\}$$

The textbook is "H_p spaces" by P. Koosis. The preliminaries are Complex analysis 1, Measure theory, Hilbert spaces and little bit of Functional analysis.

Reminder about $\mathcal{L}^p$ spaces, $(\mathcal{L}^p)^* \simeq \mathcal{L}^q$ for $1 \le p < \infty$ where $\frac{1}{p} + \frac{1}{q} = 1$. Then $\|g\|_{L^q} = \sup_{\|f\|_{L^p}} \int f \cdot g \, dt$. Another fact is that $C_c(\mathbb{R}) \subseteq \mathcal{L}^p(\mathbb{R})$ is dense. $(T_t f)(x) = f(x + t)$ is continuous in $\mathcal{L}^p$ in the sense that $t \to T_t f$ is continuous,

$$\|T_t f - f\|_{L^p} \xrightarrow{t\to 0} 0$$

the proof is left as exercise. $C(\partial\mathbb{D})^* \simeq \mathcal{M}(\partial\mathbb{D})$ and $C_0(\mathbb{R})^* \simeq \mathcal{M}(\mathbb{R})$. We can define the measure μ that takes $f \to \int_{\partial\mathbb{D}} f \, d\mu$.

1.1 Harmonic functions

Definition 1.1 (Harmonic function)**.** a function $u : \Omega \to \mathbb{R}$ is harmonic if one of the following conditions holds:

1. u is C^2 and $\Delta u = \frac{\partial^2}{\partial x^2} u + \frac{\partial^2}{\partial y^2} = 0$
2. $u \in C$ satisfies the mean-value property, namely for any disk $B(z_0, r) \subseteq \Omega$ it holds that,
$$u(z_0) = \frac{1}{2\pi} \int_{-\pi}^{\pi} u(z_0 + re^{i\theta}) \, d\theta$$
3. u is locally the real part of an analytic function.

Example 1.1. $\log|z|$ is harmonic in $\mathbb{C} \setminus \{0\}$.

We would have wanted to take $f = \log z$ but we would've encountered a problem with continuity. Note that condition 3 holds for the imaginary part as well. For the sake of convenience we will treat condition 3 as the main definition in this

course. Suppose that $u : B(0,R) \to \mathbb{R}$ is harmonic, then $u = \text{Re}\, f$ for some analytic function f, but any such function has an analytic series,

$$f(z) = \sum_{n=0}^{\infty} a_n z^n$$

such that,

$$\liminf_{n\to\infty} \frac{1}{\sqrt[n]{|a_n|}} \geq R$$

Taking real parts, we get,

$$u(rr^{i\theta}) = \sum_{n=-\infty}^{\infty} A_n r^{|n|} e^{in\theta}$$

In order for this to be real we need $A_{-n} = \overline{A_n}$ and the condition,

$$\forall n \in \mathbb{Z},\ \liminf_{n\to\infty} \frac{1}{\sqrt[|n|]{|A_n|}} \geq R$$

1.2 Poisson integral formula

Poisson formula gives us some generalization of the mean-value property in the sense that each point is valued by an harmonic function as the boundary of a consisting ball, up to weight function (kernel). Let $u : B(0,R) \to \mathbb{R}$ with $R > 1$ and $r < 1$ then,

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} u(e^{it}) \left[\sum_{-\infty}^{\infty} r^{|n|} e^{in(\theta - t)} \right] dt$$

We know that,

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} e^{int}\, dt = \begin{cases} 0 & n \neq 0 \\ 1 = 0 & \end{cases}$$

and that if u is defined as a sequence, then,

$$u(re^{it}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \sum_{-\infty}^{\infty} A_n e^{int} r^{|n|} e^{in(\theta - t)}\, dt$$

thus we get the exact formula.

Definition 1.2 (Poisson kernel)**.** For any θ we define,

$$P_r(\theta) = \sum_{-\infty}^{\infty} r^{|n|} e^{in\theta} = \frac{1 - r^2}{1 + r^2 - 2r\cos\theta} = \text{Re}\left(\frac{1 + re^{i\theta}}{1 - re^{i\theta}}\right)$$

Then,

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} u(e^{it}) P_r(\theta - t)\, dt$$

Proposition 1.3. *The following properties hold for Poisson kernel:*

1. $P_r(\theta) > 0$
2. P_r *is* 2π*-periodic*
3. $\frac{1}{2\pi} \int_{-\pi}^{\pi} P_r(\theta)\, d\theta = 1$
4. *For any* $\delta > 0$, $P_r(\theta) \to 0$ *uniformly for* $\pi > |\theta| > \delta$, *as* $r \to 1$
5. $P_r(\theta)$ *is even*

Remark. We can also denote using Poisson kernel with,

$$u(er^{i\theta}) = \frac{1}{2\pi} u(e^{it}) P_r(\theta - t)\, dt$$

Suppose that $f \in \mathcal{L}^p(\partial\mathbb{D})$ or equivalently $\in \mathcal{L}^p([-\pi,\pi])$, and let $u = P[f]$ Poisson extension of f, namely,

$$u(re^{it}) = \frac{1}{2\pi}\int_{-\pi}^{\pi} f(t)P_r(\theta - t)\,dt$$

then u is harmonic in $\mathbb{D}$. We can also consider $u + i\tilde{u}$ for $\tilde{u}$ the harmonic conjugate. So we can plug $\mathcal{L}^p$ functions into the PIF (Poisson Integration Formula) to obtain harmonic functions in the disk. What kind of functions do we get by doing so, and how are u and f related? Is $f \to u \to \tilde{u} \to \tilde{f}$ bounded in $\mathcal{L}^p$ (Singular integral operators)? That is a delicate question which we will answer during the course.

Theorem 1.4 (Extension to harmonic functions)**.** *Suppose that $u(re^{i\theta})$ is a harmonic function on the unit disk. If $1 < p < \infty$ and,*

$$\sup_{0<r<1}\int_{|z|=r} |u(re^{i\theta})|\,d\theta < \infty$$

then there is a function $f : [-\pi,\pi] \to \mathbb{R}$ in L^p such that $u(re^{i\theta}) = \frac{1}{2\pi}\int_{-\pi}^{\pi} P_r(\theta - t)f(t)\,dt$.

Proof. Compactness: We know that $(\mathcal{L}^q)^* \simeq \mathcal{L}^p$. Choose a countable dense set $G_1, G_2, \ldots \in \mathcal{L}^q$. Define,

$$\mathcal{L}G_j = \lim_{j\to\infty}\frac{1}{2\pi}\int_{-\pi}^{\pi} G_k(\theta)u((1-\frac{1}{n_j})e^{i\theta})\,d\theta$$

$G_k \in \mathcal{L}^q$ and $u((1-n_j)e^{i\theta})$ is bounded because L^p norm is bounded, then the whole integral is bounded. Remember that $|\int fg| \le \|f\|_p \cdot \|g\|_q$ for any $f \in L^p, g \in L^q$. For a general $G \in \mathcal{L}^q$, we approximate G by $G_{k_l} \in \mathcal{L}^q$ and define $\mathcal{L}G = \lim_{l\to\infty}\mathcal{L}G_{k_l}$. It follows that there exists a function $f \in \mathcal{L}^p$ such that $\mathcal{L}G = \int_{-\pi}^{\pi} fG$. Fix $-\pi \le \theta \le \pi$ and choose $G(t) = P_r(\theta - t)$, let $v = P[f]$,

$$\mathcal{L}P_r(\theta - t) = \int_{-\pi}^{\pi} f(t)P_r(\theta - t)\,dt = 2\pi v(re^{i\theta}) = \lim_{j\to\infty}\int_{-\pi}^{\pi} P_r(\theta - t)u((1-\frac{1}{n_j})e^{i\theta})$$

The dilate $u((1-\frac{1}{n_j})e^{i\theta})$ is harmonic on $B(0, \frac{1}{1-\frac{1}{n_j}}) \subseteq B(0,1)$ therefore we can use Poisson formula,

$$\lim_{j\to\infty}\int_{-\pi}^{\pi} P_r(\theta - t)u((1-\frac{1}{n_j})e^{i\theta}) = \lim_{j\to\infty} 2\pi u((1-\frac{1}{n_j})re^{i\theta}) = 2\pi\cdot u(re^{i\theta})$$

□

Theorem 1.5 (Existence of measure to harmonic functions)**.** *Suppose that $u(er^{i\theta})$ is a harmonic function on $\mathbb{D}$ such that,*

$$\sup_{0<r<1}\int_{-\pi}^{\pi} |u(re^{i\theta})|\,d\theta < \infty$$

then there exists a measure μ such that,

$$u(re^{i\theta}) = \frac{1}{2\pi}\int_{-\pi}^{\pi} P_r(\theta - t)\,d\mu$$

Example 1.2. Remember that $P_r(\theta) = \frac{1-r^2}{1+r^2-2\cos\theta}$, in the case "$r = 1$" acts like δ_1.

2 Lecture 2 — 25.10.2026

2.1 Measures and total variation

A **signed measure** is the difference of two positive finite measures, $\mu = \mu_+ - \mu_-$. The **total variation measure** $|\mu| = \mu_+ + \mu_-$, this leads to the identity,

$$\left|\int f\, d\mu\right| \leq \int |f|\, d|\mu|$$

We define $\|\mu\|_M = |\mu|(X)$. A complex measure is, in similar manner, a sum of two measures, $\mu = \lambda + i\nu$ where λ, ν are signed measures. In this case we define the total variation as $|\mu| = |\lambda| + |\nu| = \lambda_- + \lambda_- + \nu_+ + \nu_-$.

Theorem 2.1 (Banach-Alaoglu)**.** *Let X be a Banach space, The unit ball in X^* is weak-$*$ compact.*

Example 2.1. $(\mathcal{L}^q)^* \simeq \mathcal{L}^p$ for $1 <\leq \infty$ and $\frac{1}{p} + \frac{1}{q} = 1$.

Suppose $f_1, f_2, \ldots$ with $\sup\|f_j\|_{L^p} < \infty$. $\{f_j\}$ may not have a continuous subsequence in $\mathcal{L}^p$, BA (Banach-Alaoglu) implies that there are j_k such that $f_{j_k} \to f$ in weak-$*$,

$$\forall g \in \mathcal{L}^q, \quad \int f_{j_k} g\, dx \to \int fg\, dx$$

2.2 Harmonic functions in R-disk

Denote $\mathbb{D}_R = \{z \in \mathbb{C} \mid |z| < R\}$. Function $u : \mathbb{D}_R \to \mathbb{R}$ is harmonic if,

$$u(re^{i\theta}) = \sum_{n=-\infty}^{\infty} A_n r^{|n|} e^{in\theta}$$

where $A_{-n} = \bar{A}_n$ and radius $\geq R$.

We can consider complex-valued harmonic functions, in this case we can omit the first condition and say that u is analytic iff $A_{-n} = 0$ for $n > 0$,

$$u(re^{i\theta}) = \sum_{n\geq 0} A_n r^n e^{in\theta}$$

u is **anti-analytic** iff $A_n = 0$ for $n > 0$. Any harmonic function h can be written as $h = i + iv$ where u, v are real-valued harmonic functions, but also as $h = f + \bar{g}$ where f, g are analytics.

We say that v is a **harmonic-conjugate** of u if $u + iv$ is analytic.

Lemma 2.2. *v is unique up to an additive constant.*

If v_1, v_2 are such conjugates then $(u + iv_1) - (u + iv_2) = i(v_1 - v_2)$ which is constant.

Existence:

$$\tilde{u}(re^{i\theta}) = \sum_{n=0\infty}^{\infty} (-i \operatorname{sgn} n) A_n r^{|n|} e^{in\theta}$$

From this definition it follows that $\tilde{A}_n = -iA_n$ for $n > 0$ and so $\tilde{A}_{-n} = iA_{-n}$, meaning that indeed $\tilde{A}_{-n} = \bar{\tilde{A}}_n$.

When discussing Poisson integral formula we assumed that $R > 1$,

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} u(e^{it}) \left[\sum_{n=-\infty}^{\infty} r^{|n|} e^{in(\theta - t)}\right] dt$$

When $R > 1$, $|A_n| \leq C\Theta^{-n}$ for some $\Theta > 1$.

Example 2.2. The series $z + z^2 + \cdots$ has a radius $1 = \limsup \sqrt[|n|]{|A_n|}$, to enlarge it we have to set the coefficients to be much smaller. $\frac{z}{2} + \frac{z^2}{4} + \frac{z^3}{8}$ has a radius of 2.

We will define Poisson kernel as,

$$P_r(\theta) = \sum_{n=-\infty}^{\infty} r^{|n|} e^{in\theta} = \frac{1 - r^2}{1 + r^2 - 2r\cos\theta} = \operatorname{Re} \frac{1 + re^{i\theta}}{1 - re^{i\theta}}$$

Poisson kernel $P(re^{i\theta})$ is a harmonic function in $\mathbb{D}$. This is clear as it can be written as a special case of the series defining a harmonic function, or as the real part of an analytic function. In general we can achieve the harmonic conjugate with the formula,

$$\tilde{u}(re^{i\theta}) = \frac{1}{2\pi}\int_{-\pi}^{\pi} u(e^{it})\left[\sum_{n=-\infty}^{\infty} -i\,\mathrm{sgn}(n) r^{|n|}e^{in(\theta-t)}\right] dt$$

So we define the conjugate Poisson kernel as,

$$Q_r(\theta) = \sum_{n=-\infty}^{\infty} -i\,\mathrm{sgn}(n) r^{|n|}e^{in\theta} = \frac{2r\sin\theta}{1+r^2-2r\cos\theta} = \mathrm{Im}\,\frac{1+re^{i\theta}}{1-rr^{i\theta}}$$

2.3 Harmonic functions with bounded Lp means

Theorem 2.3 (Representation with Lp fuctions)**.** *Suppose u is harmonic function on the unit disk. If,*

$$\sup_{0<r<1}\int_{-\pi}^{\pi} |u(re^{i\theta})|^p\, d\theta < \infty \tag{1}$$

then,

$$u(re^{i\theta}) = \frac{1}{2\pi}\int_{-\pi}^{\pi} F(t)P_r(\theta-t)\, dt$$

where $F \in \mathcal{L}^p$.

Proof. Set $F_n(\theta) = u((1-\frac{1}{n})e^{i\theta})$ for $\theta \in [-\pi,\pi]$. (1) sways that $\|F_n\|_{L^p} \le C$ for all n. Banach-Alaoglu means that for any $G \in L^q$,

$$\int_{-\pi}^{\pi} F_{n_k}(\theta)G(\theta)\, d\theta \to \int_{-\pi}^{\pi} F(\theta)G(\theta)\, d\theta$$

Fix $0 \le r < 1$ and $t \in [-\pi,\pi]$ and take $G(\theta) = P_r(t-\theta)$, this is a smooth function.

$$2\pi u(re^{it}) = 2\pi \lim_{k\to\infty} u((1-\frac{1}{n_k})re^{it}) = \lim_{k\to\infty}\int_{-\pi}^{\pi} u((1-\frac{1}{n_k})e^{i\theta})P_r(t-\theta)\, d\theta = \int_{-\pi}^{\pi} F(\theta)P_r(t-\theta)\, d\theta$$

□

Theorem 2.4 (equivalency in the case of 1)**.** *Suppose u is harmonic function on the unit disk and $p = 1$. If,*

$$\sup_{0<r<1}\int_{-\pi}^{\pi} |u(re^{i\theta})|\, d\theta < \infty$$

then,

$$u(re^{i\theta}) = \frac{1}{2\pi}\int_{-\pi}^{\pi} P_r(\theta-t)\, d\mu(t)$$

where μ is a signed measure.

Corollary 2.5. Suppose $u : \mathbb{D} \to \mathbb{R}$ is a positive harmonic function. Then $u = P[\mu]$ for μ a positive measure, namely $du = Pd\mu$.

Example 2.3. $P(re^{i\theta}) = \frac{1-r^2}{1+r^2-2r\cos\theta}$ is a positive harmonic function,

$$P(re^{i\theta}) = \frac{1}{2\pi}\int_{-\pi}^{\pi} P_r(\theta-t)d(2\pi\delta_0(t))$$

It follows that $P = P[2\pi\delta_0]$.

Example 2.4 (Plateau's problem)**.** Suppose that $\gamma \subseteq \mathbb{R}^3$ is a closed smooth path, find a surface $S \subseteq \mathbb{R}^3$ with $\partial S = \gamma$ of minimal area.

Let $m = \inf\{\mathrm{vol}_2\, S \mid \partial S = \gamma\}$, for any $n \in \mathbb{N}$ there is a surface S_n with $\partial S_n = \gamma$ and $\mathrm{vol}_2\, S_n < m + \frac{1}{n}$.

Theorem 2.6. *Suppose $F(\theta) \in \mathcal{L}^p$, where $1 < p \le \infty$, and $u = P[F]$. Then,*

$$\sup_{0<r<1}\int_{-\pi}^{\pi} |u(re^{i\theta})|^p\, d\theta < \infty$$

Proof. Set $F_r(\theta) = u(re^{i\theta})$ for $\theta \in [-\pi, \pi]$. Then there is a function $G \in \mathcal{L}^q$ with $\|G\|_{L^q} = 1$ such that,

$$\begin{aligned}
\|F_r\|_{L^p} &= \int_{-\pi}^{\pi} F_r(\theta)G(\theta)\, d\theta \\
&= \left| \frac{1}{2\pi} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} F(\theta - t)P_r(t)\, dt\, G(\theta)\, d\theta \right| \\
&\leq \frac{1}{2\pi} \int_{-\pi}^{\pi} \|F\|_{L^p} \|G\|_{L^q} P_r(t)\, dt \\
&= \|F\|_{L^p} \|G\|_{L^q} \frac{1}{2\pi} \int_{-\pi}^{\pi} P_r(t)\, dt \\
&= \|F\|_{L^p} \|G\|_{L^q} \\
&= \|F\|_{L^p}
\end{aligned}$$

In the case $p = 1$, $G \in \mathcal{L}^\infty$ and $G = \frac{\bar{F}_r}{|F_r|}$, the inequalities are a bit different,

$$\begin{aligned}
\|F_r\|_{L^1} &= \int_{-\pi}^{\pi} F_r(\theta)G(\theta)\, d\theta \\
&= \frac{1}{2\pi} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} P_r(\theta - t) d\mu(t) G(\theta)\, d\theta \\
&\leq \frac{1}{2\pi} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} P_r(\theta - t) |d\mu(t)|\, d\theta \\
&= \int_{-\pi}^{\pi} |d\mu(\theta)| \\
&= \|\mu\|_M
\end{aligned}$$

and the claim holds. □

2.4 Elementary study of boundary behavior

Theorem 2.7 (Extension of unit function to boundary)**.** *Suppose F is a continuous function on $[-\pi, \pi]$ and $u = P[F]$. Then u extends continuously to $\overline{\mathbb{D}}$ and,*

$$\lim_{z \to e^{i\theta}} u(z) = F(\theta)$$

Notice that $P_r(\theta) > 0$, is even, 2π-periodic, $\int_{-\pi}^{\pi} P_r(\theta)\, d\theta = 2\pi$ and $\forall \delta > 0 \exists r_0,\ r > r_0 \implies \int_{|t|>\delta} P_r(\theta)\, d\theta < \delta$.

Proof. We show $\lim_{r \to 1} u(re^{i\theta}) = F(\theta)$.

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} F(\theta - t)P_r(t)\, dt$$

Then,

$$u(re^{i\theta}) - F(\theta) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} F(\theta - t)P_r(t)\, dt$$

and in particular,

$$|u(re^{i\theta}) - F(\theta)| \leq \frac{1}{2\pi} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} |F(\theta - t) - F(\theta)| P_r(t)\, dt = \frac{1}{2\pi} \left(\int_{|t|<\delta} + \int_{|t|>\delta} \right)$$

We show that these two integrals are bounded by $\frac{\varepsilon}{2}$. Since F is uniformly continuous, there exists a $\delta > 0$ such that,

$$|F(\theta - t) - F(\theta)| < \frac{\varepsilon}{2}$$

for $|t| < \delta$. Denote the two integrals by I and II, then,

$$|I| \leq \frac{1}{2\pi} \int_{|t|<\delta} \frac{\varepsilon}{2} P_r(t)\, dt \leq \frac{\varepsilon}{2}$$

Since F is continuous, it is bounded, $M = \|F\|_{L^\infty} < \infty$.

$$|II| \leq \frac{1}{2\pi} \int_{|t|>\delta} 2MP_r(t)\, dt$$

but by Poisson kernel properties, we can choose a value such that r is close to 1. □

Theorem 2.8. *Suppose $F(\theta) \in \mathcal{L}^p$ with $1 < p < \infty$ and $u = P[F]$. Set $F_r(\theta) = u(re^{i\theta})$, then $F_r \to F$ in L^p as $r \to 1$.*

Proof.

$$\|F_r(\theta) - F(\theta)\|_{L^p} = \left\| \frac{1}{2\pi} \int_{-\pi}^{\pi} (F(\theta - t) - F(\theta)) P_r(t) \right\|_{L^p} \leq \frac{1}{2\pi} \int_{-\pi}^{\pi} \|F(\theta - t) - F(\theta)\|_{L^p} P_r(t)\, dt \leq \frac{1}{2\pi} \left(\int_{|t|<\delta} + \int_{|t|>\delta} \right)$$

and we once again denote them I and II. $I < \frac{\varepsilon}{2}$ by the same claim as the last theorem. If $\|F\|_{L^p} = M$ then $\|F(\theta - t) - F(\theta)\|_{L^p} \leq 2M$ hence $II < \frac{\varepsilon}{2}$ by a choice of δ. □

theorems & definitions

Definition 1.1 (Harmonic function) . . . 3
Definition 1.2 (Poisson kernel) . . . 4
Theorem 1.4 (Extension to harmonic functions) . . . 5
Theorem 1.5 (Existence of measure to harmonic functions) . . . 5
Theorem 2.1 (Banach-Alaoglu) . . . 6
Theorem 2.3 (Representation with Lp fuctions) . . . 7
Theorem 2.4 (equivalency in the case of 1) . . . 7
Theorem 2.6 . . . 7
Theorem 2.7 (Extension of unit function to boundary) . . . 8
Theorem 2.8 . . . 9