Solution to Final Exercise — Model Theory (1), 80616

February 28, 2026

Question 1

Urysohn Space. Let $L = \{D_q \mid q \in \mathbb{Q}_{\geq 0}\}$ such that D_q is a binary relation symbol for any q. For any metric space (X, d) we can define L-structure $\mathcal{X}$ with $|\mathcal{X}| = X$ and $(x, y) \in D_q^{\mathcal{X}} \iff d(x, y) = q$.

Part a

Let $\mathcal{K}$ be the class of all finite metric spaces with rational distance.
We will show that $\mathcal{K}$ is a Fraïssé class.

Proof. We will follow the definition of Fraïssé class.

Closure under isomorphism If $\mathcal{X} \in \mathcal{K}$ and $\mathcal{Y} \cong \mathcal{X}$, then let us define $(X, d), (Y, \rho)$ the equivalent metric spaces, and let $f : \mathcal{X} \to \mathcal{Y}$ witness the isomorphism, it follows that,

$$d(x_1, x_2) = q \iff D_q^{\mathcal{X}}(x_1, x_2) \iff D_q^{\mathcal{Y}}(f(x_1), f(x_2)) \iff \rho(f(x_1), f(x_2)) = q$$

hence (Y, ρ) is a finite metric space with rational distance and therefore $\mathcal{Y} \in \mathcal{K}$ as well.

Essential countability Let $X_n = [n]$ and $d : X_n^2 \to \mathbb{Q}$ be a metric, we define $\mathcal{X}_n^d$ as the equivalent of the metric space (X_n, d). Let,

$$\mathcal{M} = \{\mathcal{X}_n^d \mid n < \omega, d \text{ is a metric}\}$$

and notice that for each $n < \omega$, the collection of metric functions defined on X_n is countable as it is a subset of $\mathbb{Q}^{n^2}$. It follows that $\mathcal{M}$ is a countable collection, and notice that $\mathcal{M} \subseteq \mathcal{K}$.
Let $\mathcal{Y} \in \mathcal{K}$, and let (Y, ρ) be the equivalent metric space, notate $n = |Y|$ and let us define $f : [n] \to Y$ a bijection, then,

$$d : X_n^2 \to \mathbb{Q}, \qquad d(x_1, x_2) = \rho(f(x_1), f(x_2))$$

is a rational metric on X_n and thus $\mathcal{X}_n^d \in \mathcal{M}$ with $\mathcal{X}_n^d \cong \mathcal{Y}$.

Hereditary property Assume that $\mathcal{X} \subseteq \mathcal{Y} \in \mathcal{K}$ and show that $\mathcal{X} \in \mathcal{K}$, we omit the requirement that $\mathcal{X}$ is finitely-generated as there are no function symbols and $|Y| < \omega$. Metric property is closed under restrictions then if (Y, ρ) a metric space and $X \subseteq Y$ then $(X, \rho \restriction X^2)$ is a metric space as well.

The joint embedding property Suppose that $\mathcal{X}, \mathcal{Y} \in \mathcal{K}$ and let $(X, d), (Y, \rho)$ be their respective equivalencies. Assume that $X \cap Y \neq \emptyset$ then let $x_0 \in X \cap Y$ be some element, and let $C = A \cup B$, define,

$$f : C^2 \to \mathbb{Q}_{\geq 0}, \qquad f(a, b) = \begin{cases} d(a, b) & a, b \in X \\ \rho(a, b) & a, b \in Y \\ d(a, x_0) + \rho(b, x_0) & a \in X, b \in Y \\ d(b, x_0) + \rho(a, x_0) & b \in X, a \in Y \end{cases}$$

Then (C, f) is a metric space and it follows that $\mathrm{id}_X, \mathrm{id}_Y$ are embeddings of X, Y into C respectively.
If $X \cap Y = \emptyset$ then denote some arbitrary elements $x_0 \in X, y_0 \in Y$, let $f : C^2 \to \mathbb{Q}_{\geq 0}$ defined by,

$$f(a, b) = \begin{cases} d(a, b) & a, b \in X \\ \rho(a, b) & a, b \in Y \\ d(x_0, a) + \rho(y_0, b) + 1 & a \in X, b \in Y \\ d(x_0, b) + \rho(y_0, a) + 1 & \text{otherwise} \end{cases}$$

Thus f is a metric over C, as it is symmetric, non-negative and fulfills triangle inequality. If $\mathcal{C}$ is the structure of (C, f) then $\mathcal{C} \in \mathcal{K}$ and $\mathrm{id}_X, \mathrm{id}_Y$ are embedding of $\mathcal{X}, \mathcal{Y}$ into $\mathcal{C}$.

The amalgamation property Let $\mathcal{X}, \mathcal{Y}, \mathcal{Z} \in \mathcal{K}$ with $(X, d_x), (Y, d_y), (Z, d_z)$ such that $g_x : \mathcal{Z} \to \mathcal{X}, g_y : \mathcal{Z} \to \mathcal{Y}$ are

embeddings. Let $\mathcal{C} \in \mathcal{K}$ as in the last statement, and assume without loss of generality that $\mathcal{C} \subseteq \mathcal{X}, \mathcal{Y}$ (otherwise we can use closure under isomorphism) and therefore g_x, g_y are both identity. Let $f_x : \mathcal{X} \to \mathcal{C}, f_y : \mathcal{Y} \to \mathcal{C}$ be embeddings, it is implied that,

$$f_y \circ g_y = f_y \circ \mathrm{id}_Z = f_x \circ \mathrm{id}_Z = f_x \circ g_x$$

and the property holds as wished. □

Let us denote $\mathbb{U}_\mathbb{Q}$ the Fraïssé limit of $\mathcal{K}$, and let $(U_\mathbb{Q}, d)$ be its equivalent metric space.

Part b

Let (X, d) be a metric space, a Katětov map is a function $f : X \to \mathbb{R}_{\geq 0}$ such that,

$$\forall x, y \in X,\ |f(x) - f(y)| \leq d(x, y) \leq f(x) + f(y)$$

A rational Katětov map is a Katětov map such that $\operatorname{Im} f \subseteq \mathbb{Q}$.
We will show that for every finite $A \subseteq U_\mathbb{Q}$, any rational Katětov map $f : A \to \mathbb{Q}_{\geq 0}$ is realized, namely there is some $b \in U_\mathbb{Q}$ such that $\forall x \in U_\mathbb{Q},\ f(x) = d(x, b)$.

Proof. Note that if $\mathcal{X} \cong \mathcal{Y}$ and $f : \mathcal{X} \to \mathcal{Y}$ isomorphism, then $R_q^{\mathcal{X}}(x, y) \iff R_q^{\mathcal{Y}}(f(x), f(y))$, meaning that f is an isometry of X, Y as metric spaces.
Let f be some rational Katětov map of A. Let $B = A \uplus \{c\}$ for some element, we define a metric on B, $\rho : B^2 \to \mathbb{Q}_{\geq 0}$ by,

$$\rho(x, y) = \begin{cases} d(x, y) & x, y \in A \\ f(x) & x \in A \\ 0 & x = y = c \end{cases}$$

ρ is symmetrical and non-negative, and fulfills the triangle inequality for elements of A, let us verify that for c as well,

$$\rho(c, c) = 0 \leq 2f(x) = \rho(c, x) + \rho(x, c)$$

and,

$$\rho(x, y) = d(x, y) \leq f(x) + f(y) = \rho(x, c) + \rho(c, y)$$

Then (B, ρ) is indeed a metric space, but B is finite and thus $\mathcal{B}$ is isomorphic to some substructure $\mathcal{C} \subseteq \mathbb{U}_\mathbb{Q}$, and let $g : \mathcal{B} \leftrightarrow \mathcal{C}$ denote the isomorphism. $\mathbb{U}_\mathbb{Q}$ is ultra-homogeneous then the isomorphism of $A \to \mathcal{C} \setminus \{g(c)\}$ can be extended to $\sigma \in \operatorname{Aut}(\mathbb{U}_\mathbb{Q})$, and let $e = \sigma^{-1}(c)$. For any $x \in A$, $f(x) = \rho(x, c) = d(g(x), g(c)) = d(\sigma^{-1}(g(x)), \sigma^{-1}(g(c))) = d(x, e)$ as required. □

We will show that $\mathbb{U}_\mathbb{Q}$ is the unique countable rational metric space such that for any $A \subseteq X$ finite and rational Katětov map on A, there is $b \in X$ realizing f.

Proof. Let (X, ρ) be a metric space with this property, we will show that $(X, \rho) \cong (U_\mathbb{Q}, d)$. We start by using induction to show that for every X_n^f (as defined in part 1) there is $Y \subseteq X$ such that $(X_n^f, f) \cong (Y, \rho \restriction Y^2)$. The case of $n = 1$ is trivial as the single metric on single valued space is 0.
Let us assume the claim holds for $m < n$ and let X_n^f be some finite and rational metric space. Take $(X_{n-1}^{f \restriction X_{n-1}^2}, f \restriction X_{n-1}^2)$, we get a sub-metric space of size $n-1$, meaning that there is some such $Y \subseteq X$ and let $\sigma : X_{n-1} \leftrightarrow Y$. Define $g : Y \to \mathbb{Q}_{\geq 0}$ by $g(x) = f(n-1, \sigma^{-1}(x))$, then g is a Katětov map, as a result of the last subpart. There is an element $b \in X$ that realizes g, therefore $\rho(x, b) = g(x) = f(n-1, \sigma^{-1}(x))$, meaning that $f(\sigma^{-1}(\cdot), \sigma^{-1}(\cdot)) = \rho \restriction (Y \cup \{b\})^2$.
We have shown that $\{X_n^f \mid n < \omega, f \text{ is a metric}\} \subseteq \operatorname{Age}(\mathcal{X})$ for $\mathcal{X}$ the equivalent of (X, ρ), therefore $\operatorname{Age}(\mathcal{X}) = \mathcal{K} = \operatorname{Age}(\mathbb{U}_\mathbb{Q})$, then $\mathcal{X} \cong \mathbb{U}_\mathbb{Q}$. □

Part c

Let $\mathbb{U}$ be the completion of $\mathbb{U}_{\mathbb{Q}}$, we will show that any finite metric space (X, d) is isometrically embedded into $\mathbb{U}$.

Proof. The proof is by induction over $n = |X|$. For $n = 1$ the proposition is trivial. Assume that the proposition holds for $m < n$ and let us show the case of n. Let us notate $X = \{x_k \mid k < n\}$ and let $Y = X \setminus \{x_0\}$ and therefore $(Y, d \restriction Y^2)$ is a finite metric space of size $n - 1$. The induction hypothesis holds for Y, then let $\varphi : (Y, d \restriction Y^2) \to \mathbb{U}$ be an isometry embedding of Y into $\mathbb{U}$.

Let $\varepsilon > 0$, For any $0 < m < n$ there is $|u_m^\varepsilon - \varphi(x_m)| < \varepsilon$ such that $u_m^\varepsilon \in U_{\mathbb{Q}}$. By the last part we also deduce that there is $u_0^\varepsilon \in U_{\mathbb{Q}}$ such that if (U, ρ) is the equivalent metric space of $\mathbb{U}$, then,

$$|d(x_i, x_j) - \rho(u_i^\varepsilon, u_j^\varepsilon)| = |\rho(\varphi(x_i), \varphi(x_j)) - \rho(u_i^\varepsilon, u_j^\varepsilon)| \leq |\rho(\varphi(x_i)u_i^\varphi) + \rho(u_i^\varepsilon, u_j^\varepsilon) + \rho(u_j^\varepsilon, \varphi(x_j)) - \rho(u_i^\varepsilon, u_j^\varepsilon)| < 2\varepsilon$$

Define the sequences $\langle x_m^{\frac{1}{k}} \mid 0 < k < \omega \rangle$, then $x_m^{\frac{1}{k}} \to x_m$ and let us define $u_0^{\frac{1}{k}}$ as well. By our definition of $\langle u_0^{\frac{1}{k}} \mid 0 < k < \omega \rangle$, it is a Cauchy sequence and therefore it converges, let us denote its limit with u_0. It follows that $|d(x_i, x_j) - \rho(u_i, u_j)| = 0$, meaning that the map $x_i \mapsto u_i$ is an isometry into $\mathbb{U}$. $\square$

Part d

We will show that for any $A \subseteq \mathbb{U}$ finite and Katětov map f on A is realized by point $b \in \mathbb{U}$.

Proof. Let us define new finite metric space $Y = \{y_n \mid n < |A| + 1\}$ and $\varphi : Y \restriction |A| \to A$ a bijection, and define Y's metric as the one induced by φ as an isometry, in other words $\rho(x, y) = d(\varphi(x), \varphi(y))$ for $x, y \in Y \setminus \{y_n\}$. Let us extend the definition of ρ to Y by setting $\rho(x, y_n) = f(\varphi(x))$ and $\rho(y_n, y_n) = 0$, by part b this is indeed a metric. (Y, ρ) is isometrically embeddable in $\mathbb{U}$ by part c, let us denote this embedding by $g : (Y, \rho) \hookrightarrow (Z, d)$, as a result of the proof of part b we get $g(Y \setminus \{y_n\}) = A$. If $c = g(y_n)$ then by our definition $\rho(x, y_n) = f(\varphi(x)) = d(\varphi(x), c)$, meaning that $\forall a \in A,\ f(a) = d(a, c)$ as required. $\square$

Part e

We will show that every complete separable metric space is isometrically embedded into $\mathbb{U}$.

Proof. By combining part c and d we get that if (X, d) is a finite metric space and $(X \cup \{y\}, f)$ is an extension of (X, d), namely f is a metric and $f \restriction X^2 = d$, then the isometry embedding of (X, d) into $\mathbb{U}$ is a sub-extension of $(X \cup \{y\}, f)$ into $\mathbb{U}$.

Let (X, d) be a countable metric space and suppose that $X = \{x_i \mid i < \omega\}$, then $(\{x_i \mid i < n\}, d \restriction (\{x_i \mid i < n\})^2)$ is isometrically embedded by σ_n into $\mathbb{U}$, and by our assumption $\sigma_i \subseteq \sigma_j$ for all $i < j < \omega$. Let us denote $\sigma = \bigcup_{i<\omega} \sigma_n$, then $\sigma : X \to \mathbb{U}$ is an isometry embedding, as for any choice of points x, y, there is n such that $x, y \in \{x_i \mid i < n\}$.

We can now move to the case of (X, d) complete separable metric space. Take $Y \subseteq X$ dense subset, and let $\sigma : Y \to \mathbb{U}$ be isometry embedding. For each $x \in X$ there is a sequence $(y_n)_{n=1}^\infty \subseteq Y$ such that $y_n \to x$, and therefore $\sigma(y_n) \to b$ for some $b \in \mathbb{U}$, we define $x \mapsto b$ for all $x \in X$. The result is an isometry embedding $\sigma' : X \to \mathbb{U}$. $\square$

Question 2

Groups.

Part a

Let $\mathcal{G}$ be a group in the language $L = \{\cdot\}$ of group theory.

i

Assume that $\mathrm{Th}(\mathcal{G})$ is ω-categorical, we will show that for every finite $A \subseteq G$, it holds that $\mathcal{A} = \langle A \rangle$ is finite, where $\langle A \rangle$ is the subgroup generated by A.

Proof. Assume that $|G| = \omega$, then $\mathcal{G}$ is saturated. We assume for the sake of contradiction that the proposition does not hold, meaning that there is $A \subseteq G$ finite with countable $\langle A \rangle$, but by ω-category $\langle A \rangle = \mathcal{G}$. We define $T : A^{<\omega} \to G$ by,

$$\forall f : [n] \to A, \quad T(f) = \prod_{i=0}^{n-1} f(i)$$

and let $p \in S_1(\mathcal{G})$ defined as completion of,

$$\{T(f) \neq x \mid f \in A^{<\omega}\}$$

p is realized in $\mathcal{G}$, but by p's definition it holds that if $c \in G$ witnessing p then $c \notin \langle A \rangle$, a contradiction. $\square$

ii

Let us assume that for every finite $A \subseteq G$, $|\langle A \rangle|$ is finite as well.
We will find a counter example to the claim that $\mathrm{Th}(G)$ is ω-categorical.
Solution. Let μ_p denote Prüfer p-group over $\mathbb{C}$, namely,

$$\mu_p = \{z \in \mathbb{C} \mid z^p = 1\}$$

and $\mu_\infty = \bigcup_{0<n} \mu_p$. Each finite $A \subseteq \mu_\infty$ equals to $\mu_{\gcd A}$, in particular $\langle A \rangle$ is finite.
Notice $x^n = 1 \iff x^{n+1} = x$, meaning that order of elements is definable, and let,

$$\varphi_n(x) = x^{n+1} = x$$

Therefore $\varphi_n^G(a) \iff o(a) = n$ for any $a \in G$, each element has a unique order ($T \models \forall x\ \varphi_n(x) \to (\neg\varphi_m(x))$ for any $n \neq m$), then $\varphi_n \not\equiv_T \varphi_m$. It implies that there are at least countably many non-equivalent formulas over single variable in T, by Ryll-Nardzewski theorem T is not ω-categorical.

Part b

Let $\mathcal{G}$ be a group and let us assume that $\mathrm{Th}(\mathcal{G})$ is totally transcendental. We will show that there is no infinite strictly decreasing sequence of definable subgroups $\langle H_n \mid n < \omega \rangle$.

Proof. We will show that if $\mathcal{G}$ has definable strictly decreasing subgroups then $\mathrm{Th}(\mathcal{G})$ has the binary tree property. Let us assume that $\langle H_n \mid n < \omega \rangle$ is such a sequence, meaning that $\mathcal{G} = \mathcal{H}_0 > \mathcal{H}_1 > \cdots$. H_n is definable and let $\varphi_n(x)$ witness it, meaning that $\mathcal{G} \models \varphi_i(x) \to \varphi_j(x)$ for any $i < j < \omega$. Let $\psi_n(x) = \varphi_n(x) \wedge (\neg\varphi_{n+1}(x))$, that is $\mathcal{M} \models \psi_n(x)$ if $x \in H_n \setminus H_{n+1}$, the sequence is strictly decreasing therefore $\mathcal{G} \models \exists x\ \psi_n(x)$ for any n.
We denote h_n as a witness of ψ_n, meaning that $h_n \in H_n \setminus H_{n+1}$, it follows that $h_n H_{n+1}$ is a proper coset of H_{n+1} in H_n (and in G). So $H_{n+1} \cap h_n H_{n+1} = \emptyset$, and define $\varphi_{n+1}^{h_n}(x) = \varphi_{n+1}(h_n^{-1}x)$, hence $\mathcal{G} \models \varphi_{n+1}^g(x) \iff x \in h_n H_{n+1}$. Given

a formula φ_n^g for some $g \in H$, we can define the following formulas,

$$\varphi_{n+1}^g(x) = \varphi_{n+1}(g^{-1}x), \qquad \varphi_{n+1}^{h_n g}(x) = \varphi(g^{-1}h_n^{-1}x)$$

By $H_{n+1} \subsetneq H_n$ we get that $gH_{n+1} \subsetneq gH_n$, therefore $\{\varphi_n^g, \varphi_{n+1}^g\}$ is consistent, We also get that $\{\varphi_n^g, \varphi_{n+1}^{h_n g}\}$ is consistent as a result of the previous statement, but $\mathcal{G} \models \varphi_{n+1}^g(x) \wedge \varphi_{n+1}^{h_n g}(x) \iff x \in gH_n, h_n gH_n$, but $gH_n \cap h_n gH_n \neq \emptyset \iff h_n H_n \cap H_n \neq \emptyset$, implying that $\varphi_{n+1}^g \wedge \varphi_{n+1}^{h_n g}$ is inconsistent.

We can now get to the construction of the binary tree. Let $T : 2^{<\omega} \to G$ be defined as,

$$T(\eta) = \prod_{n \in \eta^{-1}(\{1\})} h_n$$

and let $\Gamma : 2^{<\omega} \to \mathrm{form}_L$ be defined by,

$$\forall \eta : 2^n \to G, \quad \Gamma(\eta) = \varphi_{n+1}^{T(\eta)}$$

Γ fulfills the binary tree property, and therefore witnessing that $\mathcal{G}$ is not totally transcendental.

The proposition holds by contra positive principle. □

Part c

Let G be an infinite group and let $L = \{1, \cdot, ^{-1}\}$. We will show that if $\mathrm{Th}(G)$ admits quantifier elimination then G is Abelian.

Proof. Let $\varphi(x)$ be some formula in sent_L, then φ is equivalent to one of the formulas,

$$\bigwedge_{n \in A} x^n \neq 1, \qquad \bigvee_{n \in A} x^n = 1.$$

as a result of quantifier elimination, where $A \subseteq \omega \setminus \{0\}$ is finite. In other words $T \models \varphi(x) \iff o(x) \in A_\varphi$ such that A_φ is finite or co-finite, for any $\varphi \in \mathrm{sent}_L$.

Let $\theta(x) = \forall y\ x \cdot y = y \cdot x$, that is $\theta^G(x) \iff x \in Z(G)$. We get from $o(1) = 1$ that $1 \in A_\theta$, as well $2 \in A_\theta$ as $o(x) = 2$ if and only if $x = x^{-1}$.

We assume for contradiction that G is not Abelian.
Assume that G has an infinite order element $a \in G$. If $a \in Z(G)$ then $o(a) \neq n \iff n \notin A_\theta$ for all n, then $A_\theta = \emptyset$, implying that $o(g) = 0 \iff g \in Z(G)$ for all $g \in G$. If $g \in G$ is of finite order n, then if $a \cdot g$ is of finite order m we get,

$$(ag)^{nm} = 1 \iff a^{nm}g^{nm} = 1 \iff a^{nm} = 1.$$

then ag is of an infinite order and therefore $\in Z(G)$, then $g \in Z(G)$ as well, but $o(g) = n \notin A_\theta$ in contradiction, thus there is no element of infinite order in $Z(G)$.

$Z(G)$ does not contain infinite order element. Let us assume that $a \in G \setminus Z(G)$ is of infinite order, for the same reason as before elements are of infinite order if and only if they are not commutative. There wasn't a lot of types in this theory in the first place, now we know that the single type there is isolated, and therefore each model of T is atomic. Every countable and atomic model is prime, therefore any two countable models of $\mathrm{Th}(G)$ are mutually embedded in each other, in other words $\mathrm{Th}(G)$ is ω-categorical. By part a of the question $\langle a \rangle$ is finite in contradiction, therefore there is no such a, it follows that there are no infinite order elements in G.

Let us define $A = \{n < \omega \mid \exists g \in G,\ o(g) = n\}$. Let us define the partial type $p(x) = \{x^n = 1 \mid n \in A\}$, then the theory $\mathrm{Th}(G) \cup p(c)$ for new constant symbol c is finitely satisfiable therefore satisfiable by a countable model H. if we omit c from the language of H we get $H \models \mathrm{Th}(G)$ and if A is unbound then H has infinite order element in direct contradiction to our last claim. We deduce that A is finite, and denote $M = \max A < \omega$, then $A_\theta \subseteq A$, in particular A_θ is bounded by M as well.

Let us denote $H = Z(G)$ and $K = \langle G \setminus Z(G) \rangle = \{1\} \cup (G \setminus Z(G))$, then $HK = G$, $H \cap K = \{1\}$, $H \triangleleft G$ as its center, and $K \triangleleft G$ as if $g \in K$ then $gK = K = Kg$ and if $g \in Z(G)$ then $gK = Kg$. Therefore $G \simeq H \times K$, so for any $x \in H, y \in K$ we get $o(xy) = o((x, y)) = \gcd(o(x), o(y)) \in A_\theta$, but A_θ is closed under squares, then $A_\theta = A$. $\square$

Question 3

Part a

Let (X, cl) be a pre-geometry, we will show that there is a basis for X and that every two bases have the same cardinality.

Proof. We say that a set $B \subseteq X$ is a basis if $\mathrm{cl}(B) = X$ and $\forall b \in B,\ b \notin \mathrm{cl}(B \setminus \{b\})$ (independent). We will show that B is a basis if and only if B is maximally independent.
Assume that B is a basis, and let $x \in X \setminus B$, then $x \in \mathrm{cl}(B)$ and therefore B is maximally independent.
In the other direction, let us assume that B is maximally independent, we will show that $\mathrm{cl}(B) = X$. We assume towards contradiction that there is $x \in X \setminus \mathrm{cl}(B)$, and let $B' = B \cup \{x\}$, thus $x \in \mathrm{cl}(B')$ but $x \notin \mathrm{cl}(B' \setminus \{x\}) = \mathrm{cl}(B)$. For any $b \in B$ we already assumed that $b \notin \mathrm{cl}(B \setminus \{b\})$, therefore B' is independent and $B' \supsetneq B$, a contradiction.

Let $\mathcal{B} = \{B \subseteq X \mid B \text{ is independent}\}$, notice that $\emptyset$ is independent then $\mathcal{B}$ is not empty. In intention of using Zorn's lemma, let $C \subseteq \mathcal{B}$ be a chain under $\subseteq$, and let $B^* = \bigcup C$. If $b \in B^*$, then,
$\mathrm{cl}(B^* \setminus \{b\}) = \bigcup\{\mathrm{cl}(B_0) \mid B_0 \subseteq B^* \setminus \{b\}, |B_0| < \omega\} \overset{(1)}{=} \bigcup\{\mathrm{cl}(B_0) \mid \exists B \in C\ B_0 \subseteq B \setminus \{b\}, |B_0| < \omega\} = \bigcup\limits_{B \in C} \mathrm{cl}(B \setminus \{b\})$
where (1) is following from the fact that if B_0 is finite and $B_0 \subseteq \bigcup C$ then there is an element in the chain such that $B_0 \subseteq B \in C$. All $B \in C$ are independent then $b \notin \mathrm{cl}(B \setminus \{b\})$, it follows that $b \notin \mathrm{cl}(B^* \setminus \{b\})$ as well, B^* is independent and maximal. The conditions for Zorn's lemma are fulfilled, as a result $\mathcal{B}$ has a maximal element, in other words there is $B \in \mathcal{B}$ that is maximally independent, and thus a basis.

Let $B_1, B_2 \subseteq X$ be bases for (X, cl), we will show that $|B_1| = |B_2|$. Let $B_1 = \{b_\alpha \mid \alpha < \kappa\}$, for any α, $b_\alpha \in \mathrm{cl}(B_2)$, so there is finite set $B_0^\alpha \subseteq B_2$ such that $b_\alpha \in \mathrm{cl}(B_0^\alpha)$. Take $C = \bigcup_{\alpha<\kappa} B_0^\alpha$, then $B \subseteq \mathrm{cl}(C) \implies X = \mathrm{cl}(B) \subseteq \mathrm{cl}(\mathrm{cl}(C)) = \mathrm{cl}(C) \implies \mathrm{cl}(C) = X$ and $|C| = \kappa$ as a union of finite sets. Notice that $C \subseteq B_2$, then $|B_1| = |C| \leq |B_2|$, by symmetry it also follows that $|B_2| \leq |B_1|$ and by CSB $|B_1| = |B_2|$. $\square$

Part b

Let T be a countable complete $\aleph_1$-categorical theory. We will show that T has at most $\aleph_0$ different isomorphism classes for countable models.

Proof. We assume for the sake of contradiction that $\langle \mathcal{M}_\alpha \mid \alpha < \omega_1 \rangle$ is a sequence of non-isomorphic countable models. T is $\aleph_1$-categorical and thus its only model is saturated and $\aleph_1$-stable, we infer that T is totally transcendental and that there is a prime countable model $\mathcal{M} \models T$. We can also deduce that there is a strongly minimal formula $\varphi(x, a)$ over $\mathcal{M}$, where $a \in M$ is a parameter. $\mathcal{M}$ is prime then there is an embedding of $\mathcal{M}$ into $\mathcal{M}_\alpha$ for all α, by closure to larger models the embedding of a, $a \mapsto a_\alpha$ holds that $\varphi(x, a_\alpha)$ is strongly minimal in $\mathcal{M}_\alpha$.

We will show the lemma that if $\varphi(x)$ is strongly minimal in a structure $\mathcal{M}$, then $\mathrm{acl}_\varphi : \mathcal{P}(X) \to \mathcal{P}(X)$ defined by $\mathrm{acl}_\varphi(B) = \varphi(M) \cap \mathrm{acl}(B)$ where $X = \varphi(M)$, is a closure.
If $A \subseteq B \subseteq X$ then,

$$\mathrm{acl}_\varphi(A) = \varphi(M) \cap \mathrm{acl}(A) \subseteq \varphi(M) \cap \mathrm{acl}(B) = \mathrm{acl}_\varphi(B)$$

$A \subseteq X = \varphi(M)$ and $A \subseteq \mathrm{acl}(A)$ by acl's definition, therefore $A \subseteq \mathrm{acl}_\varphi(A)$ as well. acl_φ is also closed as we have shown in former exercise that $\mathrm{acl}(\mathrm{acl}(A)) = \mathrm{acl}(A)$. We deduce that acl_φ is a closure, and it remains to show finite character and exchange properties.
Let $A \subseteq X$ and let $b \in \mathrm{acl}_\varphi(A)$, then there is a formula $\psi(x)$ with parameters from A such that $\psi(X)$ is finite and $b \in \psi(X)$, therefore $\mathrm{acl}_\varphi(A) \subseteq \bigcup\{\mathrm{acl}_\varphi(A_0) \mid A_0 \subseteq A, A_0 \text{ is finite}\}$, the other direction is direct.
Let us assume that $a \in \mathrm{acl}_\varphi(A \cup \{b\}) \setminus \mathrm{acl}_\varphi(A)$, then there is a formula $\psi(x, y)$ such that $\psi(x, b)$ is algebraic and $\psi(a, b)$

holds. In turn, $b \in \psi(a, X)$ and φ is strongly minimal so $\psi(a, y) \wedge \varphi(y)$ is (without loss of generality) finite and witness $b \in \mathrm{acl}_\varphi(A \cup \{a\})$. This conclude our lemma.

Returning to our initial proposition, let us observe $(\varphi(M_\alpha, a_\alpha), \mathrm{acl}_{\varphi(x,a_\alpha)})$, this is a pre-geometry according to the lemma, thus it has a basis of constant size, let us denote $n_\alpha = \dim(\varphi(M_\alpha, a_\alpha), \mathrm{acl}_{\varphi(x,a_\alpha)})$, in particular $n_\alpha \leq |M_\alpha| = \omega$. By the assumption that there are ω_1 equivalency classes we get that there are $\alpha \neq \beta < \omega_1$ such that $n_\alpha = n_\beta$.
Let B_α, B_β be the respective bases, there is $f : B_\alpha \to B_\beta$ a bijection. For each $t \in M_\alpha$ there is an algebraic formula $\phi(x)$ such that $\phi^{\mathcal{M}_\alpha}(t)$ and therefore t is atomic over B_α. By our assumption $f(tp(t/B_\alpha))$ is realized and let $s \in M_\beta$ denote its realization, then f can be extended with $\langle t, s\rangle$, implying that f has an extension to the elementary injection $f' : \mathrm{acl}_{\varphi(x,a_\alpha)}(B_\alpha) \to \mathrm{acl}_{\varphi(x,a_\beta)}(B_\beta)$, that is $f' : \mathcal{M}_\alpha \to \mathcal{M}_\beta$. By using back and forth we can extend f' to a bijection, thus $\mathcal{M}_\alpha \cong \mathcal{M}_\beta$ in contradiction to our assumption that $\alpha = \beta \iff \mathcal{M}_\alpha \cong \mathcal{M}_\beta$. $\square$