

REPUBLIQUE DU CAMEROUN

*Paix-Travail-patrie*

UNIVERSITE DE YAOUNDE I

FACULTE DE MEDECINE ET DES  
SCIENCES BIOMEDICALES



REPUBLIC OF CAMEROON

*Peace-Work-Fatherland*

UNIVERSITY OF YAOUNDE I

FACULTY OF MEDICINE AND  
BIOMEDICAL SCIENCES

DEPARTMENT OF PEDIATRICS

# **PEDIATRICIANS' KNOWLEDGE, ATTITUDES AND PRACTICES TOWARDS PARAFUNCTIONAL ORAL HABITS AND DENTAL MALOCCLUSIONS IN CHILDREN**

*Thesis presented and publicly defended in view of obtaining a doctorate in dental medicine*

*by*

**AWA CLAVICE WOPONG**

*Matricule: 17M169*

## **Supervisor**

**Pr. KOKI Ndombo Paul**  
Professor, Pediatrics

## **Co-Supervisors**

**Dr. TONY NENGOM Jocelyn**  
Lecturer, Pediatrics

**Dr. LOWE Jacqueline Michele**  
Lecturer, Pediatric Dentistry


Academic year: 2023 -2024

REPUBLIQUE DU CAMEROUN

*Paix-Travail-patrie*

UNIVERSITE DE YAOUNDE I

FACULTE DE MEDECINE ET DES  
SCIENCES BIOMEDICALES



REPUBLIC OF CAMEROON

*Peace-Work-Fatherland*

UNIVERSITY OF YAOUNDE I

FACULTY OF MEDICINE AND  
BIOMEDICAL SCIENCES

DEPARTMENT OF PEDIATRICS

# **PEDIATRICIANS' KNOWLEDGE, ATTITUDES AND PRACTICES TOWARDS PARAFUNCTIONAL ORAL HABITS AND DENTAL MALOCCLUSIONS IN CHILDREN**

## **Jury:**

### **President**

Pr ZE Mikande

### **Supervisor**

Pr KOKI Ndombo Paul

### **Member**

Dr Mengong Hortense

## **Supervisor**

**Pr. KOKI Ndombo Paul**

Professor, Pediatrics

## **Co-Supervisors**

**Dr. TONY NENGOM Jocelyn**

Lecturer, Pediatrics

**Dr. LOWE Jacqueline Michele**

Lecturer, Pediatric Dentistry

**Academic year: 2023 -2024**

## TABLE OF CONTENT

TABLE OF CONTENT .....	III
DEDICATION .....	VI
ACKNOWLEDGEMENT .....	VII
THE PHYSICIAN'S OATH .....	XXVI
LIST OF TABLES .....	XXVII
LIST OF FIGURES .....	XXVIII
LIST OF ABBREVIATIONS .....	XXIX
ABSTRACT .....	XXX
RÉSUMÉ .....	XXXI
CHAPTER I: INTRODUCTION .....	- 1 -
I.1 BACKGROUND .....	- 2 -
I.2 JUSTIFICATION .....	- 3 -
I.3 RESEARCH QUESTION .....	- 3 -
I.4 OBJECTIVES .....	- 3 -
CHAPTER II: LITERATURE REVIEW .....	- 4 -
II.1 THE ORAL CAVITY .....	- 5 -
II.2 CRANIOFACIAL GROWTH AND DEVELOPMENT .....	- 8 -
II.3 TOOTH DEVELOPMENT .....	- 17 -
II.4 TOOTH ERUPTION .....	- 21 -

II.5 HUMAN DENTITION AND OCCLUSION .....	- 25 -
II.6 PARAFUNCTIONAL HABITS.....	- 27 -
II.7 MALOCCLUSION .....	- 31 -
CHAPTER III: METHODOLOGY .....	<b>ERREUR ! SIGNET NON DEFINI.</b>
III.1 STUDY TYPE.....	- 35 -
III.2 STUDY SITE .....	- 35 -
III.3 STUDY PERIOD .....	- 35 -
III.4 STUDY POPULATION .....	- 35 -
III.5 PROCEDURE .....	- 36 -
III.6 MATERIALS .....	- 36 -
III.7 ETHICAL CONSIDERATIONS .....	- 38 -
CHAPTER IV: RESULTS .....	- 39 -
IV.1. STUDY PARTICIPANTS .....	- 40 -
IV.2 SOCIODEMOGRAPHIC CHARACTERISTICS OF THE STUDY POPULATION ..	- 41 -
IV.3. RESULTS ON KNOWLEDGE OF PARAFUNCTIONAL ORAL HABITS AND DENTAL MALOCCLUSIONS AND THEIR RELATIONSHIP .....	- 43 -
IV.4. RESULTS ON ATTITUDE TOWARDS PARAFUNCTIONAL HABITS AND MALOCCLUSIONS .....	- 46 -
CHAPTER V; DISCUSSION .....	- 50 -
5.1 SOCIO-PROFESSIONAL DATA .....	- 51 -
5.2 KNOWLEDGE .....	- 52 -

5.3 ATTITUDE .....	- 53 -
5.4 PRACTICE.....	- 53 -
CHAPTER VI: CONCLUSION AND RECOMMENDATIONS .....	- 55 -
CONCLUSION .....	- 56 -
RECOMMENDATIONS .....	- 56 -
REFERENCES .....	- 57 -
APPENDIX .....	- 57 -

**DEDICATION**

This work is dedicated to my parents;

Mr. Awa Eric Fonkam and Mrs Awa Melvis Ngwi.

## ACKNOWLEDGEMENT

I wish to extend my sincere and heart felt gratitude to:

GOD Almighty, for His presence in my life as well as the grace, strength, and favor He gave me throughout my medical training and the period of this work.

Pr. KOKI Ndombo Paul, the supervisor of this research work. You are an admirable professor and researcher, an epitome of knowledge and a source of inspiration and encouragement. It was an honour working with you.

Dr. TONY Jocelyn, for accepting to co-direct and for being there from the start of this work till the end. Thank you for painstakingly going through this work every step of the way and for your ready availability and selfless support.

Dr LOWE Michele for equally co-directing and giving constructive guidance to the completion of this work. Thank you, a lot, for your patience.

The president and members of Jury for accepting to scrutinize this work in a bid to make it better.

The Dean, Pr ZE MINKANDE Jacqueline for the management and training she offers to the Faculty of Medicine and Biomedical Sciences

All the teaching and administrative staff of the FMBS for your relentless efforts and sacrifices in the correct programming and dispensation of the different lectures during our training.

All the pediatricians who consented to participate in this research, this work could be possible without your accord. Thank you for your trust.

My beloved father, Mr. AWA Eric for your support and encouragements. Your strength and unwavering confidence in me have pushed me to be everything I am today. Words will never be enough.

My sweet mother, Mrs AWA Melvis Ngwi for constantly praying for me. You have been an epitome of strength and my pillar.

My siblings, Awa Eric Angelo, Awa Praise, Awa Daizy, Awa Mark-Shalom and Afanyu Divine Favour for their love for me and for being my source of inspiration.

Mr FK Joshua for standing by me, making me a priority despite your busy schedule and always pushing me to be the best of myself.

My grandmum, Mrs Naah Rebecca- your ageless strength and wisdom constantly inspire me.

My entire family for always being there for me. For all the prayers, encouragement, trust, I am grateful.

Mrs Timah Miriam, Miss Tuku Melvis and Mr Ngwa Godwill for your constant prayers and unconditional love.

My oldest friends, Miss Fowedlung Tsonju Agafina, Miss Nya Tounda Laurie and Miss Nanje Maxmine for your strong and resilient support.

My mentors, Dr LONI Ekali Gabriel and Dr. Ekali Solange. You have been a true source of inspiration to me.

My senior doctors; Dr Ashu George, Dr Gwan Emmanuel and Dr Basambo Fabrice for your meticulous corrections and encouragement.

My classmate, friend and sister Meniemoh Ranebel. Your friendship has been a major driving force throughout this journey. I'm grateful.

My 4.9 BBC Sisters; Leyuga Senkaa, Fondzewong Larissa, Ayeni Doris, Thom Claude and Sake Jolie Coeur - I couldn't have wished for a better circle.

The entire BBC Family for your support, mentorship, and encouragements.

The 49<sup>th</sup> batch of the Faculty of Medicine and Biomedical Sciences of the University of Yaoundé 1, it has been a great experience.



## THE ADMINISTRATIVE AND TEACHING STAFF

### ADMINISTRATIVE STAFF

Dean	Pr. ZE MINKANDE Jacqueline
Vice-Dean in charge of programming and academic activities	Pr. NTSAMA ESSOMBA Claudine Mireille
Vice-Dean in charge of Students' records, follow-up, and statistics	Pr. NGANOU Chris Nadège épse GNINDJIO
Vice-Dean in charge of Co-operation and Research	Pr. ZEH Odile Fernande
Director of Academics, Student records and Research	Dr VOUNDI VOUNDI Esther
Director of Administrative and Financial Affairs:	Mme ESSONO EFFA Muriel Glawdis
General Coordinator of the Specialization Cycle	Pr NJAMNSHI Alfred KONGNYU
Chief of service, Finance	Mme NGAMLI NGOU Mireille Albertine épouse WAH
Deputy Chief of service, Finance	Mme MANDA BANA Marie Madeleine épouse ENGUENE
Chief of service, Administration and Personnel	Pr. SAMBA Odette NGANO ép. TCHOUAWOU

Chief of service, Certifications	Mrs ASSAKO Anne DOOBA
Deputy chief of service, Certifications	Dr NGONO AKAM MARGA Vanina
Chief of service, Student records and statistics	Mme BIENZA Aline
Deputy Chief of service, Student records and statistics:	Mme FAGNI MBOUOMBO AMINA épouse ONANA
Chief of service, Equipment and Maintenance	Mrs HAWA OUMAROU
Deputy Chief of service, Equipment and Maintenance	Dr NDONGO née Mpono EMENGUELE
Interim Chief Librarian	Mrs FROUISOU née MAME Marie-Claire
Stores Accountant	Mr MOUMEMIE NJOUNDIYIMOUN MAZOU
PROGRAM AND SPECIALTY TRAINING COORDINATORS	
Coordinator, Oral Medicine	Pr. BENGONDO MESSANGA Charles
Coordinator, Pharmacy	Pr. NTSAMA ESSOMBA Claudine
Coordinator, Intern Specialisation Cycle	Pr. ONGOLO ZOGO Pierre
Coordinator, Specialization in Morbid Anatomy/Pathology	Pr. SANDO Zacharie
Coordinator, Specialization in Anaesthesia and Critical Care	Pr. ZE MINKANDE Jacqueline
Coordinator, Specialization in General Surgery	Pr. NGO NONGA Bernadette
Coordinator, Specialization in Gynaecology and Obstetrics	Pr DOHBIT Julius SAMA
Coordinator, Specialization in Internal Medicine	Pr. NGANDEU Madeleine

Coordinator, Specialization in Paediatrics	Pr. MAH Evelyn MUNGYEH
Coordinator, Specialization in Clinical Laboratory Sciences	Pr. KAMGA FOUAMNO Henri Lucien
Coordinator, Specialization in Radiology and Medical Imaging	Pr. ONGOLO ZOGO Pierre
Coordinator, Specialization in Public Health: Coordinator, Post-Graduate Education Program:	Pr. TAKOUGANG Innocent Pr. KASIA Jean Marie
Point focal projet CESSI Pedagogic Manager	Pr NGOUPAYO Joseph Pr. ANKOUANE ANDOULO Firmin

#### HONORARY DIRECTORS OF CUSS (University Centre for Health sciences)

Pr. MONEKOSSO Gottlieb (1969-1978)

Pr. EBEN MOUSSI Emmanuel (1978-1983)

Pr. NGU LIFANJI Jacob (1983-1985)

Pr. CARTERET Pierre (1985-1993)

#### HONORARY DEANS OF FMBS (Faculty of Medicine and Biomedical Sciences)

Pr. SOSSO Maurice Aurélien (1993-1999)

Pr. NDUMBE Peter (1999-2006)

Pr. TETANYE EKOE Bonaventure (2006-2012)

Pr. EBANA MVOGO Côme (2012-2015)

## TEACHING STAFF

N°	NAMES	RAN	FIELD
	K		
	DEPARTMENT OF SURGERY AND SPECIALTIES		
1	SOSSO Maurice Aurélien (HD)	P	General Surgery
2	DJIENTCHEU Vincent de Paul	P	Neurosurgery
3	ESSOMBA Arthur (Interim HD)	P	General Surgery
4	HANDY EONE Daniel	P	Trauma/Orthopaedic Surgery
5	MOUAFO TAMBO Faustin	P	Paediatric Surgery
6	NGO NONGA Bernadette	P	General Surgery
7	NGOWE NGOWE Marcellin	P	General Surgery
8	OWONO ETOUNDI Paul	P	Anaesthesia-Critical Care
9	ZE MINKANDE Jacqueline	P	Anaesthesia-Critical Care
10	BAHEBECK Jean	AP	Trauma/Orthopaedic Surgery
11	BANG GUY Aristide	AP	General Surgery
12	BENGONO BENGONO Roddy	AP	Anaesthesia-Critical Care
	Stéphan		
13	FARIKOU Ibrahima	AP	Orthopaedic Surgery
14	JEMEA Bonaventure	AP	Anaesthesia-Critical Care
15	BEYIHA Gérard	AP	Anaesthesia-Critical Care
16	EYENGA Victor Claude	AP	Surgery/ Neurosurgery
17	GUIFO Marc Leroy	AP	General Surgery
18	NGO YAMBEN Marie Ange	AP	Trauma/Orthopaedic Surgery
19	TSIAGADIGI Jean Gustave	AP	Trauma/Orthopaedic Surgery
20	BELLO FIGUIM	SL	Neurosurgery
21	BIWOLE BIWOLE Daniel	SL	General Surgery
	Claude Patrick		
22	FONKOUÉ Loïc	SL	Trauma/Orthopaedic Surgery
23	KONA NGONDO François	SL	Anaesthesia-Critical Care
	Stéphane		

24	MBOUCHE Landry Oriole	SL	Urology
25	MEKEME MEKEME Junior Barthelemy	SL	Urology
26	MULUEM Olivier Kennedy	SL	Trauma/ Orthopaedic Surgery
27	SAVOM Eric Patrick	SL	General Surgery
28	AHANDA ASSIGA	L	General Surgery
29	AMENGLE Albert Ludovic	L	Anaesthesia-Critical Care
30	BIKONO ATANGANA Ernestine Renée	L	Neurosurgery
31	BWELE Georges	L	General Surgery
32	EPOUPA NGALLE Frantz Guy	L	Urology
33	FOUDA Jean Cédric	L	Urology
34	IROUME Cristella Raïssa BIFOUNA Epse NTYO'O NKOUMOU	L	Anaesthesia-Critical Care
35	MOHAMADOU GUEMSE Emmanuel	L	Trauma/ Orthopaedic Surgery
36	NDIKONTAR KWINJI Raymond	L	Anaesthesia-Critical Care
37	NWAHA MAKON Axel Stéphane	L	Urology
38	NYANIT BOB Dorcas	L	Pediatric Surgery
39	OUMAROU HAMAN NASSOUROU	L	Neurosurgery
40	ARROYE BETOU Fabrice Stéphane	L	Thoracic and Cardiovascular Surgery
41	ELA BELLA Amos Jean-Marie	L	Thoracic Surgery
42	FOLA KOPONG Olivier	L	Surgery
43	FOSSI KAMGA GACELLE	L	Pediatric Surgery
44	GOUAG	L	Anaesthesia-Critical Care
45	MBELE Richard II	L	Thoracic Surgery

46	MFOUAPON EWANE Hervé Blaise	L	Neurosurgery
47	NGOUATNA DJEUMAKOU Serge Rawlings	L	Anaesthesia-Critical Care
48	NYANKOUE MEBOUINZ Ferdinand	L	Trauma/Orthopaedic Surgery
DEPARTMENT OF INTERNAL MEDICINE AND SPECIALTIES			
49	SINGWE Madeleine Epse NGANDEU (HD)	P	Internal Medicine/Rheumatology
50	ANKOUANE ANDOULO	P	Internal Medicine/Gastroenterology and Hepatology
51	ASHUNTANTANG Gloria Enow	P	Internal Medicine/Nephrology
52	BISSEK Anne Cécile	P	Internal Medicine/Dermatology
53	KAZE FOLEFACK François	P	Internal Medicine/Nephrology
54	KUATE TEGUEU Calixte	P	Internal Medicine/Neurology
55	KOUOTOU Emmanuel Armand	P	Internal Medicine/Dermatology
56	MBANYA Jean Claude	P	Internal Medicine/Endocrinology
57	NDJITOYAP NDAM Elie Claude	P	Internal Medicine/Gastroenterology and Hepatology
58	NDOM Paul	P	Internal Medicine/Oncology
59	NJAMNSHI Alfred KONGNYU	P	Internal Medicine/Neurology
60	NJOYA OUDOU	P	Internal Medicine/Gastroenterology

61	SOBNGWI Eugène	P	Internal Medicine/Endocrinology
62	PEFURA YONE Eric Walter	P	Internal Medicine/Pneumology
63	BOOMBHI Jérôme	AP	Internal Medicine/Cardiology
64	FOUDA MENYE Hermine Danielle	AP	Internal Medicine/Nephrology
65	HAMADOU BA	AP	Internal Medicine/Cardiology
66	MENANGA Alain Patrick	AP	Internal Medicine/Cardiology
67	NGANOU Chris Nadège	AP	Internal Medicine/Cardiology
68	KOWO Mathurin Pierre	AP	Internal Medicine/Gastroenterology and Hepatology
69	KUATE née MFEUKEU KWA Liliane Claudine	AP	Internal Medicine/Cardiology
70	NDONGO AMOUGOU Sylvie	AP	Internal Medicine/Cardiology
71	DEHAYEM YEFOU Mesmin	SL	Internal Medicine/Endocrinology
72	ESSON MAPOKO Berthe Sabine Epse PAAMBOG	SL	Internal Medicine/Oncology
73	ETOA NDZIE Epse ETOGA Martine Claude	SL	Internal Medicine/Endocrinology
74	MAÏMOUNA MAHAMAT	SL	Internal Medicine/Nephrology
75	MASSONGO MASSONGO	SL	Internal Medicine/Pneumology
76	MBONDA CHIMI Paul-Cédric	SL	Internal Medicine/Neurology
77	NDJITOYAP NDAM Antonin Wilson	SL	Internal Medicine/Gastroenterology
78	NDOBO Epse KOE Juliette Valérie Danielle	SL	Internal Medicine/Cardiology
79	NGAH KOMO Elisabeth	SL	Internal Medicine/Pneumology
80	NGARKA Léonard	SL	Internal Medicine/Neurology

81	NKORO OMBEDE Grâce Anita	SL	Internal Medicine/Dermatology
82	OWONO NGABEDE Amalia Ariane	SL	Internal Medicine/Interventional Cardiology
83	NTSAMA ESSOMBA Marie Josiane Epse EBODE	SL	Internal Medicine/Gériatrics
84	ATENGUENA OBALEMBA Etienne	L	Internal Medicine/Oncology
85	FOJO TALONGONG Baudelaire	L	Internal Medicine/Rheumatology
86	KAMGA OLEN Jean Pierre Olivier	L	Internal Medicine/Psychiatry
87	MENDANE MEKOBÉ Francine Epse EKOBEA	L	Internal Medicine/Endocrinology
88	MINTOM MEDJO Pierre Didier	L	Internal Medicine/Cardiology
89	NTONE ENYIME Félicien	L	Internal Medicine/Psychiatry
90	NZANA Victorine Bandolo Epse FORKWA MBAH	L	Internal Medicine/Nephrology
91	ANABA MELINGUI Victor Yves	L	Internal Medicine/Rheumatology
92	EBENE MANON Guillaume	L	Internal Medicine/Cardiology
93	ELIMBY NGANDE Lionel Patrick Joël	L	Internal Medicine/Nephrology
94	KUABAN Alain	L	Internal Medicine/Pneumology
95	NKECK Jan René	L	Internal Medicine
96	NSOUNFON ABDOU WOUOLIYOU	L	Internal Medicine/Pneumology
97	NTYO'O NKOUMOU Arnaud Laurel	L	Internal Medicine/Pneumology



98	TCHOUANKEU KOUNGA Fabiola	L	Internal Medicine/Psychiatry
DEPARTMENT OF MEDICAL IMAGING AND RADIOLOGY			
99	ZEH Odile Fernande (HD)	P	Radiology/Medical Imaging
100	GUEGANG GOUJOU. Emilienne	P	Medical Imaging/Neuroradiology
101	MOIFO Boniface	P	Radiology/Medical Imaging
102	ONGOLO ZOGO Pierre	AP	Radiology/Medical imaging
103	SAMBA Odette NGANO	AP	Biophysics/Medical Physics
104	MBEDE Maggy Epse ENDEGUE MANGA	SL	Radiology/Medical Imaging
105	MEKA'H MAPENYA Ruth- Rosine	SL	Radiotherapy
106	NWATSOCK Joseph Francis	L	Radiology/Nuclear Medicine
107	SEME ENGOUMOU Ambroise Merci	L	Radiology/Medical Imaging
108	ABO'O MELOM Adèle Tatiana	L	Radiology/Medical Imaging
DEPARTMENT OF OBSTETRICS AND GYNECOLOGY			
109	NGO UM Esther Juliette épouse MEKA (HD)	AP	Obstetrics and Gynecology
110	FOUMANE Pascal	P	Obstetrics and Gynecology
111	KASIA Jean Marie	P	Obstetrics and Gynecology
112	KEMFANG NGOWA Jean Dupont	P	Obstetrics and Gynecology
113	MBOUDOU Émile	P	Obstetrics and Gynecology
114	MBU ENOW Robinson	P	Obstetrics and Gynecology
115	NKWABONG Elie	P	Obstetrics and Gynecology
116	TEBEU Pierre Marie	P	Obstetrics and Gynecology
117	BELINGA Etienne	AP	Obstetrics and Gynecology
118	ESSIBEN Félix	AP	Obstetrics and Gynecology

119	FOUEDJIO Jeanne Hortence	AP	Obstetrics and Gynecology
120	NOA NDOUA Claude Cyrille	AP	Obstetrics and Gynecology
121	DOHBIT Julius SAMA	AP	Obstetrics and Gynecology
122	MVE KOH Valère Salomon	AP	Obstetrics and Gynecology
123	METOGO NTSAMA Junie Annick	SL	Obstetrics and Gynecology
124	MBOUA BATOUM Véronique Sophie	L	Obstetrics and Gynecology
125	MENDOUA Michèle Florence Epse NKODO	L	Obstetrics and Gynecology
126	NSAHLAI Christiane JIVIR FOMU	L	Obstetrics and Gynecology
127	NYADA Serge Robert	L	Obstetrics and Gynecology
128	TOMPEEN Isidore	L	Obstetrics and Gynecology
129	EBONG Cliford EBONTANE	L	Obstetrics and Gynecology
130	MPONO EMENGUELE Pascale Epse NDONGO	L	Obstetrics and Gynecology
131	NGONO AKAM Marga Vanina	L	Obstetrics and Gynecology
DEPARTMENT OF OPHTHALMOLOGY, ENT AND STOMATOLOGY			
132	DJOMOU François (HD)	P	ENT
133	EBANA MVOGO Côme	P	Ophthalmology
134	ÉPÉE Émilienne Epse ONGUENE	P	Ophthalmology
135	KAGMENI Gilles	P	Ophthalmology
136	NDJOLO Alexis	P	ENT
137	NJOCK Richard	P	ENT
138	OMGBWA EBALE André	P	Ophthalmology
139	BILLONG Yannick	AP	Ophthalmology
140	DOHVOMA Andin Viola	AP	Ophthalmology
141	EBANA MVOGO Stève Robert	AP	Ophthalmology

142	KOKI Godefroy	AP	Ophthalmology
143	MINDJA EKO David	AP	ENT/Maxillo-Facial Surgery
144	NGABA Olive	AP	ENT
145	ANDJOCK NKOUO Yves Christian	SL	ENT
146	MEVA'A BIOUELE Roger Christian	SL	ENT-MFS
147	MOSSUS Yannick	SL	ENT-MFS
148	MVILONGO TSIMI Epse BENGONO Caroline	SL	Ophthalmology
149	NGO NYEKI Adèle-Rose Epse MOUAHA-BELL	SL	ENT-MFS
150	NOMO Arlette Francine	SL	Ophthalmology
151	AKONO ZOUA Epse ETEME Marie Evodie	L	Ophthalmology
152	ASMAOU BOUBA Dalil	L	ENT
153	ATANGA Léonel Christophe	L	ENT-MFS
154	BOLA SIAFA Antoine	L	ENT
155	NANFACK NGOUNE Chantal	L	Ophthalmology
DEPARTMENT OF PAEDIATRICS			
156	ONGOTSOYI Angèle Epse PONDY (HD)	P	Paediatrics
157	KOKI NDOMBO Paul	P	Paediatrics
158	ABENA OBAMA Marie Thérèse	P	Paediatrics
159	CHIABI Andreas	P	Paediatrics
160	CHELO David	P	Paediatrics
161	MAH Evelyn	P	Paediatrics
162	NGUEFACK Séraphin	P	Paediatrics

163	NGUEFACK Epse DONGMO Félicitée	P	Paediatrics
164	NGO UM KINJEL Suzanne Epse SAP	AP	Paediatrics
165	KALLA Ginette Claude Epse MBOPI KEOU	AP	Paediatrics
166	MBASSI AWA Hubert Désiré	AP	Paediatrics
167	NOUBI Nelly Epse KAMGAING MOTING	AP	Paediatrics
168	EPEE Epse NGOUE Jeannette	SL	Paediatrics
169	KAGO TAGUE Daniel Armand	SL	Paediatrics
170	MEGUIEZE Claude-Audrey	SL	Paediatrics
171	MEKONE NKWELE Isabelle	SL	Paediatrics
172	TONY NENGOM Jocelyn	SL	Paediatrics

DEPARTMENT OF MICROBIOLOGY, PARASITOLOGY, HAEMATOLOGY AND INFECTIOUS  
DISEASES

173	MBOPI KEOU François-Xavier (HD)	P	Bacteriology/ Virology
174	ADIOGO Dieudonné	P	Microbiology/Virology
175	GONSU born KAMGA Hortense	P	Bacteriology
176	LUMA Henry	P	Bacteriology/ Virology
177	MBANYA Dora	P	Haematology
178	OKOMO ASSOUMOU Marie Claire	P	Bacteriology/ Virology
179	TAYOU TAGNY Claude	P	Microbiology/Haematology
180	CHETCHA CHEMEGNI Bernard	AP	Microbiology/Haematology
181	LYONGA Emilia ENJEMA	AP	Microbiology
182	TOUKAM Michel	AP	Microbiology

183	NGANDO Laure épouse MOUDOUTE	SL	Parasitology
184	BEYALA Frédérique	L	Infectious Diseases
185	BOUM II YAP	L	Microbiology
186	ESSOMBA René Ghislain	L	Immunology
187	MEDI SIKE Christiane Ingrid	L	Infectious Diseases
188	NGOGANG Marie Paule	L	Clinical Biology
189	NDOUMBA Annick Epse MINTYA	L	Haematology
190	VOUNDI VOUNDI Esther	L	Virology
191	ANGANDJI TIPANE Prisca Epse ELLA	L	Clinical Biology/Haematology
192	Georges MONDINDE IKOMEY	L	Immunology
193	MBOUYAP Pretty Rosereine	L	Virology
DEPARTMENT OF PUBLIC HEALTH			
194	KAMGNO Joseph (HD)	P	Public Health /Epidemiology
195	ESSI Marie José	P	Public Health/Medical Anthropology
196	TAKOUGANG Innocent	P	Public Health
197	BEDIANG Georges Wylfred	AP	Medical Information Technology/Public Health
198	BILLONG Serges Clotaire	SL	Public Health
199	NGUEFACK TSAGUE	SL	Public Health /Biostatistics
200	EYEBE EYEBE Serge Bertrand	L	Public Health/Epidemiology
201	KEMBE ASSAH Félix	L	Epidemiology
202	KWEDI JIPPE Anne Sylvie	L	Epidemiology
203	MOSSUS Tatiana born ETOUNOU AKONO	L	Health Promotion Expert
204	NJOUNEMI ZAKARIAOU	L	Public Health/Health Economics

205	ABBA-KABIR Mahamat	Haamit-	L	Pharmacist
206	AMANI ADIDJA		L	Public Health
207	ESSO ENDALLE Lovet Linda Augustine Julia		L	Public Health
208	MBA MAADJHOU Berjauline Camille		L	Public Health/Nutritional Epidemiology
DEPARTMENT OF MORPHOLOGICAL SCIENCES AND MORBID ANATOMY				
209	MENDIMI NKODO Joseph (HD)		AP	Morbid Anatomy/Pathology
210	SANDO Zacharie		P	Morbid Anatomy/Pathology
211	BISSOU MAHOP Josue		AP	Sports Medicine
212	KABEYENE OKONO Angèle Clarisse		AP	Histology/Embryology
213	AKABA Désiré		AP	Human Anatomy
214	NSEME ETOUCKEY Georges Eric		AP	Legal Medicine
215	NGONGANG Gilbert Frank Olivier		SL	Legal Medicine
216	MENDOUGA MENYE Coralie Reine Bertine Epse KOUOTOU		L	Morbid Anatomy
217	ESSAME Eric Fabrice		L	Morbid Anatomy
DEPARTMENT OF BIOCHEMISTRY				
218	NDONGO EMBOLA Epse TORIMIRO Judith (HD)		P	Molecular Biology
219	PIEME Constant Anatole		P	Biochemistry
220	AMA MOOR Vicky Joceline		P	Clinical Biology/Biochemistry
221	EUSTACE BONGHAN BERINYUY		L	Biochemistry
222	GUEWO FOKENG Magellan		L	Biochemistry

223	MBONO SAMBA ELOUMBA Esther Astrid	L	Biochemistry
DEPARTMENT OF PHYSIOLOGY			
224	ETOUNDI NGOA Laurent Serges (HD)	P	Physiology
225	ASSOMO NDEMBA Peguy Brice	AP	Physiology
226	AZABJI KENFACK Marcel	L	Physiology
227	DZUDIE TAMDJIA Anastase	L	Physiology
228	EBELL'A DALLE Ernest Remy Hervé	L	Human Physiology
DEPARTMENT OF PHARMACOLOGY AND TRADITIONAL MEDICINE			
229	NGONO MBALLA Rose ABONDO (HD)	AP	African Pharmacotherapeutics
230	NDIKUM Valentine	L	Pharmacology
231	ONDOUA NGUELE Marc Olivier	L	Pharmacology
DEPARTMENT OF ORAL SURGERY, MAXILLO-FACIAL SURGERY AND PERIODONTOLOGY			
232	BENGONDO MESSANGA Charles (HD)	P	Stomatology
233	EDOUMA BOHIMBO Jacques Gérard	SL	Stomatology and Surgery
234	LOWE NANTCHOUANG Jacqueline Michèle Epse ABISSEGUE	L	Pediatric Dentistry
235	MBEDE NGA MVONDO Rose	L	Dental Medicine
236	MENGONG Epse MONEBOULOU Hortense	L	Pediatric Dentistry
237	NDJOH Jules Julien	L	Dental Medicine
238	NOKAM TAGUEMNE M.E.	L	Dental Medicine

239	GAMGNE Catherine M	GUIADEM	L	Dental Surgery
240	KWEDI Karl Guy Grégoire		L	Oral Surgery
241	NIBEYE Yannick Carine Brice		L	Bacteriology
242	NKOLO TOLO Francis Daniel		L	Dental Surgery
DEPARTMENT OF PHARMACOGNOSY AND PHARMACEUTICAL CHEMISTRY				
243	NTSAMA ESSOMBA Claudine (HD)		P	Pharmacognosy/ Pharmaceutical Chemistry
244	NGAMENI Bathélémy		P	Phytochemistry/Organic Chemistry
245	NGOUPAYO Joseph		P	Phytochemistry/Pharmacognos y
246	GUEDJE Nicole Marie		AP	Ethnopharmacology/Plant Biology
247	BAYAGA Hervé Narcisse		L	Pharmacy
DEPARTMENT OF PHARMACOTOXICOLOGY AND PHARMACOKINETICS				
248	ZINGUE Stéphane (HD)		AP	Pharmacy
249	FOKUNANG Charles		P	Molecular Biology
250	TEMBE Estella Epse FOKUNANG		AP	Clinical Pharmacology
251	ANGO Yves Patrick		L	Natural Substance Chemistry
252	NENE AHIDJO Epse NJITUNG TEM		L	Neuropharmacology
DEPARTMENT OF GALENIAL PHARMACY AND PHARMACEUTICAL LEGISLATION				
253	NNANGA NGA Emmanuel (CD)		P	Galenial Pharmacy
254	MBOLE Jeanne Mauricette Epse MVONDO		L	Quality Control and Management of Health Products and Food
255	NYANGONO NDONGO Martin		L	Pharmacy



256	SOPPO Lobe Charlotte Vanessa	L	Quality control of Drugs
257	ABA'A Marthe Dereine	L	Drug Analysis
258	FOUMANÉ MANIEPI NGOUOPIHO Jacqueline Saurelle	L	Pharmacology
259	MINYEM NGOMBI Aude Périne Epse AFUH	L	Pharmaceutical Regulation

KEY:

HD= Head of Department


P= Professor

AP= Associate Professor

SL= Senior Lecturer

L= Lecturer

## THE PHYSICIAN'S OATH



Association in Geneva, Switzerland, September 1948  
and amended by the 22nd World Medical Assembly, Sydney,  
Australia (August 1968).

*I will practice my profession with conscience and dignity;*

*The health of my patients was my first consideration;*

*I will respect secrets confided in me, even after the patient has  
died;*

*I will maintain by all the means in my power the honor and noble  
traditions of the medical profession;*

*My colleagues was my brothers and sisters;*

*I will not permit considerations of religion, nationality, race,  
political party or social standing to intervene between my duty  
and my patient;*

*I will maintain the utmost respect for human life from the time of  
conception, even under threat I will not use my medical  
knowledge contrary to the laws of humanity; I make these  
promises solemnly, freely and upon my honor.*

## LIST OF TABLES

TABLE I; BRACHIAL ARCH DERIVATIVES .....	- 9 -
TABLE II ; APPROXIMATE DATES OF TOOTH DEVELOPMENT, ERUPTION AND EXFOLIATION FOR THE PRIMARY DENTITION.....	- 23 -
TABLE III ; APPROXIMATE DATES OF TOOTH DEVELOPMENT AND ERUPTION FOR THE PERMANENT DENTITION.....	- 24 -
TABLE IV : KAP STUDIES ASSESSMENT .....	- 37 -
TABLE V; SOCIODEMOGRAPHIC REPRESENTATION OF PARTICIPANTS .....	- 41 -
TABLE VI; DISTRIBUTION ACCORDING TO KNOWLEDGE OF RISK FACTORS OF MALOCCLUSION-	43 -
TABLE VII; IDENTIFICATION OF PARAFUNCTIONAL ORAL HABITS AND MALOCCLUSIONS AND TRAINING.....	- 44 -
TABLE VIII; DISTRIBUTION ACCORDING TO ATTITUDE .....	- 46 -

## LIST OF FIGURES

FIG 1 : ANATOMY OF THE ORAL CAVITY.....	- 7 -
FIG 2; PRENATAL DEVELOPMENT OF THE FACE .....	- 12 -
FIG 3; DOWNWARD AND FORWARD GROWTH OF THE NASOMAXILLARY COMPLEX AS A RESULT OF SUTURAL GROWTH .....	- 14 -
FIG 4; SECONDARY DISPLACEMENT (A) AND REMODELING (B) .....	- 15 -
FIG 5; MANDIBULAR GROWTH PATTERN .....	- 17 -
FIGURE 6; SCHEMATIC REPRESENTATION OF OUR STUDY PARTICIPANTS .....	- 40 -
FIG 7: KNOWLEDGE OF PARTICIPANTS .....	- 45 -
FIG 8: PRACTICE OF PARTICIPANTS .....	- 49 -

## **LIST OF ABBREVIATIONS**

W.H.O	World Health Organisation
O.M.S	Organisation Mondiale de la Sante
AAP	American Academy of Pediatrics

## ABSTRACT

**Introduction.** The World Health Organization (WHO) considers malocclusion as one of the most important oral health problems, after caries and periodontal disease. Its prevalence is highly variable and is estimated to be between 39% and 93% in children and adolescents. Parafunctional habits are abnormal hyperactive functions conducted by the masticatory structures, i.e. tongue, teeth, oral muscles, and so on. They include thumb sucking, tongue thrusting, bruxism, prolonged use of feeding bottle or pacifier and so on. Parafunctional habits happen to be a major risk factor for malocclusion. Adequate knowledge of pediatricians and their collaboration with oral healthcare providers would go a long way into sensitizing patients on parafunctional habits and their effect on the developing dentition, as well as intercepting malocclusions at an early stage where they can be corrected through referrals to orthodontists or pediatric dentists

**Objective.** Our objective was to evaluate the knowledge, attitude and practice of paediatricians in Cameroon towards parafunctional oral habits and dental malocclusions in children.

**Materials and Methods.** A descriptive, cross-sectional survey was sent to a sample of pediatricians in Cameroon. The questionnaire consisted of 35 questions in four domains; socioprofessional information, knowledge, attitude and practice.

**Results.** A total of 86 pediatricians participated in the survey. In terms of knowledge, 43% of respondents reported having knowledge that children with parafunctional oral habits are more likely to develop malocclusion. Regarding their attitude, 97.7% of respondents had a good attitude towards the subject. In terms of examination of the oral cavity, a low frequency of examination amongst pediatricians for malocclusion (10.5%) and oral functional habits (76.7%) was detected. 47.7% recommend oral health examinations only if there is a problem and 53.5% refer patients to a pediatric dentists when they notice any abnormality.

**Conclusion.** Although a vast majority of the paediatricians in our survey recognised their place in identifying and preventing parafunctions and malocclusions, very few have adequate knowledge and skills to perform evaluations and referrals.

*Key words: Pediatricians, knowledge, attitude, practice, parafunctions, malocclusion*

## RÉSUMÉ

**Introduction.** L'Organisation Mondiale de la Santé (OMS) considère la malocclusion comme l'un des problèmes de santé bucco-dentaire les plus importants, après les caries et les maladies parodontales. Sa prévalence est très variable et est estimée entre 39 % et 93 % chez les enfants et adolescents. Les habitudes parafonctionnelles sont des fonctions hyperactives anormales exercées par les structures masticatoires, c'est-à-dire la langue, les dents, les muscles buccaux, etc. Ils comprennent la succion du pouce et de la langue, le bruxisme, l'utilisation prolongée d'un biberon ou d'une tétine, rongement des ongles etc. Les habitudes parafonctionnelles s'avèrent être un facteur de risque majeur de malocclusion. Une connaissance adéquate des pédiatres et leur collaboration avec les prestataires de soins bucco-dentaires contribueraient grandement à sensibiliser les patients aux habitudes parafonctionnelles et à leurs effets sur le développement de la dentition, ainsi qu'à intercepter les malocclusions à un stade précoce où elles peuvent être corrigées en les référant à des orthodontistes ou des pédodontistes.

**Objectif.** Notre objectif était d'évaluer les connaissances, l'attitude et la pratique des pédiatres au Cameroun face aux habitudes bucco-dentaires parafonctionnelles et aux malocclusions dentaires chez l'enfant.

**Matériels et méthodes.** Une enquête descriptive et transversale a été envoyée à un échantillon de pédiatres au Cameroun. Le questionnaire comprenait 35 questions dans quatre domaines ; informations socioprofessionnelles, connaissances, attitudes et pratiques.

**Résultats.** Au total, 86 pédiatres ont participé à l'enquête. En termes de connaissances, 43 % des personnes interrogées ont déclaré savoir que les enfants ayant des habitudes buccales parafonctionnelles sont plus susceptibles de développer une malocclusion. Concernant leur attitude, 97,7% des répondants avaient une bonne attitude envers le sujet. En termes d'examen de la cavité buccale, une faible fréquence d'examens chez les pédiatres pour les malocclusions (10,5%) et les habitudes fonctionnelles buccales a été détectée. 47,7 % recommandent des examens de santé buccodentaire uniquement en cas de problème et 53,5% orientent les patients vers un dentiste pédiatrique lorsqu'ils constatent une anomalie.

**Conclusion.** Bien qu'une grande majorité des pédiatres de notre enquête reconnaissent leur place dans l'identification et la prévention des parafonctions et des malocclusions, très peu possèdent les connaissances et les compétences adéquates pour effectuer des évaluations et des références.

*Mots clés : Pédiatres, connaissances, attitude, pratiques, parafonctions, malocclusion*

## **CHAPTER I: INTRODUCTION**



## **I.1 BACKGROUND**

A habit is a practice acquired by the frequent repetition of the same act, which occurs consciously at first, then unconsciously [1]. Nasal breathing, chewing and swallowing are considered physiological and functional habits. However, digital sucking, pacifier sucking, tongue-thrusting, nail-biting, lip-biting and mouth breathing, amongst others, are considered non-physiological, and therefore, deleterious or parafunctional habits [1]. Studies have shown that while parafunctional habits can interfere with the position of the teeth and normal pattern of skeletal growth causing malocclusion, the negative influence of parafunctional habits on occlusion originates in childhood. Bottle feeding and non-nutritive sucking habits have been associated with malocclusions starting from the primary dentition. Several authors have pointed out that non-nutritive sucking habits are associated with an atypical swallowing pattern, and with tongue thrusting, which may be related to the development of malocclusions such as posterior crossbites [2]. A study carried out in Cameroon by Kalla *et al.* in 2022 showed an increasing prevalence of parafunctional oral habits in children [3].

The World Health Organization (WHO) considers malocclusion one of the most important oral health problems, after caries and periodontal disease. Its prevalence is highly variable and is estimated to be between 39% and 93% in children and adolescents [4]. In the same light, malocclusions have been noted to be of huge impact on children increasing their risk of dental caries, periodontal disease, increased possible temporomandibular disease, and increased risk of trauma and equally affecting their appearance, feeding habits and even their self- esteem.

Pediatricians usually are the first healthcare providers children visit and are most likely to examine patients earlier than pediatric dentists. However, very few patients present before pediatric dentists and orthodontists with referrals from pediatricians for parafunctional habits and malocclusions [5]. In a study published by Kumar *et al.* in India showed that most pediatricians were unaware that oral habits could be responsible for malocclusion [1]. The early identification and treatment of orthodontic abnormalities in the deciduous and early mixed dentition can prevent the development of significant anomalies in late mixed and permanent dentition, as well as reduce or even eliminate the need for subsequent orthodontic treatment.

## **I.2 JUSTIFICATION**

Adequate knowledge of pediatricians on parafunctional habits and their collaboration with oral healthcare providers would go a long way in sensitizing patients and their care-givers on the long term effects of parafunctional habits on the developing dentition. Furthermore, identifying malocclusions at an early stage is essential in intercepting and correcting them through referrals to orthodontists or pediatric dentists. This study will therefore contribute to the available body of literature on this subject. It will equally raise an awareness on the role of pediatricians' in identifying and intercepting parafunctional oral habits and malocclusions. Lastly, this study will contribute to developing policy around early and widespread detection of malocclusion.

## **I.3 RESEARCH QUESTION**

Do pediatricians have adequate knowledge, attitudes and practices towards parafunctional oral habits and dental malocclusions in children?

## **I.4 OBJECTIVES**

### **I.4.i GENERAL OBJECTIVE**

To assess pediatricians' knowledge, attitudes and practices towards parafunctional habits and malocclusions in children.

### **I.4.ii SPECIFIC OBJECTIVES**

1-To assess pediatricians' knowledge of parafunctional habits and malocclusions as well as their understanding of the relationship between the two.

2-To describe pediatricians' attitude towards parafunctional habits and malocclusions.

3-To identify pediatricians' practices towards parafunctional habits and malocclusions.

## **CHAPTER II: LITERATURE REVIEW**

Our literature review will cover the anatomy of the oral cavity, craniofacial growth and development, tooth development, parafunctional habits and malocclusions. We will conclude with a review of existing literature on the subject.

## **II.1 THE ORAL CAVITY**

The mouth or oral cavity serves as the first portion of the digestive system. The oral cavity is situated anteriorly on the face, under the nasal cavities. It opens to the face through the oral fissure, while posteriorly the oral cavity communicates with the oropharynx through a narrow passage called the oropharyngeal isthmus (also termed the isthmus of the fauces). The cavity is separated into an anterior oral vestibule and the oral cavity proper located posteriorly to the teeth. Although a small compartment, the oral cavity is a unique and complex structure with several different nerves and blood vessels inside it. This intricate network is necessary for its unique and diverse role in human life [6].

### **Vestibule;**

The oral vestibule is bounded externally by the lips and the cheek mucosa and internally by the alveolar processes and the teeth. When the teeth are in occlusion, the vestibule communicates with the oral cavity proper via the intermaxillary commissure behind the last molar teeth [7]

### **Lips;**

The lips are the soft, visible part of the humans' mouth and comprise the upper and the lower lips. Together these serve as the gatekeepers of the oral cavity. Externally, the lips' skin forms a junction with the surrounding facial skin, which is marked by the vermilion border (also called the cupid's bow in the case of the upper lip). Internally, it forms a junction with the mucous membrane inside the oral cavity. The lips help in food intake and in articulation of speech. Moreover, being tactile sensory organs and due to their rich sensory innervation, they also serve as erogenous zones in acts of intimacy. Additionally, lips also contribute to the facial expressions [8].

### Oral Cavity proper;

It is filled mostly with the tongue and is bound anterolaterally by the posterior surface of the teeth and gums, superiorly by the soft and hard palate, inferiorly by the floor of the mouth and communicates posteriorly with the oropharynx.

### Tongue [9];

The tongue is predominantly muscle. There are 8 in total; 4 intrinsic muscles (They are responsible for many of the tongue's functions; such as talking, mastication and any other action that requires the tongue to move. These muscles control movements such as twirling, curling, flattening and broadening of the tongue) and 4 extrinsic muscles (assist the tongue and support it in more complex actions such as protrusion and retraction). Besides the muscles, the other important feature of the tongue is its mucosa. The dorsal tongue mucosa is covered with lingual papillae which function as the sensory receptors for taste. All of the tongue muscles are innervated by the hypoglossal nerve (CN XII), except for the palatoglossus muscle which is supplied by the vagus nerve (CN X).

### Palate (8);

The palate is a bony/muscular partition that forms the roof of the oral cavity and the floor of the nasal cavities. It consists of two main parts; the hard palate and soft palate. The hard palate is the anterior bony portion, while the soft palate is the posterior muscular part. The palate completely separates the oral cavity and nasal cavities. This division is essential for preventing pressure changes within these cavities, thus enabling important processes such as suckling and breathing. Additionally, due to its position and structure, the palate plays an important role in articulation (formation of speech), digestion and swallowing.




Fig 1 : Anatomy of the oral cavity

## **II.2 CRANIOFACIAL GROWTH AND DEVELOPMENT**

Craniofacial development is a highly coordinated process under a tight genetic control and environmental influence. The development of the craniofacial skeleton occurs as a result of a sequence of normal developmental events: brain growth and development, optic pathway development, speech and swallowing development, airway and pharyngeal development, muscle development, and tooth development and eruption [10].

### **ii.2.i Brachial Arches**

Majority of the craniofacial skeleton is of ectomesenchymal origin whereas the other skeleton of the entire body is derived from mesoderm. In specific areas of the developing embryo, the migrating and rapidly proliferating ectomesenchymal cells develop elevations between ectoderm and endoderm. In the somite period, 4th week IUL, such elevations are seen in the ventral foregut resulting in the formation of six pharyngeal arches or brachial arches bilaterally, the fifth arch perishes; finally, only five arches remain. The arches are separated by 4 branchial grooves on the ectodermal aspect and 5 pharyngeal pouches on the endodermal aspect. Mesenchymal cells migrate between the ectoderm and endoderm and around the mesodermal condensation. Branchial arches decrease in size craniocaudally. Each branchial arch gives rise to some specific derivatives and every arch has the following structures: A cartilage rod, centrally located, forms the skeleton of the arch; a branchiomere that gives rise to the muscles of the arch; a vascular component, aortic arch artery, supplying the area; a nervous element, giving rise to sensory and specific visceral motor fibers for the cranial nerve supplying specific arch[11].

Table I shows the different brachial arches as well as their derivatives.

Table I; Brachial arch derivatives

ARCH	NERVE	SKELETAL DERIVATIVE	POUCH DERIVATIVE	MUSCLE DERIVATIVE	ARCH DERIVATIVES	ARTERY
First (Mandibular)	Trigeminal	Greater wing of sphenoid, malleus, incus, maxilla, zygoma, mandible, anterior ligament of malleus, sphenomandibular ligament	Auditory tube, middle cavity	Muscle s of mastication, anterior belly of digastric, mylohyoid, tensor tympani, tensor veli palatine		Portion of external carotid arteries
Second (Hyoid)	Facial	Stapes, styloid process, lesser cornu and upper hyoid, stylohyoid ligament	Palatine tonsillar fossa	Muscle s of facial expression, posterior belly of digastric, stylohyoid, stapedius		Tympanic branch of internal carotid artery
Third	Glossopharyngeal	Greater cornu and lower part of hyoid	Inferior parathyroids, thymus	Stylopharyngeus		Common carotid and proximal internal carotid arteries
Fourth	Superior laryngeal of vagus	Laryngeal cartilages	Superior parathyroid, lateral thyroid	Pharyngeal constrictors, cricothyroid, levator veli palatine, palatoglossus		Left arch of aorta, brachiocephalic and right subclavian arteries, distal pulmonary arteries
Fifth/Sixth	Recurrent laryngeal of vagus	Laryngeal cartilages	Ultimobranchial body	Laryngeal muscles except cricothyroid, striated muscle of esophagus		Proximal pulmonary arteries, ductus arteriosus



## II.2.ii Formation of the Face

The face derives from five prominences that surround a central depression, the *stomodeum* which constitutes the future mouth. These prominences are the single median frontonasal prominence and the paired maxillary and mandibular prominences. Facial development occurs mainly between the fourth and eighth weeks of gestation as follows[12] :

- Fourth week of development

Primordia of the face appear at the cephalic end of the embryo.

Two nasal placodes cap the bulbous frontal prominence.

The optic discs appear posterolateral to the frontal prominence.

Three paired branchial arches have formed.

The first arches split into maxillary and mandibular prominences. The hyoid arches are the second pair.

Between the first arches and frontal prominence, the buccopharyngeal membrane becomes fenestrated.

- Fifth week of development

Nasal pits develop in the nasal placodes, and the rims of the placodes differentiate into medial and lateral nasal prominences.

The lens vesicles invaginate and close within the optic discs.

The mesenchyme of the mandibular arch fills in across the midline.

The caudal end of the medial nasal prominences begins to fuse with the maxillary prominences.

- At the beginning of the sixth week of development

The nasals have shifted to a more ventral, central position.

Growing and shifting subectodermal mesenchyme smooths out the furrows between prominences and arches, and the second arch becomes more massive.

Six auricular hillocks, which will become the pinna of the ears, form on the mandibular and hyoid arches.

- By the end of the sixth week of development

Medial and lateral nasal prominences fuse.

Maxillary prominences begin the formation of the upper jaw.

The midline approximation of the medial nasal prominences forms the nasal septum.

The mandible appears as a band of dense fibrous tissue known as Meckel's cartilage; this cartilage provides a framework around which the bone will form.

- At the beginning of the seventh week of development

The tip of the nose is elevated between the medial nasal prominences and is visible in profile.

Eyelids become prominent.

The pinna of the ear takes shape.

- End of the seventh week of development

The pattern of facial features has taken on a human appearance. However, facial proportions develop during the fetal period.

Bone formation commences at the mental foramen area and begins to spread backwards, forwards and upwards outlining the future body of the mandible.

The fusion of the medial nasal prominences, which forms the central axis of the nose and the philtrum of the lip, is complete.

From the beginning of the eighth week of development till birth, the final facial development occurs slowly and consists mainly of changes in the proportion and relative positions of the facial components. During the early fetal period, the nose is flat and the mandible is underdeveloped. They obtain their characteristic form while facial development is being completed. As the brain enlarges, it creates a prominent forehead, the eyes move medially, and the

external ears rise. The prenatal face is small because of the rudimentary upper and lower jaws, the unerupted primary teeth, and the small size of the nasal cavities and maxillary sinuses.




Fig 2; Prenatal development of the face

## **II.2.iii POST NATAL GROWTH OF THE MAXILLA AND MANDIBLE**

### **a- Growth of the Maxilla**

Since, the maxillary complex is attached to the cranial base, there is a strong influence of the latter on the former. Hence, while discussing the growth of nasomaxillary complex, we have to look into three aspects [13];

#### **1) The displacement in the position of the maxillary complex**

-Primary displacement occurs in a forward direction. This occurs by growth of the maxillary tuberosity in a posterior direction. This results in the whole maxilla being carried anteriorly.

-Secondary displacement- Occurs in a downward & forward direction as the cranial base grows.

#### **2) Growth at sutures:**

- Sutural connective tissue ; Proliferation, ossification, surface apposition, resorption and translation are the mechanisms for maxillary growth.

- The maxilla is related to cranium at least partially by the: Frontomaxillary suture, frontonasal suture, zygomaticomaxillary suture, zygomaticotemporal suture and pterygopalatine suture. These sutures are all oblique & more or less parallel with each other. The growth in these areas would serve to move the maxilla downward & forward.

#### **3) Surface Remodeling:**

Remodeling occurs by bone deposition & resorption to bring about increase in size, change in shape and change in functional relationship.

Bone resorption is seen on the floor of the nasal cavity. To compensate, there is bone deposition on the palatal side. Thus, a net downward shift leading to increase in maxillary height.

Bone deposition occurs along the posterior margin of the maxillary tuberosity. This causes lengthening of the dental arch and enlargement of the anteroposterior dimension of the entire maxillary body.

#### Growth timings of the maxilla

1 – 5 years of age; 45% of total growth completed

5 – 10 years of age; 65% of total growth

10 – 20 years of age; Remaining 35% completed.




Fig 3; Downward and forward growth of the nasomaxillary complex as a result of sutural growth




Fig 4; Secondary displacement (a) and remodeling (b)

### **b- Growth of mandible**

At birth the two rami of the mandible are short, condylar development is minimum and there is no articular eminence in glenoid fossa. A thin layer of fibrocartilage & connective tissue exists at the midline of symphysis to separate right & left mandibular bodies [14].

At fourth month of age and end of first year symphysial cartilage is replaced by bone.

During first year of life appositional growth is active at alveolar border, at distal & superior surfaces of the ramus, at the condyle, along the lower border of mandible and on its lateral surface.

After the first year of life these changes occur ;

- Mandibular growth becomes more selective, condyle shows considerable activities, mandible moves and grows downward & forward.
- Appositional growth occurs on posterior border of the ramus and on the alveolar process.
- Resorption occurs along the anterior border of ramus lengthening the alveolar border & maintaining the anterior- posterior dimension of ramus.
- Gonial angle changes after little muscle activity.
- Transverse dimension is mainly due to growth at posterior border in an expanding V pattern.
- The two rami also diverge outward from below to above so that additive growth at coronoid notch coronoid process & condyle also increases the superior inter-ramus dimension.
- Alveolar process of mandible grows upward & outward on an expanding arc. This permits dental arc to accommodate the larger permanent teeth.
- On lingual surface, behind the chin heavy periosteal growth occurs, with the dense lamellar bone merging and overlapping on the labial side of the chin.

- In males, the apposition of the bone at symphysis seems to be about the last change in shape during the growing period. This change is much less apparent in the females.






Fig 5; Mandibular growth pattern

## II.3 TOOTH DEVELOPMENT

### II.3.i Development of the tooth germ

The tooth germ develops in three stages: bud, cap and bell [15].

**Bud-** At 8 weeks of intrauterine life, clumps of mesenchymal cells induce the dental lamina to form swellings known as enamel organs. Each enamel organ will be responsible for the development of each tooth.

**Cap-** As the enamel organs grow and increase in size, the inner aspect becomes concave resembling skull caps. By the late cap stage, at 12 weeks of intrauterine life, cells on the inner aspect of the enamel organ change from cuboidal to columnar forming the inner enamel epithelium. The outer layer of cells remains cuboidal and is known as the outer enamel epithelium. Beneath the inner enamel epithelium the condensation of mesenchymal cells is termed the dental papilla; this will eventually become the pulp. A fibrous capsule surrounds each enamel organ and this is termed the dental follicle; this will eventually become the periodontal ligament.

### **Bell**

By 14 weeks of intrauterine life the enamel organ consists of the following:

Inner enamel epithelium:

- Cells lining the inner surface of the enamel organ which are columnar in shape.
- The inner enamel epithelium defines the shape of the crown.
- They will eventually differentiate into enamel forming cells (ameloblasts).

Stratum intermedium:

- Lies over the inner enamel epithelium.
- Consists of two to three layers of cells.



- Transports nutrients to and from the ameloblasts.

Stellate reticulum:

- Lies between the stratum intermedium and the outer enamel epithelium.
- It consists of star-shaped cells that protect the underlying dental tissues.
- It also maintains the shape of the tooth.

Outer enamel epithelium:

- Cells lining the outer surface of the enamel organ. They are cuboidal in shape.
- They maintain the shape of the enamel organ.
- The outer enamel epithelium meets with the internal enamel epithelium at the cervical loop.

Eventually the inner and outer enamel epithelium grows downwards at the cervical loop forming Hertwig's root sheath, which maps out the shape of the root.

At the late bell stage the dental lamina disintegrates and is ready for the formation of dental hard tissue. Dentine formation always precedes enamel formation.

### **II.3.ii Dentine formation (Dentinogenesis)**

- Late bell stage-the inner enamel epithelium cells have mapped out the shape of the crown.
- The inner enamel epithelium cells induce cells at the periphery of the dental papilla to form columnar odontoblast cells (dentine-forming cells).
- Odontoblast cells begin to secrete an unmineralised dentine matrix.
- As more dentine matrix is deposited, the odontoblast cells retreat in the direction of the pulp leaving an elongated process known as the odontoblast process.

-The dentine matrix formed prior to mineralisation is termed predentine. A narrow layer of predentine is always present on the surface of the pulp.

-Mineralisation of dentine begins when the predentine is approximately 5  $\mu\text{m}$  thick.

-Spherical zones of hydroxyapatite called calcospherites are formed within the dentine matrix.

-Mineralisation of the dentine matrix starts at random points and eventually these calcospherites fuse together to form mineralised dentine.

-Dentinal tubules form around each odontoblast process.

-The odontoblasts retreat in S-shaped curves towards the dental papilla.

-The first layer of mineralised dentine is called mantle dentine and the remaining bulk of the mineralised dentine is known as circumpulpal dentine.

### **II.3.iii Enamel formation (Amelogenesis)**

-Immediately after the first layer of dentine is formed, the inner enamel epithelium ameloblast cells (enamel forming cells).

-The ameloblast cell is columnar in shape with its base attached to cells of the stratum intermedium.

-At the secretory end of ameloblast cells is a pyramidal extension called the *Tomes' process*.

-The enamel matrix is secreted through the Tomes' process at the amelodentinal junction.

-Calcium and phosphate ions are secreted into the enamel matrix and mineralisation of enamel occurs immediately, forming hydroxyapatite crystallites.

-As more enamel matrix is secreted and mineralised, the ameloblast cells move away from the amelodentinal junction forming a pattern of crystallites, which are contained within enamel prisms.

-Enamel prisms are also known as enamel rods they run from the amelodentinal junction to the enamel surface.

-During maturation from pre-enamel to mature enamel, the enamel crystallites increase in size and the organic content is reduced.

-On completion of enamel formation the ameloblast cell loses the Tomes' process, flattens and becomes the reduced enamel epithelium.

-The reduced enamel epithelium protects the enamel during eruption and will eventually become the junctional epithelium.

#### **II.3.iv Formation of the root**

-Occurs when the crown has completed.

-The internal and external enamel epithelium grows downwards at the cervical loop to form a double layered epithelial wall – Hertwig's root.

-The Hertwig's root sheath grows apically mapping out the shape of the root enclosing the dental papilla.

-The dental follicle lies external to the Hertwig's root sheath.

#### **II.3.v Cementum formation (Cementogenesis)**

-The Hertwig's root sheath induces the formation of odontoblast cells.

-When root dentine has formed, Hertwig's root sheath fragments allowing adjacent cells from the dental follicle to come into contact with the root dentine.

-These cells differentiate into cementoblasts (cementum-forming cells).

-Cementoblasts are cuboidal in shape and form a single layer on the surface of the root dentine.

-The cementoblasts secrete cementum matrix and crystallites of hydroxyapatite are deposited in this matrix and mineralisation occurs.

-During formation, a thin layer of unmineralised cementum is always present on the surface; this is known as *cementoid*.

#### **II.3.vi Formation of the periodontal ligament**

- Cells within the dental follicle give rise to fibroblasts that secrete collagen.
- Once cementum formation has begun, collagen fibres within the dental follicle orientate themselves into bundles, which are perpendicular to the root surface.
- These fibres will form the principal fibres of the periodontal ligament.
- The ends of these fibres become embedded in the developing cementum and alveolar bone and are known as Sharpey's fibres.

## **II.4 TOOTH ERUPTION**

Tooth eruption is the bodily movement of a tooth from its development position into its functional position in the oral cavity. It can be broken down into two phases; the pre-functional eruptive phase and the functional eruptive phase[16].

### **Prefunctional phase**

During the prefunctional phase, crown formation has completed.

As root formation begins the developing tooth begins to erupt.

The overlying alveolar bone is resorbed by osteoclasts and gradually the tooth moves in an axial direction towards the oral cavity.

The enamel surface of the tooth is covered by the reduced enamel epithelium which fuses with the oral epithelium.

The pressure from the tip of the tooth breaks down the oral epithelium allowing the tooth to emerge into the oral cavity without any rupturing of blood vessels.

Once the tooth has emerged, the reduced enamel epithelium is known as the epithelial attachment.

Tooth eruption continues until the tooth contacts (occludes with) the opposing tooth in the opposite jaw.

### **Functional eruptive phase**

The functional eruptive phase continues throughout life due to functional changes.

The alveolar bone continuously remodels in response to tooth movement and enamel wear allowing teeth to maintain contact with each other and with opposing teeth.

### **Mechanisms of tooth eruption**

The eruptive force of tooth eruption is unclear; however, several theories have been put forward although there is little evidence to support them. These are:

Root growth generates a force beneath the tooth, elevating the tooth towards the oral cavity.

Remodelling and deposition of the bone beneath the developing tooth pushes the tooth upwards.

Traction of the periodontal fibres exerts an upward pull on the tooth.

Cellular proliferation at the base of the pulp creates pressure that pushes the tooth from the dental follicle.

An increase in tissue fluid or blood pressure generates an eruptive force on the tooth.

Table II shows the calcification, eruption and exfoliation dates of primary teeth while Table III shows the calcification and eruption dates of permanent teeth.

Table II ; Approximate dates of tooth development, eruption and exfoliation for the primary dentition.

	Approximate calcification date	Approximate eruption date	Approximate exfoliation date
Upper and lower primary incisors(A's)	3-4 months in utero	6-8 months	6-7 years
Upper and lower primary lateral incisors(B's)	4.5 months in utero	7-9 months	7-8 years
Upper and lower primary first molars (D's)	5 months in utero	16-18 months	9-12 years
Upper and lower primary canines( C's)	5 months in utero	16-18 months	9-12 years
Upper and lower primary second molars (E's)	6 months in utero	20-24 months	10-12 years

Table III ; Approximate dates of tooth development and eruption for the permanent dentition.

Tooth (in order of eruption)	Approximate initial calcification date	Approximate eruption date
Upper and lower permanent first molars (6's)	At birth	6–7 years
Upper and lower permanent central incisors (1's)	3–4 months	6–7 years
Upper and lower permanent lateral incisors (2's)	Upper 10–12 months	7–8 years
Lower	3–4 months	
Lower permanent canines (3's)	4–5 months	9–10 years
Upper and lower permanent first premolars (4's)	1–2 years	10–11 years
Upper and lower permanent second premolars (5's)	2–3 years	11–12 years
Upper permanent canines (3's)	4–5 months	11–12 years
Upper and lower permanent second molars (7's)	2–3 years	12–13 years
Upper and lower permanent third molars (8's)	7–10 years	18+ years

## **II.5 HUMAN DENTITION AND OCCLUSION**

Humans have two sets of dentition in their lifetime- temporary and permanent dentition. The temporary (primary) dentition comprises the first set of teeth to erupt in children which are 20 in number. The permanent dentition is comprised of 32 teeth. All these set of teeth are evenly distributed between the maxillary and mandibular arches. These arches are equally divided into 4 quadrants [17]. There are four main types of teeth – incisors, canines, premolars, and molars. The premolars are only present in the permanent dentition. Each tooth is made up of two parts- the crown of the tooth is the visible part of the tooth in the mouth, while the root is hidden beneath the gingiva and alveolar bone [18]. The gingiva covers the border of the alveolar process that is adjacent to the teeth. The cemento-enamel junction is the anatomical boundary between the enamel-covered crown and cementum covered root. Dentin makes up the core of the entire tooth that surrounds the pulp, which contains the neurovascular structures. The apical foramen at the root apex is where the neurovascular structures enter the tooth and travel up the root canal to the expanded pulp chamber of the crown. The roots of the tooth vary depending on the type of tooth. The molars typically have three roots: a lingual root on the lingual aspect and a mesiobuccal root and distobuccal root on the buccal aspect. The crown of the tooth has five surfaces. The surface facing the lip or cheek is called the facial surface for incisors and canines and buccal surfaces for premolars and molars. The surface facing the inside of the mouth is referred to as the palatal surface in the maxilla and the lingual surface in the mandible. The surfaces referring to the boundaries of adjacent teeth are called mesial and distal. Mesial refers to the surface closer to the midline of the face, and distal refers to the surface away from the midline of the face. The biting surface is called the occlusal surface [19].

Occlusion is the relationship between the maxillary and mandibular teeth at rest (static occlusal relationship) and in function during various jaw movements (dynamic occlusal relationship) [20].

-Centric occlusion (CO) is the occlusion the patient makes when they fit their teeth together in maximum intercuspation. It is the contact between the greatest number of opposing teeth. This



is the occlusion the patient nearly always makes when they are asked to bite together as it is the bite they are habituated to.

-Centric relation (CR) is the mandible's relationship to the maxilla when the condyles are in their most anterior superior position in the glenoid fossae.

#### II.5.i Classification of static occlusion

Key classifications are;

##### *British standards institute classification of incisor relationship*

This is the relationship between the maxillary and mandibular incisors when in centric occlusion [21].

-Class I-The lower incisal edges occlude with or lie immediately below the cingulum plateau of the upper incisors.

-Class II-The lower incisal edges occlude posterior to the cingulum plateau of the upper incisors. It is further divided into two divisions:

Division I: the upper central incisors are proclined, usually resulting in an increased overjet

Division II: the upper central incisors are retroclined, usually resulting in a decreased overjet

-Class III-The lower incisal edges occlude anterior to the cingulum plateau of the upper incisors

##### *Canine Classification [22]*

-Class I: mesial incline of the upper canine overlaps the distal slope of the lower canine (The maxillary canine occludes between the mandibular canine and 1st premolar).

-Class II: Distal slope of the maxillary canine occludes or contact the mesial slope of the lower canine.

-Class III: The mandibular canine is displaced anterior to the maxillary canine with no overlapping.

*Molar Classification (Angle's classification) [23]*

-Class 1 (Normocclusion)-The mesiobuccal cusp of the maxillary first molar occluding with the buccal groove of the mandibular first molar i.e. the maxillary first molar is slightly posteriorly positioned relative to the mandibular first molar.

-Class 2 (Distocclusion)- The mesiobuccal cusp of the maxillary first molar occluding anterior to the buccal groove of the mandibular first molar i.e. the maxillary first molar is in line with or anteriorly positioned relative to the mandibular first molar. It is further divided into 2 divisions;

Class 2 division 1 – Class 2 molars with normally inclined or proclined maxillary central incisors.

Class 2 division 2 – Class 2 molars with retroclined maxillary central incisors.

- Class 3(Mesioclusion)-The mesiobuccal cusp of the maxillary first molar occluding posterior to the buccal groove of the mandibular first molar i.e. the maxillary first molar is severely posteriorly positioned relative to the mandibular first molar.

## **II.6 PARAFUNCTIONAL HABITS**

Habits stem from the repetition of the same action over time. Repetitive behaviors are common in infantile period and most of them are started and finished spontaneously [24]. Generally, oral habits can be divided into 2 main groups [25];

-Physiological or Functional such as chewing and swallowing.

-Parafunctional (deleterious) habits, such as finger and thumb sucking, pacifier and bottle sucking etc

Persistence of parafunctional oral habits during and beyond preschool age are one of the important etiological factors in developing malocclusion and other ill effects on orofacial

structures [26]. The effect of parafunctional habits on the dentition is intensity, frequency, and duration dependent.

Deleterious oral habits can act in such a way to disrupt the muscular balance mentioned above, causing significant interferences in the normal development of the maxillary bones, correct positioning of teeth, and respiratory, deglutition and phonation processes [27].

### **II.6.i Types of Parafunctional Habits**

- **Thumb-sucking;**

Thumb sucking is one of the most common habits of children. The habit starts early in life. About 90% of newborns show some form of hand sucking by 2 hours after birth. Thumb sucking is normal in infants and young children. It should cause no permanent problems if it stops by age 5 or before the eruption of permanent incisors [17].

Sucking a thumb gives a sense of safety or joy to the infants; and helps them better understand the world around themselves. On the other hand, thumbs placed inside the mouth exert force on the tissues thus creating physical changes as dental and maxillofacial abnormalities [28].

There are 2 types [27]:

- **Active:** In this type, there is a heavy force by the muscles during the sucking and if this habit continues for a long period, the position of permanent teeth and the shape of mandible will be affected.
- **Passive:** In this type, the child puts his/her finger in mouth, but because there is no force on teeth and mandible, so this habit is not associated with skeletal changes.

Some effects of thumb sucking include anterior open bite, increased overjet, lingual inclination of lower incisor and labial inclination of upper incisors, posterior cross bite, deep palate, speech defect, finger defects (Eczema of the finger due to alternate dryness and moisture that occurs and even angulations of the finger).

- **Nail biting or onychophagia**

Nail biting is a common and untreated medical problem among children. Complications caused by nail biting include malocclusion of the anterior teeth, root resorption intestinal parasitic infections, change of oral carriage of Enterobacteriaceae, bacterial infection and so on [29].

- **Mouth breathing**

Effects of mouth breathing include large overbite and crowded teeth, poor posture, inflamed tonsils, dry cracked lips, dry mouth which leads to bad breath, periodontal disease and tooth cavities. It can equally worsen symptoms in people with asthma [30]

- **Tongue thrusting**

Tongue thrust is defined as a condition in which the tongue makes contact with anterior teeth during swallowing and even continuously at random moments. Normal mature swallow shows the positioning of the tongue high on the palate behind the maxillary incisors and no lip and cheek activity during swallowing. A transitional swallowing pattern is seen in mixed dentition, when some primary teeth are lost and permanent are yet to erupt or are erupting. This type of swallow is self-correcting. In cases of open bite mostly created by habits like thumb sucking, the tongue gets thrust ahead to achieve a lip seal. This type of tongue thrust is called a “Simple Tongue Thrust”.

“Complex tongue thrust” is seen in individuals with a diffuse open bite, seen most commonly in mouth breathers and in a child with a history of chronic nasorespiratory disease/allergies [31]. Tongue thrusting can result in increased overjet, anterior open bite, posterior crossbite, labial incompetence (when at rest an open mouth position with a forward tongue posture is noted e.g while watching television or reading a book), difficulty with speech, especially the s and z sounds etc. [32]

- **Bruxism**

Bruxism, defined as a habitual nonfunctional forceful contact between occlusal tooth surfaces, is involuntary, excessive grinding, clenching or rubbing of teeth during nonfunctional movements of the masticatory system. Bruxism may occur during the day or during sleep. Bruxism has become an increasing concern in children due to its negative effects on life quality and also for being considered an important risk factor for temporomandibular dysfunctions. It may cause

tooth wear and, in more severe cases, dental traumas. Recent studies have indicated a relation between bruxism and respiratory alterations (27).

- **Tongue sucking**

When a child has a tongue sucking habit, he constantly sucks on the tongue as though on candy. The pressure put by children between the teeth while sucking the tongue causes incorrect swallowing and speech. If this disorder persists, it may have a devastating effect on a child's dental growth provoking more commonly an open bite and oral health in general (28).

### **Management of Parafunctional habits**

Old habits are hard to break and new habits are hard to form because the behavioral patterns we repeat are imprinted in our neural pathways, but it is possible to form new habits through repetition. In order to replace the adverse oral habits by good habits, a wholistic approach is indicated, which includes patient-parent counselling, behaviour modification techniques, use of habit breaking appliances, followed by recall visits and reinforcement [29]. Prevention and interception of these deleterious oral habits at an early stage is of utmost important for the good oral health of the children.

These treatment methods include [33];

**1. Psychological therapy-** Behavior shaping and modification and stress relief.

**2. Reminder therapy**

Extraoral approach: Ace bandage and long night sleeve gown.

Intraoral approach: removable appliances with palatal cribs.

**3. Mechanotherapy-**

Palatal crib appliance

Bluegrass appliance by Haskell and Mink consists of an acrylic roller that is placed in the most superior position of the palate with no contact onto the palatal tissues so that the roller can be moved freely with the tongue and this is welded onto the molar bands. Patient is asked to roll the roller with the tongue whenever he/she feels like thumb sucking.

Hay rake appliance consists of either sharp/blunt points/rakes can also feature loops or a palatal bar with short, blunt protuberances forming a hybrid which interferes with the digit or tongue while digit sucking or tongue thrusting and serves as a negative/painful feedback.

Quad helix appliance.

Electronic devices have been used like Alarming wrist watch consisted of a new device by *Krishnappa et al* with an alarm system that gets activated when the child placed the finger in the mouth. The alarm was incorporated in a wristwatch that made it appealing for the child to wear it and thus cooperated [34]

## II.7 MALOCCLUSION

Malocclusion is a form of relationship of the upper and lower jaw which deviate from the standard form that is accepted as a normal. On the sagittal planet, the basis for assessment is the intermaxillary positions of the first molars- ANGLE'S Classification [23] as seen previously. Other manifestations of malocclusion include;

### **Open bite**

Open bite is condition in which there is the presence of occlusal or incisal space between the teeth when the upper jaw and lower jaws are in centric occlusion.

Anterior open bite is a malocclusion with no contact in the anterior region of the dental arches, and the posterior teeth in occlusion. It is called combined open bite when malocclusion extends to the posterior segment.

Posterior open bite is failure of contact between the posterior teeth when the teeth occlude in centric occlusion, there is no occlusion between the maxillary and mandibular premolars. The

maxillary and mandibular molars have a slight contact. The anterior overjet and overbite are normal [17].

### **Deep bite**

Deep bite is defined as an excessive vertical overlap of the incisors, i.e. vertically, where more than two-thirds of the buccal surfaces of the mandibular incisors are covered by the maxillary incisors. Most often, the reason for deep bite is an over-eruption of the incisors or an anterior rotation of the mandible. A deep bite occasionally manifests with contact between the edges of the mandibular incisors and the palatal mucosa behind the maxillary incisors [35].

### **Posterior Crossbite**

In a posterior crossbite, the buccal cusps of the maxillary premolars and/or molars occlude lingually to the buccal cusps of the mandibular premolars and/or molars. The posterior crossbite can be either unilateral or bilateral. Unilateral crossbites of dento- alveolar origin are caused by palatal tipping of the maxillary premolars and molars, and is most often accompanied with a forced guidance of the mandible, thus deviating the midline of the mandible to the crossbite (29).

### **Crowding**

Dental crowding is defined as an inconsistency between tooth size and arch dimension that results in malocclusion; it occurs because of a lack of coordination between tooth size and arch dimensions leading to limited space for proper eruption [36].

## **REVIEW OF LITERATURE ON THE SUBJECT**

Although the prevalence of oral parafunctional habits and dental malocclusions vary in literature, several researches carried out around the world indicate a high prevalence.

A study in Cameroon in June 2022 by Kalla *et al.* showed that the most common parafunctional habit was thumb sucking (17.4%) with the higher proportion of thumb suckers being girls (63.79%) with an overall mean age of 4.80+-5 years. The highest proportion of thumb sucking was found in firstborn children (39.65%) and in those who breastfed for less than 6 months (85.62%) [3]

A worldwide study carried out in 2020 showed that the worldwide prevalence of malocclusion was 56% without differences in gender. The highest prevalence was in Africa (81%) and Europe (72%), followed by America (53%) and Asia (48%). The distributions of Class I, Class II and Class III malocclusions in Africa were found to be 76.9%, 9.7% and 4.0% respectively. Regarding vertical malocclusions, the observed deep bite and open bite were 10.9% and 6.5% respectively. Cross bites affected 9.8% of the sample [37].

However, we could find few studies have been carried out evaluating pediatricians' knowledge, attitudes and practices towards parafunctional habits and malocclusions.

A study conducted by Maden *et al.* in Turkey evaluating pediatricians' knowledge, attitude and practices on parafunctional habits and orthodontic problems in children revealed low level of knowledge on some aspects of parafunctional habits and orthodontic problems. The practice scores were lower (61.2%) compared to attitude scores (94.0%) and knowledge scores (71.2%) [5].

Kumar *et al.* in India researched the knowledge and attitude of pediatricians towards digit sucking in children. This study showed that many respondents were unaware that oral habits could be responsible for malocclusion. A high percentage of respondents preferred not to examine oral features in digit sucking child. Among the etiological factor that may contribute to digit sucking habit in a child, most of the respondents expressed that habitual, psychological and inadequate parental care are possible causes. Most of the participants answered that parental and child counseling is essential to stop the habit. Most of the respondents did not refer the digit sucking child to the dentist for the better management of associated malocclusion [1].

As seen above, similar studies were transverse and descriptive in nature which provides a comprehensive picture of the characteristics, behaviors, and attributes of the population which can be useful in informing future research and policy decision.



## **CHAPTER III: METHODOLOGY**

### **III.1 STUDY TYPE**

This is a descriptive cross-sectional study.

### **III.2 STUDY SITE**

Cameroon.

### **III.3 STUDY PERIOD**

STUDY DURATION- The study lasted from November 2023 to June 2024

DATA COLLECTION PERIOD- February to March 2024

### **III.4 STUDY POPULATION**

#### **III.4.i Target population**

This study targeted pediatricians in Cameroon.

#### **III.4.ii Source Population**

Pediatricians in Cameroon.

#### **III.4.iii Inclusion criteria**

The criteria for inclusion is any practicing pediatrician who consents to fill the questionnaire.

#### **III.4.iv Non-inclusion criteria**

Any physician who doesn't fit the inclusion criteria or doesn't consent to fill the questionnaire

#### **III.4.v Sampling**

*Type-* Consecutive and Exhaustive Sampling

### **III.5 PROCEDURE**

#### **III.5.i-Recruitment**

Pediatricians in Cameroon have a whatsapp group called *Communication SOCAPED*. Our questionnaire was shared in that group with reminders sent every couple of days. Also, some pediatricians in Yaounde and Douala were visited in their offices and given the hard copy of the questionnaires to fill.

Questionnaires were printed and submitted to the pediatricians and collected on appointment depending on their availability.

#### **III.5.ii-Administrative procedure**

Authorization was gotten at the ethics institutional committee for research of FMBS.

### **III.6 MATERIALS**

#### **Materials for data collection**

-Questionnaires

-Google forms

- Pens

#### **Materials for data analysis**

Data entry was done using Microsoft Excel 2016.

Analysis was done with the Statistical Package for Social Sciences (SPSS) 23.0.

The data was graded by section; Knowledge, Attitude and Practice (see table IV).

We used the scale for KAP studies by ESSI Marie and Njoya Oudou [38]

Each response was allocated 1 for correct answers and 0 for wrong answers.

-Total marks per section were 21,7 and 10 for the knowledge, attitude and practice sections respectively.

Knowledge was graded as poor, inadequate, average or good.

For attitude, it could be correct, approximative, wrong or poor.

Analysis of the practice was either poor, inadequate or adequate.

Table IV : KAP studies assessment

KAP	Category	Grade
Knowledge	Poor	Less than 50%
	Inadequate	[50-65% [
	Average	[65-80% [
	Good	Above 80%
Attitude	Poor	Less than 50%
	Inadequate	[50-75% [
	Adequate	Above 75%
Practice	Poor	Less than 50%
	Erroneous	[50-65% [
	Average	[65-80% [
	Good	Above 80%

### **III.7 Ethical Considerations**

After protocol validation by thesis supervisors, we sought approval from the Ethical Committee of the Faculty of Medicine and Biomedical Sciences of the University of Yaounde I.

Timeliness and interest of our study for the nation.

Parafunctional habits have been shown to have a negative impact on the developing occlusion which probably accounts for the increasing incidence of malocclusions in children. Cooperation between pediatricians and pediatric dentists will play a fundamental role in prevention.

#### **Voluntary Participation**

Participants were not coerced into taking part in the research.

#### **Risk of Harm**

There is no risk posed to any of the participants

#### **Confidentiality**

Participants' identities were kept secret during and after the research.

#### **Conflict of interest**

No conflict of interest were identified or reported during this study

## **CHAPTER IV: RESULTS**

#### IV.1. STUDY PARTICIPANTS

The aim of our study was to determine the knowledge, attitude and practice of paediatricians in Cameroon regarding parafunctional oral habits and dental malocclusions in children. A total of 159 respondents were encountered, 86 (54.09%) completed the study, 5 (3.14%) did not fill up to 75% of the questionnaires due to lack of time according to their reasons and 68 (42.77%) did not fill the questionnaire due to their busy schedules, numerous appointments without response and other reasons not mentioned.




Figure 6; Schematic representation of our study participants

## IV.2 SOCIODEMOGRAPHIC CHARACTERISTICS OF THE STUDY POPULATION

Of all the participants, 69.8% (n=60) were female while 30.2% (n=26) were male. With regards to the age distribution, the largest group of our participants 51.2% (n=44) were between the ages 40-49 years old with 44.2% (n=38) practising in both public and private setting. With regards to subspecialty, 30.2% (n=26) had a subspecialty, while 69.8% (n=60) do not.

When considering years of clinical practice, 50.0% (n=43), has 10-19 years of experience and 77.1% (n=67) of the sample spend 40 or more hours in patient care per week with the largest group, 53.5% (n=46) seeing 10-19 patients per day. Geographically, the regions with the highest representation are the Centre and the Littoral with 45.3% (n=39) and 38.4% (n=33) responses respectively, with all other regions making up smaller percentage (Table V)

Table V; Sociodemographic representation of participants

Variable (n=86)	Frequency	Percentage
Age(years)		
30-39	28	32.6
40-49	44	51.2
50+	16	16.3
Gender		
Male	26	30.2
Female	60	69.8
Subspecialty		
Yes	26	30.2
No	60	69.8
Years of clinical practice		
0-9	37	43.0
10-19	43	50.0
20+	6	7.0
Type of practice		
Public practice	31	36.0
Private practice	17	19.8



Both	38	44.2
Hours of patient care per week		
<40	19	22.9
>=40	67	77.1
Number of patients seen per day		
<=10	34	39.5
10-19	46	53.5
20-29	6	7.0
Region of practice		
North west	3	3.5
South west	2	2.3
Centre	39	45.3
Littoral	33	38.4
East	2	2.3
West	5	5.8
Far North	1	1.2
Adamawa	1	1.2

#### IV.3. RESULTS ON KNOWLEDGE OF PARAFUNCTIONAL ORAL HABITS AND DENTAL MALOCCLUSIONS AND THEIR RELATIONSHIP

According to the results, 43% (n=37) of respondents reported having knowledge that children with parafunctional oral habits are more likely to develop malocclusion, while 57% (n=49) do not. Additionally, 18.6% (n=16) indicated that long-term bottle feeding increases the possibility of malocclusion. Only 2.3% (n=2) of respondents believed that children with snoring/breathing problems are more prone to crowding. 51.2% (n=44) knew that prolonged pacifier use increases the risk of open-bite. Regarding the impact of supporting healthy chewing/swallowing, 43% (n=37) said it can help prevent malocclusion. Furthermore, 9.3% (n=8) reported that nail biting increases the risk of malocclusion, 52.3% (n=45) knew that thumb sucking increases the risk, and 32.6% (n=28) knew that tongue-sucking increases the risk (Table VI).

Table VI; Distribution according to knowledge of risk factors of malocclusion

Variable (N=86)	Frequency	Percentage
Children with parafunctional oral habits are more likely to develop malocclusion.	37	43
Long-term feeding of breast milk or formula with a bottle increases the possibility of developing parafunctional oral habits and malocclusion.	16	18.6
Children with snoring and breathing problems (mouth breathing) are more prone to crowding.	2	2.3
If the pacifier use time is prolonged, the possibility of open-bite increases?	44	51.2
Supporting healthy chewing and swallowing function in early childhood can help prevent malocclusion.	37	43
Nail biting or Onychophagia increases the risk of malocclusion.	8	9.3
Thumb sucking increases the risk of malocclusion.	45	52.3
Tongue-sucking increases the risk of malocclusion.	28	32.6

The most commonly known parafunctional habits were thumb sucking (95.3%, n=82), tongue thrusting (81.4%, n=70), and nail biting (58.1%, n=50). The most commonly known malocclusion was open bite (45.3%, n=39) with 41.9% (n=36) unable to identify any. Notably, 83.7% (n=72) had not received any training on parafunctional habits, and all respondents (100%) had not received training on orthodontic problems in children. When asked about recommended methods to receive such training, 52.3% (n=45) suggested doing so during medical school, 24.4% (n=21) during specialization, and 23.3% (n=20) through workshops

Table VII; Identification of parafunctional oral habits and malocclusions and training

Variable (N=86)	Frequency	Percentage
Parafunctional habits you know and can identify		
Thumb sucking	82	95.3
Lip biting	32	37.2
Tongue thrusting	70	81.4
Nail biting	50	58.1
Mouth breathing	26	30.2
Clenching of teeth	18	20.9
None	1	1.2
Manifestations of malocclusions you can identify		
Open bite	39	45.3
Deep bite	10	11.6
Crowding	20	23.3
Reverse bite	10	11.6
Increased overjet	3	3.5
None	36	41.9
Have you received any training on parafunctional habits (bad oral habits) and approaches to be applied in children?		
No	72	83.7

Yes, during medical school education.	11	12.8
Yes, during specialization.	3	3.5
Have you received any training on orthodontic problems in children and the approaches to be applied?		
No	86	100
Yes, during medical school education.	0	0
Yes, during specialization.	0	0
What methods to receive education and training on parafunctional oral habits and malocclusions would you recommend?		
Workshops	20	23.3
During medical school	45	52.3
During specialization	21	24.4




Fig 7: Knowledge of participants

40.7% of our participants had poor knowledge while 22.2% were average.

#### IV.4. RESULTS ON ATTITUDE TOWARDS PARAFUNCTIONAL HABITS AND MALOCCLUSIONS

98.8% (n=85) of our respondents believe that paediatricians have to examine the oral cavity, and the same percentage also believe that dental malocclusions can be prevented. Additionally, 97.7% (n=84) of respondents indicated that dental examination is important in the prevention of parafunctional oral habits and malocclusions. 96.5% (n=83) of the respondents agreed that paediatricians play an important role in identifying and diagnosing parafunctional oral habits and malocclusions in children, while 90.7% (n=78) believe paediatricians have an important part to play in preventing these issues. Table VIII shows information about their attitude.

Notably, 100% (n=86) of our respondents stated that they need more education on parafunctional habits and malocclusion prevention..

Table VIII; Distribution according to attitude

Variable (N=86)	Frequency	Percentage
Pediatricians have to examine the oral cavity.	85	98.8
Dental malocclusions can be prevented.	85	98.8
Dental examination is important in the prevention of parafunctional oral habits and malocclusions.	84	97.7
Pediatricians play an important role in the pointing out and diagnosing of parafunctional oral habits and malocclusions in children.	83	96.5
Pediatricians play an important role in the prevention of parafunctional oral habits and malocclusion in children.	78	90.7
I need more education on parafunctional habits and malocclusion prevention.	86	100

No respondents (0%) scored in the Poor range. The vast majority, 84 respondents or 97.7%, achieved a Good score.

#### IV.5. PRACTICE

According to our results, 76.7% (n=66) of respondents evaluate the oral functional habits of their patients. Only 10.5% (n=9) perform oral health examinations for malocclusion. Regarding the timing of the first oral health examination, 12.8% (n=11) do it immediately after birth, 37.7% (n=32) do it between 0-3 years, and 11.6% n=10) do it at 3 years or older. In terms of frequency, 47.7% (n=41) recommend oral health examinations only if there is a problem. Finally, 53.5% (n=46) refer patients to a pediatric dentist when they notice parafunctional oral habits or malocclusion.

Table IX; Practice

Variable (N=86)	Frequency	Percentage
You evaluate the oral functional habits of your patients?	66	76.7
You perform oral health examinations for malocclusion for your patients.	9	10.5
When do you perform the first oral health examination for your patients?		
I don't	33	38.4
Immediately after birth	11	12.8
0-3 years	32	37.7
>=3 years	10	11.6
When would you recommend the first oral health examination to your patients?		
Immediately after birth	17	19.8
0-3 years	62	72.1
>=3 years	7	8.1
How often do you recommend oral health examination?		
If any problem	41	47.7
Per 6 months	7	8.1

Per year	25	29.1
You refer your patient to a pediatric dentist when you notice any parafunctional oral habit or malocclusion	46	53.5

Of the 46 who refer patients, 60.9% (n=28) do it by advising families to visit the pediatric dentist, 36.9% (n=17) send them with a referral form and 2.2% (n=1) work directly with the pediatric dentist. On the other hand, only 60.9% (n=28) check if the visit was done.

Table X; Distribution according to referral methods

Variable (N=46)	Frequency	Percentage
If your answer to the previous question is yes, how do you do the referral?		
I advise families to go to a pediatric dentist.	28	60.9
I send them with a referral form to the pediatric dentist.	17	36.9
I work directly with the pediatric dentist	1	2.2
When your patient, whom you referred to the pediatric dentist comes to you for a check- up, you check whether the pediatric dentist examination is performed.	28	60.9




Fig 8: Practice of participants

60 individuals or 69.8%, scored in the Poor range while a significant portion, 25 respondents or 29.1%, scored in the Average range. Only 1.1% of the total, achieved a Good score of over 80.



## **Chapter V; DISCUSSION**

The aim of our study was to determine the knowledge, attitude and practice of pediatricians in Cameroon towards parafunctional oral habits and dental malocclusions. Out of the 159 pediatricians we approached, 86 were retained giving a response rate of 54.09%

### **LIMITATIONS OF STUDY**

- Our response rate was relatively low; probably due to the fact that our questionnaire was mostly administered online.
- Not all regions in Cameroon were represented.
- Pediatricians are not the only healthcare providers who work with children.
- Lastly, due to the self-reporting aspect of the study, we could not determine whether reported practices reflected actual clinical practices, so it is possible that respondents may over- or under report attitude and practice.

### **STRENGTHS**

- A major percentage of our sample live in the urban setting where there is relatively higher exposure.
- Eight out of the ten regions of Cameroon were represented which made it fairly representative.

## **5.1 SOCIO-PROFESSIONAL DATA**

In our study, 69.8% of the respondents were female. These results are similar to those of *Marianna et al* in 2019 in Greece where 67% of their respondents were female [39]. This could be explained by the fact that women have a relatively higher affinity for children than men which makes more women to go in for this specialty than men. Also, 51.2% of our respondents were between the ages of 40-49 years old.

53.5% reported consulting over 10 patients a day. This could be a contributing factor to them not carrying out oral examinations because of limited time.

45.3% and 38.4% of our participants were from the Centre and Littoral regions respectively with the other regions making up smaller percentages. This is probably due to the higher population

density in these regions. Also, the fact that the older faculties of medicine are found in these regions could explain the higher population of physicians in these regions.

With regards to years of clinical practice, 50.0% had 10-19 years of experience and 77.1% of the sample spend 40 or more hours in patient care per week. This can be explained by the relatively high pediatrician-patient ratio which makes it necessary for them to work long hours.

## 5.2 KNOWLEDGE

43% of respondents reported having knowledge that children with parafunctional oral habits are more likely to develop malocclusion, and 52.3% knew that thumb sucking increases the risk of malocclusion. These results are similar to those of *Kumar et al* in India where 56% of the respondents were aware of oral habits and 55% were aware of the adverse effects of thumb sucking on dentition [1].

Also, in our study, 18.6% of respondents indicated that long-term bottle feeding increases the possibility of malocclusion. These are significantly different from *Kumar et al*'s results where 86.1% had knowledge regarding the fact that long-term feeding of breast milk or formula with a bottle increases the likelihood of parafunctional oral habit and malocclusion [1].

Only 2.3% of respondents believed that children with snoring/breathing problems are more prone to crowding. 51.2% knew that prolonged pacifier use increases the risk of open-bite. Regarding the impact of supporting healthy chewing/swallowing, 43% said it can help prevent malocclusion and 9.3% reported that nail biting increases the risk of malocclusion. On the contrary, *Maden et al* had different results. 81.5% of the participants had sufficient knowledge regarding the fact that if the duration of pacifier use is prolonged, the possibility of open-bite increases and 71.7% of the participants answered the question that supporting healthy chewing function in early childhood can help to prevent malocclusion correctly. For the possibility of malocclusion in children with snoring and respiratory problems (mouth breathing), 71.7% of the participants answered this question correctly [5]. This difference might be related to the lack of required dental training and education of the majority of our participants.

Our respondents have an overall insufficient knowledge of parafunctional habits and malocclusions which could be explained by the fact that the field of dental medicine is relatively recent in our context.

In our study, 83.7% had not received any training on parafunctional habits, and 100% had not received training on orthodontic problems in children. This is similar to the results obtained by *Sezer et al* in 2013 where 89.2% reported having no training on oral health during residency and/or medical school [41] . This could be justified by the fact that dental courses have not fully been integrated in the curriculum of their training.

However, we can infer that the percentage of our participants who were able to identify malocclusions or parafunctional habits got the information through other means or through interactions with dentists given that most had not received any formal training on the subject.

### **5.3 ATTITUDE**

98.8% of our respondents believe that pediatricians have to examine the oral cavity, and the same percentage also believe that dental malocclusions can be prevented. Additionally, 97.7% of respondents indicated that dental examination is important in the prevention of parafunctional oral habits and malocclusions. 96.5% of the respondents agreed that pediatricians play an important role in identifying and diagnosing parafunctional oral habits and malocclusions in children. *Farsi et al* had very similar results during their study in Jeddah where a vast majority of pediatricians portrayed an overall good attitude towards oral health [42].

The high attitude scores in our study implied that majority of our respondents acknowledge their role in diagnosing and managing parafunctional oral habits and malocclusions.

### **5.4 PRACTICE**

According to our results, 76.7% of respondents evaluate the functional habits of their patients. Regarding the timing of the first oral health examination, 38.4% don't remember to do it. In a national survey with post-residency members of the American Academy of Pediatrics, 54%

claimed to examine the teeth of their 0–3year-old patients [43]. Lack of training was the most common reason for not performing an oral examination which applies in our context.

Demir *et al* found out that 19.3% of pediatricians in their study in Turkey don't perform the first oral examination [40].

In terms of frequency, 8.1% recommend oral health examinations every 6 months. Finally, 53.5% refer patients to a pediatric dentist when they notice parafunctional oral habits or malocclusion. These results are contrary to those of Maden *et al* where regarding how often an oral health examination was recommended, 49.4% of the participants indicated it as six months. Moreover, 97.6% said they would refer a child to pediatric dentists when they identified a child with malocclusion or any parafunctional habits [5]. This is somehow contradictory to the lower knowledge scores and their response to oral examination so we would assume that they notice these parafunctional habits without actually seeking them out.

In our study, practice scores were quite poor indicating no discrepancy with the knowledge scores and still could be justified by their relative lack of training in the field of dentistry.

## **Chapter VI: CONCLUSION and RECOMMENDATIONS**

## **CONCLUSION**

The purpose of this study was to assess the knowledge, attitude and practice of pediatricians regarding parafunctional oral habits and dental malocclusions in children. A cross sectional study was conducted among 86 pediatricians in Cameroon. The results showed that;

Pediatricians had poor knowledge regarding parafunctional oral habits and dental malocclusions in children. They could however identify certain parafunctional habits but had little knowledge regarding its effect on the developing dentition.

Pediatricians had a good attitude towards the subject. This was likely due to the fact that they had received little or no training in parafunctional oral habits and dental malocclusions.

Pediatricians practice regarding parafunctional oral habits and dental malocclusions was equally poor for a majority and average for a few.

## **RECOMMENDATIONS**

To pediatricians

To actively take part in trainings in the domain of parafunctional oral habits and dental malocclusions.

To do personal studies on the subject.

To pediatric and dental associations

To organize joint seminars in order to foster collaboration between both fields.

To researchers

To carry out related studies on a larger scale.

To the Faculty of Medicine and Biomedical Sciences

To organize seminars, workshops and conferences to train students on parafunctional habits and dental malocclusions.

To include courses on this subject in the curriculum of general medicine and specialization in pediatrics.



## REFERENCES

1. Kumar V, Shivanna V, Kopuri RC. Knowledge and attitude of pediatricians toward digit sucking habit in children. *J Indian Soc Pedod Prev Dent*. 2019 Mar;37(1):18.
2. Grippaudo C, Paolantonio EG, Antonini G, Saulle R, La Torre G, Deli R. *ACTA OTORHINOLARYNGOLOGICA ITALICA*. *Acta Otorhinolaryngol Ital*. 2016 Oct;36(5):386–94.
3. Kalla GCM, Tiomo EDM, Onana J, Mbopi-Keou FX, Messanga CB. Morphological and functional abnormalities of the orofacial sphere associated with thumb sucking in children aged 3 to 10 years old in Yaounde, Cameroon. *Pan Afr Med J [Internet]*. 2022 Jun 9 [cited 2024 Jun 5];42(107). Available from: <https://www.panafrican-med-journal.com/content/article/42/107/full>
4. Cenzato N, Nobili A, Maspero C. Prevalence of Dental Malocclusions in Different Geographical Areas: Scoping Review. *Dent J*. 2021 Oct 11;9(10):117.
5. Maden EA, Eker İ. Pediatricians' Knowledges, Attitudes and Practices on Parafunctional Oral Habits and Orthodontic Problems in Children. *Clin Exp Health Sci*. 2021 Oct 26;11(4):834–41.
6. Kamrani P, Sadiq NM. Anatomy, Head and Neck, Oral Cavity (Mouth). In: *StatPearls [Internet]*. Treasure Island (FL): StatPearls Publishing; 2024 [cited 2024 Jun 2]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK545271/>
7. Medscape Registration [Internet]. [cited 2024 Jun 2]. Available from: <https://emedicine.medscape.com/article/1899122-overview?form=fpf>
8. Lips (Left) | Complete Anatomy [Internet]. [cited 2024 Jun 2]. Available from: <https://www.elsevier.com/resources/anatomy/integument/skin/lips-left/19742>
9. Oral cavity: Anatomy, tongue muscles, nerves and vessels | Kenhub [Internet]. [cited 2024 Jun 2]. Available from: <https://www.kenhub.com/en/library/anatomy/the-oral-cavity>
10. Craniofacial Growth: Current Theories and Influence on Management - PubMed [Internet]. [cited 2024 Jun 2]. Available from: <https://pubmed.ncbi.nlm.nih.gov/32151371/>

11. Textbook of Craniofacial Growth - 1st Ed PDF | PDF | Collagen | Bone [Internet]. [cited 2024 Jun 2]. Available from: <https://www.scribd.com/document/410518665/Textbook-of-Craniofacial-Growth-1st-ed-pdf>
12. Face Embryology: Overview, Facial Morphogenesis, Early Development of the Face [Internet]. [cited 2024 Jun 2]. Available from: <https://emedicine.medscape.com/article/844962-overview>
13. GROWTH OF MAXILLA.
14. growth\_of\_mandible\_\_ortho\_.pdf [Internet]. [cited 2024 Jun 5]. Available from: [https://is.muni.cz/www/379606/26759304/growth\\_of\\_mandible\\_\\_ortho\\_.pdf](https://is.muni.cz/www/379606/26759304/growth_of_mandible__ortho_.pdf)
15. Child and Adult Dentition (Teeth) - Structure - Primary - Permanent - TeachMeAnatomy [Internet]. [cited 2024 Jun 2]. Available from: <https://teachmeanatomy.info/head/other/child-adult-dentition/>
16. Themes UFO. 1 Oral embryology, histology and anatomy [Internet]. Pocket Dentistry. 2015 [cited 2024 Apr 1]. Available from: <https://pocketdentistry.com/1-oral-embryology-histology-and-anatomy/>
17. Child and Adult Dentition (Teeth) - Structure - Primary - Permanent - TeachMeAnatomy [Internet]. [cited 2023 Dec 11]. Available from: <https://teachmeanatomy.info/head/other/child-adult-dentition/>
18. Anatomy of the tooth [Internet]. Kenhub. [cited 2023 Dec 11]. Available from: <https://www.kenhub.com/en/library/anatomy/anatomy-of-the-tooth>
19. Morris AL, Tadi P. Anatomy, Head and Neck, Teeth. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 [cited 2023 Dec 11]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK557543/>
20. Hedayat-Kelishadi·Dentistry· M. Dental Occlusion | Geeky Medics [Internet]. 2021 [cited 2023 Dec 11]. Available from: <https://geekymedics.com/occlusion/>
21. Ireland RI. incisor relationship. In: Ireland R, editor. A Dictionary of Dentistry [Internet]. Oxford University Press; 2010 [cited 2023 Dec 11]. Available from:


<https://www.oxfordreference.com/display/10.1093/acref/9780199533015.001.0001/acref-9780199533015-e-2195>

22. Mageet A. Classification of Skeletal and Dental Malocclusion: Revisited. *Stomatol EDU J*. 2016 Jan 1;3:205–11.
23. Biyani P. Angle's Classification for Malocclusions [Internet]. *dentalnotebook*. 2014 [cited 2023 Dec 11]. Available from: <https://www.dentalnotebook.com/angles-classification/>
24. Yassaei S, Rafieian M, Ghafari R. Abnormal oral habits in the children of war veterans. *J Clin Pediatr Dent*. 2005;29(3):189–92.
25. Rebouças P, Oliveira L, Lima H, Rodrigues L, Ferreira R. Prevalence of oral habits in children. *Revistas*. 2017 Dec 27;74:272.
26. S Dhull K, Verma T, Dutta B. Prevalence of Deleterious Oral Habits among 3- to 5-year-old Preschool Children in Bhubaneswar, Odisha, India. *Int J Clin Pediatr Dent*. 2018;11(3):210–3.
27. Massignan C, Santos J, Ximenes M, Cardoso M, Bolan M. Malocclusion traits and sucking habits in preschool children: a cross-sectional study. 2019 Jan 15;3:67–74.
28. Achmad H, Riyanti E, Djais AI, Rieuwpassa IE, Saptarini R, Arief SM. The Effect of Thumb Sucking Habit on Children's Dentomaxillofacial Development: A Systematic Review. *Syst Rev Pharm*. 2021;12(2).
29. C B S. ORAL PARA-FUNCTIONAL HABITS – A HAWK-EYE VIEW. *Int J Recent Adv Multidiscip Res*. 2017 Feb 28;4:2263–5.
30. Mouth Breathing [Internet]. *Healthline*. 2017 [cited 2023 Dec 11]. Available from: <https://www.healthline.com/health/mouth-breathing>
31. Shah SS, Nankar MY, Bendgude VD, Shetty BR. Orofacial Myofunctional Therapy in Tongue Thrust Habit: A Narrative Review. *Int J Clin Pediatr Dent*. 2021;14(2):298–303.
32. Ali F, Soni S, Kaur G, Bagga MK. Oral Habits in Relation to Malocclusions: A Review. *Int J Health Sci*. 2021 Dec 31;230–8.

33. Greenleaf S, Dds JM. A Retrospective Study of the Use of the Bluegrass Appliance in the Cessation of Thumb Habits. *Pediatr Dent*. 2003;
34. Thumb Sucking - StatPearls - NCBI Bookshelf [Internet]. [cited 2024 Apr 22]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK556112/>
35. Themes UFO. Classification of malocclusions [Internet]. Pocket Dentistry. 2017 [cited 2023 Dec 11]. Available from: <https://pocketdentistry.com/classification-of-malocclusions/>
36. Patano A, Malcangi G, Inchingolo AD, Garofoli G, De Leonardis N, Azzollini D, et al. Mandibular Crowding: Diagnosis and Management—A Scoping Review. *J Pers Med*. 2023 Apr 29;13(5):774.
37. Lombardo G, Vena F, Negri P, Pagano S, Barilotti C, Paglia L, et al. Worldwide prevalence of malocclusion in the different stages of dentition: A systematic review and meta-analysis. *Eur J Paediatr Dent*. 2020 Jun;21(2):115–22.
38. Essi MJ, Njoya O. L'enquête CAP en recherche médicale. *Health Sci Dis* [Internet]. 2013 Aug 9 [cited 2024 Jun 5];14(2). Available from: <https://www.hsd-fmsb.org/index.php/hsd/article/view/183>
39. Paediatricians' awareness on orthodontic problems and related conditions—a national survey - PMC [Internet]. [cited 2024 Jun 2]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6702498/>
40. DemiR P, BiLmez Selen M. Knowledge and Management of Pediatricians About Children's Oral Health. *Clin Exp Health Sci*. 2021 Sep 27;11(3):582–8.
41. Sezer RG, Paketci C, Bozaykut A. Paediatricians' awareness of children's oral health: Knowledge, training, attitudes and practices among Turkish paediatricians. *Paediatr Child Health*. 2013 Apr;18(4):e15-19.
42. Farsi D, Alagili D. Oral Health Knowledge, Attitudes, and Clinical Practices of Pediatricians and Pediatric Residents: A Cross-Sectional Study. *Cureus*. 15(12):e50785.
43. Oral health and pediatricians: results of a national survey - PubMed [Internet]. [cited 2024 Jun 5]. Available from: <https://pubmed.ncbi.nlm.nih.gov/19945080/>

**APPENDIX**

**UNIVERSITÉ DE YAOUNDE II**  
**FACULTÉ DE MÉDECINE ET DES SCIENCES BIOMÉDICALES**  
**COMITÉ INSTITUTIONNEL D'ÉTHIQUE DE LA RECHERCHE**  
 Tel/ fax : 22 31-05-86 22 311224  
 Email: decanatfmsb@hotmail.com



**THE UNIVERSITY OF YAOUNDE I**  
**FACULTY OF MEDICINE AND BIOMEDICAL SCIENCES**  
**INSTITUTIONAL ETHICAL REVIEW BOARD**

Ref. : N° 0710 /UY1/FMSB/VRRC/D/ASR/CSD

**CLAIRANCE ÉTHIQUE**

10 JUIN 2017

**Le COMITÉ INSTITUTIONNEL D'ÉTHIQUE DE LA RECHERCHE (CIER) de la FMSB a examiné**  
 La demande de la clairance éthique soumise par :

**M.Mme : AWA CLAVICE WOPONG**                      **Matricule: 17M169**

**Travaillant sous la direction de :**

- ♦ Pr KOKI NDOMBO Paul
- ♦ Dr TONY NENGOM Jocelyn
- ♦ Dr LOWE NANTCHOUANG Jacqueline Michèle Epse ABISSEGUE

**Concernant le projet de recherche intitulé :** Pediatricians' knowleges, attitudes and practices regarding parafunctional habits and malocclusions in children


**Les principales observations sont les suivantes**

Evaluation scientifique	
Evaluation de la convenance institutionnelle/valeur sociale	
Equilibre des risques et des bénéfices	
Respect du consentement libre et éclairé	
Respect de la vie privée et des renseignements personnels (confidentialité) :	
Respect de la justice dans le choix des sujets	
Respect des personnes vulnérables :	
Réduction des inconvénients/optimalisation des avantages	
Gestion des compensations financières des sujets	
Gestion des conflits d'intérêt impliquant le chercheur	

Pour toutes ces raisons, le **CIER** émet un avis **favorable** sous réserve des modifications recommandées dans la grille d'évaluation scientifique.

L'équipe de recherche est responsable du respect du protocole approuvé et ne devra pas y apporter d'amendement sans avis favorable du **CIER**. Elle devra collaborer avec le **CIER** lorsque nécessaire, pour le suivi de la mise en œuvre dudit protocole. La clairance éthique peut être retirée en cas de non - respect de la réglementation ou des recommandations sus évoquées. En foi de quoi la présente clairance éthique est délivrée pour servir et valoir ce que de droit

**LE PRÉSIDENT DU COMITE ETHIQUE**



## Questionnaire

### KNOWLEDGE, ATTITUDES AND PRACTICES OF PEDIATRICIANS TOWARDS PARAFUNCTIONAL ORAL HABITS AND DENTAL MALOCCLUSIONS IN CHILDREN

No	Question	Answer	Score
<b>SOCIODEMOGRAPHIC DATA</b>			
1	Age	30-39 40-49 50+	
2	Gender	Male Female	
3	Subspecialty	Yes No	
4	Years of clinical experience	0-9 10-19 20+	
5	Type of practice	Government Hospital Private Hospital Both	
6	Hours of patient care per week	<40 ≥40	
7	Number of patients seen per day	≤10 10-19	



		20-29	
8	Region of practice	North West South West Centre Littoral East West Far North North Adamawa South	
KNOWLEDGE			
9	Children with parafunctional oral habits are more likely to develop malocclusion. Do you have any information about this?	Yes No	
10	Long-term feeding of breast milk or formula with a bottle increases the possibility of developing parafunctional oral habits and malocclusion. Do you have any information about this?	Yes No	
11	Children with snoring and breathing problems (mouth breathing) are more prone to crowding. Do you have any information about this?	Yes No	

12	If the pacifier use time is prolonged, the possibility of open-bite increases. Do you have any information about this?	Yes No	
13	Supporting healthy chewing and swallowing function in early childhood can help prevent malocclusion. Do you have any information about this?	Yes No	
14	Nail biting or Onychophagia increases the risk of malocclusion. Do you have any information about this?	Yes No	
15	Thumb sucking increases the risk of malocclusion. Do you have any information about this?	Yes No	
16	Tongue-sucking increases the risk of malocclusion. Do you have any information about this?	Yes No	
17	Which of the following parafunctional habits do you know and can identify? (Multiple answers)	Thumb sucking Lip biting Tongue thrusting Nail biting Mouth breathing Clenching of teeth	

18	Which of the following manifestations of malocclusions can you identify? (multiple answers)	Open bite Deep bite Crowding Reverse bite Increased Overjet None	
19	Have you received any training on parafunctional habits (bad oral habits) and approaches to be applied in children?	No Yes, I got it in my medical education Yes, I got it in my specialist training	
20	Have you received any training on orthodontic problems in children and the approaches to be applied?	No Yes, I got it in my medical school training Yes, I got it in my specialist training	
ATTITUDE			
21	Pediatricians have to examine the oral cavity.	Yes No	
22	Dental malocclusions can be prevented.	Yes	

		No	
23	Dental examination is important in the prevention of parafunctional oral habits and malocclusions.	Yes No	
24	Pediatricians play an important role in the pointing out and diagnosing of parafunctional oral habits and malocclusions in children.	Yes No	
25	Pediatricians play an important role in the prevention of parafunctional oral habits and malocclusion in children.	Yes Non	
26	I need more education on parafunctional habits and malocclusion prevention.	Yes Non	
27	If your previous answer is yes, what methods to receive education and training on parafunctional oral habits and malocclusions would you prefer?	Workshops and colleagues During medical school During specialisation	
PRACTICE			
28	Do you evaluate the functional habits of your patients?	No Yes	

29	Do you perform oral health examinations for malocclusion for your patients?	Yes No	
30	When do you perform the first oral health examination for your patients?	I don't Immediately after birth 0-3 years ≥3 years	
31	When would you recommend the first oral health examination to your patients?	Immediately after birth 0-3 years ≥3 years	
32	How often do you recommend oral health examination?	If any problem per 6 months per year I don't	
33	Do you refer your patient to a pediatric dentist when you notice any parafunctional oral habit or malocclusion?	Yes No	
34	If your answer to the previous question is yes, how do you do the referral? (multiple answers possible)	I advise families to go to a pediatric dentist. I send them with a referral form to the pediatric dentist. I work directly with the pediatric dentist by giving information myself.	

35	When your patient, whom you referred to the pediatric dentist comes to you for a check-up, do you check whether the pediatric dentist examination is performed?	Yes No	

### Questionnaire

#### CONNAISSANCES, ATTITUDES ET PRATIQUES DES PÉDIATRES À L'ÉGARD DES HABITUDES BUCCALES PARAFONCTIONNELLES ET DES MALOCCLUSIONS DENTAIRES CHEZ L'ENFANT

No	Question	Reponse	Note
DONNÉES SOCIO-PROFESSIONNELLES			
1	Âge	30-39 40-49 50+	
2	Genre	Mâle Femelle	
3	Sous spécialité	Yes No	
4	Années d'expérience clinique	0-9 10-19 20+	
5	Lieu de pratique	Hopital public	

		Hopital prive Les deux	
6	Heures de soins aux patients par semaine	<40 ≥40	
7	Nombre de patients vus par jour	≤10 10-19 20-29	
8	Région de pratique	Nord Ouest Sud-Ouest Centre Littoral Est Ouest Extrême nord Nord Adamaoua Sud	
Section II ; CONNAISSANCES			
9	Les enfants ayant des habitudes buccales parafunctionnelles sont plus susceptibles de développer une malocclusion. Avez-vous des informations à ce sujet ?	Oui Non	

10	L'alimentation à long terme avec du lait maternel ou du lait maternisé au biberon augmente la possibilité de développer des habitudes buccales parafunctionnelles et une malocclusion. Avez-vous des informations à ce sujet ?	Oui Non	
11	Les enfants souffrant de ronflements et de problèmes respiratoires (respiration buccale) sont plus sujets à la promiscuité. Avez-vous des informations à ce sujet ?	Oui Non	
12	Si la durée d'utilisation de la tétine est prolongée, la possibilité de béance augmente. Avez-vous des informations à ce sujet ?	Oui Non	
13	Soutenir une fonction saine de mastication et de déglutition dès la petite enfance peut aider à prévenir les malocclusions. Avez-vous des informations à ce sujet ?	Oui Non	
14	Le rongement des ongles ou Onychophagie augmente le risque de malocclusion. Avez-vous des informations à ce sujet ?	Oui Non	
15	La succion du pouce augmente le risque de malocclusion. Avez-vous des informations à ce sujet ?	Oui Non	
16	La succion de la langue augmente le risque de malocclusion. Avez-vous des informations à ce sujet ?	Oui Non	



17	Parmi les habitudes parafonctionnelles suivantes, lesquelles connaissez-vous et pouvez-vous identifier ? (Plusieurs réponses)	<p>Succion du pouce</p> <p>Succion de la langue</p> <p>Rongement des ongles</p> <p>Respiration buccale</p> <p>grincement des dents</p> <p>Aucun</p>	
18	Parmi les manifestations suivantes de malocclusions, lesquelles pouvez-vous identifier ? (plusieurs réponses)	<p>Béance</p> <p>Encombrement</p> <p>Occlusion inversée</p> <p>Augmentation du surplomb</p> <p>Aucun</p>	
19	Avez-vous reçu une formation sur les habitudes parafonctionnelles chez les enfants et les approches à appliquer ?	<p>Non</p> <p>Oui, je l'ai obtenu lors de ma formation en médecine</p> <p>Oui, je l'ai obtenu dans ma spécialisation</p>	

20	Avez-vous reçu une formation sur les problèmes orthodontiques chez les enfants et les approches à appliquer ?	Non  Oui, je l'ai obtenu lors de ma formation en médecine  Oui, je l'ai obtenu dans ma spécialisation	
21	Quelles méthodes préféreriez-vous pour recevoir une éducation et une formation sur les habitudes buccales parafunctionnelles et les malocclusions ?	<ul style="list-style-type: none"> <li>• Ateliers et collègues</li> <li>• Pendant les études de médecine</li> <li>• Pendant la spécialisation</li> </ul>	
ATTITUDE			
22	Les pédiatres doivent examiner la cavité buccale.	Oui  Non	
23	Les malocclusions dentaires peuvent être évitées.	Oui  Non	
24	L'examen dentaire est important dans la prévention des habitudes buccales parafunctionnelles et des malocclusions.	Oui  Non	

25	Les pédiatres jouent un rôle important dans la détection et le diagnostic des habitudes buccales parafunctionnelles et des malocclusions chez les enfants.	Oui Non	
26	Les pédiatres jouent un rôle important dans la prévention des habitudes buccales parafunctionnelles et des malocclusions chez les enfants.	Oui Non	
27	J'ai besoin de plus d'éducation sur les habitudes parafunctionnelles et la prévention des malocclusions.	Oui Non	
PRATIQUES			
28	Évaluez-vous les habitudes fonctionnelles de vos patients ?	Non Oui	
29	Effectuez-vous des examens de santé bucco-dentaire pour détecter une malocclusion chez vos patients ?	Oui Non	
30	Quand effectuez-vous le premier examen de santé bucco-dentaire de vos patients ? •	Je ne le fais pas Immédiatement après la naissance 0-3 ans ≥3 ans	

31	Quand recommanderiez-vous le premier examen de santé bucco-dentaire à vos patients ?	Immédiate ment après la naissance  0-3 ans  ≥3 years	
32	À quelle fréquence recommandez-vous un examen de santé bucco-dentaire ?	En cas de probleme  Tout les 6 mois  Par an  Je ne le fais pas	
33	Orientez-vous votre patient vers un dentiste pédiatrique lorsque vous constatez une habitude buccale parafunctionnelle ou une malocclusion ?	Oui  Non	
34	Si votre réponse à la question précédente est oui, comment procédez-vous au référencement ? (plusieurs réponses possibles)	Je conseille aux familles d'aller chez un pedodontiste  Je les envoie avec un formulaire de référence au pedodontiste pédiatrique.  Je travaille directement avec le pedodontiste en	

		donnant moi-même les informations.	
35	Lorsque votre patient, que vous avez adressé au dentiste pédiatrique, vient chez vous pour un contrôle, vérifiez-vous si l'examen du dentiste pédiatrique a été effectué ?	Oui Non	

## ANTI-PLAGIARISM REPORT

---

### PEDIATRICIANS' KNOWLEDGE, ATTITUDES AND PRACTICES TOWARDS PARAFUNCTIONAL ORAL HABITS AND DENTAL MALOCCLUSIONS IN CHILDREN

#### ABSTRACT

**Introduction.** The World Health Organization (WHO) considers malocclusion as one of the most important oral health problems, after caries and periodontal disease. Its prevalence is highly variable and is estimated to be between 39% and 93% in children and adolescents. Parafunctional habits are abnormal hyperactive functions conducted by the masticatory structures, i.e. tongue, teeth, oral muscles, and so on. They include thumb sucking, tongue thrusting, bruxism, prolonged use of feeding bottle or pacifier and so on. Parafunctional habits happen to be a major risk factor for malocclusion. Adequate knowledge of pediatricians and their collaboration with oral healthcare providers would go a long way into sensitizing patients on parafunctional habits and their effect on the developing dentition, as well as intercepting malocclusions at an early stage where they can be corrected through referrals to orthodontists or pediatric dentists.

**Objective.** Our objective was to evaluate the knowledge, attitude and practice of paediatricians in Cameroon towards parafunctional oral habits and dental malocclusions in children.

**Materials and Methods.** A descriptive, cross-sectional survey was sent to a sample of pediatricians in Cameroon. The questionnaire consisted of 35 questions in four domains; socioprofessional information, knowledge, attitude and practice.

**Results.** A total of 86 pediatricians participated in the survey. In terms of knowledge, 43% of respondents reported having knowledge that children with parafunctional oral habits are more likely to develop malocclusion. Regarding their attitude, 97.7% of respondents had a good attitude towards the subject. In terms of examination of the oral cavity, a low frequency of examination amongst pediatricians for malocclusion (10.5%) and oral functional habits (76.7%) was detected. 47.7% recommend oral health examinations only if there is a problem and 53.5% refer patients to a pediatric dentist when they notice any abnormality.

**Conclusion.** Although a vast majority of the paediatricians in our survey recognised their place in identifying and preventing parafunctions and malocclusions, very few have adequate knowledge and skills to perform evaluations and referrals.

*Key words: Pediatricians, knowledge, attitude, practice, parafunctions, malocclusion*

#### CHAPTER I: INTRODUCTION










##### 1.1 BACKGROUND














## Dupli Checker Plagiarism Report

Originality Report

 <b>16%</b> <b>Plagiarized Content</b>	 <b>84%</b> <b>Unique Content</b>	<b>4328</b> Words	<b>30760</b> Characters
--	---	----------------------	----------------------------











## Primary Sources

1	<a href="https://www.ncbi.nlm.nih.gov/pmc/arti...">https://www.ncbi.nlm.nih.gov/pmc/arti...</a> by N Cenzato · 2021 · Cited by 83 — The World Health Organization (WHO) considers malocclusion one of the most...	 <b>1.30 %</b>
2	<b>Plagiarized Content Link</b> WEB · The World Health Organization (WHO) considers malocclusion one of the most important oral health problems,...	 <b>0.33 %</b>
3	<a href="https://www.sciencedirect.com/science...">https://www.sciencedirect.com/science...</a> A cross-sectional survey to ascertain the prevalence of oral ...	 <b>0.33 %</b>
4	<a href="https://www.jrmds.in/articles/knowled...">https://www.jrmds.in/articles/knowled...</a> Knowledge and Awareness on Habits and Habit Breaking ...	 <b>0.33 %</b>
5	<a href="https://www.ncbi.nlm.nih.gov/pmc/arti...">https://www.ncbi.nlm.nih.gov/pmc/arti...</a> Jul 10, 2015 · The negative influence of bad habits on occlusion originates in childhood. Bottle feeding and nonnutritive sucking...	 <b>0.65 %</b>
6	<a href="https://pubmed.ncbi.nlm.nih.gov/3467...">https://pubmed.ncbi.nlm.nih.gov/3467...</a> Oct 11, 2021 · The World Health Organization (WHO) considers malocclusion one of the most important oral health problems,...	 <b>0.33 %</b>
7	<a href="https://www.ncbi.nlm.nih.gov/pmc/arti...">https://www.ncbi.nlm.nih.gov/pmc/arti...</a> by NE Mueller · 2018 · Cited by 29 — For example, in a study published by Kumar et al) a majority of their potential participa...	 <b>0.33 %</b>
8	<a href="https://pubmed.ncbi.nlm.nih.gov/3080...">https://pubmed.ncbi.nlm.nih.gov/3080...</a> Thumb Sucking - StatPearls - NCBI BookshelfKnowledge and attitude of pediatricians toward digit sucking ...	 <b>0.33 %</b>
9	<a href="https://go.gale.com/ps/i.do?id=GALE%...">https://go.gale.com/ps/i.do?id=GALE%...</a> by EA Maden · 2021 · Cited by 2 — ... prevent the development of significant anomalies in late mixed and permanent dentition to...	 <b>1.63 %</b>

10	<a href="https://pubmed.ncbi.nlm.nih.gov/2797...">https://pubmed.ncbi.nlm.nih.gov/2797...</a> Dec 15, 2016 · Methods: This is a descriptive cross-sectional study via an anonymous web-based survey administered between Apr...	 0.33 %
11	<a href="https://www.ncbi.nlm.nih.gov/pmc/arti...">https://www.ncbi.nlm.nih.gov/pmc/arti...</a> by AZ Fatani · 2021 · Cited by 5 — Data entry was done using Microsoft Excel 2016 (Microsoft Corporation, Redmond, WA) an...	 0.33 %
12	<a href="https://pubmed.ncbi.nlm.nih.gov/3858...">https://pubmed.ncbi.nlm.nih.gov/3858...</a> by R Mahalie · 2024 — The analysis was done with the Statistical Package for Social Sciences (SPSS) version 25. Pearson Chi-Squa...	 0.33 %
13	<a href="https://www.ncbi.nlm.nih.gov/pmc/arti...">https://www.ncbi.nlm.nih.gov/pmc/arti...</a> by SN Amougou · 2019 · Cited by 13 — Ethical considerations: the study was approved by the Institutional Ethics Committee of the—	 0.33 %
14	<a href="https://www.tandfonline.com/doi/full/1...">https://www.tandfonline.com/doi/full/1...</a> by MT Macheke · 2023 — Participants were not coerced into taking part in the research and they were fully informed about...	 0.33 %
15	<a href="https://pubmed.ncbi.nlm.nih.gov/3325...">https://pubmed.ncbi.nlm.nih.gov/3325...</a> The aim of our study was to determine the knowledge and skills of such persons who bring victims of road traffic crashes to...	 0.65 %
16	<a href="https://www.academia.edu/93582304/...">https://www.academia.edu/93582304/...</a> ... Children with parafunctional oral habits are more likely to develop malocclusion. Do you have any information about this?...	 2.61 %
17	<a href="https://dergipark.org.tr/tr/download/a...">https://dergipark.org.tr/tr/download/a...</a> by EA MADEN · Cited by 2 — Dental examination is important in the prevention of malocclusion and parafunctional oral habits...	 0.65 %
18	<a href="https://pubmed.ncbi.nlm.nih.gov/3608...">https://pubmed.ncbi.nlm.nih.gov/3608...</a> Dec 1, 2022 · Summary: Pediatricians play an important role in the prevention of teratogenic exposures. Ensuring optimal...	 0.33 %
19	<a href="https://dergipark.org.tr/tr/download/a...">https://dergipark.org.tr/tr/download/a...</a> by EA MADEN — Do you evaluate the oral functional habits of your patients with ATH? Yes. 84. 68.3. No. 13. 10.6. If any proble...	 0.33 %
20	<a href="https://www.supremecourt.ohio.gov/B...">https://www.supremecourt.ohio.gov/B...</a> Jun 2, 2017 — Q44 If your answer to the previous question is "yes", how many eligible defendants were released last year? Q4...	 0.33 %
21	<b>Plagiarized Content Link</b> WEB · Not all regions in Cameroon were represented making it difficult to generalize the findings of this review. All sources of...	 0.33 %
22	<a href="https://en.wikipedia.org/wiki/Languag...">https://en.wikipedia.org/wiki/Languag...</a> Eight out of the ten regions of Cameroon are primarily francophone and two are anglophone. The official percentage o...	 0.33 %

1



23	<a href="https://ludwig.guru/s/are+similar+to+t...">https://ludwig.guru/s/are+similar+to+t...</a> The part of the sentence "are similar to those of previous" is correct and usable in written English. You can use it to compare...	 0.33 %
24	<a href="https://www.reddit.com/r/barstoolspor...">https://www.reddit.com/r/barstoolspor...</a> This could be a contributing factor to them not bringing in as many new personalities as they could/should. You bring a guy in...	 0.33 %
25	<a href="https://www.researchgate.net/figure/P...">https://www.researchgate.net/figure/P...</a> This is probably due to the higher population sizes of domesticated populations compared to wild populations, as wel...	 0.33 %
26	<a href="https://ludwig.guru/s/by+the+relative...">https://ludwig.guru/s/by+the+relative...</a> This can be explained by the relatively smaller Mo2C crystallites, which give higher Mo2C dispersion over the catalysts. 1. Applied...	 0.33 %
27	<b>Plagiarized Content Link</b> WEBSimilarly, 81.5% of the participants had sufficient knowledge regarding that if the duration of pacifier use is prolonged, the...	 0.33 %
28	<a href="https://www.ncbi.nlm.nih.gov/pmc/arti...">https://www.ncbi.nlm.nih.gov/pmc/arti...</a> by AR Bowers · 2016 · Cited by 16 — It is possible that this difference might be related to the lack of prior driving experien...	 0.33 %
29	<a href="https://pubmed.ncbi.nlm.nih.gov/3591...">https://pubmed.ncbi.nlm.nih.gov/3591...</a> A survey of Indian corneal surgeons	 0.33 %
30	<a href="https://moneder.market/en/blog/news...">https://moneder.market/en/blog/news...</a> Feb 9, 2024 — The importance of training in the field of dentistry is undeniable and crucial to achieving excellent results and...	 0.33 %
31	<a href="https://pubmed.ncbi.nlm.nih.gov/3360...">https://pubmed.ncbi.nlm.nih.gov/3360...</a> Abstract. Background COVID-19 is significantly affecting the healthcare system globally. As a result, healthcare workers need...	 0.33 %
32	<a href="https://twitter.com/jaiswalmdphd/statu...">https://twitter.com/jaiswalmdphd/statu...</a> This was likely due to the fact that 1 out of every 18 exonic mutations in HSCs led to a fitness advantage, a number that is...	 0.33 %