

Applied Partial Differential Equations
with Fourier Series and Boundary Value Problems
Richard Haberman
Solutions by David A. Lee

All errors, typographical and substantive, and other offenses, are entirely my own.

Table of Contents

3 - Fourier Series	3
3.2 - Statement of Convergence Theorem	3
3.3 - Fourier Cosine and Sine Series	12
3.4 - Term-by-Term Differentiation of Fourier Series	39

3 - Fourier Series

3.2 - Statement of Convergence Theorem

Question: 3.2.1

For the following functions, sketch the Fourier series of $f(x)$ (on the interval $-L \leq x \leq L$). Compare $f(x)$ to its Fourier series:

(a) $f(x) = 1$

(b) $f(x) = x^2$



(c) $f(x) = 1 + x$






(d) $f(x) = e^x$

(e) $f(x) = \begin{cases} x & x < 0 \\ 2x & x > 0 \end{cases}$

(f) $f(x) = \begin{cases} 0 & x < 0 \\ 1 + x & x > 0 \end{cases}$

(g) $f(x) = \begin{cases} x & x < L/2 \\ 0 & x > L/2 \end{cases}$





Question: 3.2.2

For the following functions, sketch the Fourier series of $f(x)$ (on the interval $-L \leq x \leq L$ and determine the Fourier coefficients:

(a) $f(x) = x$

(b) $f(x) = e^{-x}$

(c) $f(x) = \sin\left(\frac{\pi x}{L}\right)$

(d) $f(x) = \begin{cases} 0 & x < 0 \\ x & x > 0 \end{cases}$


(e) $f(x) = \begin{cases} 1 & |x| < L/2 \\ 0 & |x| > L/2 \end{cases}$

(f) $f(x) = \begin{cases} 0 & x < 0 \\ 1 & x > 0 \end{cases}$

(g) $f(x) = \begin{cases} 1 & x < 0 \\ 2 & x > 0 \end{cases}$


- (a) Since $f(x)$ is odd, we only have non-zero coefficients for the Fourier sine series:

$$\begin{aligned} b_n &= \frac{1}{L} \int_{-L}^L x \sin\left(\frac{n\pi x}{L}\right) dx \\ &= \frac{1}{L} \left[-\frac{L}{n\pi} x \cos\left(\frac{n\pi x}{L}\right) \Big|_{-L}^L + \frac{L}{n\pi} \int_{-L}^L \cos\left(\frac{n\pi x}{L}\right) dx \right] \\ &= -\frac{2L}{n\pi} \cos(n\pi) \\ &= \frac{2L}{n\pi} (-1)^{n+1} \end{aligned}$$




(b)

$$\begin{aligned}
a_0 &= \frac{1}{2L} \int_{-L}^L e^{-x} dx \\
&= \frac{1}{2L} (e^L - e^{-L}) \\
&= \frac{1}{L} \sinh(L) \\
a_n &= \frac{1}{L} \int_{-L}^L e^{-x} \cos\left(\frac{n\pi x}{L}\right) dx \\
&= \frac{1}{L} \left[\frac{e^{-x} \left(\frac{n\pi}{L} \sin\left(\frac{n\pi x}{L}\right) - \cos\left(\frac{n\pi x}{L}\right) \right)}{1 + \frac{n^2\pi^2}{L^2}} \right]_{-L}^L \\
&= \frac{(-1)^n (e^L - e^{-L})}{L \left(1 + \frac{n^2\pi^2}{L^2}\right)} \\
&= \frac{2(-1)^n L \sinh(L)}{L^2 + n^2\pi^2} \\
b_n &= \frac{1}{L} \int_{-L}^L e^{-x} \sin\left(\frac{n\pi x}{L}\right) dx \\
&= -\frac{1}{L} \left[\frac{e^{-x} \left(\sin\left(\frac{n\pi x}{L}\right) + \frac{n\pi}{L} \cos\left(\frac{n\pi x}{L}\right) \right)}{1 + \frac{n^2\pi^2}{L^2}} \right]_{-L}^L \\
&= \frac{n\pi(-1)^n (e^L - e^{-L})}{L^2 \left(1 + \frac{n^2\pi^2}{L^2}\right)} \\
&= \frac{2(-1)^n n\pi \sinh(L)}{L^2 + n^2\pi^2}
\end{aligned}$$




(c) Since $f(x)$ is odd, $a_0 = 0$. By orthogonality, $a_n = 0$ for all n , and $b_n = 0$, $n \neq 1$. We are left with

$$\begin{aligned}
b_1 &= \frac{1}{L} \int_{-L}^L \sin^2\left(\frac{\pi x}{L}\right) dx \\
&= 1
\end{aligned}$$



(d)


$$\begin{aligned}
 a_0 &= \frac{1}{2L} \int_0^L x \, dx \\
 &= \frac{L}{4} \\
 a_n &= \frac{1}{L} \int_0^L x \cos\left(\frac{n\pi x}{L}\right) dx \\
 &= \frac{1}{L} \left[\frac{L}{n\pi} x \sin\left(\frac{n\pi x}{L}\right) \Big|_0^L - \frac{L}{n\pi} \int_0^L \sin\left(\frac{n\pi x}{L}\right) dx \right] \\
 &= \frac{1}{L} \cdot \frac{L^2}{n^2\pi^2} \cos\left(\frac{n\pi x}{L}\right) \Big|_0^L \\
 &= \frac{L}{n^2\pi^2} ((-1)^n - 1) \\
 b_n &= \frac{1}{L} \int_0^L x \sin\left(\frac{n\pi x}{L}\right) dx \\
 &= \frac{1}{L} \left[-\frac{L}{n\pi} x \cos\left(\frac{n\pi x}{L}\right) \Big|_0^L + \frac{L}{n\pi} \int_0^L \cos\left(\frac{n\pi x}{L}\right) dx \right] \\
 &= \frac{1}{L} \left[-\frac{L^2(-1)^n}{n\pi} + \frac{L^2}{n^2\pi^2} \sin\left(\frac{n\pi x}{L}\right) \Big|_0^L \right] \\
 &= \frac{L(-1)^{n+1}}{n\pi}
 \end{aligned}$$



(e) Since $f(x)$ is even, the b_n coefficients vanish.

$$\begin{aligned}
 a_0 &= \frac{1}{2L} \int_{-L/2}^{L/2} dx \\
 &= \frac{1}{2} \\
 a_n &= \frac{1}{L} \int_{-L/2}^{L/2} \cos\left(\frac{n\pi x}{L}\right) dx \\
 &= \frac{1}{n\pi} \sin\left(\frac{n\pi x}{L}\right) \Big|_{-L/2}^{L/2} \\
 &= \frac{2}{n\pi} \sin\left(\frac{n\pi}{2}\right) \\
 a_{2m-1} &= \frac{2}{(2m-1)\pi} (-1)^{m+1}
 \end{aligned}$$

where in the last line we change the index to reflect the fact that $a_n = 0$ for even n .




(f) One intuitive observation is to see that $f(x)$ is equal to

$$f(x) = \frac{1}{2} + \frac{1}{2} \text{sgn}(x)$$

which adds an odd function to a constant, rendering the a_n coefficients for $n \geq 1$ zero.

$$\begin{aligned}
 a_0 &= \frac{1}{2L} \int_0^L dx \\
 &= \frac{1}{2} \\
 b_n &= \frac{1}{L} \int_0^L \sin\left(\frac{n\pi x}{L}\right) dx \\
 &= -\frac{1}{n\pi} \cos\left(\frac{n\pi x}{L}\right) \Big|_0^L \\
 &= \frac{1 - (-1)^n}{n\pi}
 \end{aligned}$$



- (g) Here, the step function $f(x)$ differs from the previous example only by an upward vertical shift of 1. We can equivalently write


$$f(x) = \frac{3}{2} + \frac{1}{2}\text{sgn}(x)$$

which, once more, is a constant added by an odd function. Therefore the a_n coefficients remain zero, while the a_0 coefficient simply is added by 1, which we can verify by

$$a_0 = \frac{1}{2L} \left[\int_{-L}^0 dx + \int_0^L 2 dx \right] = \frac{3}{2}$$

The b_n coefficients remain unchanged. The quick way to see this is by exploiting the oddness of sine:

$$\begin{aligned} b_n &= \frac{1}{L} \left[\underbrace{\int_{-L}^0 \sin\left(\frac{n\pi x}{L}\right) dx}_{= -\int_0^L \sin\left(\frac{n\pi x}{L}\right) dx} + 2 \int_0^L \sin\left(\frac{n\pi x}{L}\right) dx \right] \\ &= \frac{1}{L} \int_0^L \sin\left(\frac{n\pi x}{L}\right) dx \end{aligned}$$



Question: 3.2.3

Show that the Fourier series operation is linear: that is, show that the Fourier series of $c_1 f(x) + c_2 g(x)$ is the sum of c_1 times the Fourier series of $f(x)$ and c_2 times the Fourier series of $g(x)$.

Suppose that f and g have Fourier series defined over $[-L, L]$ with period $2L$. The respective coefficients for any function $h(x)$ on the aforementioned interval are defined by

$$\begin{aligned} [a_0]_h &= \frac{1}{2L} \int_{-L}^L h(x) \, dx \\ [a_n]_h &= \frac{1}{L} \int_{-L}^L h(x) \cos\left(\frac{n\pi x}{L}\right) \, dx \\ [b_n]_h &= \frac{1}{L} \int_{-L}^L h(x) \sin\left(\frac{n\pi x}{L}\right) \, dx \end{aligned}$$

By linearity of integration, the coefficients for the linear combination are

$$\begin{aligned} [a_n]_{c_1 f + c_2 g} &= \frac{1}{L} \int_{-L}^L [c_1 f(x) + c_2 g(x)] \cos\left(\frac{n\pi x}{L}\right) \, dx = c_1 [a_n]_f + c_2 [a_n]_g \\ [b_n]_{c_1 f + c_2 g} &= \frac{1}{L} \int_{-L}^L [c_1 f(x) + c_2 g(x)] \sin\left(\frac{n\pi x}{L}\right) \, dx = c_1 [b_n]_f + c_2 [b_n]_g \end{aligned}$$

with an analogous calculation for $[a_0]_h$. Since the coefficients of the linear combination are the sum of c_1 times the Fourier coefficients of f and c_2 times the Fourier coefficients of g , it follows that the Fourier series itself is composed of c_1 times that of f plus c_2 times that of g , establishing that the Fourier series operation is linear.

Question: 3.2.4

Suppose that $f(x)$ is piecewise smooth. What value does the Fourier series of $f(x)$ converge to at the endpoint $x = -L$? at $x = L$?

One subtlety to be mindful of is $f(x)$ is defined over $[-L, L]$. Let the superscript $+$ and $-$ in $f(x^+)$ and $f(x^-)$ be defined as approaching x from above (the right) and below (the left). From the construction of our interval, however, it is impossible to have either $f(-L^-)$ or $f(L^+)$, for L and $-L$ are the absolute maximum and minimum values of the interval.

Fortunately, there is a workaround. Let F be the periodic extension of f . Note that there is a difference in where F and f are defined; the former is over all x , and the latter only on $[-L, L]$. By Fourier's theorem, the Fourier series of f converges to $\frac{1}{2}[F(x^+) + F(x^-)]$; by periodicity, this means that approaching the left boundary from below is akin to approaching the right boundary from below (and vice-versa for the other boundary). Equivalently,

$$F(-L^-) = f(L^-) \quad \text{and} \quad F(L^+) = f(-L^+)$$

From this logic, it turns out that the point of convergence is the same at $x = -L$ and $x = L$, which is

$$\frac{1}{2}[f(-L^+) + f(L^-)]$$

3.3 - Fourier Cosine and Sine Series

Question: 3.3.1

For the following functions, sketch $f(x)$, the Fourier series of $f(x)$, the Fourier sine series of $f(x)$, and the Fourier cosine series of $f(x)$.


(a) $f(x) = 1$



(b) $f(x) = 1 + x$



(c) $f(x) = \begin{cases} x & x < 0 \\ 1 + x & x > 0 \end{cases}$

(d) $f(x) = e^x$

(e) $f(x) = \begin{cases} 2 & x < 0 \\ e^{-x} & x > 0 \end{cases}$







Question: 3.3.2

For the following functions, sketch the Fourier sine series of $f(x)$ and determine its Fourier coefficients:

(a) $f(x) = \cos(\pi x/L)$ [Verify formula (3.3.13).]

(b) $f(x) = \begin{cases} 1 & x < L/6 \\ 3 & L/6 < x < L/2 \\ 0 & x > L/2 \end{cases}$

(c) $f(x) = \begin{cases} 0 & x < L/2 \\ x & x > L/2 \end{cases}$

(d) $f(x) = \begin{cases} 1 & x < L/2 \\ 0 & x > L/2 \end{cases}$


(a)

$$\begin{aligned}
 B_1 &= \frac{2}{L} \int_0^L \cos\left(\frac{\pi x}{L}\right) \sin\left(\frac{\pi x}{L}\right) dx \\
 &= \frac{1}{L} \int_0^L \sin\left(\frac{2\pi x}{L}\right) dx \\
 &= -\frac{1}{2\pi} \cos\left(\frac{2\pi x}{L}\right) \Big|_0^L \\
 &= 0 \\
 B_n &= \frac{2}{L} \int_0^L \cos\left(\frac{\pi x}{L}\right) \sin\left(\frac{n\pi x}{L}\right) dx \\
 &= \frac{1}{L} \int_0^L \left[\sin\left(\frac{(n+1)\pi x}{L}\right) + \sin\left(\frac{(n-1)\pi x}{L}\right) \right] dx \\
 &= \frac{1}{\pi} (1 - (-1)^{n+1}) \left[\frac{1}{n+1} + \frac{1}{n-1} \right] \\
 &= \frac{2n(1 + (-1)^n)}{\pi(n^2 - 1)}, \quad n \neq 1
 \end{aligned}$$

The B_n coefficients split into even and odd cases:


$$B_n = \begin{cases} 0 & n \text{ odd} \\ \frac{4n}{\pi(n^2 - 1)} & n \text{ even} \end{cases}$$

verifying equation (3.3.13).




(b)

$$\begin{aligned}
 B_n &= \frac{2}{L} \left[\int_0^{L/6} \sin\left(\frac{n\pi x}{L}\right) dx + 3 \int_{L/6}^{L/2} \sin\left(\frac{n\pi x}{L}\right) dx \right] \\
 &= \frac{2}{L} \cdot \frac{L}{n\pi} \left[-\cos\left(\frac{n\pi x}{L}\right) \Big|_0^{L/6} - 3 \cos\left(\frac{n\pi x}{L}\right) \Big|_{L/6}^{L/2} \right] \\
 &= \frac{2}{n\pi} \left[\left(1 - \cos\left(\frac{n\pi}{6}\right)\right) + 3 \left(\cos\left(\frac{n\pi}{6}\right) - \cos\left(\frac{n\pi}{2}\right)\right) \right] \\
 &= \frac{2}{n\pi} \left[1 + 2 \cos\left(\frac{n\pi}{6}\right) - 3 \cos\left(\frac{n\pi}{2}\right) \right]
 \end{aligned}$$



(c)


$$\begin{aligned}
B_n &= \frac{2}{L} \int_{L/2}^L x \sin\left(\frac{n\pi x}{L}\right) dx \\
&= \frac{2}{L} \left[-\frac{L}{n\pi} x \cos\left(\frac{n\pi x}{L}\right) \Big|_{L/2}^L + \frac{L}{n\pi} \int_{L/2}^L \cos\left(\frac{n\pi x}{L}\right) dx \right] \\
&= \frac{2}{L} \left[-\frac{L}{n\pi} \left(L(-1)^n - \frac{L}{2} \cos\left(\frac{n\pi}{2}\right) \right) + \frac{L^2}{n^2\pi^2} \sin\left(\frac{n\pi x}{L}\right) \Big|_{L/2}^L \right] \\
&= \frac{2L}{n\pi} \left[\frac{1}{2} \cos\left(\frac{n\pi}{2}\right) - (-1)^n \right] - \frac{2L}{n^2\pi^2} \sin\left(\frac{n\pi}{2}\right)
\end{aligned}$$



(d)

$$\begin{aligned}
B_n &= \frac{2}{L} \int_0^{L/2} \sin\left(\frac{n\pi x}{L}\right) dx \\
&= -\frac{2}{n\pi} \cos\left(\frac{n\pi x}{L}\right) \Big|_0^{L/2} \\
&= \frac{2}{n\pi} \left[1 - \cos\left(\frac{n\pi}{2}\right) \right] \\
&= \begin{cases} \frac{2}{n\pi} & n \text{ odd} \\ \frac{1}{m\pi} (1 - (-1)^m) & n = 2m \end{cases}
\end{aligned}$$

where we substitute $n = 2m$.




Question: 3.3.3



For the following functions, sketch the Fourier sine series of $f(x)$. Also, roughly sketch the sum of a *finite* number of nonzero terms (at least the first two) of the Fourier sine series:

(a) $f(x) = \cos(\pi x/L)$ [Use formula (3.3.13).]

(b) $f(x) = \begin{cases} 1 & x < L/2 \\ 0 & x > L/2 \end{cases}$


(c) $f(x) = x$ [Use formula (3.3.12).]





Question: 3.3.4

Sketch the Fourier cosine series of $f(x) = \sin(\pi x/L)$. Briefly discuss.



The Fourier cosine series looks exactly like $|\sin(\pi x/L)|$, having period L . If we were to actually derive the Fourier series (and thus the even extension of $f(x)$), we would find

$$\begin{aligned}
 A_0 &= \frac{1}{L} \int_0^L \sin\left(\frac{\pi x}{L}\right) dx \\
 &= -\frac{1}{\pi} \cos\left(\frac{\pi x}{L}\right) \Big|_0^L \\
 &= \frac{2}{\pi} \\
 A_1 &= \frac{2}{L} \int_0^L \sin\left(\frac{\pi x}{L}\right) \cos\left(\frac{\pi x}{L}\right) dx \\
 &= \frac{1}{L} \int_0^L \sin\left(\frac{2\pi x}{L}\right) dx \\
 &= -\frac{1}{2\pi} \cos\left(\frac{2\pi x}{L}\right) \Big|_0^L \\
 &= 0 \\
 A_n &= \frac{2}{L} \int_0^L \sin\left(\frac{\pi x}{L}\right) \cos\left(\frac{n\pi x}{L}\right) dx \\
 &= \frac{1}{L} \int_0^L \left[\sin\left(\frac{(n+1)\pi x}{L}\right) - \sin\left(\frac{(n-1)\pi x}{L}\right) \right] dx \\
 &= \frac{1}{\pi} (1 + (-1)^n) \left[\frac{1}{n+1} - \frac{1}{n-1} \right] \\
 &= -\frac{2(1 + (-1)^n)}{\pi(n^2 - 1)}, \quad n \neq 1
 \end{aligned}$$

for the A_n , let $n = 2m$ and reindex as

$$A_m = -\frac{4}{\pi(4m^2 - 1)}$$

which combine to produce

$$f(x) \sim \frac{2}{\pi} - \frac{4}{\pi} \sum_{m=1}^{\infty} \frac{\cos(2m\pi x/L)}{4m^2 - 1}$$

Observe the pattern: *the Fourier cosine series consists only of even terms.* Moreover, since $\sin(\pi x/L) = 0$ at $x = 0$ and $x = L$, the Fourier series converges to $f(x)$ for *all* x over $[0, L]$, not only $(0, L)$. Crucially, **there is no Gibbs overshoot at the boundary**, as we have no jump discontinuity. **This is a crucial point and one I would advise the reader to internalize!**

Question: 3.3.5

For the following functions, sketch the Fourier cosine series of $f(x)$ and determine its Fourier coefficients:


(a) $f(x) = x^2$

(b) $f(x) = \begin{cases} 1 & x < L/6 \\ 3 & L/6 < x < L/2 \\ 0 & x > L/2 \end{cases}$

(c) $f(x) = \begin{cases} 0 & x < L/2 \\ x & x > L/2 \end{cases}$


(a)

$$\begin{aligned} A_0 &= \frac{1}{L} \int_0^L x^2 dx \\ &= \frac{L^2}{3} \\ A_n &= \frac{2}{L} \int_0^L x^2 \cos\left(\frac{n\pi x}{L}\right) dx \\ &= \frac{2}{L} \left[\frac{L}{n\pi} x^2 \sin\left(\frac{n\pi x}{L}\right) \Big|_0^L - \frac{2L}{n\pi} \int_0^L x \sin\left(\frac{n\pi x}{L}\right) dx \right] \\ &= -\frac{4}{n\pi} \left[-\frac{L}{n\pi} x \cos\left(\frac{n\pi x}{L}\right) \Big|_0^L + \frac{L}{n\pi} \int_0^L \cos\left(\frac{n\pi x}{L}\right) dx \right] \\ &= -\frac{4}{n\pi} \left[-\frac{L^2(-1)^n}{n\pi} + 0 \right] \\ &= \frac{4L^2(-1)^n}{n^2\pi^2} \end{aligned}$$




(b)

$$\begin{aligned}
 A_0 &= \frac{1}{L} \left[\int_0^{L/6} dx + \int_{L/6}^{L/2} 3 dx \right] \\
 &= \frac{1}{L} \left(\frac{L}{6} + L \right) \\
 &= \frac{7}{6} \\
 A_n &= \frac{2}{L} \left[\int_0^{L/6} \cos\left(\frac{n\pi x}{L}\right) dx + 3 \int_{L/6}^{L/2} \cos\left(\frac{n\pi x}{L}\right) dx \right] \\
 &= \frac{2}{n\pi} \left[\sin\left(\frac{n\pi x}{L}\right) \Big|_0^{L/6} + 3 \sin\left(\frac{n\pi x}{L}\right) \Big|_{L/6}^{L/2} \right] \\
 &= \frac{2}{n\pi} \left[3 \sin\left(\frac{n\pi}{2}\right) - 2 \sin\left(\frac{n\pi}{6}\right) \right]
 \end{aligned}$$



(c)

$$\begin{aligned}
 A_0 &= \frac{1}{L} \int_{L/2}^L x \, dx \\
 &= \frac{1}{L} \left(\frac{L^2}{2} - \frac{L^2}{8} \right) \\
 &= \frac{3L}{8} \\
 A_n &= \frac{2}{L} \int_{L/2}^L x \cos\left(\frac{n\pi x}{L}\right) \, dx \\
 &= \frac{2}{L} \left[\frac{L}{n\pi} x \sin\left(\frac{n\pi x}{L}\right) \Big|_{L/2}^L - \frac{L}{n\pi} \int_{L/2}^L \sin\left(\frac{n\pi x}{L}\right) \, dx \right] \\
 &= \frac{2}{L} \left[-\frac{L^2}{2n\pi} \sin\left(\frac{n\pi}{2}\right) + \frac{L^2}{n^2\pi^2} \cos\left(\frac{n\pi x}{L}\right) \Big|_{L/2}^L \right] \\
 &= -\frac{L}{n\pi} \sin\left(\frac{n\pi}{2}\right) + \frac{2L}{n^2\pi^2} \left[(-1)^n - \cos\left(\frac{n\pi}{2}\right) \right]
 \end{aligned}$$



Question: 3.3.6

For the following functions, sketch the Fourier cosine series of $f(x)$. Also, roughly sketch the sum of a finite number of nonzero terms (at least the first two) of the Fourier cosine series:

(a) $f(x) = x$ [Use formulas (3.3.22) and (3.3.23)].

(b) $f(x) = \begin{cases} 0 & x < L/2 \\ 1 & x > L/2 \end{cases}$ [Use carefully formulas (3.2.6) and (3.2.7).]

(c) $f(x) = \begin{cases} 1 & x < L/2 \\ 0 & x > L/2 \end{cases}$ [Hint: Add the functions in parts (b) and (c).]


Note that there is a typographical error in the text; what I have written here is the correct problem statement.

(a)

$$\begin{aligned}
 A_0 &= \frac{1}{L} \int_0^L x \, dx \\
 &= \frac{1}{L} \left(\frac{L^2}{2} \right) \\
 &= \frac{L}{2} \\
 A_n &= \frac{2}{L} \int_0^L x \cos\left(\frac{n\pi x}{L}\right) \, dx \\
 &= \frac{2}{L} \frac{L^2}{n^2 \pi^2} \left[\cos\left(\frac{n\pi x}{L}\right) + \frac{n\pi x}{L} \sin\left(\frac{n\pi x}{L}\right) \right]_0^L \\
 &= \frac{2L}{(n\pi)^2} [\cos(n\pi) - 1]
 \end{aligned}$$


Since $\cos(n\pi) - 1$ is 0 for even n and -2 for odd n , the even coefficients vanish and, writing $n = 2m - 1$,

$$A_{2m-1} = -\frac{4L}{\pi^2(2m-1)^2}$$



(b)


$$\begin{aligned}
 A_0 &= \frac{1}{L} \int_{L/2}^L dx \\
 &= \frac{1}{2} \\
 A_n &= \frac{2}{L} \int_{L/2}^L \cos\left(\frac{n\pi x}{L}\right) dx \\
 &= \frac{2}{n\pi} \sin\left(\frac{n\pi x}{L}\right) \Big|_{L/2}^L \\
 &= -\frac{2}{n\pi} \sin\left(\frac{n\pi}{2}\right)
 \end{aligned}$$



(c)

$$\begin{aligned}
 A_0 &= \frac{1}{L} \int_0^{L/2} dx \\
 &= \frac{1}{2} \\
 A_n &= \frac{2}{L} \int_0^{L/2} \cos\left(\frac{n\pi x}{L}\right) dx \\
 &= \frac{2}{n\pi} \sin\left(\frac{n\pi x}{L}\right) \Big|_0^{L/2} \\
 &= \frac{2}{n\pi} \sin\left(\frac{n\pi}{2}\right)
 \end{aligned}$$

Observe that adding the two Fourier series cancels the A_n terms from each, leaving only the A_0 terms: $1/2 + 1/2 = 1$.

**Question: 3.3.7**

Show that e^x is the sum of an even and an odd function.

$$e^x = \underbrace{\frac{1}{2}(e^x + e^{-x})}_{f_{\text{even}}} + \underbrace{\frac{1}{2}(e^x - e^{-x})}_{f_{\text{odd}}} = \cosh(x) + \sinh(x)$$

Observe that

$$f_{\text{even}}(x) = \frac{1}{2}(e^x + e^{-x}) = \frac{1}{2}(e^{-x} + e^x) = f_{\text{even}}(-x)$$

and

$$f_{\text{odd}}(x) = \frac{1}{2}(e^x - e^{-x}) = -\frac{1}{2}(e^{-x} - e^x) = -f_{\text{odd}}(-x)$$

Question: 3.3.8

- (a) Determine formulas for the even extension of any $f(x)$. Compare to the formula for the even part of $f(x)$.
- (b) Do the same for the odd extension of $f(x)$ and the odd part of $f(x)$.
- (c) Calculate and sketch the four functions of parts (a) and (b) if

$$f(x) = \begin{cases} x & x > 0 \\ x^2 & x < 0 \end{cases}$$

Graphically add the even and odd parts of $f(x)$. What occurs? Similarly, add the even and odd extensions. What occurs then?

- (a) The even extension of $f(x)$ is given by

$$f_{\text{even}}(x) = \begin{cases} f(x) & x > 0 \\ f(-x) & x < 0 \end{cases}$$

As Haberman is emphatic about, do **not** confuse the even *part* of $f(x)$ with its *extension*. They are equal, however, when $f(x)$ is itself an even function. The part and extension differ in their construction: the extension performs a mirror reflection of $f(x)$ about the vertical axis, while the part takes an arithmetic mean of $f(x)$ evaluated at *both* positive and negative x .

- (b) The odd extension of $f(x)$ is given by

$$f_{\text{odd}}(x) = \begin{cases} f(x) & x > 0 \\ -f(-x) & x < 0 \end{cases}$$

Similarly, when $f(x)$ is itself odd, the extension and part are equal. In their construction, the extension is a reflection through the origin (i.e. reflecting across the vertical axis *and then* the horizontal axis), while the part is an arithmetic mean of $f(x)$ and $-f(-x)$.

- (c) First, for clarity in evaluating $f(-x)$, keep the following in mind: define $y = -x$. When $x < 0$, then $y > 0$. When that is the case, $f(y) = y = -x$. Alternatively, when $x > 0$, $y < 0$, and $f(y) = y^2 = (-x)^2 = x^2$.

It is critical to not simply substitute in a negative sign in each branch. Instead, the sign of $-x$ directs us to which case of $f(x)$ we should be on.

$$f(x) = \begin{cases} x & x > 0 \\ x^2 & x < 0 \end{cases}, \quad f(-x) = \begin{cases} x^2 & x > 0 \\ -x & x < 0 \end{cases}, \quad -f(-x) = \begin{cases} -x^2 & x > 0 \\ x & x < 0 \end{cases}$$

Then we derive

$$f_{\text{even part}}(x) = \frac{1}{2}(f(x) + f(-x)) = \begin{cases} \frac{1}{2}(x + x^2) & x > 0 \\ \frac{1}{2}(-x + x^2) & x < 0 \end{cases}$$

$$f_{\text{odd part}}(x) = \frac{1}{2}(f(x) - f(-x)) = \begin{cases} \frac{1}{2}(x - x^2) & x > 0 \\ \frac{1}{2}(x + x^2) & x < 0 \end{cases}$$



$$f_{\text{even ext.}}(x) = \begin{cases} x & x > 0 \\ -x & x < 0 \end{cases} = |x|$$

$$f_{\text{odd ext.}}(x) = \begin{cases} x & x > 0 \\ x & x < 0 \end{cases} = x$$

Adding the parts, we simply get back $f(x)$, as expected. Adding the extensions produces

$$f_{\text{even ext.}}(x) + f_{\text{odd ext.}}(x) = \begin{cases} 2x & x > 0 \\ 0 & x < 0 \end{cases}$$

which we expect, as both extensions will agree on the $x > 0$ portion (in a sense, they double each other in the sum), but cancel each other out on the $x < 0$ portion, being mirror images of one another about the horizontal axis. See the graph below for all four equations and the sums of the parts and extensions graphed out alongside $f(x)$.



Question: 3.3.9

What is the sum of the Fourier sine series of $f(x)$ and the Fourier cosine series of $f(x)$? [What is the sum of the even and odd extensions of $f(x)$?]

Akin to our solution from 3.3.8(c), the sum of the Fourier sine and cosine series will converge to

$$g(x) = \begin{cases} 2f(x) & 0 < x < L \\ 0 & -L < x < 0 \end{cases}$$

and extended with period $2L$. This parallels the $x < 0$ segment vanishing in the previous problem. From the convergence theorem, we know that at the endpoints, the series converges to the arithmetic mean of $g(x^+)$ and $g(x^-)$.

Question: 3.3.10

If $f(x) = \begin{cases} x^2 & x < 0 \\ e^{-x} & x > 0 \end{cases}$, what are the even and odd parts of $f(x)$?

When $x < 0$, then $y = -x > 0$, and $f(y) = e^y = e^x$. Additionally, when $x > 0$, $y = -x < 0$, and $f(y) = y^2 = (-x)^2 = x^2$. With these insights we can conclude

$$f(-x) = \begin{cases} e^x & x < 0 \\ x^2 & x > 0 \end{cases}, \quad -f(-x) = \begin{cases} -e^x & x < 0 \\ -x^2 & x > 0 \end{cases}$$

Then we can compute

$$\begin{aligned} f_{\text{even}}(x) &= \begin{cases} \frac{1}{2}(x^2 + e^x) & x < 0 \\ \frac{1}{2}(x^2 + e^{-x}) & x > 0 \end{cases} = \frac{1}{2}(x^2 + e^{-|x|}) \\ f_{\text{odd}}(x) &= \begin{cases} \frac{1}{2}(x^2 - e^x) & x < 0 \\ \frac{1}{2}(-x^2 + e^{-x}) & x > 0 \end{cases} = \frac{1}{2}\text{sgn}(x)(e^{-|x|} - x^2) \end{aligned}$$

As we anticipate, $f_{\text{even}} + f_{\text{odd}} = f$.

Question: 3.3.11

Given a sketch of $f(x)$, describe a procedure to sketch the even and odd parts of $f(x)$.

For the even part:


- (1) Draw the mirror image of $f(x)$ about the vertical axis.
- (2) Draw the even part, which is the vertical average of the original function and its mirror.

For the odd part:

- (1) Rotate the graph of $f(x)$ by 180° ; that is what the mirror image of $f(x)$ looks like on the original plot.
- (2) Draw the odd part, which is the vertical average of the original function and its mirror.

Question: 3.3.12

- (a) Graphically show that the even terms (n even) of the Fourier sine series of any function on $0 \leq x \leq L$ are odd (antisymmetric) around $x = L/2$.
- (b) Consider a function $f(x)$ that is odd around $x = L/2$. Show that the odd coefficients (n odd) of the Fourier sine series of $f(x)$ on $0 \leq x \leq L$ are zero.



- (b) The coefficients are given by

$$B_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

By premise, over $[0, L]$, $f(x)$ is odd about $x = L/2$. We wish to establish that for n odd, $\sin(n\pi x/L)$ is even about $L/2$, as that would mean the integrand $f(x) \sin(n\pi x/L)$ is odd about $L/2$ and vanishes when integrated over the symmetric interval $[0, L]$. Equivalently, we can demonstrate that

$$g(x) = \sin\left[\frac{n\pi}{L}\left(x + \frac{L}{2}\right)\right]$$

is even about the origin. Let $n = 2m + 1$, $m = 0, 1, \dots$ and derive

$$\begin{aligned} g(x) &= \sin\left[\frac{(2m+1)\pi}{L}\left(x + \frac{L}{2}\right)\right] = \sin\left[\frac{(2m+1)\pi x}{L} + \frac{(2m+1)\pi}{2}\right] \\ &= (-1)^m \cos\left[\frac{(2m+1)\pi x}{L}\right] \end{aligned}$$

Since cosine is even about the origin, we must also have $g(x)$ is even, proving the result.

Question: 3.3.13

Consider a function $f(x)$ that is even around $x = L/2$. Show that the even coefficients (n even) of the Fourier sine series of $f(x)$ on $0 \leq x \leq L$ are zero.

The coefficients are given by

$$B_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

By premise, over $[0, L]$, $f(x)$ is even about $x = L/2$. The objective is to demonstrate that when n is even, $f(x) \sin(n\pi x/L)$ is odd about $x = L/2$. Equivalently, we need only prove that $\sin(n\pi x/L)$ is odd about that point (since an even times an odd function is odd), and an even easier way to frame this is to show that

$$g(x) = \sin\left[\frac{n\pi}{L}\left(x + \frac{L}{2}\right)\right]$$

is odd about the origin. Let $n = 2m$ and rewrite as

$$g(x) = \sin\left[\frac{2m\pi}{L}\left(x + \frac{L}{2}\right)\right] = \sin\left(\frac{2m\pi x}{L} + m\pi\right) = (-1)^m \sin\left(\frac{2m\pi x}{L}\right)$$

Immediately we can deduce

$$g(-x) = (-1)^{m+1} \sin\left(\frac{2m\pi x}{L}\right) \implies -g(-x) = (-1)^m \sin\left(\frac{2m\pi x}{L}\right) = g(x)$$

Proving that $g(x)$ is odd. Given that the integrand is established to be odd about $x = L/2$ for even n , by symmetry of $[0, L]$ about $L/2$, we have proven that $B_n = 0$.

Question: 3.3.14

- (a) Consider a function $f(x)$ that is even around $x = L/2$. Show that the odd coefficients (n odd) of the Fourier cosine series of $f(x)$ on $0 \leq x \leq L$ are zero.
- (b) Explain the result of part (a) by considering a Fourier cosine series of $f(x)$ on the interval $0 \leq x \leq L/2$.

- (a) The coefficients are given by

$$A_n = \frac{2}{L} \int_0^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx$$

By premise, over $[0, L]$, $f(x)$ is even about $x = L/2$. We wish to prove that when n is odd, $f(x) \cos(n\pi x/L)$ is odd about $L/2$. Put differently, we must prove that $\cos(n\pi x/L)$ is odd about that point, or equivalently

$$g(x) = \cos\left[\frac{n\pi}{L}\left(x + \frac{L}{2}\right)\right]$$

is odd about the origin. Let $n = 2m + 1$, $m = 0, 1, \dots$ and deduce

$$\begin{aligned} g(x) &= \cos\left[\frac{(2m+1)\pi}{L}\left(x + \frac{L}{2}\right)\right] = \cos\left[\frac{(2m+1)\pi x}{L} + \frac{(2m+1)\pi}{2}\right] \\ &= (-1)^{m+1} \sin\left[\frac{(2m+1)\pi x}{L}\right] \end{aligned}$$

and since sine is odd, we must also have $g(x)$ is odd, establishing the result.

- (b) It is good to motivate some geometric intuition: since $f(x)$ is even about $L/2$, we know that the segments on $[0, L/2]$ and $[L/2, L]$ are mirror images of each other around $x = L/2$. If we replace the interval with $[0, L/2]$, our coefficients are written as

$$A_n = \frac{4}{L} \int_0^{L/2} f(x) \cos\left(\frac{2n\pi x}{L}\right) dx$$

In particular, observe that the basis functions $\cos(2n\pi x/L)$ are only the even modes. The odd modes are nowhere to be found, which comports with our result from part (a). The Fourier cosine series built from these coefficients converges to $f(x)$ on $[0, L/2]$. But we get two for the price of one: convince yourself that when we horizontally shift the cosine to the left by $L/2$, we can see that

$$g(x) = \cos\left[\frac{2n\pi}{L}\left(x + \frac{L}{2}\right)\right] = \cos\left(\frac{2n\pi x}{L} + n\pi\right) = (-1)^n \cos\left(\frac{2n\pi x}{L}\right)$$

is even about the origin; by the translation, we can immediately deduce that $\cos(2n\pi x/L)$ is even about $L/2$. Since both $f(x)$ and the cosine basis functions are even about that point, A_n is exactly the same over $[0, L/2]$ and $[L/2, L]$, and by even symmetry, $\int_0^L = 2 \int_0^{L/2}$. This establishes

$$\begin{aligned} \frac{2}{L} \int_0^L f(x) \cos\left(\frac{2n\pi x}{L}\right) dx &= \frac{4}{L} \int_0^{L/2} f(x) \cos\left(\frac{2n\pi x}{L}\right) dx \\ &= \frac{4}{L} \int_{L/2}^L f(x) \cos\left(\frac{2n\pi x}{L}\right) dx \end{aligned}$$

where the left side includes only the even modes that match term-by-term with A_n , proving that the odd modes must vanish as we established in part (a).

Question: 3.3.15

Consider a function $f(x)$ that is odd around $x = L/2$. Show that the even coefficients (n even) of the Fourier cosine series of $f(x)$ on $0 \leq x \leq L$ are zero.

The coefficients are given by

$$A_n = \frac{2}{L} \int_0^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx$$

Our objective is to prove that $f(x) \cos(n\pi x/L)$ is odd about $L/2$. By premise, $f(x)$ is odd around that point, so we need only prove that $\cos(n\pi x/L)$ is even about $L/2$. Equivalently, prove

$$g(x) = \cos\left[\frac{n\pi}{L}\left(x + \frac{L}{2}\right)\right]$$

is even about the origin. Let $n = 2m$ and deduce

$$g(x) = \cos\left(\frac{2m\pi x}{L} + m\pi\right) = (-1)^m \cos\left(\frac{2m\pi x}{L}\right)$$

Since cosine is even about the origin, so is $g(x)$. Then $f(x) \cos(n\pi x/L)$ is odd about $L/2$, and by symmetry of $[0, L]$ around that point, A_n vanishes for all even n .

Question: 3.3.16

Fourier series can be defined on other intervals besides $-L \leq x \leq L$. Suppose that $g(y)$ is defined for $a \leq y \leq b$. Represent $g(y)$ using periodic trigonometric functions with period $b - a$. Determine formulas for coefficients. [Hint: Use the linear transformation

$$y = \frac{a+b}{2} + \frac{b-a}{2L}x.]$$

First, let us experiment with the book's hint for y . Observe that $y(L) = b$ and $y(-L) = a$. Structurally, it is a sum of the average of the y -endpoints and a scaling term transforming between the $2L$ and $b - a$ periods.

Expressing in terms of x , do some algebraic manipulation to derive

$$x = \left(\frac{a+b}{b-a}\right)L + \frac{2L}{b-a}y$$

where the differentials are related by

$$dx = \frac{2L}{b-a}dy$$

Substituting into our full Fourier series, we have

$$\begin{aligned} g(y) &\sim A_0 + \sum_{n=1}^{\infty} A_n \cos\left[n\pi\left(\frac{a+b}{b-a} + \frac{2}{b-a}y\right)\right] + \sum_{n=1}^{\infty} B_n \sin\left[n\pi\left(\frac{a+b}{b-a} + \frac{2}{b-a}y\right)\right] \\ &\sim A_0 + \sum_{n=1}^{\infty} A_n \cos\left[\frac{2n\pi}{b-a}\left(y - \frac{a+b}{2}\right)\right] + \sum_{n=1}^{\infty} B_n \sin\left[\frac{2n\pi}{b-a}\left(y - \frac{a+b}{2}\right)\right] \end{aligned}$$

where the coefficients are

$$\begin{aligned} A_0 &= \frac{1}{b-a} \int_a^b g(y) dy \\ A_n &= \frac{2}{b-a} \int_a^b g(y) \cos\left[\frac{2n\pi}{b-a}\left(y - \frac{a+b}{2}\right)\right] dy \\ B_n &= \frac{2}{b-a} \int_a^b g(y) \sin\left[\frac{2n\pi}{b-a}\left(y - \frac{a+b}{2}\right)\right] dy \end{aligned}$$

But since the coefficients and basis functions have the same phase shift, we can effectively get rid of the $-(a+b)/2$ terms everywhere. All that matters is that the integrals in the coefficients cycle through the entire period over $[a, b]$.

Question: 3.3.17

Consider

$$\int_0^1 \frac{dx}{1+x^2}$$

- (a) Evaluate explicitly.
- (b) Use the Taylor series of $1/(1+x^2)$ (itself a geometric series) to obtain an infinite series for the integral.
- (c) Equate part (a) to part (b) in order to derive a formula for π .

- (a) Let $x = \tan(u)$. Then $dx = \sec^2(u) du$. The bounds change to $[0, \pi/4]$. Substituting, we get

$$\int_0^{\pi/4} \frac{\sec^2(u)}{1+\tan^2(u)} du$$

Using the fact that $1 + \tan^2(u) = \sec^2(u)$, we have

$$\int_0^{\pi/4} du = \frac{\pi}{4}$$

- (b) The Taylor series is

$$\frac{1}{1+x^2} = \sum_{n=0}^{\infty} (-1)^n x^{2n}$$

Note that a crucial point that Haberman omits is that this sum is valid only for $|x| < 1$. We will handwave our way through this for now. Then the integral can be rewritten as

$$\begin{aligned} \int_0^1 \sum_{n=0}^{\infty} (-1)^n x^{2n} dx &= \int_0^1 (1 - x^2 + x^4 - \dots) dx = \left(x - \frac{x^3}{3} + \frac{x^5}{5} - \dots \right) \Big|_0^1 \\ &= \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{2n-1} \end{aligned}$$

- (c) Combining the two results, derive

$$\pi = 4 \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{2n-1}$$

Question: 3.3.18

For continuous functions,

- (a) Under what conditions does $f(x)$ equal its Fourier series for all x , $-L \leq x \leq L$?
- (b) Under what conditions does $f(x)$ equal its Fourier sine series for all x , $0 \leq x \leq L$?
- (c) Under what conditions does $f(x)$ equal its Fourier cosine series for all x , $0 \leq x \leq L$?

In all cases, $f(x)$ must be continuous and piecewise smooth.

- (a) $f(-L) = f(L)$
- (b) $f(0) = f(L) = 0$
- (c) No extra condition required.

3.4 - Term-by-Term Differentiation of Fourier Series

Question: 3.4.1

The integration-by-parts formula

$$\int_a^b u \frac{dv}{dx} dx = uv \Big|_a^b - \int_a^b v \frac{du}{dx} dx$$

is known to be valid for functions $u(x)$ and $v(x)$, which are continuous and have continuous first derivatives. However, we will assume that u , v , du/dx , and dv/dx are continuous only for $a \leq x \leq c$ and $c \leq x \leq b$; we assume that all quantities may have a jump discontinuity at $x = c$.

- Derive an expression for $\int_a^b u dv/dx dx$ in terms of $\int_a^b v du/dx dx$.
- Show that this reduces to the integration-by-parts formula if u and v are continuous across $x = c$. It is *not* necessary for du/dx and dv/dx to be continuous at $x = c$.

- Split the interval $[a, b]$ as partitioned by c and evaluate over both sub-intervals:

$$\int_a^c u \frac{dv}{dx} dx + \int_c^b u \frac{dv}{dx} dx$$

Since u and v and their first derivatives are continuous within each sub-interval, we can apply the integration-by-parts formula normally in each:

$$\begin{aligned} \int_a^c u \frac{dv}{dx} dx &= uv \Big|_a^{c^-} - \int_a^c v \frac{du}{dx} dx \\ \int_c^b u \frac{dv}{dx} dx &= uv \Big|_{c^+}^b - \int_c^b v \frac{du}{dx} dx \end{aligned}$$

Now add these expressions to find

$$\begin{aligned} \int_a^c u \frac{dv}{dx} dx + \int_c^b u \frac{dv}{dx} dx &= \left(uv \Big|_a^{c^-} + uv \Big|_{c^+}^b \right) - \left(\int_a^c v \frac{du}{dx} dx + \int_c^b v \frac{du}{dx} dx \right) \\ &= uv \Big|_a^b + uv \Big|_{c^+}^{c^-} - \left(\int_a^c v \frac{du}{dx} dx + \int_c^b v \frac{du}{dx} dx \right) \end{aligned}$$

As for the $\int v du/dx dx$ terms, note that the discontinuity of du/dx does *not* affect the evaluation of the definite integral at all. In geometric terms, we evaluate the area underneath $v du/dx$, which is not a function of the exact value of $\lim_{x \rightarrow c^-} v du/dx$ or $\lim_{x \rightarrow c^+} v du/dx$. Put differently, the value of the integrand at a single point does not change the definite integral. Thus we have

$$\int_a^b v \frac{du}{dx} dx = \int_a^c v \frac{du}{dx} dx + \int_c^b v \frac{du}{dx} dx$$

Leaving us with the final expression

$$uv \Big|_a^b + uv \Big|_{c^+}^{c^-} - \int_a^b v \frac{du}{dx} dx$$

- (b) With a potential discontinuity, it is not generally true that $u(c^-)v(c^-) = u(c^+)v(c^+)$. But with the assumption of continuity of u and v at $x = c$, then equality holds, and the uv terms collapse into pointwise evaluation at the endpoints $[a, b]$ as before:

$$\int_a^b u \frac{dv}{dx} dx = uv \Big|_a^b + \underbrace{uv \Big|_{c^+}^{c^-}}_{=0} - \int_a^b v \frac{du}{dx} dx = uv \Big|_a^b - \int_a^b v \frac{du}{dx} dx$$

All throughout, never once did we require continuity of the derivatives.

Question: 3.4.2

Suppose that $f(x)$ and df/dx are piecewise smooth. Prove that the Fourier series of $f(x)$ can be differentiated term by term if the Fourier series of $f(x)$ is continuous.

Suppose that the Fourier series of $f(x)$ is continuous. Since the Fourier series has period $2L$, we will have $f(L) = f(-L)$ on each period.