

Module 4A


Supervised Machine Learning

Different flavors of REGRESSION and General Linear Models


Review: Correlation:

- Measures the amount/degree of linear association between two **numerical** variables
- Estimate the degree to which variables **covary**
 - With no attempt to interpret the causality of the association

Example: arm length and leg length covary together (individuals with longer arms often have longer legs) but they are influenced by other underlying variables **not** each other (longer legs do not cause longer arms)



Scientists be like




Regression:

- Statistics is about prediction
- Used to **predict** value of one numerical variable from the value of another
 - predicting dependent/response variable, Y from independent/predictor X
- Linear regression assumes that the relationship between X and Y can be described by a line
 - Fits a straight line to a (messy) scatterplot

Example: ambient temperature may impact growth rate of a plant species, but the reverse is probably not true

A common point of confusion: Pearson's correlation is the slope of the best-fit line **after standardizing both** variables. The regression slope is the **unstandardized** version that tells you how much Y changes per unit X.




- Other kinds:
 - Lasso
 - Variable selection (weighting a predictor variable by 0)
 - Ridge
 - Allows analysis even in the face of **collinearity**


Regression:

- Linear regression assumes that the relationship between X and Y can be described by a line
 - Fits a straight line to a (messy) scatterplot
- Homoscedasticity: Y is normally distributed with equal variance for all values of X


Example: ambient temperature may impact growth rate of a plant species, but the reverse is probably not true




correlation vs regression



$r = 1; Y = 3 + 0.8X$



$r = 0.6; Y = 3 + 0.8X$



$r = 0.6; Y = 3 + 2X$

←
Different correlation;
same slope

→
Same correlation;
different slope

The parameters of linear regression





$$Y = \alpha + \beta X + \varepsilon_1$$

intercept

Slope*

* Very similar to Pearson's correlation (but normalized only wrt to X)

Regression Overview



Estimating a regression line

$$Y = a + bX + \varepsilon_1$$

Quick Review:

$$Z = \frac{\bar{Y} - \mu}{\sigma_{\bar{Y}}} = \frac{Y - \mu}{\sigma / \sqrt{n}}$$


$$t = \frac{\bar{Y} - \mu}{SE_{\bar{Y}}} = \frac{\bar{Y} - \mu}{s / \sqrt{n}}$$

$$\text{F-value} = \frac{\text{MS}_{\text{group}}}{\text{MS}_{\text{error}}} \quad \begin{array}{l} \text{SIGNAL} \\ \text{NOISE} \end{array}$$

$$r = \frac{\sum (X - \bar{X})(Y - \bar{Y})}{\sqrt{\sum (X - \bar{X})^2} \sqrt{\sum (Y - \bar{Y})^2}} = \frac{\text{Covariance}(X, Y)}{s_x s_y}$$

(Ordinary) Least Squares:

- Best fitting line through a scatterplot
 - Line that minimized spread of y values
- Minimize $SS_{\text{residuals}}$
 - Measurement of how much the line's predicted y_i deviate from actual data values






Figure 20.1: Whitlock and Schluter, Fig 15.1.2 – Illustrating the partitioning of sum of squares into MS_{group} and MS_{error} components.


Regression Overview

Least Squares:

- What are the elements of this equation?

$$SS_{residual} = \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

$$\hat{y}_i = a + bx_i$$



- Residuals:


- Residuals measure the scatter of points above and below the least squares regression line
- MS_{residual} is the variance of the residuals, $residual = Y_i - \hat{Y}_i$

$$MS_{\text{residual}} = \frac{\sum (Y_i - \hat{Y}_i)^2}{n - 2}$$

- $MS_{\text{regression}}$ is the variance of the regression

$$MS_{\text{regression}} = \frac{\sum (\hat{Y}_i - \bar{Y})^2}{n - 2}$$

- Coefficient of determination (r^2) = SSR/SST






R^2 predicts the amount of variance in Y explained by the regression line

We saw this in ANOVA where R^2 gave 'precision' of model (i.e. Ability of the model to explain variation)




- The coefficient of determination
- Sometimes written as r^2
- Square of the correlation coefficient, r

$$R^2 = \frac{SS_{regression}}{SS_{Total}}$$

Y



Y



X

X

X

Best estimate of slope:

b = Sum of cross products


Sum of squares of X

$$b = \frac{\sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y})}{\sum_{i=1}^n (X_i - \bar{X})^2}$$


Best estimate of slope:

b = Sum of cross products

Sum of squares of X

$$b = \frac{\sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y})}{\sum_{i=1}^n (X_i - \bar{X})^2}$$


Denominator ONLY normalizes based on X (Independent/Explanatory variable)

$$r = \frac{\sum (X - \bar{X})(Y - \bar{Y})}{\sqrt{\sum (X - \bar{X})^2} \sqrt{\sum (Y - \bar{Y})^2}} = \frac{\text{Covariance}(X, Y)}{s_x s_y}$$


Denominator normalizes based on **both** X and Y variables

Finding **a**:

$$\bar{Y} = a + b\bar{X}$$

OR

$$a = \bar{Y} - b\bar{X}$$