

# Selected Topics in Mathematics of Learning

## High-Dimensional Statistics

Lecturer: Marius Yamakou

Winter Semester 2024/25  
Department of Data Science, FAU

December 17, 2024

## Part V: Large Inverse Covariance Matrices continued ...

## 2. Estimation : Why not $\hat{\Sigma}^{-1}$ ?

### First Natural Thought:

- Use the **sample covariance matrix** and compute its inverse.
- Recall the sample covariance matrix:

$$\hat{\Sigma} = \frac{1}{n} \sum_{k=1}^n (X_k - \bar{X})(X_k - \bar{X})^\top =: (\hat{\sigma}_{ij})_{i,j=1,\dots,p}.$$

## 2. Estimation : Why not $\hat{\Sigma}^{-1}$ ?

### First Natural Thought:

- Use the **sample covariance matrix** and compute its inverse.
- Recall the sample covariance matrix:

$$\hat{\Sigma} = \frac{1}{n} \sum_{k=1}^n (X_k - \bar{X})(X_k - \bar{X})^\top =: (\hat{\sigma}_{ij})_{i,j=1,\dots,p}.$$

- The inverse of the sample covariance matrix provides an estimator for  $\Theta$ :

$$\hat{\Theta} = \hat{\Sigma}^{-1}.$$

## 2. Estimation : Why not $\hat{\Sigma}^{-1}$ ?

### First Natural Thought:

- Use the **sample covariance matrix** and compute its inverse.
- Recall the sample covariance matrix:

$$\hat{\Sigma} = \frac{1}{n} \sum_{k=1}^n (X_k - \bar{X})(X_k - \bar{X})^\top =: (\hat{\sigma}_{ij})_{i,j=1,\dots,p}.$$

- The inverse of the sample covariance matrix provides an estimator for  $\Theta$ :

$$\hat{\Theta} = \hat{\Sigma}^{-1}.$$

### Challenges with This Approach:

- **Non-Invertibility:**  $\hat{\Sigma}$  is not invertible when  $n \ll p$ .

## 2. Estimation : Why not $\hat{\Sigma}^{-1}$ ?

### First Natural Thought:

- Use the **sample covariance matrix** and compute its inverse.
- Recall the sample covariance matrix:

$$\hat{\Sigma} = \frac{1}{n} \sum_{k=1}^n (X_k - \bar{X})(X_k - \bar{X})^\top =: (\hat{\sigma}_{ij})_{i,j=1,\dots,p}.$$

- The inverse of the sample covariance matrix provides an estimator for  $\Theta$ :

$$\hat{\Theta} = \hat{\Sigma}^{-1}.$$

### Challenges with This Approach:

- **Non-Invertibility:**  $\hat{\Sigma}$  is not invertible when  $n \ll p$ .
- **Non-Sparsity:** Even if invertible,  $\hat{\Sigma}$  typically lacks exact zeros.

## 2. Estimation : Why not $\hat{\Sigma}^{-1}$ ?

### First Natural Thought:

- Use the **sample covariance matrix** and compute its inverse.
- Recall the sample covariance matrix:

$$\hat{\Sigma} = \frac{1}{n} \sum_{k=1}^n (X_k - \bar{X})(X_k - \bar{X})^\top =: (\hat{\sigma}_{ij})_{i,j=1,\dots,p}.$$

- The inverse of the sample covariance matrix provides an estimator for  $\Theta$ :

$$\hat{\Theta} = \hat{\Sigma}^{-1}.$$

### Challenges with This Approach:

- **Non-Invertibility:**  $\hat{\Sigma}$  is not invertible when  $n \ll p$ .
- **Non-Sparsity:** Even if invertible,  $\hat{\Sigma}$  typically lacks exact zeros.
- **Numerical Instability:** Inversion can lead to large estimation errors, especially in high dimensions.

## 2. Estimation : Why not $\hat{\Sigma}^{-1}$ ?

### First Natural Thought:

- Use the **sample covariance matrix** and compute its inverse.
- Recall the sample covariance matrix:

$$\hat{\Sigma} = \frac{1}{n} \sum_{k=1}^n (X_k - \bar{X})(X_k - \bar{X})^\top =: (\hat{\sigma}_{ij})_{i,j=1,\dots,p}.$$

- The inverse of the sample covariance matrix provides an estimator for  $\Theta$ :

$$\hat{\Theta} = \hat{\Sigma}^{-1}.$$

### Challenges with This Approach:

- **Non-Invertibility:**  $\hat{\Sigma}$  is not invertible when  $n \ll p$ .
- **Non-Sparsity:** Even if invertible,  $\hat{\Sigma}$  typically lacks exact zeros.
- **Numerical Instability:** Inversion can lead to large estimation errors, especially in high dimensions.

### Proposed Solution:

- Re-parameterize the maximum likelihood estimation (MLE) in terms of the **precision matrix**  $\Theta$ .



## 2. Estimation : Why not $\hat{\Sigma}^{-1}$ ?

### First Natural Thought:

- Use the **sample covariance matrix** and compute its inverse.
- Recall the sample covariance matrix:

$$\hat{\Sigma} = \frac{1}{n} \sum_{k=1}^n (X_k - \bar{X})(X_k - \bar{X})^\top =: (\hat{\sigma}_{ij})_{i,j=1,\dots,p}.$$

- The inverse of the sample covariance matrix provides an estimator for  $\Theta$ :

$$\hat{\Theta} = \hat{\Sigma}^{-1}.$$

### Challenges with This Approach:

- **Non-Invertibility:**  $\hat{\Sigma}$  is not invertible when  $n \ll p$ .
- **Non-Sparsity:** Even if invertible,  $\hat{\Sigma}$  typically lacks exact zeros.
- **Numerical Instability:** Inversion can lead to large estimation errors, especially in high dimensions.

### Proposed Solution:

- Re-parameterize the maximum likelihood estimation (MLE) in terms of the **precision matrix**  $\Theta$ .
- Introduce a **penalty term** to enforce sparsity or improve stability.

## 2. Estimation: Recall Maximum Likelihood Estimation

### What is Maximum Likelihood Estimation (MLE)?

- A method to estimate the parameters of a probability distribution by maximizing the **likelihood function**.

## 2. Estimation: Recall Maximum Likelihood Estimation

### What is Maximum Likelihood Estimation (MLE)?

- A method to estimate the parameters of a probability distribution by maximizing the **likelihood function**.
- Ensures that, under the assumed model, the observed data is most probable.

## 2. Estimation: Recall Maximum Likelihood Estimation

### What is Maximum Likelihood Estimation (MLE)?

- A method to estimate the parameters of a probability distribution by maximizing the **likelihood function**.
- Ensures that, under the assumed model, the observed data is most probable.

### Steps in Maximum Likelihood Estimation:

- Assume a random sample from an unknown joint probability distribution, parameterized by a set of parameters.

## 2. Estimation: Recall Maximum Likelihood Estimation

### What is Maximum Likelihood Estimation (MLE)?

- A method to estimate the parameters of a probability distribution by maximizing the **likelihood function**.
- Ensures that, under the assumed model, the observed data is most probable.

### Steps in Maximum Likelihood Estimation:

- Assume a random sample from an unknown joint probability distribution, parameterized by a set of parameters.
- Define the joint density function for the observed data:
  - If random vectors are independent, the joint density equals the product of their individual densities.

## 2. Estimation: Recall Maximum Likelihood Estimation

### What is Maximum Likelihood Estimation (MLE)?

- A method to estimate the parameters of a probability distribution by maximizing the **likelihood function**.
- Ensures that, under the assumed model, the observed data is most probable.

### Steps in Maximum Likelihood Estimation:

- Assume a random sample from an unknown joint probability distribution, parameterized by a set of parameters.
- Define the joint density function for the observed data:
  - If random vectors are independent, the joint density equals the product of their individual densities.
- Evaluate the likelihood function:  $L(\theta) = \prod_{k=1}^n f(X_k; \theta)$ , where  $f(X_k; \theta)$  is the density function.

## 2. Estimation: Recall Maximum Likelihood Estimation

### What is Maximum Likelihood Estimation (MLE)?

- A method to estimate the parameters of a probability distribution by maximizing the **likelihood function**.
- Ensures that, under the assumed model, the observed data is most probable.

### Steps in Maximum Likelihood Estimation:

- Assume a random sample from an unknown joint probability distribution, parameterized by a set of parameters.
- Define the joint density function for the observed data:
  - If random vectors are independent, the joint density equals the product of their individual densities.
- Evaluate the likelihood function:  $L(\theta) = \prod_{k=1}^n f(X_k; \theta)$ , where  $f(X_k; \theta)$  is the density function.
- Maximize the likelihood function to find parameter values that best explain the observed data.

## 2. Estimation: Recall Maximum Likelihood Estimation

### What is Maximum Likelihood Estimation (MLE)?

- A method to estimate the parameters of a probability distribution by maximizing the **likelihood function**.
- Ensures that, under the assumed model, the observed data is most probable.

### Steps in Maximum Likelihood Estimation:

- Assume a random sample from an unknown joint probability distribution, parameterized by a set of parameters.
- Define the joint density function for the observed data:
  - If random vectors are independent, the joint density equals the product of their individual densities.
- Evaluate the likelihood function:  $L(\theta) = \prod_{k=1}^n f(X_k; \theta)$ , where  $f(X_k; \theta)$  is the density function.
- Maximize the likelihood function to find parameter values that best explain the observed data.

### Intuition:

- MLE identifies the parameter values that make the observed data **most probable** under the given statistical model.



## 2. Estimation: Likelihood estimation

**Multivariate normal:** A real random vector  $X = (X_1, \dots, X_p)^T$  is called a normal random vector if there exists a random  $p$ -vector  $Z$ , which is a standard normal random vector, a  $p$ -vector  $\mu$ , and a matrix  $A$ , such that  $X = A^T Z + \mu$ .

## 2. Estimation: Likelihood estimation

**Multivariate normal:** A real random vector  $X = (X_1, \dots, X_p)^T$  is called a normal random vector if there exists a random  $p$ -vector  $Z$ , which is a standard normal random vector, a  $p$ -vector  $\mu$ , and a matrix  $A$ , such that  $X = A^T Z + \mu$ .

Suppose  $X \sim \mathcal{N}_p(\mu, \Sigma)$  and that  $\Sigma$  is full rank; then  $X$  has a density:

$$f_{\mathbf{X}}(x_1, \dots, x_p) = \frac{1}{\sqrt{(2\pi)^p |\Sigma|}} \exp \left( -\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^T \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right)$$

where  $|\Sigma|$  denotes the determinant of  $\Sigma$ .

## 2. Estimation: Likelihood estimation

**Multivariate normal:** A real random vector  $X = (X_1, \dots, X_p)^T$  is called a normal random vector if there exists a random  $p$ -vector  $Z$ , which is a standard normal random vector, a  $p$ -vector  $\mu$ , and a matrix  $A$ , such that  $X = A^T Z + \mu$ .

Suppose  $X \sim \mathcal{N}_p(\mu, \Sigma)$  and that  $\Sigma$  is full rank; then  $X$  has a density:

$$f_{\mathbf{X}}(x_1, \dots, x_p) = \frac{1}{\sqrt{(2\pi)^p |\Sigma|}} \exp \left( -\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^T \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right)$$

where  $|\Sigma|$  denotes the determinant of  $\Sigma$ .

### Key Properties:

- The density is symmetric and bell-shaped, extending into  $p$ -dimensional space.

## 2. Estimation: Likelihood estimation

**Multivariate normal:** A real random vector  $X = (X_1, \dots, X_p)^T$  is called a normal random vector if there exists a random  $p$ -vector  $Z$ , which is a standard normal random vector, a  $p$ -vector  $\mu$ , and a matrix  $A$ , such that  $X = A^T Z + \mu$ .

Suppose  $X \sim \mathcal{N}_p(\mu, \Sigma)$  and that  $\Sigma$  is full rank; then  $X$  has a density:

$$f_{\mathbf{X}}(x_1, \dots, x_p) = \frac{1}{\sqrt{(2\pi)^p |\Sigma|}} \exp \left( -\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^T \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right)$$

where  $|\Sigma|$  denotes the determinant of  $\Sigma$ .

### Key Properties:

- The density is symmetric and bell-shaped, extending into  $p$ -dimensional space.
- The covariance matrix  $\Sigma$  controls the shape and orientation of the distribution.

## 2. Estimation: Likelihood Estimation

### Setup:

- We have random vectors  $X_n = (X_{n,1}, \dots, X_{n,p})^\top$ , for  $n = 1, \dots, N$ , where  $X_n \sim \mathcal{N}_p(\mu, \Sigma)$ .
- The mean vector  $\mu \in \mathbb{R}^p$  and covariance matrix  $\Sigma \in \mathbb{R}^{p \times p}$  are unknown.

## 2. Estimation: Likelihood Estimation

### Setup:

- We have random vectors  $X_n = (X_{n,1}, \dots, X_{n,p})^\top$ , for  $n = 1, \dots, N$ , where  $X_n \sim \mathcal{N}_p(\mu, \Sigma)$ .
- The mean vector  $\mu \in \mathbb{R}^p$  and covariance matrix  $\Sigma \in \mathbb{R}^{p \times p}$  are unknown.

**Joint Density:** The joint density of the observations  $X_1, \dots, X_N$  is:

$$\prod_{n=1}^N f_{\mathbf{X}_n}(x_1, \dots, x_p) = \prod_{n=1}^N \frac{1}{\sqrt{(2\pi)^p |\Sigma|}} \exp \left( -\frac{1}{2} (\mathbf{x}_n - \boldsymbol{\mu})^\top \Sigma^{-1} (\mathbf{x}_n - \boldsymbol{\mu}) \right).$$

## 2. Estimation: Likelihood Estimation

**Negative Log-Likelihood:** Taking the negative log of the joint density:

## 2. Estimation: Likelihood Estimation

**Negative Log-Likelihood:** Taking the negative log of the joint density:

$$\begin{aligned} -\log \prod_{n=1}^N f_{\mathbf{X}_n}(x_1, \dots, x_p) &= -\log \prod_{n=1}^N \frac{1}{\sqrt{(2\pi)^p |\Sigma|}} \exp \left( -\frac{1}{2} (\mathbf{x}_n - \boldsymbol{\mu})^\top \Sigma^{-1} (\mathbf{x}_n - \boldsymbol{\mu}) \right) \\ &= -\log \left( \frac{1}{((2\pi)^p |\Sigma|)^{N/2}} \exp \left( -\frac{1}{2} \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu})^\top \Sigma^{-1} (\mathbf{x}_n - \boldsymbol{\mu}) \right) \right) \\ &= \frac{N}{2} \log ((2\pi)^p |\Sigma|) + \frac{1}{2} \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu})^\top \Sigma^{-1} (\mathbf{x}_n - \boldsymbol{\mu}) \\ &= \frac{pN}{2} \log(2\pi) + \frac{N}{2} \log |\Sigma| + \frac{1}{2} \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu})^\top \Sigma^{-1} (\mathbf{x}_n - \boldsymbol{\mu}) \end{aligned}$$



## 2. Estimation: Maximum Likelihood Estimators

**Negative Log-Likelihood:**

$$L(\mu, \Sigma) = \frac{pN}{2} \log(2\pi) + \frac{N}{2} \log |\Sigma| + \frac{1}{2} \sum_{n=1}^N (X_n - \mu)^\top \Sigma^{-1} (X_n - \mu).$$

## 2. Estimation: Maximum Likelihood Estimators

### Negative Log-Likelihood:

$$L(\mu, \Sigma) = \frac{pN}{2} \log(2\pi) + \frac{N}{2} \log |\Sigma| + \frac{1}{2} \sum_{n=1}^N (X_n - \mu)^\top \Sigma^{-1} (X_n - \mu).$$

**Estimating  $\mu$  and  $\Sigma$ :** To minimize  $L(\mu, \Sigma)$ , the Maximum Likelihood Estimators (MLEs) are:

$$\operatorname{argmin}_{\mu} L(\mu, \Sigma) = \frac{1}{N} \sum_{n=1}^N X_n,$$

$$\operatorname{argmin}_{\Sigma} L(\mu, \Sigma) = \frac{1}{N} \sum_{n=1}^N (X_n - \mu)(X_n - \mu)^\top.$$

## 2. Estimation: Maximum Likelihood Estimators

### Negative Log-Likelihood:

$$L(\mu, \Sigma) = \frac{pN}{2} \log(2\pi) + \frac{N}{2} \log |\Sigma| + \frac{1}{2} \sum_{n=1}^N (X_n - \mu)^\top \Sigma^{-1} (X_n - \mu).$$

**Estimating  $\mu$  and  $\Sigma$ :** To minimize  $L(\mu, \Sigma)$ , the Maximum Likelihood Estimators (MLEs) are:

$$\operatorname{argmin}_{\mu} L(\mu, \Sigma) = \frac{1}{N} \sum_{n=1}^N X_n,$$

$$\operatorname{argmin}_{\Sigma} L(\mu, \Sigma) = \frac{1}{N} \sum_{n=1}^N (X_n - \mu)(X_n - \mu)^\top.$$

**Key Idea:** - The sample mean  $\hat{\mu}$  and sample covariance matrix  $\hat{\Sigma}$  are natural estimators under the MLE framework.

## 2. Estimation: MLE for Precision Matrix $\Theta$

**Re-Parameterization:** To get MLE for  $\Theta$  we can re-parameterize in terms of precision matrix  $\Theta = \Sigma^{-1}$ .

## 2. Estimation: MLE for Precision Matrix $\Theta$

**Re-Parameterization:** To get MLE for  $\Theta$  we can re-parameterize in terms of precision matrix  $\Theta = \Sigma^{-1}$ .

Rewriting the negative log-likelihood:

## 2. Estimation: MLE for Precision Matrix $\Theta$

**Re-Parameterization:** To get MLE for  $\Theta$  we can re-parameterize in terms of precision matrix  $\Theta = \Sigma^{-1}$ .

Rewriting the negative log-likelihood:

$$\begin{aligned} L(\mu, \Sigma) &= \frac{pN}{2} \log(2\pi) + \frac{N}{2} \log |\Sigma| + \frac{1}{2} \sum_{n=1}^N (X_n - \mu)^\top \Sigma^{-1} (X_n - \mu) \\ &= \frac{pN}{2} \log(2\pi) + \frac{N}{2} \underbrace{\log |\Sigma|}_{-\log |\Sigma^{-1}|} + \frac{1}{2} \underbrace{\sum_{n=1}^N (X_n - \mu)^\top \underbrace{\Sigma^{-1}}_{\Theta} (X_n - \mu)}_{\substack{\text{tr} \left( \sum_{n=1}^N (X_n - \mu)^\top \Theta (X_n - \mu) \right) \\ \stackrel{(*)}{=} \text{tr} \left( \sum_{n=1}^N (X_n - \mu) (X_n - \mu)^\top \Theta \right) \\ = \text{tr} \left( N \hat{\Sigma} \Theta \right) \\ = N \text{tr} \left( \hat{\Sigma} \Theta \right)}} \end{aligned}$$

## 2. Estimation: MLE for Precision Matrix $\Theta$

**Re-Parameterization:** To get MLE for  $\Theta$  we can re-parameterize in terms of precision matrix  $\Theta = \Sigma^{-1}$ .

Rewriting the negative log-likelihood:

$$\begin{aligned} L(\mu, \Sigma) &= \frac{pN}{2} \log(2\pi) + \frac{N}{2} \log |\Sigma| + \frac{1}{2} \sum_{n=1}^N (X_n - \mu)^\top \Sigma^{-1} (X_n - \mu) \\ &= \frac{pN}{2} \log(2\pi) + \frac{N}{2} \underbrace{\log |\Sigma|}_{-\log |\Sigma^{-1}|} + \frac{1}{2} \underbrace{\sum_{n=1}^N (X_n - \mu)^\top \underbrace{\Sigma^{-1}}_{\Theta} (X_n - \mu)}_{\substack{\text{tr} \left( \sum_{n=1}^N (X_n - \mu)^\top \Theta (X_n - \mu) \right) \\ \stackrel{(*)}{=} \text{tr} \left( \sum_{n=1}^N (X_n - \mu) (X_n - \mu)^\top \Theta \right) \\ = \text{tr} \left( N \hat{\Sigma} \Theta \right) \\ = N \text{tr} \left( \hat{\Sigma} \Theta \right)}} \end{aligned}$$

$$\implies L(\mu, \Sigma) = \frac{pN}{2} \log(2\pi) - \frac{N}{2} \log |\Theta| + \frac{1}{2} N \text{tr}(\hat{\Sigma} \Theta)$$

## 2. Estimation: MLE for Precision Matrix $\Theta$

where:

- $\hat{\Sigma} = \frac{1}{N} \sum_{n=1}^N (X_n - \mu)(X_n - \mu)^\top$  is the sample covariance matrix.



## 2. Estimation: MLE for Precision Matrix $\Theta$

where:

- $\hat{\Sigma} = \frac{1}{N} \sum_{n=1}^N (X_n - \mu)(X_n - \mu)^\top$  is the sample covariance matrix.
- (\*) exploit the cyclic property of the trace operator  $\text{tr}(\cdot)$  given by:

$$\text{tr}(ABC) = \text{tr}(CAB).$$

## 2. Estimation: MLE for Precision Matrix $\Theta$

where:

- $\hat{\Sigma} = \frac{1}{N} \sum_{n=1}^N (X_n - \mu)(X_n - \mu)^\top$  is the sample covariance matrix.
- (\*) exploit the cyclic property of the trace operator  $\text{tr}(\cdot)$  given by:

$$\text{tr}(ABC) = \text{tr}(CAB).$$

**Summary:** The MLE for  $\Theta$  minimizes:

$$L(\mu, \Sigma) = \frac{pN}{2} \log(2\pi) - \frac{N}{2} \log |\Theta| + \frac{N}{2} \text{tr}(\hat{\Sigma}\Theta).$$

## 2.2 Graphical LASSO

**Motivation:** The graphical LASSO introduces sparsity into the estimation of the precision matrix  $\Theta = \Sigma^{-1}$ , leading to interpretable graphical models by penalizing small off-diagonal entries.

## 2.2 Graphical LASSO

**Motivation:** The graphical LASSO introduces sparsity into the estimation of the precision matrix  $\Theta = \Sigma^{-1}$ , leading to interpretable graphical models by penalizing small off-diagonal entries.

**Maximum Likelihood Estimator (MLE):** Based on the sample covariance matrix  $\hat{\Sigma}$ , the MLE of  $\Theta$  is:

$$\hat{\Theta} = \underset{\Theta \succ 0, \Theta^T = \Theta}{\operatorname{argmin}} \left( \operatorname{tr}(\Theta \hat{\Sigma}) - \log |\Theta| \right),$$

subject to two constraints:

## 2.2 Graphical LASSO

**Motivation:** The graphical LASSO introduces sparsity into the estimation of the precision matrix  $\Theta = \Sigma^{-1}$ , leading to interpretable graphical models by penalizing small off-diagonal entries.

**Maximum Likelihood Estimator (MLE):** Based on the sample covariance matrix  $\hat{\Sigma}$ , the MLE of  $\Theta$  is:

$$\hat{\Theta} = \underset{\Theta \succ 0, \Theta^T = \Theta}{\operatorname{argmin}} \left( \operatorname{tr}(\Theta \hat{\Sigma}) - \log |\Theta| \right),$$

subject to two constraints:

- $\Theta \succ 0$ : Positive definiteness of the precision matrix.
- $\Theta^T = \Theta$ : Symmetry of the precision matrix.

## 2.2 Graphical LASSO

**Why the Objective:**  $\hat{\Theta} = \underset{\Theta \succ 0, \Theta^{\top} = \Theta}{\operatorname{argmin}} \left( \operatorname{tr}(\Theta \hat{\Sigma}) - \log |\Theta| \right) ?$

## 2.2 Graphical LASSO

**Why the Objective:**  $\hat{\Theta} = \underset{\Theta \succ 0, \Theta^\top = \Theta}{\operatorname{argmin}} \left( \operatorname{tr}(\Theta \hat{\Sigma}) - \log |\Theta| \right) ?$

Because:

- $\operatorname{tr}(\Theta \hat{\Sigma})$  ensures that the precision matrix  $\Theta$  aligns with the empirical covariance matrix  $\hat{\Sigma}$ .
- It penalizes deviations of  $\Theta$  from matching  $\hat{\Sigma}$ .

## 2.2 Graphical LASSO

**Why the Objective:**  $\hat{\Theta} = \underset{\Theta \succ 0, \Theta^\top = \Theta}{\operatorname{argmin}} \left( \operatorname{tr}(\Theta \hat{\Sigma}) - \log |\Theta| \right) ?$

Because:

- $\operatorname{tr}(\Theta \hat{\Sigma})$  ensures that the precision matrix  $\Theta$  aligns with the empirical covariance matrix  $\hat{\Sigma}$ .
- It penalizes deviations of  $\Theta$  from matching  $\hat{\Sigma}$ .
- $-\log |\Theta|$  represents the log-determinant of  $\Theta$ , which has two key roles:



## 2.2 Graphical LASSO

**Why the Objective:**  $\hat{\Theta} = \underset{\Theta \succ 0, \Theta^\top = \Theta}{\operatorname{argmin}} \left( \operatorname{tr}(\Theta \hat{\Sigma}) - \log |\Theta| \right) ?$

Because:

- $\operatorname{tr}(\Theta \hat{\Sigma})$  ensures that the precision matrix  $\Theta$  aligns with the empirical covariance matrix  $\hat{\Sigma}$ .
  - It penalizes deviations of  $\Theta$  from matching  $\hat{\Sigma}$ .
- $-\log |\Theta|$  represents the log-determinant of  $\Theta$ , which has two key roles:
  - Ensures the matrix  $\Theta$  is invertible.

## 2.2 Graphical LASSO

**Why the Objective:**  $\hat{\Theta} = \underset{\Theta \succ 0, \Theta^\top = \Theta}{\operatorname{argmin}} \left( \operatorname{tr}(\Theta \hat{\Sigma}) - \log |\Theta| \right) ?$

Because:

- $\operatorname{tr}(\Theta \hat{\Sigma})$  ensures that the precision matrix  $\Theta$  aligns with the empirical covariance matrix  $\hat{\Sigma}$ .
  - It penalizes deviations of  $\Theta$  from matching  $\hat{\Sigma}$ .
- $-\log |\Theta|$  represents the log-determinant of  $\Theta$ , which has two key roles:
  - Ensures the matrix  $\Theta$  is invertible.
  - Penalizes overly large or overly small eigenvalues, maintaining numerical stability.

## 2.2 Graphical LASSO

**Why the Objective:**  $\hat{\Theta} = \underset{\Theta \succ 0, \Theta^\top = \Theta}{\operatorname{argmin}} \left( \operatorname{tr}(\Theta \hat{\Sigma}) - \log |\Theta| \right) ?$

Because:

- $\operatorname{tr}(\Theta \hat{\Sigma})$  ensures that the precision matrix  $\Theta$  aligns with the empirical covariance matrix  $\hat{\Sigma}$ .
  - It penalizes deviations of  $\Theta$  from matching  $\hat{\Sigma}$ .
- $-\log |\Theta|$  represents the log-determinant of  $\Theta$ , which has two key roles:
  - Ensures the matrix  $\Theta$  is invertible.
  - Penalizes overly large or overly small eigenvalues, maintaining numerical stability.

**Intuition:** The objective function balances:

- **Fidelity:** The trace term  $\operatorname{tr}(\Theta \hat{\Sigma})$  ensures consistency with the observed data.

## 2.2 Graphical LASSO

**Why the Objective:**  $\hat{\Theta} = \underset{\Theta \succ 0, \Theta^\top = \Theta}{\operatorname{argmin}} \left( \operatorname{tr}(\Theta \hat{\Sigma}) - \log |\Theta| \right) ?$

Because:

- $\operatorname{tr}(\Theta \hat{\Sigma})$  ensures that the precision matrix  $\Theta$  aligns with the empirical covariance matrix  $\hat{\Sigma}$ .
  - It penalizes deviations of  $\Theta$  from matching  $\hat{\Sigma}$ .
- $-\log |\Theta|$  represents the log-determinant of  $\Theta$ , which has two key roles:
  - Ensures the matrix  $\Theta$  is invertible.
  - Penalizes overly large or overly small eigenvalues, maintaining numerical stability.

**Intuition:** The objective function balances:

- **Fidelity:** The trace term  $\operatorname{tr}(\Theta \hat{\Sigma})$  ensures consistency with the observed data.
- **Regularization:** The log-determinant term  $-\log |\Theta|$  prevents degenerate solutions and stabilizes the optimization.

## 2.2 Graphical LASSO

**Graphical LASSO: Introducing Sparsity** The graphical LASSO estimator adds an  $\ell_1$ -penalty on the off-diagonal entries of  $\Theta$  to induce sparsity:

## 2.2 Graphical LASSO

**Graphical LASSO: Introducing Sparsity** The graphical LASSO estimator adds an  $\ell_1$ -penalty on the off-diagonal entries of  $\Theta$  to induce sparsity:

$$\hat{\Theta}_\lambda = \underset{\Theta \succ 0, \Theta^\top = \Theta}{\operatorname{argmin}} \left( \operatorname{tr}(\Theta \hat{\Sigma}) - \log |\Theta| + \lambda \|\Theta\|_{1,\text{off}} \right),$$

where:

## 2.2 Graphical LASSO

**Graphical LASSO: Introducing Sparsity** The graphical LASSO estimator adds an  $\ell_1$ -penalty on the off-diagonal entries of  $\Theta$  to induce sparsity:

$$\hat{\Theta}_\lambda = \underset{\Theta \succ 0, \Theta^\top = \Theta}{\operatorname{argmin}} \left( \operatorname{tr}(\Theta \hat{\Sigma}) - \log |\Theta| + \lambda \|\Theta\|_{1,\text{off}} \right),$$

where:

$$\|\Theta\|_{1,\text{off}} = \sum_{\substack{i,j=1 \\ i \neq j}}^p |\Theta_{ij}|.$$

## 2.2 Graphical LASSO

**Graphical LASSO: Introducing Sparsity** The graphical LASSO estimator adds an  $\ell_1$ -penalty on the off-diagonal entries of  $\Theta$  to induce sparsity:

$$\hat{\Theta}_\lambda = \underset{\Theta \succ 0, \Theta^\top = \Theta}{\operatorname{argmin}} \left( \operatorname{tr}(\Theta \hat{\Sigma}) - \log |\Theta| + \lambda \|\Theta\|_{1,\text{off}} \right),$$

where:

$$\|\Theta\|_{1,\text{off}} = \sum_{\substack{i,j=1 \\ i \neq j}}^p |\Theta_{ij}|.$$

**Key Question:**

- Is the penalized likelihood a convex function in  $\Theta$ ?



## 2.2 Graphical LASSO

**Graphical LASSO: Introducing Sparsity** The graphical LASSO estimator adds an  $\ell_1$ -penalty on the off-diagonal entries of  $\Theta$  to induce sparsity:

$$\hat{\Theta}_\lambda = \underset{\Theta \succ 0, \Theta^\top = \Theta}{\operatorname{argmin}} \left( \operatorname{tr}(\Theta \hat{\Sigma}) - \log |\Theta| + \lambda \|\Theta\|_{1,\text{off}} \right),$$

where:

$$\|\Theta\|_{1,\text{off}} = \sum_{\substack{i,j=1 \\ i \neq j}}^p |\Theta_{ij}|.$$

**Key Question:**

- Is the penalized likelihood a convex function in  $\Theta$ ?

**Key Insight:**

- The graphical LASSO objective is a combination of a convex negative log-likelihood function and a convex  $\ell_1$ -norm penalty term.

## 2.2 Graphical LASSO

**Graphical LASSO: Introducing Sparsity** The graphical LASSO estimator adds an  $\ell_1$ -penalty on the off-diagonal entries of  $\Theta$  to induce sparsity:

$$\hat{\Theta}_\lambda = \underset{\Theta \succ 0, \Theta^\top = \Theta}{\operatorname{argmin}} \left( \operatorname{tr}(\Theta \hat{\Sigma}) - \log |\Theta| + \lambda \|\Theta\|_{1,\text{off}} \right),$$

where:

$$\|\Theta\|_{1,\text{off}} = \sum_{\substack{i,j=1 \\ i \neq j}}^p |\Theta_{ij}|.$$

### Key Question:

- Is the penalized likelihood a convex function in  $\Theta$ ?

### Key Insight:

- The graphical LASSO objective is a combination of a convex negative log-likelihood function and a convex  $\ell_1$ -norm penalty term.
- Therefore, the overall problem remains convex in  $\Theta$ . And the answer to the above question is Yes!