

# Selected Topics in Mathematics of Learning

## **High-Dimensional Statistics**

Lecturer: Marius Yamakou

Winter Semester 2024/25  
Department of Data Science, FAU

January 21, 2025

## Part VI

### Sparse vector autoregressive models

#### Motivation

- Traditional assumptions:
  - Data are multivariate Gaussian
  - Observations are independent and identically distributed (i.i.d.)

## Part VI

### Sparse vector autoregressive models

#### Motivation

- Traditional assumptions:
  - Data are multivariate Gaussian
  - Observations are independent and identically distributed (i.i.d.)
- Real-world data often deviate from these assumptions:
  - Examples: fMRI data, ETFs
  - Temporal correlations are common

## Part VI

### Sparse vector autoregressive models

#### Motivation

- Traditional assumptions:
  - Data are multivariate Gaussian
  - Observations are independent and identically distributed (i.i.d.)
- Real-world data often deviate from these assumptions:
  - Examples: fMRI data, ETFs
  - Temporal correlations are common

**Key Idea:** Vector Autoregressive Models (VARs) are powerful tools for capturing temporal correlations in high-dimensional data.

# Outline

- Preliminaries of VAR
- Estimation of Sparse VAR using LASSO
- LASSO: Estimation
- LASSO: Properties

# 1. Preliminaries of VAR

A time series  $\{X_t\}_{t \in \mathbb{Z}} = \{(X_{j,t})_{j=1,\dots,d}\}_{t \in \mathbb{Z}}$  follows a **VAR**( $p$ ) model if:

$$X_t = \Phi_1 X_{t-1} + \dots + \Phi_p X_{t-p} + \epsilon_t, \quad t \in \mathbb{Z},$$

# 1. Preliminaries of VAR

A time series  $\{X_t\}_{t \in \mathbb{Z}} = \{(X_{j,t})_{j=1,\dots,d}\}_{t \in \mathbb{Z}}$  follows a **VAR**( $p$ ) model if:

$$X_t = \Phi_1 X_{t-1} + \dots + \Phi_p X_{t-p} + \epsilon_t, \quad t \in \mathbb{Z},$$

where:

- $\Phi_1, \dots, \Phi_p$  are  $d \times d$  matrices.
- $\{\epsilon_t\}_{t \in \mathbb{Z}}$  is a white noise series with the following statistics:

$$\mathbb{E}[\epsilon_t] = 0, \quad \mathbb{E}[\epsilon_t \epsilon_t'] = \Sigma_\epsilon, \quad \mathbb{E}[\epsilon_s \epsilon_t'] = 0 \quad \text{for } s \neq t.$$

# 1. Preliminaries of VAR

A time series  $\{X_t\}_{t \in \mathbb{Z}} = \{(X_{j,t})_{j=1,\dots,d}\}_{t \in \mathbb{Z}}$  follows a **VAR**( $p$ ) model if:

$$X_t = \Phi_1 X_{t-1} + \dots + \Phi_p X_{t-p} + \epsilon_t, \quad t \in \mathbb{Z},$$

where:

- $\Phi_1, \dots, \Phi_p$  are  $d \times d$  matrices.
- $\{\epsilon_t\}_{t \in \mathbb{Z}}$  is a white noise series with the following statistics:

$$\mathbb{E}[\epsilon_t] = 0, \quad \mathbb{E}[\epsilon_t \epsilon_t'] = \Sigma_\epsilon, \quad \mathbb{E}[\epsilon_s \epsilon_t'] = 0 \quad \text{for } s \neq t.$$

**Example:** VAR(1) with  $d = 2$

$$\Phi_1 = (\Phi_{1,jk})_{j,k=1,2} = \begin{bmatrix} \Phi_{1,11} & \Phi_{1,12} \\ \Phi_{1,21} & \Phi_{1,22} \end{bmatrix}$$



# 1. Preliminaries of VAR

A time series  $\{X_t\}_{t \in \mathbb{Z}} = \{(X_{j,t})_{j=1,\dots,d}\}_{t \in \mathbb{Z}}$  follows a **VAR**( $p$ ) model if:

$$X_t = \Phi_1 X_{t-1} + \dots + \Phi_p X_{t-p} + \epsilon_t, \quad t \in \mathbb{Z},$$

where:

- $\Phi_1, \dots, \Phi_p$  are  $d \times d$  matrices.
- $\{\epsilon_t\}_{t \in \mathbb{Z}}$  is a white noise series with the following statistics:

$$\mathbb{E}[\epsilon_t] = 0, \quad \mathbb{E}[\epsilon_t \epsilon_t'] = \Sigma_\epsilon, \quad \mathbb{E}[\epsilon_s \epsilon_t'] = 0 \quad \text{for } s \neq t.$$

**Example:** VAR(1) with  $d = 2$

$$\Phi_1 = (\Phi_{1,jk})_{j,k=1,2} = \begin{bmatrix} \Phi_{1,11} & \Phi_{1,12} \\ \Phi_{1,21} & \Phi_{1,22} \end{bmatrix}$$

The equations are:

$$X_{1,t} = \Phi_{1,11} X_{1,t-1} + \Phi_{1,12} X_{2,t-1} + \epsilon_{1,t},$$

$$X_{2,t} = \Phi_{1,21} X_{1,t-1} + \Phi_{1,22} X_{2,t-1} + \epsilon_{2,t}.$$

# 1. Preliminaries of VAR

Please note the following:

- $\mathbb{E}[\epsilon_t] = 0$ : This indicates that the white noise has a mean (or expected value) of 0 for all time points  $t$ . Essentially, the noise fluctuates around zero on average.
- $\mathbb{E}[\epsilon_t \epsilon_t'] = \Sigma_\epsilon$ : Here,  $\Sigma_\epsilon$  is the covariance matrix of  $\epsilon_t$ , which characterizes the variance (for scalar  $\epsilon_t$ ) or the relationships between components (for vector-valued  $\epsilon_t$ ). For scalar white noise, this reduces to  $\mathbb{E}[\epsilon_t^2] = \sigma_\epsilon^2$ , where  $\sigma_\epsilon^2$  is the variance.
- $\mathbb{E}[\epsilon_s \epsilon_t'] = 0$  for  $s \neq t$ : This indicates that  $\epsilon_t$  values at different time points  $t$  are uncorrelated. That is, the noise at time  $t$  does not depend on or influence the noise at any other time  $s$ . For vector-valued  $\epsilon_t$ , this implies that the cross-covariance between vectors at different times is 0.

# 1. Preliminaries of VAR

- If the **VAR(1)** model has  $d = 1$  (i.e., a univariate time series) and  $p = 1$ , the equation simplifies significantly because there is only one time series and only one lag, making it one of the simplest forms of a VAR model.

# 1. Preliminaries of VAR

- If the **VAR(1)** model has  $d = 1$  (i.e., a univariate time series) and  $p = 1$ , the equation simplifies significantly because there is only one time series and only one lag, making it one of the simplest forms of a VAR model.
- The univariate time series  $\{X_t\}_{t \in \mathbb{Z}}$ , a **VAR(1)** model with  $d = 1$  and  $p = 1$  simplifies to:

$$X_t = \Phi_1 X_{t-1} + \epsilon_t,$$

# 1. Preliminaries of VAR

- If the **VAR(1)** model has  $d = 1$  (i.e., a univariate time series) and  $p = 1$ , the equation simplifies significantly because there is only one time series and only one lag, making it one of the simplest forms of a VAR model.
- The univariate time series  $\{X_t\}_{t \in \mathbb{Z}}$ , a **VAR(1)** model with  $d = 1$  and  $p = 1$  simplifies to:

$$X_t = \Phi_1 X_{t-1} + \epsilon_t,$$

where:

- $\Phi_1$  is a scalar (instead of a matrix since  $d = 1$ ).
- $\epsilon_t$  is white noise:

$$\mathbb{E}[\epsilon_t] = 0, \quad \mathbb{E}[\epsilon_t^2] = \sigma_\epsilon^2, \quad \mathbb{E}[\epsilon_s \epsilon_t] = 0 \text{ for } s \neq t.$$

# 1.1 Stability

- A VAR(1) model is:
  - **Strictly stable** if:

$$\det(I_d - \Phi_1 z) \neq 0, \quad \text{for } |z| < 1,$$

which is equivalent to requiring that all eigenvalues of  $\Phi_1$  satisfy  $|\lambda| < 1$ .

# 1.1 Stability

- A VAR(1) model is:

- **Strictly stable** if:

$$\det(I_d - \Phi_1 z) \neq 0, \quad \text{for } |z| < 1,$$

which is equivalent to requiring that all eigenvalues of  $\Phi_1$  satisfy  $|\lambda| < 1$ .

- **Marginally stable** if:

$$\det(I_d - \Phi_1 z) = 0 \quad \text{for } |z| = 1,$$

but the system does not diverge over time. This typically occurs when some eigenvalues lie exactly on the unit circle (e.g.,  $|\lambda| = 1$ ).

# 1.1 Stability

- A VAR(1) model is:

- **Strictly stable** if:

$$\det(I_d - \Phi_1 z) \neq 0, \quad \text{for } |z| < 1,$$

which is equivalent to requiring that all eigenvalues of  $\Phi_1$  satisfy  $|\lambda| < 1$ .

- **Marginally stable** if:

$$\det(I_d - \Phi_1 z) = 0 \quad \text{for } |z| = 1,$$

but the system does not diverge over time. This typically occurs when some eigenvalues lie exactly on the unit circle (e.g.,  $|\lambda| = 1$ ).

- For  $d = 1$ , the model  $X_t = \Phi_1 X_{t-1} + \epsilon_t$ , is:

- **Strictly stable** if  $|\Phi_1| < 1$ ,



# 1.1 Stability

- A VAR(1) model is:

- **Strictly stable** if:

$$\det(I_d - \Phi_1 z) \neq 0, \quad \text{for } |z| < 1,$$

which is equivalent to requiring that all eigenvalues of  $\Phi_1$  satisfy  $|\lambda| < 1$ .

- **Marginally stable** if:

$$\det(I_d - \Phi_1 z) = 0 \quad \text{for } |z| = 1,$$

but the system does not diverge over time. This typically occurs when some eigenvalues lie exactly on the unit circle (e.g.,  $|\lambda| = 1$ ).

- For  $d = 1$ , the model  $X_t = \Phi_1 X_{t-1} + \epsilon_t$ , is:

- **Strictly stable** if  $|\Phi_1| < 1$ ,
  - **Marginally stable** if  $|\Phi_1| = 1$ ,

# 1.1 Stability

- A VAR(1) model is:

- **Strictly stable** if:

$$\det(I_d - \Phi_1 z) \neq 0, \quad \text{for } |z| < 1,$$

which is equivalent to requiring that all eigenvalues of  $\Phi_1$  satisfy  $|\lambda| < 1$ .

- **Marginally stable** if:

$$\det(I_d - \Phi_1 z) = 0 \quad \text{for } |z| = 1,$$

but the system does not diverge over time. This typically occurs when some eigenvalues lie exactly on the unit circle (e.g.,  $|\lambda| = 1$ ).

- For  $d = 1$ , the model  $X_t = \Phi_1 X_{t-1} + \epsilon_t$ , is:

- **Strictly stable** if  $|\Phi_1| < 1$ ,
  - **Marginally stable** if  $|\Phi_1| = 1$ ,
  - **Unstable** if  $|\Phi_1| > 1$ .

# 1.1 Stability

**General Stability Condition:** For a  $\text{VAR}(p)$  model, stability requires that its characteristic polynomial satisfies:

$$\det(\Phi(z)) \neq 0, \quad \text{for } |z| \leq 1, z \in \mathbb{C},$$

# 1.1 Stability

**General Stability Condition:** For a VAR( $p$ ) model, stability requires that its characteristic polynomial satisfies:

$$\det(\Phi(z)) \neq 0, \quad \text{for } |z| \leq 1, \quad z \in \mathbb{C},$$

where:

$$\Phi(z) = I_d - \Phi_1 z - \Phi_2 z^2 - \dots - \Phi_p z^p.$$

# 1.1 Stability

**General Stability Condition:** For a VAR( $p$ ) model, stability requires that its characteristic polynomial satisfies:

$$\det(\Phi(z)) \neq 0, \quad \text{for } |z| \leq 1, \quad z \in \mathbb{C},$$

where:

$$\Phi(z) = I_d - \Phi_1 z - \Phi_2 z^2 - \dots - \Phi_p z^p.$$

**Assumption:** In this part of the course, we assume all VAR models are strictly stable unless otherwise stated.

## 1.2 Stationarity

A VAR model is (second-order) **Stationarity** if:

- 1 The mean of the process is constant:

$$\mathbb{E}(X_t) = 0 \quad (\text{or constant over time}).$$

## 1.2 Stationarity

A VAR model is (second-order) **Stationarity** if:

- 1 The mean of the process is constant:

$$\mathbb{E}(X_t) = 0 \quad (\text{or constant over time}).$$

- 2 The **autocovariance function (ACVF)** depends only on the lag  $\ell$  and not on  $t$ , that is:

$$\Gamma_X(\ell) = \mathbb{E}(X_{t+\ell}X_t') = \mathbb{E}(X_\ell X_0').$$

## 1.2 Stationarity

A VAR model is (second-order) **Stationarity** if:

- 1 The mean of the process is constant:

$$\mathbb{E}(X_t) = 0 \quad (\text{or constant over time}).$$

- 2 The **autocovariance function (ACVF)** depends only on the lag  $\ell$  and not on  $t$ , that is:

$$\Gamma_X(\ell) = \mathbb{E}(X_{t+\ell}X_t') = \mathbb{E}(X_\ell X_0').$$

**Remark:** When  $\ell = 0$ , we get the **covariance**:  $\Gamma_X(0) = \Sigma_X = \mathbb{E}(X_tX_t')$



## 1.2 Stationarity

A VAR model is (second-order) **Stationarity** if:

- 1 The mean of the process is constant:

$$\mathbb{E}(X_t) = 0 \quad (\text{or constant over time}).$$

- 2 The **autocovariance function (ACVF)** depends only on the lag  $\ell$  and not on  $t$ , that is:

$$\Gamma_X(\ell) = \mathbb{E}(X_{t+\ell}X_t') = \mathbb{E}(X_\ell X_0').$$

**Remark:** When  $\ell = 0$ , we get the **covariance**:  $\Gamma_X(0) = \Sigma_X = \mathbb{E}(X_tX_t')$

**Implications of stationarity:**

- The statistical properties of the time series do not change over time.

## 1.2 Stationarity

A VAR model is (second-order) **Stationarity** if:

- 1 The mean of the process is constant:

$$\mathbb{E}(X_t) = 0 \quad (\text{or constant over time}).$$

- 2 The **autocovariance function (ACVF)** depends only on the lag  $\ell$  and not on  $t$ , that is:


$$\Gamma_X(\ell) = \mathbb{E}(X_{t+\ell}X_t') = \mathbb{E}(X_\ell X_0').$$

**Remark:** When  $\ell = 0$ , we get the **covariance**:  $\Gamma_X(0) = \Sigma_X = \mathbb{E}(X_tX_t')$

**Implications of stationarity:**

- The statistical properties of the time series do not change over time.
- Stationarity ensures that the VAR model's behavior is predictable and consistent over time, making it suitable for modeling and forecasting.


## 1.2 Stationarity: Long-range dependence (LRD)



### Left: Time Series:

- The time series shows a pattern with persistent, slowly decaying behavior over time, indicating that observations are strongly correlated even at long time lags.

## 1.2 Stationarity: Long-range dependence (LRD)




### Right: Autocorrelation Function (ACF):

- The autocorrelation function (ACF)  $\Gamma_X(\ell)/\Gamma_X(0)$  decreases very slowly as the lag  $\ell$  increases.

### Left: Time Series:

- The time series shows a pattern with persistent, slowly decaying behavior over time, indicating that observations are strongly correlated even at long time lags.

## 1.2 Stationarity: Long-range dependence (LRD)




### Left: Time Series:

- The time series shows a pattern with persistent, slowly decaying behavior over time, indicating that observations are strongly correlated even at long time lags.

### Right: Autocorrelation Function (ACF):

- The autocorrelation function (ACF)  $\Gamma_X(\ell)/\Gamma_X(0)$  decreases very slowly as the lag  $\ell$  increases.
- This behavior reflects **long-range dependence (LRD)**, where correlations persist over long time periods.

## 1.2 Stationarity: Long-range dependence (LRD)



### Left: Time Series:

- The time series shows a pattern with persistent, slowly decaying behavior over time, indicating that observations are strongly correlated even at long time lags.


### Right: Autocorrelation Function (ACF):

- The autocorrelation function (ACF)  $\Gamma_X(\ell)/\Gamma_X(0)$  decreases very slowly as the lag  $\ell$  increases.
- This behavior reflects **long-range dependence (LRD)**, where correlations persist over long time periods.

### Stationarity Implications:

- The slow decay in the ACF suggests that the time series is **not stationary**, as stationarity requires the ACF to decay rapidly (e.g., exponentially or geometrically—decreases by a constant ratio in each time step).


## 1.2 Stationarity: Short-range dependence (SRD)



### Left: Time Series:

- The time series exhibits more erratic behavior compared to LRD. While there are small dependencies between consecutive observations, they dissipate quickly over time.

## 1.2 Stationarity: Short-range dependence (SRD)



### Right: Autocorrelation Function (ACF):


- The ACF decreases rapidly as  $\ell$  increases and eventually becomes close to zero, indicating weak or no correlation at large lags. This is characteristic of **short-range dependence (SRD)**.

### Left: Time Series:

- The time series exhibits more erratic behavior compared to LRD. While there are small dependencies between consecutive observations, they dissipate quickly over time.



## 1.2 Stationarity: Short-range dependence (SRD)



### Left: Time Series:

- The time series exhibits more erratic behavior compared to LRD. While there are small dependencies between consecutive observations, they dissipate quickly over time.


### Right: Autocorrelation Function (ACF):

- The ACF decreases rapidly as  $\ell$  increases and eventually becomes close to zero, indicating weak or no correlation at large lags. This is characteristic of **short-range dependence (SRD)**.
- The rapid decay indicates that the time series is likely stationary, as the ACF does not persist indefinitely.

### Stationarity Implications:

- The rapid decay in the ACF suggests that the time series is likely **stationary**.

## 1.2 Stationarity: Short-range dependence (SRD)



### Left: Time Series:

- The time series exhibits more erratic behavior compared to LRD. While there are small dependencies between consecutive observations, they dissipate quickly over time.


### Right: Autocorrelation Function (ACF):

- The ACF decreases rapidly as  $\ell$  increases and eventually becomes close to zero, indicating weak or no correlation at large lags. This is characteristic of **short-range dependence (SRD)**.
- The rapid decay indicates that the time series is likely stationary, as the ACF does not persist indefinitely.

### Stationarity Implications:

- The rapid decay in the ACF suggests that the time series is likely **stationary**.
- Stationary time series exhibit consistent statistical properties over time.


## 1.2 Stationarity: In Functional Magnetic Resonance Imaging (fMRI).



### Left: Time Series:

- The fMRI signal shows fluctuations over time, typical of neuro-physiological processes, with moderate temporal correlations.

## 1.2 Stationarity: In Functional Magnetic Resonance Imaging (fMRI).




### Right: Autocorrelation Function (ACF):

- The ACF decreases gradually but not as rapidly as in SRD. The slower decay suggests some degree of correlation over time, though it is weaker than in LRD.

### Left: Time Series:

- The fMRI signal shows fluctuations over time, typical of neuro-physiological processes, with moderate temporal correlations.

## 1.2 Stationarity: In Functional Magnetic Resonance Imaging (fMRI).



### Left: Time Series:

- The fMRI signal shows fluctuations over time, typical of neuro-physiological processes, with moderate temporal correlations.


### Right: Autocorrelation Function (ACF):

- The ACF decreases gradually but not as rapidly as in SRD. The slower decay suggests some degree of correlation over time, though it is weaker than in LRD.

### Stationarity Implications:

- The fMRI data may exhibit **weak stationarity**, as the mean and variance appear constant over time.

## 1.2 Stationarity: In Functional Magnetic Resonance Imaging (fMRI).



### Left: Time Series:

- The fMRI signal shows fluctuations over time, typical of neuro-physiological processes, with moderate temporal correlations.


### Right: Autocorrelation Function (ACF):

- The ACF decreases gradually but not as rapidly as in SRD. The slower decay suggests some degree of correlation over time, though it is weaker than in LRD.

### Stationarity Implications:

- The fMRI data may exhibit **weak stationarity**, as the mean and variance appear constant over time.
- However, correlations persist over moderate lags, requiring further investigation to confirm stationarity.


## 1.2 Stationarity: In Stock Prices



### Left: Time Series:

- The stock price time series appears highly volatile, with no clear trend or periodicity.

## 1.2 Stationarity: In Stock Prices




### Left: Time Series:

- The stock price time series appears highly volatile, with no clear trend or periodicity.
- The variations are large and erratic, consistent with typical financial time series.



## 1.2 Stationarity: In Stock Prices




### Right: Autocorrelation Function (ACF):

- The autocorrelation function (ACF)  $\Gamma_X(\ell)/\Gamma_X(0)$  does not decay as rapidly as expected for short-range dependence.

### Left: Time Series:

- The stock price time series appears highly volatile, with no clear trend or periodicity.
- The variations are large and erratic, consistent with typical financial time series.

## 1.2 Stationarity: In Stock Prices




### Right: Autocorrelation Function (ACF):

- The autocorrelation function (ACF)  $\Gamma_X(\ell)/\Gamma_X(0)$  does not decay as rapidly as expected for short-range dependence.
- This behavior suggests that correlations persist over time, indicating potential long-range dependence, weak or even non-stationarity.

### Left: Time Series:

- The stock price time series appears highly volatile, with no clear trend or periodicity.
- The variations are large and erratic, consistent with typical financial time series.

## 1.2 Stationarity: In Stock Prices



### Left: Time Series:

- The stock price time series appears highly volatile, with no clear trend or periodicity.
- The variations are large and erratic, consistent with typical financial time series.


### Right: Autocorrelation Function (ACF):

- The autocorrelation function (ACF)  $\Gamma_X(\ell)/\Gamma_X(0)$  does not decay as rapidly as expected for short-range dependence.
- This behavior suggests that correlations persist over time, indicating potential long-range dependence, weak or even non-stationarity.

### Stationarity Implications:

- The slow decay in the ACF indicates that the time series may be **weakly stationary**.

## 1.2 Stationarity: In Stock Prices



### Left: Time Series:

- The stock price time series appears highly volatile, with no clear trend or periodicity.
- The variations are large and erratic, consistent with typical financial time series.

### Right: Autocorrelation Function (ACF):

- The autocorrelation function (ACF)  $\Gamma_X(\ell)/\Gamma_X(0)$  does not decay as rapidly as expected for short-range dependence.
- This behavior suggests that correlations persist over time, indicating potential long-range dependence, weak or even non-stationarity.

### Stationarity Implications:

- The slow decay in the ACF indicates that the time series may be **weakly stationary**.
- Financial data can exhibit behaviors such as long-range dependence or structural breaks that violate stationarity assumptions.

## 1.2. Key Points on Stationarity and ACF

### 1. Stationarity:

- A time series is stationary if its mean and variance are constant over time, and the autocovariance  $\Gamma_X(\ell)$  depends only on the lag  $\ell$ , not on time  $t$ .

## 1.2. Key Points on Stationarity and ACF

### 1. Stationarity:

- A time series is stationary if its mean and variance are constant over time, and the autocovariance  $\Gamma_X(\ell)$  depends only on the lag  $\ell$ , not on time  $t$ .
- In the figures:
  - SRD, fMRI, and stock data are examples of stationary (or weakly stationary) time series.

## 1.2. Key Points on Stationarity and ACF

### 1. Stationarity:

- A time series is stationary if its mean and variance are constant over time, and the autocovariance  $\Gamma_X(\ell)$  depends only on the lag  $\ell$ , not on time  $t$ .
- In the figures:
  - SRD, fMRI, and stock data are examples of stationary (or weakly stationary) time series.
  - LRD may not be stationary due to the slow decay of the ACF, which suggests that correlations persist across large lags.

### 2. Autocorrelation Function (ACF):

- The ACF quantifies the dependence of the time series on its past values at various lags  $\ell$ .

## 1.2. Key Points on Stationarity and ACF

### 1. Stationarity:

- A time series is stationary if its mean and variance are constant over time, and the autocovariance  $\Gamma_X(\ell)$  depends only on the lag  $\ell$ , not on time  $t$ .
- In the figures:
  - SRD, fMRI, and stock data are examples of stationary (or weakly stationary) time series.
  - LRD may not be stationary due to the slow decay of the ACF, which suggests that correlations persist across large lags.

### 2. Autocorrelation Function (ACF):

- The ACF quantifies the dependence of the time series on its past values at various lags  $\ell$ .
- Stationary time series typically exhibit ACFs that decay rapidly (e.g., exponentially or geometrically) as  $\ell$  increases.



## 1.2. Key Points on Stationarity and ACF

### 1. Stationarity:

- A time series is stationary if its mean and variance are constant over time, and the autocovariance  $\Gamma_X(\ell)$  depends only on the lag  $\ell$ , not on time  $t$ .
- In the figures:
  - SRD, fMRI, and stock data are examples of stationary (or weakly stationary) time series.
  - LRD may not be stationary due to the slow decay of the ACF, which suggests that correlations persist across large lags.

### 2. Autocorrelation Function (ACF):

- The ACF quantifies the dependence of the time series on its past values at various lags  $\ell$ .
- Stationary time series typically exhibit ACFs that decay rapidly (e.g., exponentially or geometrically) as  $\ell$  increases.
- Non-stationary time series may show very slow or no decay in the ACF.

## 1.2. Note1 : The Autocovariance function (ACVF) $\neq$ Autocorrelation function (ACF)

### 1 Autocovariance Function (ACVF):

- ACVF measures the covariance of a time series with its lagged version.

## 1.2. Note1 : The Autocovariance function (ACVF) $\neq$ Autocorrelation function (ACF)

### 1 Autocovariance Function (ACVF):

- ACVF measures the covariance of a time series with its lagged version.
- For a time series  $X_t$ , the ACVF at lag  $\ell$  is defined as:

$$\Gamma_X(\ell) = \text{Cov}(X_t, X_{t+\ell}) = \mathbb{E}[(X_t - \mu)(X_{t+\ell} - \mu)],$$

where  $\mu = \mathbb{E}[X_t]$  is the mean of the time series (assuming stationarity).

## 1.2. Note1 : The Autocovariance function (ACVF) $\neq$ Autocorrelation function (ACF)

### 1 Autocovariance Function (ACVF):

- ACVF measures the covariance of a time series with its lagged version.
- For a time series  $X_t$ , the ACVF at lag  $\ell$  is defined as:

$$\Gamma_X(\ell) = \text{Cov}(X_t, X_{t+\ell}) = \mathbb{E}[(X_t - \mu)(X_{t+\ell} - \mu)],$$

where  $\mu = \mathbb{E}[X_t]$  is the mean of the time series (assuming stationarity).

- **Units:** The autocovariance is measured in the square of the units of  $X_t$ , so it depends on the scale of the time series.

1.2. Note 1: The Autocovariance function (ACVF)  $\neq$  Autocorrelation function (ACF)

## 2 Autocorrelation Function (ACF):

1.2. Note 1: The Autocovariance function (ACVF)  $\neq$  Autocorrelation function (ACF)

## 2 Autocorrelation Function (ACF):

- The ACF is the normalized version of the ACVF.

## 1.2. Note 1: The Autocovariance function (ACVF) $\neq$ Autocorrelation function (ACF)

### 2 Autocorrelation Function (ACF):

- The ACF is the normalized version of the ACVF.
- It measures the linear dependence between  $X_t$  and  $X_{t+h}$ , scaled to lie between  $-1$  and  $1$ . It is defined as:

$$\rho_X(\ell) = \frac{\Gamma_X(\ell)}{\Gamma_X(0)},$$

where  $\Gamma_X(0)$  is the variance of the time series ( $\Gamma_X(0) = \text{Var}(X_t)$ ).

## 1.2. Note 1: The Autocovariance function (ACVF) $\neq$ Autocorrelation function (ACF)

### 2 Autocorrelation Function (ACF):

- The ACF is the normalized version of the ACVF.
- It measures the linear dependence between  $X_t$  and  $X_{t+h}$ , scaled to lie between  $-1$  and  $1$ . It is defined as:

$$\rho_X(\ell) = \frac{\Gamma_X(\ell)}{\Gamma_X(0)},$$

where  $\Gamma_X(0)$  is the variance of the time series ( $\Gamma_X(0) = \text{Var}(X_t)$ ).

- **Units:** The autocorrelation is dimensionless and does not depend on the scale of the time series.



## 1.2 Note 2: Covariance and Variance Relationship at lag $\ell$

The covariance,  $\text{Cov}(X_t, X_{t+\ell}) = \mathbb{E}[(X_t - \mu)(X_{t+\ell} - \mu)]$  will be equal to the variance of the time series,  $\text{Var}(X_t) = \mathbb{E}[(X_t - \mu)^2]$ , when the lag  $\ell = 0$ .

## 1.2 Note 2: Covariance and Variance Relationship at lag $\ell$

The covariance,  $\text{Cov}(X_t, X_{t+\ell}) = \mathbb{E}[(X_t - \mu)(X_{t+\ell} - \mu)]$  will be equal to the variance of the time series,  $\text{Var}(X_t) = \mathbb{E}[(X_t - \mu)^2]$ , when the lag  $\ell = 0$ .

### Explanation:

- **Variance:** The variance of a random variable  $X_t$  is a special case of the covariance where the two random variables are identical, i.e., the lag is  $\ell = 0$ :

$$\text{Var}(X_t) = \text{Cov}(X_t, X_t) = \mathbb{E}[(X_t - \mu)(X_t - \mu)].$$

## 1.2 Note 2: Covariance and Variance Relationship at lag $\ell$

The covariance,  $\text{Cov}(X_t, X_{t+\ell}) = \mathbb{E}[(X_t - \mu)(X_{t+\ell} - \mu)]$  will be equal to the variance of the time series,  $\text{Var}(X_t) = \mathbb{E}[(X_t - \mu)^2]$ , when the lag  $\ell = 0$ .

### Explanation:

- **Variance:** The variance of a random variable  $X_t$  is a special case of the covariance where the two random variables are identical, i.e., the lag is  $\ell = 0$ :

$$\text{Var}(X_t) = \text{Cov}(X_t, X_t) = \mathbb{E}[(X_t - \mu)(X_t - \mu)].$$

- **Covariance at  $\ell = 0$ :** When  $\ell = 0$ , the covariance reduces to:

$$\text{Cov}(X_t, X_{t+0}) = \mathbb{E}[(X_t - \mu)(X_t - \mu)] = \text{Var}(X_t).$$

## 1.2 Note 2: Covariance and Variance Relationship at lag $\ell$

The covariance,  $\text{Cov}(X_t, X_{t+\ell}) = \mathbb{E}[(X_t - \mu)(X_{t+\ell} - \mu)]$  will be equal to the variance of the time series,  $\text{Var}(X_t) = \mathbb{E}[(X_t - \mu)^2]$ , when the lag  $\ell = 0$ .

### Explanation:

- **Variance:** The variance of a random variable  $X_t$  is a special case of the covariance where the two random variables are identical, i.e., the lag is  $\ell = 0$ :

$$\text{Var}(X_t) = \text{Cov}(X_t, X_t) = \mathbb{E}[(X_t - \mu)(X_t - \mu)].$$

- **Covariance at  $\ell = 0$ :** When  $\ell = 0$ , the covariance reduces to:

$$\text{Cov}(X_t, X_{t+0}) = \mathbb{E}[(X_t - \mu)(X_t - \mu)] = \text{Var}(X_t).$$

- For  $\ell \neq 0$ , the covariance involves the dependence between  $X_t$  and its lagged value  $X_{t+\ell}$ . Unless the time series is perfectly correlated for that lag ( $X_{t+\ell} = X_t$ ), the covariance at lag  $\ell$  will **not** equal the variance.

## 1.2 Note 2: Covariance and Variance Relationship at lag $\ell$

The covariance,  $\text{Cov}(X_t, X_{t+\ell}) = \mathbb{E}[(X_t - \mu)(X_{t+\ell} - \mu)]$  will be equal to the variance of the time series,  $\text{Var}(X_t) = \mathbb{E}[(X_t - \mu)^2]$ , when the lag  $\ell = 0$ .

### Explanation:

- **Variance:** The variance of a random variable  $X_t$  is a special case of the covariance where the two random variables are identical, i.e., the lag is  $\ell = 0$ :

$$\text{Var}(X_t) = \text{Cov}(X_t, X_t) = \mathbb{E}[(X_t - \mu)(X_t - \mu)].$$

- **Covariance at  $\ell = 0$ :** When  $\ell = 0$ , the covariance reduces to:

$$\text{Cov}(X_t, X_{t+0}) = \mathbb{E}[(X_t - \mu)(X_t - \mu)] = \text{Var}(X_t).$$

- For  $\ell \neq 0$ , the covariance involves the dependence between  $X_t$  and its lagged value  $X_{t+\ell}$ . Unless the time series is perfectly correlated for that lag ( $X_{t+\ell} = X_t$ ), the covariance at lag  $\ell$  will **not** equal the variance.

**Conclusion:**  $\text{Cov}(X_t, X_{t+\ell}) = \text{Var}(X_t)$  if and only if  $\ell = 0$ .