

Introduction to R and RStudio

The goal of this lab is to introduce you to R and RStudio, which you'll be using throughout the course both to learn the statistical concepts discussed in the textbook and also to analyze real data and come to informed conclusions. To straighten out which is which: R is the name of the programming language itself and RStudio is a convenient interface.

As the labs progress, you are encouraged to explore beyond what the labs dictate; a willingness to experiment will make you a much better programmer. Before we get to that stage, however, you need to build some basic fluency in R. Today we begin with the fundamental building blocks of R and RStudio: the interface, reading in data, and basic commands.

The panel in the upper right contains your *workspace* as well as a history of the commands that you've previously entered. Any plots that you generate will show up in the panel in the lower right corner.

The panel on the left is where the action happens. It's called the *console*. Everytime you launch RStudio, it will have the same text at the top of the console telling you the version of R that you're running. Below that information is the *prompt*. As its name suggests, this prompt is really a request, a request for a command. Initially, interacting with R is all about typing commands and interpreting the output. These commands and their syntax have evolved over decades (literally) and now provide what many users feel is a fairly natural way to access data and organize, describe, and invoke statistical computations.

To get you started, enter the following command at the R prompt (i.e. right after `>` on the console). You can either type it in manually or copy and paste it from this document.

```
source("more/arbuthnot.R")
```

This command instructs R to access the OpenIntro website and fetch some data: the Arbuthnot baptism counts for boys and girls. You should see that the workspace area in the upper righthand corner of the RStudio window now lists a data set called `arbuthnot` that has 82 observations on 3 variables. As you interact with R, you will create a series of objects. Sometimes you load them as we have done here, and sometimes you create them yourself as the byproduct of a computation or some analysis you have performed. Note that because you are accessing data from the web, this command (and the entire assignment) will work in a computer lab, in the library, or in your dorm room; anywhere you have access to the Internet.

The Data: Dr. Arbuthnot's Baptism Records

The Arbuthnot data set refers to Dr. John Arbuthnot, an 18th century physician, writer, and mathematician. He was interested in the ratio of newborn boys to newborn girls, so he gathered the baptism records for children born in London for every year from 1629 to 1710. We can take a look at the data by typing its name into the console.

```
arbuthnot
```

```
##   year boys girls
## 1  1629 5218 4683
## 2  1630 4858 4457
## 3  1631 4422 4102
## 4  1632 4994 4590
## 5  1633 5158 4839
## 6  1634 5035 4820
## 7  1635 5106 4928
## 8  1636 4917 4605
```




Figure 1: rinterface

```
## 9 1637 4703 4457
## 10 1638 5359 4952
## 11 1639 5366 4784
## 12 1640 5518 5332
## 13 1641 5470 5200
## 14 1642 5460 4910
## 15 1643 4793 4617
## 16 1644 4107 3997
## 17 1645 4047 3919
## 18 1646 3768 3395
## 19 1647 3796 3536
## 20 1648 3363 3181
## 21 1649 3079 2746
## 22 1650 2890 2722
## 23 1651 3231 2840
## 24 1652 3220 2908
## 25 1653 3196 2959
## 26 1654 3441 3179
## 27 1655 3655 3349
## 28 1656 3668 3382
## 29 1657 3396 3289
## 30 1658 3157 3013
## 31 1659 3209 2781
## 32 1660 3724 3247
## 33 1661 4748 4107
## 34 1662 5216 4803
## 35 1663 5411 4881
```

```
## 36 1664 6041 5681
## 37 1665 5114 4858
## 38 1666 4678 4319
## 39 1667 5616 5322
## 40 1668 6073 5560
## 41 1669 6506 5829
## 42 1670 6278 5719
## 43 1671 6449 6061
## 44 1672 6443 6120
## 45 1673 6073 5822
## 46 1674 6113 5738
## 47 1675 6058 5717
## 48 1676 6552 5847
## 49 1677 6423 6203
## 50 1678 6568 6033
## 51 1679 6247 6041
## 52 1680 6548 6299
## 53 1681 6822 6533
## 54 1682 6909 6744
## 55 1683 7577 7158
## 56 1684 7575 7127
## 57 1685 7484 7246
## 58 1686 7575 7119
## 59 1687 7737 7214
## 60 1688 7487 7101
## 61 1689 7604 7167
## 62 1690 7909 7302
## 63 1691 7662 7392
## 64 1692 7602 7316
## 65 1693 7676 7483
## 66 1694 6985 6647
## 67 1695 7263 6713
## 68 1696 7632 7229
## 69 1697 8062 7767
## 70 1698 8426 7626
## 71 1699 7911 7452
## 72 1700 7578 7061
## 73 1701 8102 7514
## 74 1702 8031 7656
## 75 1703 7765 7683
## 76 1704 6113 5738
## 77 1705 8366 7779
## 78 1706 7952 7417
## 79 1707 8379 7687
## 80 1708 8239 7623
## 81 1709 7840 7380
## 82 1710 7640 7288
```

What you should see are four columns of numbers, each row representing a different year: the first entry in each row is simply the row number (an index we can use to access the data from individual years if we want), the second is the year, and the third and fourth are the numbers of boys and girls baptized that year, respectively. Use the scrollbar on the right side of the console window to examine the complete data set.

Note that the row numbers in the first column are not part of Arbuthnot's data. R adds them as part of its printout to help you make visual comparisons. You can think of them as the index that you see on the

left side of a spreadsheet. In fact, the comparison to a spreadsheet will generally be helpful. R has stored Arbuthnot's data in a kind of spreadsheet or table called a *data frame*.

You can see the dimensions of this data frame by typing:

```
dim(arbuthnot)
```

```
## [1] 82  3
```

This command should output `[1] 82 3`, indicating that there are 82 rows and 3 columns (we'll get to what the `[1]` means in a bit), just as it says next to the object in your workspace. You can see the names of these columns (or variables) by typing:

```
names(arbuthnot)
```

```
## [1] "year" "boys" "girls"
```

You should see that the data frame contains the columns `year`, `boys`, and `girls`. At this point, you might notice that many of the commands in R look a lot like functions from math class; that is, invoking R commands means supplying a function with some number of arguments. The `dim` and `names` commands, for example, each took a single argument, the name of a data frame.

One advantage of RStudio is that it comes with a built-in data viewer. Click on the name `arbuthnot` in the *Environment* pane (upper right window) that lists the objects in your workspace. This will bring up an alternative display of the data set in the *Data Viewer* (upper left window). You can close the data viewer by clicking on the *x* in the upper lefthand corner.

Some Exploration

Let's start to examine the data a little more closely. We can access the data in a single column of a data frame separately using a command like

```
arbuthnot$boys
```

```
## [1] 5218 4858 4422 4994 5158 5035 5106 4917 4703 5359 5366 5518 5470 5460
## [15] 4793 4107 4047 3768 3796 3363 3079 2890 3231 3220 3196 3441 3655 3668
## [29] 3396 3157 3209 3724 4748 5216 5411 6041 5114 4678 5616 6073 6506 6278
## [43] 6449 6443 6073 6113 6058 6552 6423 6568 6247 6548 6822 6909 7577 7575
## [57] 7484 7575 7737 7487 7604 7909 7662 7602 7676 6985 7263 7632 8062 8426
## [71] 7911 7578 8102 8031 7765 6113 8366 7952 8379 8239 7840 7640
```

This command will only show the number of boys baptized each year.

1. What command would you use to extract just the counts of girls baptized? Try it! **Answer 1. The command I would use to extract just the counts of girls baptized is below**


```
arbuthnot$girls
```

```
## [1] 4683 4457 4102 4590 4839 4820 4928 4605 4457 4952 4784 5332 5200 4910
## [15] 4617 3997 3919 3395 3536 3181 2746 2722 2840 2908 2959 3179 3349 3382
## [29] 3289 3013 2781 3247 4107 4803 4881 5681 4858 4319 5322 5560 5829 5719
## [43] 6061 6120 5822 5738 5717 5847 6203 6033 6041 6299 6533 6744 7158 7127
## [57] 7246 7119 7214 7101 7167 7302 7392 7316 7483 6647 6713 7229 7767 7626
## [71] 7452 7061 7514 7656 7683 5738 7779 7417 7687 7623 7380 7288
```

Notice that the way R has printed these data is different. When we looked at the complete data frame, we saw 82 rows, one on each line of the display. These data are no longer structured in a table with other variables, so they are displayed one right after another. Objects that print out in this way are called *vectors*; they represent a set of numbers. R has added numbers in [brackets] along the left side of the printout to indicate locations within the vector. For example, 5218 follows [1], indicating that 5218 is the first entry in the vector. And if [43] starts a line, then that would mean the first number on that line would represent the 43rd entry in the vector.


R has some powerful functions for making graphics. We can create a simple plot of the number of girls baptized per year with the command

```
plot(x = arbuthnot$year, y = arbuthnot$girls)
```



By default, R creates a scatterplot with each x,y pair indicated by an open circle. The plot itself should appear under the *Plots* tab of the lower right panel of RStudio. Notice that the command above again looks like a function, this time with two arguments separated by a comma. The first argument in the plot function specifies the variable for the x-axis and the second for the y-axis. If we wanted to connect the data points with lines, we could add a third argument, the letter *l* for line.

```
plot(x = arbuthnot$year, y = arbuthnot$girls, type = "l")
```



You might wonder how you are supposed to know that it was possible to add that third argument. Thankfully, R documents all of its functions extensively. To read what a function does and learn the arguments that are available to you, just type in a question mark followed by the name of the function that you're interested in. Try the following.

```
?plot
```

```
## starting httpd help server ... done
```

Notice that the help file replaces the plot in the lower right panel. You can toggle between plots and help files using the tabs at the top of that panel.

2. Is there an apparent trend in the number of girls baptized over the years?
How would you describe it?

Answer 2. Trend in the number of girls baptized over the years? Starting from year ~ 1660, there seems to be a positive linear correlation and trend between years and girls baptized. As the years are passing, the number of girls baptized are increasing starting from year 1660. However it looks like the relationship has changed over the years as prior to year 1660; there wasn't a definite trend between over the years. Between, years 1640 and 1650, there seems to be a negative linear relationship over the years as the amount of girls baptized declines within these years.

Now, suppose we want to plot the total number of baptisms. To compute this, we could use the fact that R is really just a big calculator. We can type in mathematical expressions like

```
5218 + 4683
```

```
## [1] 9901
```

to see the total number of baptisms in 1629. We could repeat this once for each year, but there is a faster way. If we add the vector for baptisms for boys and girls, R will compute all sums simultaneously.

```
arbuthnot$boys + arbuthnot$girls
```

```
## [1] 9901 9315 8524 9584 9997 9855 10034 9522 9160 10311 10150
## [12] 10850 10670 10370 9410 8104 7966 7163 7332 6544 5825 5612
## [23] 6071 6128 6155 6620 7004 7050 6685 6170 5990 6971 8855
## [34] 10019 10292 11722 9972 8997 10938 11633 12335 11997 12510 12563
## [45] 11895 11851 11775 12399 12626 12601 12288 12847 13355 13653 14735
## [56] 14702 14730 14694 14951 14588 14771 15211 15054 14918 15159 13632
## [67] 13976 14861 15829 16052 15363 14639 15616 15687 15448 11851 16145
## [78] 15369 16066 15862 15220 14928
```

What you will see are 82 numbers (in that packed display, because we aren't looking at a data frame here), each one representing the sum we're after. Take a look at a few of them and verify that they are right. Therefore, we can make a plot of the total number of baptisms per year with the command

```
plot(arbuthnot$year, arbuthnot$boys + arbuthnot$girls, type = "l")
```



This time, note that we left out the names of the first two arguments. We can do this because the help file shows that the default for `plot` is for the first argument to be the x-variable and the second argument to be the y-variable.

Similarly to how we computed the proportion of boys, we can compute the ratio of the number of boys to the number of girls baptized in 1629 with

```
5218 / 4683
```

```
## [1] 1.114243
```

or we can act on the complete vectors with the expression

```
arbuthnot$boys / arbuthnot$girls
```

```
## [1] 1.114243 1.089971 1.078011 1.088017 1.065923 1.044606 1.036120
## [8] 1.067752 1.055194 1.082189 1.121656 1.034884 1.051923 1.112016
## [15] 1.038120 1.027521 1.032661 1.109867 1.073529 1.057215 1.121267
## [22] 1.061719 1.137676 1.107290 1.080095 1.082416 1.091371 1.084565
## [29] 1.032533 1.047793 1.153901 1.146905 1.156075 1.085988 1.108584
## [36] 1.063369 1.052697 1.083121 1.055242 1.092266 1.116143 1.097744
## [43] 1.064016 1.052778 1.043112 1.065354 1.059647 1.120575 1.035467
## [50] 1.088679 1.034100 1.039530 1.044237 1.024466 1.058536 1.062860
## [57] 1.032846 1.064054 1.072498 1.054359 1.060974 1.083128 1.036526
## [64] 1.039092 1.025792 1.050850 1.081931 1.055748 1.037981 1.104904
## [71] 1.061594 1.073219 1.078254 1.048981 1.010673 1.065354 1.075460
## [78] 1.072132 1.090022 1.080808 1.062331 1.048299
```

The proportion of newborns that are boys

```
5218 / (5218 + 4683)
```

```
## [1] 0.5270175
```

or this may also be computed for all years simultaneously:

```
arbuthnot$boys / (arbuthnot$boys + arbuthnot$girls)
```


```
## [1] 0.5270175 0.5215244 0.5187705 0.5210768 0.5159548 0.5109082 0.5088698
## [8] 0.5163831 0.5134279 0.5197362 0.5286700 0.5085714 0.5126523 0.5265188
## [15] 0.5093518 0.5067868 0.5080341 0.5260366 0.5177305 0.5139059 0.5285837
## [22] 0.5149679 0.5322023 0.5254569 0.5192526 0.5197885 0.5218447 0.5202837
## [29] 0.5080030 0.5116694 0.5357262 0.5342132 0.5361942 0.5206108 0.5257482
## [36] 0.5153557 0.5128359 0.5199511 0.5134394 0.5220493 0.5274422 0.5232975
## [43] 0.5155076 0.5128552 0.5105507 0.5158214 0.5144798 0.5284297 0.5087122
## [50] 0.5212285 0.5083822 0.5096910 0.5108199 0.5060426 0.5142178 0.5152360
## [57] 0.5080788 0.5155165 0.5174905 0.5132301 0.5147925 0.5199527 0.5089677
## [64] 0.5095857 0.5063659 0.5123973 0.5196766 0.5135590 0.5093183 0.5249190
## [71] 0.5149385 0.5176583 0.5188268 0.5119526 0.5026541 0.5158214 0.5181790
## [78] 0.5174052 0.5215362 0.5194175 0.5151117 0.5117899
```


Note that with R as with your calculator, you need to be conscious of the order of operations. Here, we want to divide the number of boys by the total number of newborns, so we have to use parentheses. Without them, R will first do the division, then the addition, giving you something that is not a proportion.

- Now, make a plot of the proportion of boys over time. What do you see? Tip: If you use the up and down arrow keys, you can scroll through your previous commands, your so-called command history. You can also access it by clicking on the history tab in the upper right panel. This will save you a lot of typing in the future.

Answer 3. Proportion of Boys over time

```
plot(arbuthnot$year, arbuthnot$boys / (arbuthnot$boys + arbuthnot$girls), type = "l")
```



Finally, in addition to simple mathematical operators like subtraction and division, you can ask R to make comparisons like greater than, >, less than, <, and equality, ==. For example, we can ask if boys outnumber girls in each year with the expression

```
arbuthnot$boys > arbuthnot$girls
```

```
## [1] TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE
## [15] TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE
## [29] TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE
## [43] TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE
## [57] TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE
## [71] TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE TRUE
```

This command returns 82 values of either **TRUE** if that year had more boys than girls, or **FALSE** if that year did not (the answer may surprise you). This output shows a different kind of data than we have considered so far. In the **arbuthnot** data frame our values are numerical (the year, the number of boys and girls). Here, we've asked R to create *logical* data, data where the values are either **TRUE** or **FALSE**. In general, data analysis will involve many different kinds of data types, and one reason for using R is that it is able to represent and compute with many of them.

This seems like a fair bit for your first lab, so let's stop here. To exit RStudio you can click the *x* in the upper right corner of the whole window.

You will be prompted to save your workspace. If you click *save*, RStudio will save the history of your commands and all the objects in your workspace so that the next time you launch RStudio, you will see **arbuthnot** and you will have access to the commands you typed in your previous session. For now, click *save*, then start up RStudio again.

On Your Own

In the previous few pages, you recreated some of the displays and preliminary analysis of Arbuthnot's baptism data. Your assignment involves repeating these steps, but for present day birth records in the United States. Load up the present day data with the following command.

```
source("more/present.R")
```

Answers. Last Section

First view the dataset.

```
present
```

```
##   year   boys  girls
## 1  1940 1211684 1148715
## 2  1941 1289734 1223693
## 3  1942 1444365 1364631
## 4  1943 1508959 1427901
## 5  1944 1435301 1359499
## 6  1945 1404587 1330869
## 7  1946 1691220 1597452
## 8  1947 1899876 1800064
## 9  1948 1813852 1721216
## 10 1949 1826352 1733177
## 11 1950 1823555 1730594
## 12 1951 1923020 1827830
## 13 1952 1971262 1875724
## 14 1953 2001798 1900322
## 15 1954 2059068 1958294
## 16 1955 2073719 1973576
## 17 1956 2133588 2029502
## 18 1957 2179960 2074824
## 19 1958 2152546 2051266
## 20 1959 2173638 2071158
## 21 1960 2179708 2078142
## 22 1961 2186274 2082052
```

```
## 23 1962 2132466 2034896
## 24 1963 2101632 1996388
## 25 1964 2060162 1967328
## 26 1965 1927054 1833304
## 27 1966 1845862 1760412
## 28 1967 1803388 1717571
## 29 1968 1796326 1705238
## 30 1969 1846572 1753634
## 31 1970 1915378 1816008
## 32 1971 1822910 1733060
## 33 1972 1669927 1588484
## 34 1973 1608326 1528639
## 35 1974 1622114 1537844
## 36 1975 1613135 1531063
## 37 1976 1624436 1543352
## 38 1977 1705916 1620716
## 39 1978 1709394 1623885
## 40 1979 1791267 1703131
## 41 1980 1852616 1759642
## 42 1981 1860272 1768966
## 43 1982 1885676 1794861
## 44 1983 1865553 1773380
## 45 1984 1879490 1789651
## 46 1985 1927983 1832578
## 47 1986 1924868 1831679
## 48 1987 1951153 1858241
## 49 1988 2002424 1907086
## 50 1989 2069490 1971468
## 51 1990 2129495 2028717
## 52 1991 2101518 2009389
## 53 1992 2082097 1982917
## 54 1993 2048861 1951379
## 55 1994 2022589 1930178
## 56 1995 1996355 1903234
## 57 1996 1990480 1901014
## 58 1997 1985596 1895298
## 59 1998 2016205 1925348
## 60 1999 2026854 1932563
## 61 2000 2076969 1981845
## 62 2001 2057922 1968011
## 63 2002 2057979 1963747
```

The data are stored in a data frame called **present**.

- What years are included in this data set? What are the dimensions of the data frame and what are the variable or column names?

Answer

The years included in the data set is below - 1940 to 2002-

```
present$year
```

```
## [1] 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953
```

```
## [15] 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967
## [29] 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981
## [43] 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995
## [57] 1996 1997 1998 1999 2000 2001 2002
```

Dimensions of the data frame is 63 rows and 3 columns

```
dim(present)
```

```
## [1] 63  3
```

Column names are , “years”, “boys”, “girls”

```
names(present)
```

```
## [1] "year" "boys" "girls"
```

- How do these counts compare to Arbuthnot’s? Are they on a similar scale?

Answer


Yes they are. The arbuthnot’s data set had 82 rows with 3 columns, present data set has the same amount of columns, with same column values with 63 rows. They are similar scale.

- Make a plot that displays the boy-to-girl ratio for every year in the data set. What do you see? Does Arbuthnot’s observation about boys being born in greater proportion than girls hold up in the U.S.? Include the plot in your response.

Answer


Present data set boy to girl ratio plot below.

```
plot(present$year, (present$boys / present$girls), type = "l")
```



Arbuthnot data set boy to girl ration plot below.

```
plot(arbuthnot$year, (arbuthnot$boys / arbuthnot$girls), type = "l")
```



What do you see? Does Arbuthnot's observation about boys being born in greater proportion than girls hold up in the U.S.? Include the plot in your response.

Comparing the two plots, we see that both Arbuthnot's observation and Present data set has boys being born in greater proportions than girls. See two plots.

- In what year did we see the most total number of births in the U.S.? You can refer to the help files or the R reference card <http://cran.r-project.org/doc/contrib/Short-refcard.pdf> to find helpful commands.

Answer

First we can add the total births per year and create the total column.

```
total <- present$boys + present$girls
cbind(present, total)
```

```
##   year   boys   girls  total
## 1  1940 1211684 1148715 2360399
## 2  1941 1289734 1223693 2513427
## 3  1942 1444365 1364631 2808996
## 4  1943 1508959 1427901 2936860
## 5  1944 1435301 1359499 2794800
## 6  1945 1404587 1330869 2735456
## 7  1946 1691220 1597452 3288672
## 8  1947 1899876 1800064 3699940
## 9  1948 1813852 1721216 3535068
```

10 1949 1826352 1733177 3559529
 ## 11 1950 1823555 1730594 3554149
 ## 12 1951 1923020 1827830 3750850
 ## 13 1952 1971262 1875724 3846986
 ## 14 1953 2001798 1900322 3902120
 ## 15 1954 2059068 1958294 4017362
 ## 16 1955 2073719 1973576 4047295
 ## 17 1956 2133588 2029502 4163090
 ## 18 1957 2179960 2074824 4254784
 ## 19 1958 2152546 2051266 4203812
 ## 20 1959 2173638 2071158 4244796
 ## 21 1960 2179708 2078142 4257850
 ## 22 1961 2186274 2082052 4268326
 ## 23 1962 2132466 2034896 4167362
 ## 24 1963 2101632 1996388 4098020
 ## 25 1964 2060162 1967328 4027490
 ## 26 1965 1927054 1833304 3760358
 ## 27 1966 1845862 1760412 3606274
 ## 28 1967 1803388 1717571 3520959
 ## 29 1968 1796326 1705238 3501564
 ## 30 1969 1846572 1753634 3600206
 ## 31 1970 1915378 1816008 3731386
 ## 32 1971 1822910 1733060 3555970
 ## 33 1972 1669927 1588484 3258411
 ## 34 1973 1608326 1528639 3136965
 ## 35 1974 1622114 1537844 3159958
 ## 36 1975 1613135 1531063 3144198
 ## 37 1976 1624436 1543352 3167788
 ## 38 1977 1705916 1620716 3326632
 ## 39 1978 1709394 1623885 3333279
 ## 40 1979 1791267 1703131 3494398
 ## 41 1980 1852616 1759642 3612258
 ## 42 1981 1860272 1768966 3629238
 ## 43 1982 1885676 1794861 3680537
 ## 44 1983 1865553 1773380 3638933
 ## 45 1984 1879490 1789651 3669141
 ## 46 1985 1927983 1832578 3760561
 ## 47 1986 1924868 1831679 3756547
 ## 48 1987 1951153 1858241 3809394
 ## 49 1988 2002424 1907086 3909510
 ## 50 1989 2069490 1971468 4040958
 ## 51 1990 2129495 2028717 4158212
 ## 52 1991 2101518 2009389 4110907
 ## 53 1992 2082097 1982917 4065014
 ## 54 1993 2048861 1951379 4000240
 ## 55 1994 2022589 1930178 3952767
 ## 56 1995 1996355 1903234 3899589
 ## 57 1996 1990480 1901014 3891494
 ## 58 1997 1985596 1895298 3880894
 ## 59 1998 2016205 1925348 3941553
 ## 60 1999 2026854 1932563 3959417
 ## 61 2000 2076969 1981845 4058814
 ## 62 2001 2057922 1968011 4025933
 ## 63 2002 2057979 1963747 4021726

present

##	year	boys	girls
## 1	1940	1211684	1148715
## 2	1941	1289734	1223693
## 3	1942	1444365	1364631
## 4	1943	1508959	1427901
## 5	1944	1435301	1359499
## 6	1945	1404587	1330869
## 7	1946	1691220	1597452
## 8	1947	1899876	1800064
## 9	1948	1813852	1721216
## 10	1949	1826352	1733177
## 11	1950	1823555	1730594
## 12	1951	1923020	1827830
## 13	1952	1971262	1875724
## 14	1953	2001798	1900322
## 15	1954	2059068	1958294
## 16	1955	2073719	1973576
## 17	1956	2133588	2029502
## 18	1957	2179960	2074824
## 19	1958	2152546	2051266
## 20	1959	2173638	2071158
## 21	1960	2179708	2078142
## 22	1961	2186274	2082052
## 23	1962	2132466	2034896
## 24	1963	2101632	1996388
## 25	1964	2060162	1967328
## 26	1965	1927054	1833304
## 27	1966	1845862	1760412
## 28	1967	1803388	1717571
## 29	1968	1796326	1705238
## 30	1969	1846572	1753634
## 31	1970	1915378	1816008
## 32	1971	1822910	1733060
## 33	1972	1669927	1588484
## 34	1973	1608326	1528639
## 35	1974	1622114	1537844
## 36	1975	1613135	1531063
## 37	1976	1624436	1543352
## 38	1977	1705916	1620716
## 39	1978	1709394	1623885
## 40	1979	1791267	1703131
## 41	1980	1852616	1759642
## 42	1981	1860272	1768966
## 43	1982	1885676	1794861
## 44	1983	1865553	1773380
## 45	1984	1879490	1789651
## 46	1985	1927983	1832578
## 47	1986	1924868	1831679
## 48	1987	1951153	1858241
## 49	1988	2002424	1907086
## 50	1989	2069490	1971468


```
## 51 1990 2129495 2028717
## 52 1991 2101518 2009389
## 53 1992 2082097 1982917
## 54 1993 2048861 1951379
## 55 1994 2022589 1930178
## 56 1995 1996355 1903234
## 57 1996 1990480 1901014
## 58 1997 1985596 1895298
## 59 1998 2016205 1925348
## 60 1999 2026854 1932563
## 61 2000 2076969 1981845
## 62 2001 2057922 1968011
## 63 2002 2057979 1963747
```

We can further filter the present data set to see the row that has the max birth. That is year 1961.

```
filter(present, total==max(present$total))
```

```
## Warning in max(present$total): no non-missing arguments to max; returning -
## Inf
```

```
## Time Series:
## Start = 1
## End = 63
## Frequency = 1
##      [,1] [,2] [,3]
##  1    NA   NA   NA
##  2    NA   NA   NA
##  3    NA   NA   NA
##  4    NA   NA   NA
##  5    NA   NA   NA
##  6    NA   NA   NA
##  7    NA   NA   NA
##  8    NA   NA   NA
##  9    NA   NA   NA
## 10    NA   NA   NA
## 11    NA   NA   NA
## 12    NA   NA   NA
## 13    NA   NA   NA
## 14    NA   NA   NA
## 15    NA   NA   NA
## 16    NA   NA   NA
## 17    NA   NA   NA
## 18    NA   NA   NA
## 19    NA   NA   NA
## 20    NA   NA   NA
## 21    NA   NA   NA
## 22    NA   NA   NA
## 23    NA   NA   NA
## 24    NA   NA   NA
## 25    NA   NA   NA
## 26    NA   NA   NA
## 27    NA   NA   NA
```

```
## 28    NA    NA    NA
## 29    NA    NA    NA
## 30    NA    NA    NA
## 31    NA    NA    NA
## 32     0     0     0
## 33    NA    NA    NA
## 34    NA    NA    NA
## 35    NA    NA    NA
## 36    NA    NA    NA
## 37    NA    NA    NA
## 38    NA    NA    NA
## 39    NA    NA    NA
## 40    NA    NA    NA
## 41    NA    NA    NA
## 42    NA    NA    NA
## 43    NA    NA    NA
## 44    NA    NA    NA
## 45    NA    NA    NA
## 46    NA    NA    NA
## 47    NA    NA    NA
## 48    NA    NA    NA
## 49    NA    NA    NA
## 50    NA    NA    NA
## 51    NA    NA    NA
## 52    NA    NA    NA
## 53    NA    NA    NA
## 54    NA    NA    NA
## 55    NA    NA    NA
## 56    NA    NA    NA
## 57    NA    NA    NA
## 58    NA    NA    NA
## 59    NA    NA    NA
## 60    NA    NA    NA
## 61    NA    NA    NA
## 62    NA    NA    NA
## 63    NA    NA    NA
```

These data come from a report by the Centers for Disease Control http://www.cdc.gov/nchs/data/nvsr/nvsr53/nvsr53_20.pdf. Check it out if you would like to read more about an analysis of sex ratios at birth in the United States.

That was a short introduction to R and RStudio, but we will provide you with more functions and a more complete sense of the language as the course progresses. Feel free to browse around the websites for R and RStudio if you're interested in learning more, or find more labs for practice at <http://openintro.org>.