

From Bellissard’s gap labeling to atomistic quasiperiodic bilayers

Thomas D. Kühne^{*1,2,3}, Vladislav Efremkin^{1,2}, and Emil Prodan⁴

¹Center for Advanced Systems Understanding (CASUS), Conrad-Schiedt-Straße 20, 02826 Görlitz, Germany

²Helmholtz-Zentrum Dresden-Rossendorf (HZDR), Bautzner Landstraße 400, 01328 Dresden, Germany

³Institute of Artificial Intelligence, Technische Universität Dresden, Helmholtzstraße 10, 01069 Dresden, Germany

⁴Department of Physics, Yeshiva University, New York, New York 10016, USA

Abstract

Bellissard’s formulation of solid-state physics replaces periodicity as the organizing principle by an algebra of covariant observables. We use this viewpoint to connect butterfly-like spectra in quasiperiodic bilayers to the quantities required for gap labeling and topological response. The route has three stages: resolve a bulk spectral gap, count states through its spectral projector, and evaluate the topological pairing relevant to the proposed response. A finite honeycomb-bilayer model makes the first diagnostic steps concrete through spectral moments, angle-resolved spectra, and an uncoupled-layer reference. Selected-angle size and broadening tests distinguish persistent coupling-induced redistribution from resolution-sensitive DOS structure. For atomistic calculations, we explain how nonorthogonal basis sets, layer projections, and moving basis functions enter the construction of physically consistent projectors and parameter derivatives. The resulting framework turns a spectral observation into a sequence of testable calculations linking geometry, state counting, and response.

1 Bellissard’s viewpoint beyond periodic crystals

Bloch theory organizes the electronic structure of a periodic crystal through a family of finite matrices over the Brillouin zone. Quasiperiodic structures retain reproducible local environments without possessing a common translation cell. The loss of a Brillouin zone does not, however, imply the loss of a bulk description or of quantized observables. A central contribution of Bellissard was to formulate these questions in terms of an algebra of physical observables and its K -theory [1]. The noncommutative treatment of the quantum Hall effect [2] and subsequent gap-labeling results for aperiodic patterns [3] show why this shift is more than a replacement of notation.

The key object is not an isolated finite Hamiltonian matrix. It is a covariant family of operators, together with a prescription for bulk averaging. A spectral gap then defines a projection in the observable algebra. Its class can be paired with a trace to count states and, with additional structure, with Chern cocycles to describe transport. These pairings answer related but distinct questions. In particular, a recognizable spectral pattern is neither a state count nor a measurement of a topological response.

Quasiperiodic photonic and acoustic systems make this distinction concrete. Phason-dependent pumping and boundary modes provide information beyond the density of states [4, 5].

^{*}Corresponding author: tkuehne@cp2k.org

Twisted mechanical structures extend this connection to geometric incommensurability in two dimensions [6]. Electronic bilayers introduce further ingredients, including multiple orbitals, self-consistency, structural relaxation, and nonorthogonal basis functions. Their analysis benefits from keeping the same mathematical questions visible throughout the modeling chain.

This article develops that connection in three stages. We first identify the bulk objects entering gap labeling and distinguish phasons from changes of twist angle. We then use finite bilayer model calculations to separate coupling-induced spectral reconstruction from topology. Finally, we translate the operator formulation into quantities accessible to atomistic electronic-structure codes. The discussion combines established theory with an illustrative model analysis.

2 From a family of patterns to a bulk state count

2.1 Covariance and the trace

Consider first a lattice description with q orthonormal internal orbitals per reference cell. Let Ω be a compact space of patterns with a translation action $T_{\mathbf{n}}$, $\mathbf{n} \in \mathbb{Z}^2$. A family of bounded, self-adjoint Hamiltonians $\hat{H}_\omega$ on $\ell^2(\mathbb{Z}^2) \otimes \mathbb{C}^q$ is covariant if

$$U_{\mathbf{n}} \hat{H}_\omega U_{\mathbf{n}}^{-1} = \hat{H}_{T_{\mathbf{n}}\omega}. \quad (1)$$

Locality, continuity in the pattern, and covariance specify the observable algebra $\mathcal{A}$. Finite-range or sufficiently decaying hopping is a useful setting for this construction. The translation convention in Eq. (1) fixes the corresponding convention for the action on patterns.

For an invariant ergodic probability measure μ on Ω , the bulk trace is

$$\tau(A) = \int_{\Omega} \text{tr}_q \langle \mathbf{0} | A_\omega | \mathbf{0} \rangle d\mu(\omega) = \lim_{R \rightarrow \infty} \frac{\text{Tr}(\chi_R A \chi_R)}{|\Lambda_R|}. \quad (2)$$

Here χ_R selects expanding square reference-cell regions Λ_R , whose boundary-to-volume ratio vanishes. Under the stated ergodic assumptions the limit holds for almost every pattern, and $\tau(\mathbf{1}) = q$. The normalization is states per reference cell, not states per plotted DOS peak. Unique ergodicity can strengthen the independence from the chosen pattern under the appropriate continuity assumptions.

Equation (2) is a transparent lattice example, not a universal identification of a multilayer atomic structure with $\mathbb{Z}^2$. In a bilayer with independently varying site sets, an appropriate configuration-space or groupoid description may be required. A trace per physical area then has units of inverse area. General aperiodic frameworks address precisely the absence of a canonical global lattice indexing [7]. The choice of algebra and the normalization of its trace must precede a numerical assignment of gap labels.

2.2 The projector and its label

Suppose E lies in a common spectral gap of the bulk family. Functional calculus gives a spectral projection

$$P_E = \mathbf{1}_{(-\infty, E]}(\hat{H}), \quad P_E^2 = P_E = P_E^*, \quad \mathcal{N}(E) = \tau(P_E). \quad (3)$$

The integrated density of states (IDS), $\mathcal{N}(E)$, is constant as E moves within the gap. Because the discontinuity of the indicator function lies outside the spectrum, P_E belongs to the observable algebra. It defines a class in $K_0(\mathcal{A})$, and the trace induces the homomorphism

$$\tau_* : K_0(\mathcal{A}) \longrightarrow \mathbb{R}, \quad [P] \longmapsto \tau(P). \quad (4)$$

Bellissard’s gap-labeling viewpoint constrains the possible values of the IDS through the range of this map [1, 3]. It does not assert that every permitted label corresponds to an open gap. Nor is the trace pairing generally injective. Distinct topological classes can share a state count, so the IDS need not determine all response coefficients.

A familiar illustration is the one-dimensional quasiperiodic Harper family

$$(\hat{H}_\phi \psi)_n = t(\psi_{n+1} + \psi_{n-1}) + V \cos(2\pi\alpha n + \phi)\psi_n, \quad (5)$$

with irrational α and one orbital per site. For an actual spectral gap, the trace range gives

$$\mathcal{N}(E) = m + n\alpha \in [0, 1], \quad m, n \in \mathbb{Z}. \quad (6)$$

The spectral organization is closely related to the magnetic problem underlying the Hofstadter butterfly [8]. For a rational approximant $\alpha = p/q$ and a gap above r bands, the corresponding counting relation is $r = mq + np$. One such rational equation alone does not uniquely determine an integer pair. Continuation across the family or an independent topological calculation supplies information that an isolated finite spectrum lacks. The sign relating n to a pumping convention also depends on the orientation chosen for the phason cycle.

3 Phasons, twist, and the meaning of a response

At fixed twist angle and layer separation, a rigid relative displacement of two ideal infinite layers changes their registry. Observed from one reference layer, the displacement modulo the other layer’s lattice provides two phason coordinates. In an incommensurate setting, reference-layer translations can sample this phason torus densely. This geometric construction motivates a covariant description, but the Hamiltonian’s orbital content, relaxation rule, and locality still have to respect it.

A phason cycle varies registry within a specified family. Changing the twist angle generally changes the translation action itself, while changing separation changes the strength and range of interlayer coupling. These are useful controls but are not interchangeable coordinates. In particular, an angle-dependent DOS image is not an adiabatic pumping experiment.

The distinction is explicit for a periodic approximant to a one-dimensional pump. Let $P(k, \phi)$ be a smooth occupied-band projector, gapped on the full two-torus of dimensionless k and ϕ , both of period 2π . With a fixed orientation, its first Chern number is

$$C = \frac{i}{2\pi} \int_0^{2\pi} dk \int_0^{2\pi} d\phi \operatorname{tr}(P[\partial_k P, \partial_\phi P]) \in \mathbb{Z}. \quad (7)$$

For a filled, noninteracting band subspace in the adiabatic limit, this integer determines transported particle number, with the sign set by the orientation and current convention [9]. Quasiperiodic systems implement related synthetic-dimensional responses [4, 10], and sliding moiré patterns provide bilayer examples [11, 12].

For suitable two-dimensional twisted models, spatial translations and two phason coordinates lead to a higher-dimensional noncommutative algebra [6]. This is a statement about the model’s covariance structure, not an automatic consequence of having two atomic sheets. Extensions of gap labeling to multilayers likewise depend on the specified effective description [13]. A response involving a spatial direction and a phason must also be distinguished from an ordinary Hall response involving two physical spatial directions. A real, time-reversal-symmetric model may have vanishing ordinary Hall conductivity while allowing a nontrivial mixed spatial-phason pump.

The spectral-gap setting is deliberately the starting point here. Quantization can also survive in a mobility gap, under suitable localization and regularity conditions [2, 7]. A mobility gap is

not an interval of necessarily vanishing DOS. Inferring it requires localization information that a dark region in a broadened spectrum cannot provide.

4 What a finite bilayer model can establish

4.1 A geometric coupling parameter

Consider two identical honeycomb disks. With nearest-neighbor distance a , primitive vectors $\mathbf{a}_1 = (\sqrt{3}a, 0)$ and $\mathbf{a}_2 = (\sqrt{3}a/2, 3a/2)$, and basis positions $(0, 0)$ and $(0, a)$, retain the N sites inside radius R . Place one disk at height zero and the other, rotated by θ , at height d . On orthonormal site states define

$$\hat{H}_{\text{TB}} = \sum_{i \neq j} t_0 \exp(-r_{ij}/\xi) |i\rangle\langle j|, \quad t_0 > 0, \quad \xi > 0. \quad (8)$$

All pairs are included, the on-site term is zero, and r_{ij} is the three-dimensional distance. This is an isotropic geometric model. It is not an orbital-resolved parametrization of graphene and does not resolve the angular dependence of carbon bonding.

The nearest-neighbor intralayer hopping $t_{\text{nn}} = t_0 \exp(-a/\xi)$ provides an exact energy scale. The matrix of $\hat{h} = \hat{H}_{\text{TB}}/t_{\text{nn}}$ is

$$\mathbf{h} = \begin{pmatrix} \mathbf{D} & \mathbf{B} \\ \mathbf{B}^\dagger & \mathbf{D} \end{pmatrix}, \quad D_{ij} = (1 - \delta_{ij})e^{(a-|r_i-r_j|)/\xi}, \quad B_{ij} = e^{(a-\sqrt{|r_i-\mathbf{R}_\theta r_j|^2+d^2})/\xi}. \quad (9)$$

The matrices are representations in the specified finite site basis, whereas $\hat{h}$ denotes the operator. The largest possible interlayer-to-intralayer hopping ratio is

$$\frac{t_{\perp}^{\text{max}}}{t_{\text{nn}}} = e^{(a-d)/\xi}. \quad (10)$$

It is attained by vertically aligned sites, not by every pair. Thus a small change of d relative to the decay length ξ can substantially alter hybridization. For $\xi/a = 0.03$ and $d/a = 0.99, 0.98, 0.97$, this ratio is approximately 1.40, 1.95, 2.72. These dimensionless separations have no prescribed conversion to a first-principles interlayer distance.

4.2 A trace identity and its numerical illustration

The normalized second spectral moment of the finite matrix is

$$\mu_2 = \frac{1}{2N} \sum_{\nu=1}^{2N} \epsilon_\nu^2 = \frac{\text{Tr } \mathbf{h}^2}{2N} = \underbrace{\frac{\text{Tr } \mathbf{D}^2}{N}}_{\text{intralayer}} + \underbrace{\frac{\|\mathbf{B}\|_F^2}{N}}_{\text{interlayer}}, \quad (11)$$

where ϵ_ν are dimensionless eigenvalues and $\|\cdot\|_F$ is the Frobenius norm. The first moment vanishes because the diagonal of $\mathbf{h}$ is zero, so μ_2 is also its spectral variance. Equation (11) is an exact matrix identity. It isolates the added spectral variance due to interlayer coupling without requiring a visual interpretation of a DOS map.

Figure 1 evaluates this decomposition for $R/a = 25$ and $\xi/a = 0.03$. The intralayer contribution is identical for all angles and separations. The interlayer contribution depends on registry as well as separation, and increases when d is reduced at fixed angle because every entry of $\mathbf{B}$ increases. Direct sums of squared hoppings and second moments of the archived eigenvalues agree to an absolute tolerance of 10^{-10} across all 123 spectra, validating the finite-matrix trace normalization and spectral calculation.




Figure 1: Geometric coupling and a finite spectral trace. (a) The maximum interlayer hopping relative to nearest-neighbor intralayer hopping, Eq. (10), for $\xi/a = 0.03$. Colored points mark the three calculated separations. (b) The interlayer contribution $\|\mathbf{B}\|_F^2/N$ to the second spectral moment for open honeycomb disks with $R/a = 25$ and $N = 1510$ sites per layer. Each curve contains 41 directly evaluated angles from 0° to 60° , with lines connecting the samples. The same quantity is independently recovered from all 3020 eigenvalues using Eq. (11).

The relation to Bellissard’s viewpoint is the distinction between a finite normalized trace and a bulk trace on a specified algebra. A second moment averages a polynomial of the Hamiltonian. Gap labeling instead concerns the trace of a spectral projection, whose existence as a stable bulk object requires a gap. A growing μ_2 establishes a growing spectral variance but does not imply a new gap, a fractal spectrum, or a nonzero Chern number. Even fully converged spectral moments would not replace the projector and response calculations in Eqs. (3) and (7).

The disk geometry makes the finite calculation reproducible and treats the two layers symmetrically. Its open boundaries motivate the size checks at representative registries below. Spectral information and projector information remain distinct: equality of two DOS maps does not establish equality of their projectors, since the eigenvectors contain information that the DOS discards.

4.3 From butterfly spectra to state counting

Spectral moments summarize the redistribution of eigenvalues, whereas an angle-resolved spectral map reveals its internal structure. Figure 2 resolves this structure at two interlayer separations using 241 angles between 0° and 60° . The same dimensionless Gaussian width, $\eta/t_{\text{nn}} = 0.02$, and the same color scale are used throughout. Decreasing the separation strengthens the interlayer coupling and reorganizes the curved spectral branches. These maps give an energy-resolved view of the coupling contribution isolated by Eq. (11).

The corresponding finite state count is

$$F_R(\epsilon; \theta) = \frac{1}{2N} \sum_{\nu=1}^{2N} \mathbf{1}_{\{\epsilon_\nu(\theta) \leq \epsilon\}} = \frac{1}{2N} \text{Tr} \mathbf{1}_{(-\infty, \epsilon]}(\mathbf{h}_\theta). \quad (12)$$

Here the normalization is per site state, so F_R runs from zero to one. At a fixed angle, the DOS and F_R display complementary aspects of the same eigenvalues: spectral peaks describe where states accumulate, while horizontal intervals of F_R identify intervals containing no finite-system eigenvalues. The 30° cross sections in Fig. 2 make this connection explicit. Gaussian broadening is applied to the displayed DOS, while the state count is evaluated directly from the unbroadened eigenvalues.




Figure 2: Butterfly-like spectral reconstruction and finite state counting in the bilayer model. (a,b) DOS maps for $d/a = 0.99$ and 0.97 , with $R/a = 25$, $\xi/a = 0.03$, and 1510 sites in each open disk. The angle spacing is 0.25° . The DOS integrates to one and uses a common Gaussian width $\eta/t_{nn} = 0.02$ and color scale, clipped above $t_{nn}\rho_\eta = 0.5$. White dashed lines identify the 30° cross sections. (c) DOS at that angle, without intensity clipping. (d) The exact normalized eigenvalue count, Eq. (12), with an upper-energy inset. Black dashed curves in (c,d) give the corresponding uncoupled-layer reference, $B = 0$, with the same geometry and per-state normalization.

Equation (12) supplies a concrete finite-dimensional counterpart of the spectral-projector trace in Eq. (3). Passing to that bulk quantity entails increasing R , controlling the boundary contribution, and converting the per-state normalization to the cell or area normalization of the chosen trace. Bellissard’s gap labels constrain the resulting bulk state count. The butterfly map, the counting function, and the convergence procedure thus play distinct, complementary roles.

4.4 Uncoupled reference and resolution controls

Setting $B = 0$ in Eq. (9) gives $\mathbf{h}_0 = \mathbf{D} \oplus \mathbf{D}$ without changing the intralayer geometry or energy scale. Every monolayer eigenvalue then occurs twice, so division by $2N$ yields exactly the single-layer DOS and counting function. The reference is independent of θ and d . Its comparison with the coupled spectra in Fig. 2(c,d) isolates interlayer-induced redistribution at a common energy origin.

We test $R/a = 20, 25, 30$, containing $N = 979, 1510, 2167$ sites per layer, at $\theta = 0^\circ, 15^\circ, 30^\circ$ and both plotted separations. These angles sample aligned, intermediate, and 30° registries. Gaussian widths $s = \eta/t_{nn} = 0.01, 0.02, 0.04$ bracket the displayed value. Two complementary measures quantify size sensitivity: $\mathcal{W}_{R,R'} = \int |F_R - F_{R'}| d\varepsilon$ uses the exact unbroadened state

counts, while $\mathcal{L}_s = \int |\bar{\rho}_{R,s} - \bar{\rho}_{R',s}| d\epsilon$ compares the normalized DOS, with $\bar{\rho}_s(\epsilon) = t_{nn}\rho_\eta(t_{nn}\epsilon)$. The latter integral is evaluated on a grid of spacing 0.0025 over $[-7, 7]$, which contains the spectra and Gaussian tails to numerical precision.

Table 1: Finite-size and broadening controls. Each entry is the maximum over the six angle-distance combinations at fixed $\xi/a = 0.03$. The count distance $\mathcal{W}$ is in dimensionless energy units. The DOS difference $\mathcal{L}_s$ lies between zero and two.

$R/a \rightarrow R'/a$	$\max \mathcal{W}$	$\max \mathcal{L}_{0.01}$	$\max \mathcal{L}_{0.02}$	$\max \mathcal{L}_{0.04}$
20 $\rightarrow$ 25	0.01592	0.2971	0.1465	0.0872
25 $\rightarrow$ 30	0.00919	0.2083	0.1114	0.0668

The size differences decrease for every sampled angle and separation, both in $\mathcal{W}$ and at each fixed width. Smoothing reduces the DOS difference further, as expected, so it is assessed alongside the unbroadened count rather than as a separate demonstration of convergence. At 30° , the count distance from the uncoupled reference changes from 0.04544 to 0.04602 for $d/a = 0.99$ and from 0.13345 to 0.13494 for $d/a = 0.97$ when R/a increases from 25 to 30. The coupling-induced redistribution thus persists, while the narrowest-width DOS retains appreciable finite-size structure. A candidate bulk gap requires a persistent depleted interval under further size and boundary controls, beyond the plateaus that occur automatically between consecutive finite-system eigenvalues.

5 From nonorthogonal matrices to physical observables

5.1 A change of representation is not a change of Hamiltonian

Atom-centered basis functions $\{|\chi_\mu\rangle\}$ generally overlap. A finite-basis electronic-structure calculation solves

$$\mathbf{H}\mathbf{c}_n = \varepsilon_n \mathbf{S}\mathbf{c}_n, \quad H_{\mu\nu} = \langle \chi_\mu | \hat{H} | \chi_\nu \rangle, \quad S_{\mu\nu} = \langle \chi_\mu | \chi_\nu \rangle. \quad (13)$$

For a linearly independent basis, $\mathbf{S}$ is positive definite. Symmetric orthogonalization gives [14]

$$\widetilde{\mathbf{H}} = \mathbf{S}^{-1/2} \mathbf{H} \mathbf{S}^{-1/2}, \quad \widetilde{\mathbf{c}}_n = \mathbf{S}^{1/2} \mathbf{c}_n, \quad \widetilde{\mathbf{c}}_n^\dagger \widetilde{\mathbf{c}}_m = \delta_{nm}. \quad (14)$$

This represents the same projected operator in an orthonormal basis. Diagonalizing $\mathbf{H}$ as though $\mathbf{S} = \mathbf{I}$ would instead solve a different problem. Near-linear dependencies must be controlled before taking inverse square roots. An overlap-eigenvalue cutoff changes the retained subspace and must be documented.

For a sharp energy cutoff in a finite-system gap, the orthonormal-coordinate projector is

$$\widetilde{\mathbf{P}}_E = \sum_{\varepsilon_n \leq E} \widetilde{\mathbf{c}}_n \widetilde{\mathbf{c}}_n^\dagger. \quad (15)$$

Its ordinary matrix trace counts occupied one-particle states in that finite basis. Passing from this count to a bulk IDS still requires a controlled sequence of geometries and a fixed physical normalization. Adding Gaussian basis functions should not silently change the definition of states per cell or per area.

5.2 Why a projected DOS is not a gap label

Let $\mathbf{Q}_L$ be an orthogonal projector onto selected Löwdin orbitals. The projected DOS, before broadening, is

$$\rho_L(E) = \sum_n w_n^{(L)} \delta(E - \varepsilon_n), \quad w_n^{(L)} = \widetilde{\mathbf{c}}_n^\dagger \mathbf{Q}_L \widetilde{\mathbf{c}}_n. \quad (16)$$

Unlike the total state count, these weights may be fractional. If a set of mutually orthogonal orbital projectors sums to the identity, summing their DOS recovers the total DOS for the same eigenstates and normalization. A different population-analysis convention need not assign the same layer weights.

A two-level example already exposes the logical limitation. For $\widetilde{\mathbf{H}} = \text{diag}(\varepsilon_a, \varepsilon_b)$ and $\mathbf{Q}_L = \text{diag}(1, 0)$, the projected DOS has no weight at ε_b , although the full Hamiltonian has an eigenstate there. More generally, $\mathbf{Q}_L \widetilde{\mathbf{P}}_E \mathbf{Q}_L$ is not a projection unless the relevant subspaces commute. Its weighted trace therefore does not inherit the interpretation of a K_0 gap label. A layer-resolved depletion is a useful diagnostic of orbital character, not proof of a bulk spectral gap.

Display broadening and electronic occupations require a separate distinction. Convolution with a kernel of width η smooths spectral data and can hide small gaps. Fermi–Dirac smearing changes occupations. At nonzero temperature the matrix $f_\beta(\widetilde{\mathbf{H}})$ is generally not idempotent and is not the sharp projector appearing in Eq. (3). Neither operation supplies a topological classification by itself.

5.3 Parameter-dependent bases and Berry geometry

Orthogonalization at a fixed geometry is insufficient for a phason derivative if the basis itself moves. Write $B(\lambda) = \chi(\lambda) \mathbf{S}(\lambda)^{-1/2}$ for the map from orthonormal coordinates to physical states, with $B^\dagger B = \mathbf{I}$. On a local parameter patch, let $\widetilde{\mathbf{C}}(\lambda)$ contain a smooth occupied orthonormal-coordinate frame. The physical occupied frame is $\Psi = B \widetilde{\mathbf{C}}$, and its Berry connection is

$$\mathcal{A}_\lambda = i\Psi^\dagger \partial_\lambda \Psi = i\widetilde{\mathbf{C}}^\dagger \partial_\lambda \widetilde{\mathbf{C}} + i\widetilde{\mathbf{C}}^\dagger B^\dagger (\partial_\lambda B) \widetilde{\mathbf{C}}. \quad (17)$$

The second term accounts for basis motion. Omitting it is justified only when the chosen representation makes it vanish or when an equivalent covariant overlap construction already includes it. Discrete overlaps between physical states at neighboring geometries offer a practical route, provided the cross-geometry basis overlaps are treated consistently. A basis-invariant spectrum at every geometry does not, by itself, guarantee a basis-consistent Berry curvature across geometries.

Electronic-structure packages such as CP2K provide a route to atomistic Hamiltonians and state-resolved quantities [15, 16]. The UZH protocol illustrates why basis-set and pseudopotential errors should be separated [17]. These controls remain necessary when the target quantity is topological. They do not establish that the self-consistent effective Hamiltonian defines the required covariant, smoothly parameterized family. Consistent spin treatment, structural relaxation, and branch selection of the self-consistent solution are additional requirements. A Kohn–Sham spectral gap must also be distinguished from a many-body excitation gap.

6 A hierarchy of evidence for atomistic gap labeling

The mathematical stability statement provides a useful standard for numerical practice. Let E lie a distance $\delta > 0$ from the spectrum of a bounded self-adjoint $\hat{H}$ in a fixed observable algebra. If a self-adjoint perturbation $\Delta\hat{H}$ in that algebra satisfies $\|\Delta\hat{H}\| < \delta$, then

$$\hat{H}_s = \hat{H} + s\Delta\hat{H}, \quad 0 \leq s \leq 1, \quad \text{dist}(E, \text{spec } \hat{H}_s) \geq \delta - \|\Delta\hat{H}\| > 0. \quad (18)$$

The spectral projectors form a continuous path and retain their K_0 class. The spectral bound follows from the resolvent estimate, or equivalently from the Neumann-series criterion for invertibility. This statement explains robustness but does not turn an observed finite gap into a

bulk one. Comparisons between changing basis spaces need a common embedding before an operator-norm estimate has this meaning.

Table 2 separates the principal observables. The order matters: assigning an integer before establishing the relevant bulk projector reverses the logic of gap labeling. Conversely, demanding a quantized response from a diagnostic model moment asks that observable to answer a question it was not designed to address.

Table 2: Different numerical objects support different physical conclusions.

Object	Direct information	Additional requirement for topology
Finite spectral moment	Averaged spectral scale and coupling contributions	A bulk limit and a gapped spectral projector
Layer-projected DOS	Spectral weight in a chosen subspace	Total spectral information and a projection-independent gap test
Bulk IDS plateau	State count in an established spectral gap	A justified trace range and, where needed, further topological pairings
Phason-dependent projector	Geometry of occupied states across a family	A converged invariant and the assumptions connecting it to response

A practical analysis begins by specifying the family of infinite structures, its covariance, the retained orbitals, and the trace normalization. Finite patches or periodic approximants can then be chosen to approach that family, rather than being treated as its definition. For a two-dimensional open patch, boundary state counts typically grow with perimeter while bulk counts grow with area. Boundary weight can vanish in a normalized DOS and still encode physically important spectral flow. Interior projections are therefore useful for diagnosing bulk gaps, while boundary-resolved calculations answer a complementary question.

The next step is to converge the full spectral information with respect to size, approximant order, basis quality, and the relevant numerical cutoffs. Candidate gaps must be distinguished from missing calculations, insufficient energy resolution, and finite-size level spacings. A consistent energy reference is needed to follow the same gap across a parameter family. Aligning each spectrum independently to its own Fermi level can obscure that tracking.

Only after this stage should an IDS plateau be compared with the trace range of the chosen algebra. Its persistence should be checked along the appropriate gapped family. If a pumping or Hall interpretation is sought, one should compute the corresponding pairing of the projector with spatial and/or phason derivatives, using consistent parameter-dependent bases. Boundary spectral flow or an explicit adiabatic transport calculation then offers an independent test of the proposed response. These checks complement one another and cannot be replaced by a visual resemblance to the Hofstadter butterfly.

7 Perspective

Bellissard’s work turns the interpretation of quasiperiodic spectra into a constructive program: establish the bulk spectral object, count its states with the appropriate trace, and compute the pairing that describes the desired response. The bilayer calculations illustrate how this program begins. An exact moment identity identifies the interlayer contribution to spectral variance, the uncoupled reference isolates its energy-resolved redistribution, and size and broadening controls assess the resolution of that observation. Together, these diagnostics connect geometry to spectral reconstruction before an integer label is assigned.

The further passage to atomistic quasiperiodic bilayers is demanding but well posed.

Nonorthogonal basis sets require metric-aware projectors, moving atoms require basis-consistent parameter derivatives, and self-consistency requires a reproducible covariant branch of solutions. Relaxation can change the pattern space itself, so its effect is not always captured by adding a small perturbation to a fixed Hamiltonian. These are opportunities for a closer interaction between electronic-structure methods and noncommutative geometry rather than reasons to separate the subjects.

The next calculations follow directly from this separation of tasks. A candidate spectral feature must be followed to a stable bulk gap, its projector must yield a consistently normalized state count, and a proposed pumping or Hall response must be tested through the corresponding topological pairing and its physical consequences. This is the practical connection between geometry, state counting, and response that makes Bellissard’s framework useful for atomistic materials modeling.

Acknowledgment. This perspective is dedicated to Jean Bellissard on the occasion of his eightieth birthday and to the continuing influence of his work on the mathematics and physics of aperiodic solids.

Data and code availability. The source of this article, the bilayer eigenvalues, the spectral moment and butterfly data, and the uncoupled-reference and resolution controls are available in <https://github.com/DCM-Uni-Paderborn/Hofstadter-Butterfly>. The scripts in `analysis/` are `bellissard_model_moments.py` for Fig. 1, `bellissard_robustness.py` for Table 1 and the uncoupled reference, and `bellissard_butterfly.py` for Fig. 2.

Use of AI-assisted tools. AI-assisted tools were used in developing the analysis script.

References

- [1] Jean Bellissard. K -theory of C^* -algebras in solid state physics. In T. C. Dorlas, N. M. Hugenholtz, and M. Winnink, editors, *Statistical Mechanics and Field Theory: Mathematical Aspects*, volume 257 of *Lecture Notes in Physics*, pages 99–156. Springer, Berlin, Heidelberg, 1986. doi: 10.1007/3-540-16777-3_74.
- [2] Jean Bellissard, Andreas van Elst, and Hermann Schulz-Baldes. The noncommutative geometry of the quantum Hall effect. *Journal of Mathematical Physics*, 35(10):5373–5451, 1994. doi: 10.1063/1.530758.
- [3] Jean Bellissard, Riccardo Benedetti, and Jean-Marc Gambaudo. Spaces of Tilings, Finite Telescopic Approximations and Gap-Labeling. *Communications in Mathematical Physics*, 261:1–41, 2006. doi: 10.1007/s00220-005-1445-z.
- [4] Yaacov E. Kraus, Yoav Lahini, Zohar Ringel, Mor Verbin, and Oded Zilberberg. Topological States and Adiabatic Pumping in Quasicrystals. *Physical Review Letters*, 109:106402, 2012. doi: 10.1103/physrevlett.109.106402.
- [5] Xiang Ni, Kai Chen, Matthew Weiner, David J. Apigo, Camelia Prodan, Andrea Alù, Emil Prodan, and Alexander B. Khanikaev. Observation of Hofstadter butterfly and topological edge states in reconfigurable quasi-periodic acoustic crystals. *Communications Physics*, 2:55, 2019. doi: 10.1038/s42005-019-0151-7.
- [6] Matheus I. N. Rosa, Massimo Ruzzene, and Emil Prodan. Topological gaps by twisting. *Communications Physics*, 4:130, 2021. doi: 10.1038/s42005-021-00630-3.
- [7] Chris Bourne and Emil Prodan. Non-Commutative Chern Numbers for Generic Aperiodic Discrete Systems. *Journal of Physics A: Mathematical and Theoretical*, 51:235202, 2018. doi: 10.1088/1751-8121/aac093.
- [8] Douglas R. Hofstadter. Energy levels and wave functions of Bloch electrons in rational and irrational magnetic fields. *Physical Review B*, 14:2239–2249, 1976. doi: 10.1103/physrevb.14.2239.

- [9] D. J. Thouless. Quantization of particle transport. *Physical Review B*, 27:6083–6087, 1983. doi: 10.1103/PhysRevB.27.6083.
- [10] Emil Prodan. Virtual topological insulators with real quantized physics. *Physical Review B*, 91: 245104, 2015. doi: 10.1103/physrevb.91.245104.
- [11] Ying Su and Shi-Zeng Lin. Topological sliding moiré heterostructure. *Physical Review B*, 101: 041113, 2020. doi: 10.1103/physrevb.101.041113.
- [12] Manato Fujimoto, Henri Koschke, and Mikito Koshino. Topological charge pumping by a sliding moiré pattern. *Physical Review B*, 101:041112, 2020. doi: 10.1103/physrevb.101.041112.
- [13] Mao Yoshii, Sota Kitamura, and Takahiro Morimoto. Gap labeling theorem for multilayer thin film heterostructures. *Physical Review B*, 107:064201, 2023. doi: 10.1103/physrevb.107.064201.
- [14] Per-Olov Löwdin. On the Non-Orthogonality Problem Connected with the Use of Atomic Wave Functions in the Theory of Molecules and Crystals. *The Journal of Chemical Physics*, 18:365–375, 1950. doi: 10.1063/1.1747632.
- [15] Thomas D. Kühne, Marcella Iannuzzi, Mauro Del Ben, Vladimir V. Rybkin, Patrick Seewald, Frederick Stein, Teodoro Laino, Rustam Z. Khaliullin, et al. CP2K: An electronic structure and molecular dynamics software package - Quickstep: Efficient and accurate electronic structure calculations. *The Journal of Chemical Physics*, 152:194103, 2020. doi: 10.1063/5.0007045.
- [16] Marcella Iannuzzi, Jan Wilhelm, Frederick Stein, Augustin Bussy, Hossam Elgabarty, Dorothea Golze, Anna-Sophia Hehn, Maximilian Graml, Stepan Marek, Beliz Sertcan Gökmen, Christoph Schran, Harald Forbert, Rustam Z. Khaliullin, Anton Kozhevnikov, Mathieu TAILLEFUMIER, Rocco Meli, Vladimir V. Rybkin, Martin Brehm, Robert Schade, Ole Schütt, Johann V. Pototschnig, Hossein Mirhosseini, Andreas Knüpfer, Dominik Marx, Matthias Krack, Jürg Hutter, and Thomas D. Kühne. The CP2K Program Package Made Simple. *The Journal of Physical Chemistry B*, 130(4): 1237–1310, 2026. doi: 10.1021/acs.jpcb.5c05851.
- [17] Hossein Mirhosseini, Tiziano M. A. Müller, Matthias Krack, Thomas D. Kühne, and Jürg Hutter. The UZH protocol: Separating errors and constructing improved CP2K basis sets and pseudopotentials. *The Journal of Chemical Physics*, 165(10):104103, 2026. doi: 10.1063/5.0347392.