

Topological Systems with CP2K

Thomas D. Kühne^{a,b,c,*}, Alexander Cerjan^d,
Vladislav Efremkin^{a,b}, Hermann Schulz-Baldes^e, and Emil Prodan^f

^aCenter for Advanced Systems Understanding (CASUS),
Untermarkt 20, D-02826 Görlitz, Germany

^bHelmholtz-Zentrum Dresden-Rossendorf (HZDR),
Bautzner Landstrasse 400, D-01328 Dresden, Germany

^cChair of Computational Systems Science, Institute of Artificial Intelligence,
Technische Universität Dresden, D-01069 Dresden, Germany

^dCenter for Integrated Nanotechnologies, Sandia National Laboratories,
Albuquerque, New Mexico 87185, USA

^eDepartment Mathematik, Friedrich-Alexander-Universität Erlangen-Nürnberg,
Cauerstrasse 11, D-91058 Erlangen, Germany

^fDepartment of Physics, Yeshiva University, New York, USA

*Corresponding author: tkuehne@cp2k.org

Abstract

Electronic transport and topology probe information that cannot be inferred from band energies alone. Current operators, inter-state overlaps, symmetry representations, and the spatial structure of a Hamiltonian must be treated consistently. We formulate these quantities in the nonorthogonal Gaussian atomic-orbital representation of CP2K, connecting finite-temperature Kubo transport, Wilson-loop and Chern analysis, crystalline symmetry representations, spectral localizers, and the quadratic pseudospectrum. The overlap metric is retained in operator products, symmetry transformations, and spectral-gap estimates. Real-space and reciprocal-space constructions share Hamiltonian, position, and spin-orbit matrix elements without requiring a fitted Wannier Hamiltonian. Dense reference calculations are complemented by distributed sparse assembly, symmetric-indefinite inertia calculations, and an optional sparse Pfaffian back-end. We explain how the finite matrix formulations relate to noncommutative geometry and distinguish local topological protection from an electronic band gap or a mobility gap. Wilson and inversion calculations, native projective little-group and antiunitary corepresentation tests, and serial/MPI operator comparisons provide complementary validation. The implementation scope is stated explicitly: the current Kubo path evaluates symmetric dissipative DC response. Crystalline analysis includes nonsymmorphic endpoint-to-line compatibility, but a complete space-group EBR classifier and universal symmetry reduction of topological properties remain development targets.

Keywords: CP2K, Kubo transport, noncommutative geometry, spectral localizer, topological quantum chemistry, spin-orbit coupling

Working manuscript for Computer Physics Communications. Implementation and validation status: 24 September 2026.

1 Introduction

Electronic-structure theory supplies a Hamiltonian and a Hilbert-space representation; extracting transport or topological information requires additional structure. Two calculations can agree in their eigenvalues while disagreeing in matrix elements of position, current, or a symmetry operator. This distinction is especially important in a localized nonorthogonal basis, where ordinary matrix multiplication is not the representation of operator composition.

The QUICKSTEP module of CP2K combines localized Gaussian orbitals with plane-wave and multigrid representations of densities and electrostatics [1, 2, 3, 4]. This representation provides access to spatially resolved operators without first fitting a tight-binding Hamiltonian. It also creates specific obligations: the AO metric, periodic image indices, Bloch convention, nonlocal pseudopotentials, and spinor ordering must remain consistent across every analysis.

The first-principles transport programme of Kühne, Heske and Prodan [5] supplies an important starting point: thermally disordered structures are treated as configurations of a covariant family of Hamiltonians, and dissipation is included in the finite-temperature response. Bellissard’s operator-algebraic framework connects this viewpoint to topology without assuming exact translation symmetry of each configuration [6, 7]. The present manuscript combines this motivation with reciprocal-space invariants and spatially resolved finite-matrix diagnostics.

The scope is deliberately broader than a band-topology interface. Wilson loops resolve the geometry of an isolated band subspace; crystalline representations resolve symmetry constraints; a localizer tests an energy-position query; a quadratic pseudospectrum produces approximately localized states; and Kubo response determines a transport coefficient at specified temperature and dissipation. None of these outputs can simply be substituted for another.

The companion study on symmetry-reduced k-point sampling and Wannier analysis develops the underlying Bloch reconstruction and full-mesh export. We repeat the operator conventions needed to make the present article self-contained, but keep the detailed topology, transport, and localizer analysis here. In particular, Wannier90 is an optional route to a reduced representation, not a prerequisite for native AO analysis. Sections below distinguish implemented algorithms from the more general theory and from extensions still requiring validation. The materials-discovery applications discussed later are distinguished from our small test and demonstration calculations: the former motivate the scientific use of these methods, while the latter establish specific numerical identities and illustrate their operation.

2 Hamiltonians and operators in a nonorthogonal basis

2.1 Covariant matrices and the physical metric

Let Φ denote the row of basis functions $|\phi_\mu\rangle$, and $|\psi\rangle = \Phi c$. Define

$$S = \Phi^\dagger \Phi, \quad H = \Phi^\dagger \hat{H} \Phi, \quad X_a = \Phi^\dagger \hat{r}_a \Phi. \quad (2.1)$$

The overlap must be positive definite on the retained basis, and the normalization is $c^\dagger S c = 1$. The generalized eigenproblem is

$$H C = S C \varepsilon, \quad C^\dagger S C = I. \quad (2.2)$$

A covariant matrix A acts on coefficient vectors through $S^{-1} A$. Consequently, composition of two *projected* operators is represented by $A S^{-1} B$, and the projected commutator by $A S^{-1} B - B S^{-1} A$. An ordinary product AB is generally incorrect.

For a nonsingular change of basis $\Phi' = \Phi T$,

$$S' = T^\dagger S T, \quad A' = T^\dagger A T, \quad c' = T^{-1} c. \quad (2.3)$$

The expectation $c^\dagger A c$ and the generalized spectrum are unchanged. The orthonormal representation $\tilde{A} = S^{-1/2} A S^{-1/2}$ is useful for proofs and dense reference calculations. It need not be formed in a sparse implementation: products can be evaluated by solving linear systems with S .

A finite-basis projection is not an algebra homomorphism for arbitrary continuum operators. With $\Pi = \Phi S^{-1} \Phi^\dagger$, $\Phi^\dagger \hat{A} \Pi \hat{B} \Phi$ differs in general from $\Phi^\dagger \hat{A} \hat{B} \Phi$. Thus a commutator or a variance of projected positions is an observable of the projected model, not automatically its complete-basis continuum counterpart. This distinction is monitored through basis convergence rather than hidden in a numerical tolerance.

2.2 Finite positions and AO export

For an isolated system, analytic Gaussian first moments provide X_a in the same basis ordering as H and S . Translation of the coordinate origin by x_a gives $X_a - x_a S$, not $X_a - x_a I$. Likewise an energy query uses $H - E S$. These two simple identities are essential for origin and energy-shift invariance.

The `AO_MATRICES` print section can export overlap, scalar KS Hamiltonian, and the three position matrices. They permit an independent finite-Hamiltonian calculation. Activating SOC in a property section does not silently turn the separate scalar KS export into a spinor Hamiltonian export. Model Hamiltonians can use the low-level matrix kernels when the metric, positions, and symmetry structure are supplied consistently; this does not imply automatic adapters for every electronic-structure method.

2.3 Bloch operators and directed overlaps

In the cell Fourier convention,

$$H_{\mu\nu}(\mathbf{k}) = \sum_{\mathbf{R}} e^{i\mathbf{k}\cdot\mathbf{R}} H_{\mu\nu}(\mathbf{R}), \quad S_{\mu\nu}(\mathbf{k}) = \sum_{\mathbf{R}} e^{i\mathbf{k}\cdot\mathbf{R}} S_{\mu\nu}(\mathbf{R}). \quad (2.4)$$

Here $\mathbf{k}$ is Cartesian and includes inverse-length units; fractional reciprocal coordinates instead require 2π in the phase. Image-cell labels cannot be discarded before forming derivatives or inter-k-point connections.

For cell-periodic states and a directed step $\mathbf{b}$, the occupied-state overlap is

$$M(\mathbf{k}, \mathbf{b}) = C^\dagger(\mathbf{k}) O(\mathbf{k} + \mathbf{b}; \mathbf{b}) C(\mathbf{k} + \mathbf{b}), \quad O_{\mu\nu} = \sum_{\mathbf{R}} e^{i(\mathbf{k}+\mathbf{b})\cdot\mathbf{R}} \langle \phi_{\mu 0} | e^{-i\mathbf{b}\cdot\mathbf{r}} | \phi_{\nu \mathbf{R}} \rangle. \quad (2.5)$$

The zero-step limit is $S(\mathbf{k})$, but a finite link is not obtained by replacing O with S , or by simply contracting coefficient vectors. Neighbor lists, reciprocal-boundary phases, and the selected band rank are part of the observable. Wannier90's file and library interfaces can consume such data [8]; native Wilson calculations need no Wannier localization or interpolation.

2.4 Spin-orbit coupling and symmetry

The implementations considered here start from a converged scalar restricted SCF potential and add GTH spin-orbit matrix elements. The localizer and Bloch-Kubo paths use the complete AO spinor space, whereas a second-variational band-topology calculation uses a selected scalar-state space whose size must be converged separately. Neither construction is self-consistent noncollinear SOC DFT.

In spin-major ordering, the overlap is $\mathbb{S} = \text{diag}(S, S)$ and physical time reversal is $\Theta = JK$, where

$$J = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}, \quad \Theta^2 = -I. \quad (2.6)$$

Time-reversal relations must be checked for the Hamiltonian, metric, and relevant operators; the presence of an SOC pseudopotential alone does not establish them. A Cartesian spatial rotation R

acts on spin through an $SU(2)$ lift of $\det(R)R$, with inversion acting trivially on spin and separately on orbital parity. The sign ambiguity of this lift is retained in the double-group multiplication law.

3 Noncommutative geometry and physical interpretation

3.1 Covariance rather than exact order

For a family of configurations $\omega \in \Omega$, a spatial action τ_a is represented by

$$U_a H_\omega U_a^{-1} = H_{\tau_a \omega}. \quad (3.1)$$

This covariance is weaker than $[U_a, H_\omega] = 0$: a displaced thermal configuration need not have the symmetries of its probability measure. Bellissard’s formulation replaces the Brillouin-zone description by an algebra of such covariant observables with a trace per physical volume [6]. The operator-algebraic viewpoint therefore remains meaningful when a single Bloch momentum is not a good quantum number.

For sufficiently local observables, the trace per volume has the schematic realization

$$\mathcal{T}(A) = \lim_{V \rightarrow \infty} \frac{1}{V} \text{Tr}(\chi_V A \chi_V), \quad (3.2)$$

with the appropriate orbital trace and a physical-volume normalization. Ergodicity supplies the conditions under which a typical large configuration and an ensemble average agree. It does not guarantee that a particular small simulation cell has self-averaged.

The first-principles construction must preserve covariance while passing from continuum KS operators to AO matrices [5]. The metric is part of that transformation. In a finite AO space, traces of projected covariant matrices use $\text{Tr}(S^{-1}A)$; traces of products use the intervening metric factors. For local traces the spatial cutoff itself is an operator, not merely an arbitrary subset of coefficient indices.

3.2 Derivations, Fermi projections, and Chern pairings

Choose spatial derivations $\nabla_a A = -i[X_a, A]$ in an orthonormal representation. For the Fermi projection $P = \mathbf{1}_{(-\infty, E_F)}(H)$, a two-dimensional Chern pairing can be written, with a fixed orientation, as

$$C = 2\pi i \mathcal{T}(P[\nabla_1 P, \nabla_2 P]) = -2\pi i \mathcal{T}(P[[X_1, P], [X_2, P]]). \quad (3.3)$$

The trace normalization and oriented coordinates are chosen so that the periodic limit agrees with the Berry-curvature convention used below. Reversing that orientation reverses C . Index theory relates such pairings to stable integers under the required locality and gap or localization assumptions [7]. Equation (3.3) is a theoretical connection, not a claim that the current implementation evaluates a general thermodynamic-limit noncommutative Chern formula.

A spectral gap excludes eigenvalues in an interval. A mobility gap instead concerns localization and transport even when states are present. A localizer gap is the smallest singular value of a composite energy-position operator. These three gaps answer different questions. The localizer is useful precisely because spatially resolved information can remain available where a bulk band classification is not directly applicable [9]. Its numerical invertibility does not by itself prove a mobility-gap theorem for a DFT material.

3.3 Thermal disorder and convergence

An AIMD trajectory can supply configurations for a disorder average. The nuclear sampling temperature, the electronic Fermi temperature, and the dissipative relaxation scale are conceptually distinct. A calculation should record all three rather than infer them from a single input

temperature. Configuration correlations determine how many effectively independent samples have been obtained; a small standard error computed from strongly correlated snapshots is not a substitute for a correlation analysis.

The thermodynamic limit, zero-temperature limit, and vanishing dissipation limit cannot generally be interchanged. At fixed finite volume the spectrum is discrete, so taking the broadening to zero first can give a qualitatively misleading DC response. For a material prediction we therefore require size and sampling convergence at a specified positive dissipation, followed by a physically justified study of its dependence. The present validation examples establish numerical identities at fixed inputs, not a microscopic scattering model or a disorder-converged transport prediction.

4 Finite-temperature Kubo transport

4.1 Liouvillian formulation and dissipation

Let $f(H) = [1 + \exp((H - \mu)/k_B T)]^{-1}$ and define the energy-valued Liouvillian $\mathcal{L}_H(A) = i[H, A]$. In the relaxation-time approximation, a positive energy broadening Γ replaces a more general dissipative superoperator. With $\nabla_a = -i[X_a, \cdot]$, a convenient conductivity convention is

$$\sigma_{ab} = -\frac{e^2}{\hbar} \mathcal{T} \left[(\nabla_a H) (\Gamma + \mathcal{L}_H)^{-1} \nabla_b f(H) \right]. \quad (4.1)$$

The real symmetric part gives the dissipative DC response considered here. The factor $e > 0$ is the magnitude of the electron charge. This expression states the operator structure behind the finite-volume calculation, including the role of dissipation [5]. A constant Γ is an approximation to scattering; it is not obtained from the SCF convergence tolerance or from a plotted density-of-states width.

Let $a_{nm}^b = \langle n | \nabla_b H | m \rangle = \hbar v_{nm}^b$, and define the nonnegative Fermi divided difference

$$F_{nm} = \begin{cases} -\frac{f(\varepsilon_n) - f(\varepsilon_m)}{\varepsilon_n - \varepsilon_m}, & \varepsilon_n \neq \varepsilon_m, \\ -f'(\varepsilon_n) = \frac{f(\varepsilon_n)[1 - f(\varepsilon_n)]}{k_B T}, & \varepsilon_n = \varepsilon_m. \end{cases} \quad (4.2)$$

A spectral expansion of Eq. (4.1) gives

$$\sigma_{ab}^{\text{diss}} = \frac{e^2}{\hbar V_d} \sum_{\mathbf{k}, s} w_{\mathbf{k}} g_s \sum_{nm} F_{nmks} \frac{\Gamma}{\Gamma^2 + (\varepsilon_{nks} - \varepsilon_{mks})^2} \Re \left[a_{nmks}^a a_{mnks}^b \right]. \quad (4.3)$$

Here $\sum_{\mathbf{k}} w_{\mathbf{k}} = 1$, V_d is the periodic length, area, or volume, and g_s accounts for spin occupancy without double counting. For explicit SOC spinors, each state has unit maximum occupation. The tensor is projected onto the periodic Cartesian subspace before reporting a reduced-dimensional conductivity.

Equation (4.3) is positive semidefinite for $T, \Gamma > 0$. In particular, its diagonal elements are nonnegative. The units are conductance times length^{2-d}: S m in one dimension, S in two dimensions, and S/m in three dimensions, with the latter optionally printed as S/cm. Vacuum padding must not change a two-dimensional sheet conductivity through an arbitrary three-dimensional volume factor. An isolated finite-system response is not automatically a measurable bulk conductivity.

4.2 Degenerate levels and intraband response

The equal-energy branch of Eq. (4.2) applies to all matrix elements within a degenerate subspace, not just $n = m$. In a periodic system, the diagonal band velocity can be nonzero and provides an intraband term proportional to $-f'/\Gamma$. Discarding diagonal terms would remove this contribution. Likewise, dividing current matrix elements by $\varepsilon_n - \varepsilon_m$ cannot be used to define a stable position operator in a degenerate block.

A unitary change of basis inside an exactly degenerate subspace rotates the complete current block. Because the scalar kernel is constant on that block, the summed contraction is unchanged. This is why comparisons should involve block traces or projectors rather than phases of individual eigenvectors. Near equal energies, the implementation evaluates the divided difference with a smooth finite-temperature expansion instead of subtracting nearly identical Fermi functions.

4.3 Three current-operator choices

The historical atom-embedding approximation associates each AO with its nuclear center. It provides an economical representation but does not contain all intra-atomic Gaussian position matrix elements. It remains a distinct, explicitly selected approximation.

For isolated systems, the projected-AO alternative uses

$$J_a = HS^{-1}X_a - X_aS^{-1}H, \quad \hbar v_a = iJ_a. \quad (4.4)$$

J_a is anti-Hermitian. In an S -orthonormal eigenbasis it becomes $(J_a)_{nm} = (\varepsilon_n - \varepsilon_m)(X_a^{\text{MO}})_{nm}$. The complex conjugation in a conductivity contraction must be retained when SOC is present. This construction is exact for the projected finite AO model; the difference between projected and continuum commutators remains a basis-convergence question.

For periodic calculations, a bounded Cartesian position in one simulation cell is not a valid replacement for the periodic current. The Bloch path uses derivatives of the image Hamiltonian and overlap together with analytic first-moment integrals. In atomic units its matrix elements are

$$\begin{aligned} v_{nm}^a(\mathbf{k}) = & [C^\dagger \partial_a H C]_{nm} - \varepsilon_n [C^\dagger \partial_a S C]_{nm} \\ & + i(\varepsilon_n - \varepsilon_m) [C^\dagger D_a C]_{nm}, \end{aligned} \quad (4.5)$$

where ∂_a differentiates Cartesian k_a . The cell-gauge connection obeys

$$D_a - D_a^\dagger = -i \partial_a S. \quad (4.6)$$

These identities make explicit why the derivative of H alone, the canonical momentum, and the full velocity are not interchangeable in a local-orbital calculation [10].

Hermiticity can be checked directly: the anti-Hermitian part from the energy-weighted overlap derivative is cancelled by Eq. (4.6). For $n = m$, Eq. (4.5) reduces to the generalized Hellmann–Feynman derivative $c_n^\dagger (\partial_a H - \varepsilon_n \partial_a S) c_n$. No eigenvector derivative or energy-denominator division is required.

For distinct band energies, the same operator determines the interband position (dipole) matrix elements in atomic units,

$$d_{nm}^a(\mathbf{k}) = -\frac{i v_{nm}^a(\mathbf{k})}{\varepsilon_{n\mathbf{k}} - \varepsilon_{m\mathbf{k}}}, \quad \varepsilon_{n\mathbf{k}} \neq \varepsilon_{m\mathbf{k}}. \quad (4.7)$$

Here d_{nm}^a has units of length and does not include the electron charge. Substitution of Eq. (4.5) gives the Hamiltonian-derivative term, the energy-weighted overlap-derivative term, and $[C^\dagger D_a C]_{nm}$. All three are required in a general nonorthogonal local-orbital basis [10]. The $\mathbf{k}$ -point moment calculation constructs these matrices from real-space image-cell integrals and can use the SCF mesh, a separate KPOINTS set, or a KPOINT_SET path in the moments print section. Equation (4.7) is only an interband relation. It neither supplies diagonal positions in a periodic system nor replaces the undivided velocity blocks needed for degenerate and intraband transport.

4.4 Gauge covariance and nonlocal operators

Under a differentiable, nonsingular AO basis change $T(\mathbf{k})$, $H' = T^\dagger H T$, $S' = T^\dagger S T$, and $C' = T^{-1} C$. The connection must transform as

$$D'_a = T^\dagger D_a T - i(\partial_a T)^\dagger S T. \quad (4.8)$$

Substitution into Eq. (4.5) cancels all derivatives of T by using $HC = SC\varepsilon$. The physical velocity is therefore unchanged even under a k -dependent nonunitary AO transformation. This is a stronger test than comparing band energies.

The Hamiltonian derivatives include all retained image contributions, including nonlocal pseudopotential and GTH-SOC terms. The connection term is still required; including a derivative of the SOC Hamiltonian does not make the bare Peierls derivative complete. SCF-potential convergence, scalar or spinor basis convergence, and analysis-mesh convergence must be checked independently.

4.5 Property symmetry and practical scope

The optional Bloch-Kubo symmetry path diagonalizes only representatives of its own analysis mesh. Atom permutations, AO rotations, lattice-image phases, physical spin rotations, and antiunitary conjugation reconstruct the remaining eigenframes. Each candidate frame is tested through $C^\dagger SC = I$, $C^\dagger HC = \varepsilon$, and target/source current covariance. An unacceptable map is not silently inserted into the conductivity.

For a spatial Cartesian operation R , velocities rotate as a polar vector. Time reversal adds the reversal of velocity as well as complex conjugation. In the anti-Hermitian current convention used internally, the extra factor of i must also be conjugated. The response sums the full tensor over the full mesh; it is not an irreducible scalar-weight substitution for anisotropic currents.

The current implementation provides symmetric dissipative DC conductivity with a prescribed relaxation scale. It does not yet provide anomalous Hall, spin Hall, a general frequency-dependent response, or microscopic scattering superoperators. Its Bloch matrices and eigenframes are a bounded dense reference path, although eigensystems are distributed over MPI ranks. Fewer diagonalizations do not imply proportional memory or runtime savings when target operators are still explicitly evaluated.

5 Band topology: Wilson loops, Chern numbers, and time reversal

The overlap matrices used for Wannier construction also contain the geometry of an isolated band subspace. Following established hybrid-Wannier and Wilson-loop formulations [11, 12], the native analysis in CP2K evaluates Berry phases, hybrid Wannier charge centers, the $\mathbb{Z}_2$ invariant of a time-reversal-invariant plane, and the first Chern number C_1 . The native analysis evaluates Wilson loops and topological invariants directly from Gaussian-basis Bloch-state overlaps, without requiring Z2Pack or a maximally localized Wannier representation. Z2Pack remains an optional external analysis route and a complementary validation tool. Its adapter supplies the same first-principles overlaps to an external implementation with its own line and surface sampling. Neither route requires a fitted tight-binding Hamiltonian.

5.1 Interband geometry and Berry curvature

The velocity and dipole matrix elements in Eqs. (4.5) and (4.7) also connect the local-orbital representation to band geometry. For a nondegenerate band, the moment implementation evaluates the interband expression

$$\begin{aligned}\Omega_n^\gamma(\mathbf{k}) &= 2 \operatorname{Im} \sum_{m \neq n} d_{nm}^\alpha(\mathbf{k}) d_{mn}^\beta(\mathbf{k}) \\ &= 2 \operatorname{Im} \sum_{m \neq n} \frac{v_{nm}^\alpha(\mathbf{k}) v_{mn}^\beta(\mathbf{k})}{(\varepsilon_{n\mathbf{k}} - \varepsilon_{m\mathbf{k}})^2},\end{aligned}\tag{5.1}$$

where (α, β, γ) is a cyclic permutation of (x, y, z) . This equation uses the sign convention of the moment output. Comparisons with overlap-based phases require matching the phase and

surface-orientation conventions. With Cartesian reciprocal coordinates, Ω_n^γ has units of length squared. The formula reveals why small avoided crossings can produce sharply localized curvature and why band-energy convergence alone does not establish convergence of this observable. The band sum, AO basis, and reciprocal-space sampling require separate convergence checks.

The product in Eq. (5.1) is invariant under independent phase changes of nondegenerate eigenstates. At an exact degeneracy, however, the individual-band assignment is not invariant under rotations within the degenerate subspace, and division by the vanishing energy difference is not defined. An isolated group of bands must instead be treated through its projector or matrix-valued connection. The Wilson construction below implements this subspace viewpoint without resolving arbitrary eigenvectors within the group. The interband expression describes the implemented finite-basis moment representation and is not by itself a proof of complete-basis curvature or an integer Chern number. Nor does this geometric quantity constitute an implementation of anomalous Hall transport, which is distinct from the dissipative response in Section 4.

5.2 Directed overlaps and Wilson loops

Consider a closed directed path $\mathbf{k}_0, \dots, \mathbf{k}_L$ with $\mathbf{k}_L = \mathbf{k}_0 + \mathbf{G}$ and links $\mathbf{b}_j = \mathbf{k}_{j+1} - \mathbf{k}_j$. For N_{occ} occupied states, each link is the square matrix in Eq. (2.5). In the Gaussian representation, its AO operator is explicitly

$$O_{\mu\nu}(\mathbf{k} + \mathbf{b}; \mathbf{b}) = \sum_{\mathbf{R}} e^{i(\mathbf{k}+\mathbf{b})\cdot\mathbf{R}} \langle \phi_{\mu\mathbf{0}} | e^{-i\mathbf{b}\cdot\mathbf{r}} | \phi_{\nu\mathbf{R}} \rangle. \quad (5.2)$$

Here $\phi_{\mu\mathbf{R}}(\mathbf{r}) = \phi_{\mu}(\mathbf{r} - \mathbf{R})$ denotes an individual localized orbital, not a Bloch sum. The zero-step limit recovers $S(\mathbf{k})$. At finite displacement, however, this is a directed cross-k operator, not a Hermitian matrix at one k-point. Its construction retains ordered atom pairs, image-cell phases, and the same periodic atomic images as the Hamiltonian. Consequently, neither $C^\dagger(\mathbf{k})C(\mathbf{k} + \mathbf{b})$ nor replacement of the operator by $S(\mathbf{k})$ represents the required overlap. Reciprocal-boundary links retain their integer $\mathbf{G}$ shift, implementing $u_{n,\mathbf{k}+\mathbf{G}}(\mathbf{r}) = e^{-i\mathbf{G}\cdot\mathbf{r}} u_{n\mathbf{k}}(\mathbf{r})$ in a consistent boundary gauge.

At finite sampling the occupied-subspace overlap need not be unitary. Each link is factorized and replaced by its unitary polar factor,

$$M_j = U_j \Sigma_j V_j^\dagger, \quad Q_j = U_j V_j^\dagger, \quad \mathcal{W} = Q_0 Q_1 \cdots Q_{L-1}. \quad (5.3)$$

The smallest singular value of M_j diagnoses loss of continuity or an inadequately resolved subspace. A rank-deficient link is rejected. For a unitary change of occupied-band basis $C_j \mapsto C_j G_j$, the link and closed product transform as

$$Q_j \mapsto G_j^\dagger Q_j G_{j+1}, \quad \mathcal{W} \mapsto G_0^\dagger \mathcal{W} G_0. \quad (5.4)$$

Thus the Wilson spectrum is gauge invariant although individual overlaps are not. With the path and phase convention used here,

$$\lambda_n(\mathcal{W}) = e^{2\pi i \bar{x}_n}, \quad \bar{x}_n = \frac{\arg \lambda_n}{2\pi} \pmod{1}, \quad \gamma = \arg \det \mathcal{W}. \quad (5.5)$$

The $\bar{x}_n$ are the hybrid Wannier charge centers in reduced units, and γ is the total loop phase in this convention. Reversing the loop reverses the eigenphases. The absolute center origin follows the atomic and cell-origin convention, whereas the topological classification does not.

For spinful time-reversal symmetry, loops spanning half a reciprocal plane connect two time-reversal-invariant boundaries. The parity of partner exchange in the center spectrum determines the two-dimensional invariant. The largest-gap construction tracks the center z_ℓ of the largest

gap between adjacent centers on each sampled loop and counts crossings between successive loops [11, 12], giving

$$\nu = N_{\text{cross}} \pmod{2}. \quad (5.6)$$

Loop resolution and transverse sampling are refined together. Acceptance uses center displacements, neighboring-loop gap separation, boundary Kramers pairing, and stable parity. The occupied rank is fixed and separated from conduction states throughout the sampled surface.

On a closed full reciprocal-space surface, C_1 is obtained instead from the winding of the determinant Wilson phase. With $\gamma_\ell = \arg \det \mathcal{W}_\ell$ and a periodic transverse index, the discrete estimate in the present orientation convention is

$$C_1^{\text{mesh}} = \frac{1}{2\pi} \sum_\ell \text{Arg} \exp[i(\gamma_{\ell+1} - \gamma_\ell)]. \quad (5.7)$$

The phase increments must be resolved under refinement, and the final transverse boundary must be sewn to the initial one. Unlike the time-reversal half-plane construction, this calculation covers a full closed surface and does not require time-reversal symmetry. The native implementation checks closure and stable winding under refinement and retains the unrounded winding alongside the integer candidate.

In CP2K, explicit fractional k-points and directed connections are read from a `.nnkp` file with `KPOINTS_SOURCE NNKP`, or generated internally with `KPOINTS_SOURCE WILSON`. `WILSON_LOOP` evaluates a loop, `Z2` selects the half-plane analysis, and `CHERN` selects the full-surface C_1 analysis. The converged SCF potential is held fixed while the post-SCF points are diagonalized. The integration mesh defining the density and the ordered paths probing its band topology are distinct objects. Spin-orbit coupling (SOC) enters through a second-variational spinor Hamiltonian built from the scalar states and pseudopotential SOC matrix elements. For spinor coefficients the link is

$$M_j = C_{j,\uparrow}^\dagger O_j C_{j+1,\uparrow} + C_{j,\downarrow}^\dagger O_j C_{j+1,\downarrow}, \quad (5.8)$$

and band selection refers to this spinor spectrum.

This application makes the role of symmetry particularly clear. Scalar irreducible-mesh weights cannot replace an ordered Wilson product. A path requires explicit states at its points or their symmetry reconstruction with all orbital rotations, spin transformations, and boundary phases. The native explicit-path calculation obtains these states by diagonalizing the fixed-potential problem at the requested points. It therefore complements symmetry-reduced SCF quadrature while retaining the complete neighbor connectivity needed for a gauge-covariant observable.

Figure 1 demonstrates the calculation using small test systems: a periodic neon reference and buckled stanene, a representative quantum spin Hall system [13]. The computed invariants are $\nu = 0$ and $\nu = 1$, respectively. The sampling-converged stanene surface contains 193 loops with 192 points per loop. Native and Z2Pack analysis of the same polar-unitarized overlaps agree in the Wilson eigenvalues to 8×10^{-14} . A separately adaptive Z2Pack-driven calculation also yields $\nu = 1$. Operator-level checks independently test the zero-step metric, reverse-link adjoint identity, and displaced atomic centers. These complementary tests connect the local Gaussian integral representation to the global topology of the occupied bands. Inputs, sampling parameters, and figure data are given in the Supporting Information and accompanying calculation records.

6 Crystalline representations and topological quantum chemistry

6.1 Implemented inversion analysis

An independently versioned native extension evaluates inversion representations at time-reversal-invariant momenta (TRIM). This analysis requires spatial inversion of the crystal, unlike the time-reversal pairing used to reduce a scalar SCF mesh. The TRIM are diagonalized at the converged SCF potential independently of the integration mesh, without a Wannier90 execution



Figure 1: Wilson-loop test and demonstration calculations for (a) a periodic neon reference and (b) buckled stanene. Points are the computed hybrid Wannier centers in reduced units. Degenerate centers coincide. Each loop winds once along the first reciprocal coordinate, while the transverse coordinate spans $0 \leq k_2 \leq 1/2$ at $k_3 = 0$. The neon and stanene surfaces contain five and 193 loops, respectively, and yield $\nu = 0$ and $\nu = 1$. All eight occupied spinor centers are shown.

or a Wannier fit. If $T_{\mathcal{I}}(\mathbf{k})$ is the coefficient-space action of inversion, including atom permutations, orbital parities and lattice phases, its representation in the selected subspace is

$$D_{\mathcal{I}}(\mathbf{k}) = C^\dagger(\mathbf{k})S(\mathbf{k})T_{\mathcal{I}}(\mathbf{k})C(\mathbf{k}). \quad (6.1)$$

The overlap metric distinguishes the coefficient transformation $T_{\mathcal{I}}$ from the covariant AO matrix of the inversion operator. Character decomposition gives even and odd multiplicities without assigning gauge-dependent labels to individual vectors within a mixed degenerate eigenspace. For spinful time-reversal symmetry, the number of odd occupied Kramers pairs yields the two-dimensional plane index or the three-dimensional strong and weak Fu–Kane indices [14], together with an inversion z_4 indicator in three dimensions. An exact integer comparison with the analytic inversion-subgroup elementary-band-representation (EBR) signatures also tests membership in their integer lattice and the existence of a nonnegative decomposition [15, 16]. The implementation uses inversion rather than a complete space-group EBR catalogue, and an atomic-compatible signature does not establish trivial topology. Metric, subspace, energy-commutator and time-reversal checks precede classification. Additional parity calculations recover $\nu = 0$ for neon and $\nu = 1$ for stanene. The separately versioned tests are described in the Supporting Information and leave the retained Wilson spectra unchanged.

6.2 Little groups, nonsymmorphic phases, and spin

The extension from inversion to general crystalline symmetry needs more than a list of point rotations. Write a space-group operation as $g = \{W_g|\tau_g\}$ in fractional direct coordinates, so that $x \mapsto W_g x + \tau_g$. For cell matrix A , the Cartesian operation is $R_g = AW_g A^{-1}$. Reciprocal fractional vectors transform with W_g^{-T} , not in general with W_g . An antiunitary operation $g\Theta$ additionally reverses $\mathbf{k}$.

At a fixed fractional $\mathbf{k}$, the little group contains operations mapping $\mathbf{k}$ to itself modulo a reciprocal vector. Choose one representative per coset modulo integer translations of the cell. Composition gives

$$W_{gh} = W_g W_h, \quad L(g, h) = \tau_g + W_g \tau_h - \tau_{gh} \in \mathbb{Z}^3. \quad (6.2)$$

For spinors, the $SU(2)$ lifts additionally give a sign $z(g, h)$. The matrices acting in a fixed Bloch subspace must satisfy

$$D_g {}^{a_g}D_h = \omega(g, h) D_{gh}, \quad \omega(g, h) = e^{-2\pi i \mathbf{k} \cdot L(g, h)} z(g, h), \quad (6.3)$$

where $a_g = 1$ denotes antiunitarity and ${}^1D_h = \overline{D_h}$. Physical $\Theta^2 = -1$ contributes to the spin factor. Associativity requires the twisted cocycle identity

$$\omega(g, h) \omega(gh, j) = {}^{a_g}\omega(h, j) \omega(g, hj). \quad (6.4)$$

This distinguishes screw/glide operations from a point-group approximation and double-valued representations from scalar ones [17, 19].

A conventional centered cell has distinct cosets with the same rotation and different translations. Collapsing them by rotation alone loses folded translation sectors. Conversely, character data calculated in that conventional quotient cannot be matched blindly to a primitive-cell EBR table. Both the cell transformation and representation convention must be recorded.

6.3 Metric sewing and degeneracies

Let T_g contain the atom/AO action, Bloch-image phases and spin action. The sewing matrix between selected frames is

$$B_g(\mathbf{k}) = C^\dagger(g\mathbf{k}) S(g\mathbf{k}) T_g(\mathbf{k}) {}^{a_g}C(\mathbf{k}). \quad (6.5)$$

For a closed selected subspace it is unitary and reconstructs the transformed frame. Metric normalization, closure, and energy compatibility must be checked before extracting characters. Degeneracy is not resolved by labelling individual eigenvectors: unitary rotations within a degenerate block are physically arbitrary.

For a unitary little group and a complete character table with the same factor system, multiplicities follow from

$$n_\alpha = \frac{1}{|G_{\mathbf{k}}|} \sum_{g \in G_{\mathbf{k}}} \overline{\chi_\alpha(g)} \chi(g). \quad (6.6)$$

The result must be a nonnegative integer, with consistent reconstructed character and total dimension. Antiunitary operations require corepresentation theory, not unmodified use of this character formula. Characters alone also do not determine the sewing of directed Wilson links between different $\mathbf{k}$ -points.

The native implementation constructs a complete projective character table for the supplied unitary little group, rather than importing standard-label tables. Actual GPW/GAPW Gaussian coefficient actions include the atom permutation, AO rotation, cell-image phase and, where requested, the post-SCF GTH spinor action. The selected band sewing matrices are checked against Eq. (6.3) before characters are decomposed. Failed covariance or noninteger multiplicities are rejected, not repaired by rotating eigenvectors or rounding a defective representation.

6.4 Antiunitary corepresentations

Let the little group be $M = H \cup aH$, where H is its unitary subgroup. An irreducible projective character χ_α of H has antiunitary partner

$$\chi_\alpha^a(h) = \frac{\omega(h, a)}{\omega(a, a^{-1}ha)} \overline{\chi_\alpha(a^{-1}ha)}. \quad (6.7)$$

The Wigner indicator is [20]

$$w_\alpha = \frac{1}{|H|} \sum_{b \in aH} \omega(b, b) \chi_\alpha(b^2) \in \{+1, -1, 0\}. \quad (6.8)$$

Type +1 retains one copy of the unitary irrep, type -1 requires two copies, and type 0 pairs two inequivalent partner irreps. The native implementation determines these types and their integer unitary restriction matrix. Odd type- -1 multiplicities or unequal type-0 partner counts invalidate a selected band space. Spin and nonsymmorphic phases enter the same factor system and cannot be tested separately after those phases have been discarded.

These signs are neither topological indices nor antiunitary eigenvalues. Type +1 does not exclude Kramers degeneracy: an already multidimensional spin irrep can contain the required doublet without additional doubling. At a generic centrosymmetric SOC point, in contrast, the identity-only unitary subgroup may be doubled by the combined inversion-time-reversal operation. No canonical antiunitary eigenvector basis is assigned. The electronic-structure path currently constructs the nonmagnetic grey extension of the spatial group. The underlying algebra also accepts supplied magnetic factor systems, but this is not a magnetic space-group search or self-consistent noncollinear SOC calculation.

6.5 Character correspondence and explicit lift conventions

Equal dimensions do not identify irreducible representations. To compare a generated table with named reference characters, let $p(g)$ be a supplied operation isomorphism and let η_g relate the two representation lifts. With $z^{[a_g]} = z$ for a unitary operation and $z^{[a_g]} = \bar{z}$ for an antiunitary operation, consistency requires

$$\omega_t(g, h) = \eta_g \eta_h^{[a_g]} \bar{\eta}_{gh} \omega_r(p(g), p(h)), \quad (6.9)$$

$$\chi_t(g) = \eta_g \chi_r(p(g)), \quad g \in H. \quad (6.10)$$

The operation map must preserve products and antiunitary flags. The phases must be supplied from the actual spatial, Bloch and spin conventions: solving a factor-system coboundary alone cannot fix names, since an additional one-dimensional character can preserve the factors while permuting equal-dimensional irreps. In particular, the spin lifts returned by the factor construction are reused; independently recomputing a lift at a quaternion branch tie can choose the other $SU(2)$ representative through rounding. This includes some cubic rotations as well as π rotations. The Gaussian SOC sewing path uses these same stored lifts, not a second Cartesian-to-spin conversion. For an antiunitary operation the stored spin factor already contains $J = i\sigma_y$; this factor is removed before the frame transformation applies JK , so that physical time reversal is included exactly once. The conjugate spin convention of the CP2K SOC Hamiltonian is retained.

Orthogonality alone is also insufficient to certify a reference table. Let L_g denote the left regular action of the twisted unitary group algebra, $d_\alpha = \chi_\alpha(e)$, and $\nu_\alpha = |H|^{-1} \sum_{g \in H} |\chi_\alpha(g)|^2$. The native matcher constructs

$$P_\alpha = \frac{d_\alpha}{\nu_\alpha |H|} \sum_{g \in H} \overline{\chi_\alpha(g)} L_g \quad (6.11)$$

and checks Hermiticity, idempotence and commutation with every L_g , in addition to character orthogonality and $\sum_\alpha d_\alpha^2 / \nu_\alpha = |H|$. For ordinary unitary irreps $\nu_\alpha = 1$; paired and doubled Wigner restrictions have norms 2 and 4, respectively. These tests certify the unitary restrictions, not their extension to antiunitary corepresentations, which is checked separately as described above. Only a unique bijection of the complete character columns is accepted. The reference interface does not by itself standardize an arbitrary input cell, name a complete EBR catalogue, or establish global band connectivity.

6.6 Nonsymmorphic band compatibility on explicit segments

For a directed connection from $\mathbf{k}_a$ to $\mathbf{k}_b + \mathbf{G}$, define $\mathbf{d} = \mathbf{k}_b + \mathbf{G} - \mathbf{k}_a$. A segment symmetry must satisfy

$$\sigma_g W_g^T \mathbf{d} = \mathbf{d}, \quad \sigma_g = (-1)^{a_g}, \quad (6.12)$$

within the fractional-coordinate tolerance, not merely modulo a reciprocal vector. Intersection of the endpoint little groups is insufficient: ordinary time reversal can fix two TRIM endpoints without preserving any interior point of their connecting segment.

To compare representations at the endpoints, remove the continuously varying translational part using

$$\eta_g(\mathbf{k}) = e^{2\pi i \mathbf{k} \cdot \boldsymbol{\tau}_g}, \quad \bar{\omega}(g, h) = \eta_g^{a_g} \eta_h \eta_{gh}^* \omega(g, h). \quad (6.13)$$

The conjugation for an antiunitary first factor is essential. Both endpoints must produce the same rephased factor on the segment subgroup. Its unitary projective characters provide a common basis for integer endpoint-to-line branching matrices B_a, B_b . With selected endpoint multiplicities m_a, m_b , compatibility requires

$$B_a m_a = B_b m_b. \quad (6.14)$$

The destination rephasing uses $\mathbf{k}_b + \mathbf{G}$, not just $\mathbf{k}_b$. For example, a screw eigenvalue branch may exchange with another after one reciprocal period. Ignoring $\mathbf{G}$ would erase this monodromy. Integer changes of coset representatives cancel between the characters and the rephasing.

Endpoint corepresentations are restricted as whole representations. If R_a and R_ℓ give the unitary restrictions of endpoint and line corepresentations, their branching C_a satisfies

$$R_\ell C_a = B_a R_a. \quad (6.15)$$

If the line has no antiunitary stabilizer, $B_a R_a$ instead describes the branching into unitary line irreps. This retains endpoint partner constraints even when time reversal is absent on the interior. A spinful glide model illustrates the distinction: opposite glide partners at Gamma become same-branch pairs at the boundary, preventing an isolated two-band connection but allowing a four-band one. For valid corepresentations, unitary and corepresentation line-count compatibility are equivalent; the latter resolves the partner structure rather than introducing a new invariant.

These operations are implemented on explicit NNKP or generated Wilson links through `LITTLE_GROUP_COMPATIBILITY`, which implies `LITTLE_GROUP_IRREPS`. Compatible endpoints do not prove a gap between sampled points, a trivial Bloch bundle, or an atomic-band decomposition. Automatic complete Brillouin-zone connectivity and standard crystallographic irrep labels remain separate work.

6.7 Site-induced atomic band representations

The native induction kernel starts from a supplied fractional site q and constructs its orbit $q_j = t_j q$ modulo integer translations. Representatives t_j are adjusted by lattice translations to reach the chosen positions exactly. Localized orbitals carry irreps of the site stabilizer; an antiunitary stabilizer instead requires the complete corepresentations described above. Their unitary restriction characters are denoted by χ_α .

For a unitary little-group operation g that fixes site j modulo translations, set

$$L_{gj} = W_g q_j + \boldsymbol{\tau}_g - q_j \in \mathbb{Z}^3, \quad h_j = t_j^{-1} \{I - L_{gj}\} g t_j. \quad (6.16)$$

The induced character is

$$\chi_\alpha^\uparrow(g; \mathbf{k}) = \sum_{j: gq_j = q_j \bmod \mathbb{Z}^3} e^{-2\pi i \mathbf{k} \cdot L_{gj}} \frac{z(g, t_j)}{z(t_j, h_j)} a_{t_j} \chi_\alpha(h_j). \quad (6.17)$$

This is the Bloch form of site induction [36], with the spin factor in the same convention as Eq. (6.3). Antiunitary orbit representatives conjugate the onsite character even though g itself is unitary. Omitting this conjugation loses complex orbital partners.

The implementation retains conventional-cell centering cosets and keeps each local representation column fixed across $\mathbf{k}$ points. Its band rank is the orbit multiplicity times the local dimension. No full band diagonalization or orbital gauge choice is needed to generate these atomic reference

characters. Validation compares their little-group decompositions and segment restrictions with independent Cartesian s/p/d orbital and spin traces. Site induction alone neither enumerates all Wyckoff positions nor establishes that an induced representation is elementary.

6.8 Generating site-symmetry families

The possible atomic reference positions need not be occupied by atoms in the input structure. A separate geometric kernel obtains them from the complete supplied space group. For each operation it solves

$$(W_g - I)q = -\tau_g \pmod{\mathbb{Z}^3} \quad (6.18)$$

and closes the resulting connected fixed sets under nonempty intersection. Saturating the integer row lattice yields primitive normals B for each connected affine torus,

$$Bq = c \pmod{\mathbb{Z}^r}, \quad d = 3 - r. \quad (6.19)$$

Distinct components are retained: for example, $2q_x = 0 \pmod{1}$ contains both $q_x = 0$ and $q_x = 1/2$. Integer right-hand sides are enumerated from the bounds of the unit cube rather than by sampling a coordinate mesh. Subsequent space-group equivalence identifies the families, and a generic representative is checked against its entire site stabilizer.

Closure containment supplies specialization relations between families. A site group is maximal when its family has no higher-symmetry specialization. This provides the geometric input to site induction, but does not replace the representation-level elementarity tests needed for an EBR catalogue. The construction retains the supplied cell and origin conventions and does not assign standard Wyckoff letters.

A separate native matching layer can attach an independently supplied affine reference catalogue to these families. For a reference $q = q_0 + D\alpha$ in the same cell and origin convention, a group operation must satisfy

$$BW_g D = 0, \quad B(W_g q_0 + \tau_g) - c \in \mathbb{Z}^r, \quad \text{rank } D = 3 - r. \quad (6.20)$$

The tangent identity uses bounded integer arithmetic. Orbit multiplicities must also agree, and the complete correspondence must be bijective. Checking the whole affine family avoids treating a nongeneric reference origin as a generic site. Independent generators and Wyckoff coordinates provide a convention-aware validation of this layer; standard letters remain supplied metadata rather than labels inferred by the Gaussian analysis. This geometric correspondence does not identify onsite irreps with equal dimension.

6.9 Real-space specialization and composite witnesses

Matching little-group labels at sampled momenta is weaker than equivalence of band representations. A stronger constructive test uses a symmetry-preserving displacement of Wannier centers and transitivity of induction [37]. Let q lie in a generic site family with stabilizer H , and let q' be a higher-symmetry endpoint with stabilizer H' . If the lifted path from q to q' is fixed by H , then induction of an onsite representation σ gives

$$\sigma \uparrow H' = \bigoplus_{\beta} B_{\beta\sigma} \rho_{\beta}, \quad \text{BR}_{q,\sigma} \simeq \bigoplus_{\beta} B_{\beta\sigma} \text{BR}_{q',\beta}. \quad (6.21)$$

Unlike a fit of reciprocal signatures, this supplies a real-space equivalence witness. A maximal-site representation can still be composite if it is equivalent through such paths to a direct sum at another site; maximal site symmetry alone is insufficient.

The native implementation solves for an original operation g and integer lift L such that $y = W_g q' + \tau_g + L$ lies in the source's lifted affine closure. Every source stabilizer is checked against

$$(W_h - I)(q - y) = 0, \quad h \in H, \quad (6.22)$$

without reduction modulo lattice vectors. The destination stabilizer is represented as a finite group fixing y exactly, using translations $y - W_h y$. Its orbit of q gives the finite induced representation. Physical spin factors and antiunitary operations are retained. Transport to a different destination orbit image preserves the original seed's representation-column labels.

Decomposition uses complete onsite irreps or corepresentations. For the latter, unitary restrictions are orthogonal but need not have unit character norm. The implementation therefore divides each character inner product by the reference norm and checks the full reconstructed character, integer multiplicities and band dimension. Singleton inductions define equivalence classes; multi-summand inductions provide explicit composite witnesses. Recursive expansion terminates because every summand has strictly smaller positive whole-band dimension. For the resulting nonnegative matrix E , independently generated reciprocal signatures and ranks obey

$$AE = A, \quad N^T E = N^T. \quad (6.23)$$

The output retains lifted endpoints, branching matrices, equivalent representatives and decomposition provenance. It does not identify columns solely from equality in A . An unreduced generator is not, by itself, a certificate of canonical EBR identity or a verdict on the topology of the actual electronic bands.

6.10 Reciprocal fixed families and primitive cycles

The same affine-torus machinery also constructs reciprocal-space symmetry families, consistent with the relation between wave-vector types and reciprocal-space Wyckoff geometry [18]. For an original operation with antiunitary flag a_g , the effective fractional reciprocal action is

$$K_g = (-1)^{a_g} W_g^{-T}, \quad (K_g - I)\mathbf{k} = 0 \pmod{\mathbb{Z}^3}. \quad (6.24)$$

Identical K_g are identified only for geometric enumeration. The full original stabilizer is restored at each generic seed: centering cosets, nonsymmorphic translations and a possible antiunitary PT stabilizer still enter the representation factors. Translation-free reciprocal geometry does not justify discarding those operations.

Each connected closure has saturated normals N and equation $N\mathbf{k} = c \pmod{\mathbb{Z}^r}$. A Smith certificate $UNV = D$ gives a primitive basis $V_{:,r+1:3}$ of its integer tangent lattice. These integer vectors generate torus cycles; an arbitrary floating-point nullspace need not generate the full lattice. For each specialization, an original operation g and integer shift G place its seed in the parent's lifted closure,

$$N_{\text{parent}}(K_g \mathbf{k}_{\text{child}} + G - \mathbf{k}_{\text{parent}}) = 0. \quad (6.25)$$

The code checks closure containment for every child tangent direction and solves the lift as an exact integer equation. It returns explicit points and directed segments, including self-edges with nonzero reciprocal shifts for primitive cycles. Symmetry-related child images are evaluated as additional points rather than matched through assumed irrep labels.

The input mode `KPOINTS_SOURCE SYMMETRY` connects this geometry directly to the Gaussian post-SCF analysis. It diagonalizes the frozen Hamiltonian at the generated points and enables both little-group and segment-compatibility checks, without changing the SCF density or mesh. Uniform interior sampling is controlled by `LITTLE_GROUP_PATH_POINTS`, including the endpoints. If successive lifted samples are $\mathbf{k}_j + G_j$, the stored subedge uses $G_{j+1} - G_j$, not just the target offset. Thus wrapping and deduplicating the sampled points preserve the full directed path and its reciprocal winding, including nonsymmorphic monodromy.

At every retained sample, the selected bands must be separated from computed excluded bands and pass the metric, subspace, Hamiltonian and sewing-law tests. Optional `REQUIRE_GLOBAL_GAP` additionally requires a positive sampled indirect gap for a lowest-band prefix. The name does not imply a rigorous gap certificate between samples; sampling convergence and independent BZ-wide isolation remain necessary. The output retains the geometry, eigenvalues, representations and

directed restrictions. These paths are not a Wannier90 neighbour mesh or Wilson surface, so no overlap file is synthesized for them.

Removing higher-symmetry subsets can disconnect a fixed torus, so connected closures are not a complete decomposition into open cells. Their incidence graph and primitive cycles do not by themselves establish complete equivariant band connectivity, primitive-cell equivalence or EBR elementarity.

6.11 Atomic signature matrices on explicit reciprocal sets

Combining the generated sites with the induction kernel gives a native integer signature matrix for a supplied reciprocal set $\mathcal{K}$,

$$A_{(\mathbf{k},\beta),(q,\alpha)} = n_\beta \left[\chi_{q,\alpha}^\uparrow(\cdot; \mathbf{k}) \right], \quad \mathbf{k} \in \mathcal{K}. \quad (6.26)$$

The column (q, α) retains one Wyckoff-family representative and one onsite irrep or corepresentation at every point. Rows use unitary irreps unless the little group has an antiunitary coset; in that case they count complete corepresentations after checking the required unitary partner multiplicities. Consequently,

$$\sum_{\beta} d_{\mathbf{k}\beta} A_{(\mathbf{k},\beta),(q,\alpha)} = N_{q\alpha} \quad (6.27)$$

is the same physical band rank for every $\mathbf{k}$. Counts are not generally numbers of Kramers pairs: corepresentation dimensions and pairing types are retained explicitly.

The implementation preserves all sites, including nonmaximal ones, and does not identify columns merely because their sampled signatures coincide. An incoming unitary character vector can be decomposed in the same row basis after matching its operation indices, provided it uses the same spin/Bloch convention. This algebraic check does not replace Hamiltonian covariance or selected-subspace closure for actual Gaussian states. The matrix remains a redundant atomic reference for the explicit set $\mathcal{K}$, not an automatically complete EBR matrix: reciprocal-stratum completeness and representation elementarity remain separate from the integer decomposition of this supplied matrix.

6.12 Exact signed and nonnegative signature decompositions

For an integer signature matrix $A \in \mathbb{Z}^{m \times n}$, the native integer kernel constructs a Smith form [36],

$$UAV = D, \quad D = \text{diag}(d_1, \dots, d_r, 0, \dots), \quad d_i > 0, \quad d_i \mid d_{i+1}, \quad (6.28)$$

where U and V are unimodular integer matrices. With $c = Ub$, the equation $Ax = b$ has an integer solution precisely when

$$c_i = 0 \quad (i > r), \quad c_i \equiv 0 \pmod{d_i} \quad (i \leq r). \quad (6.29)$$

A witness is obtained as $x = Vy$, with $y_i = c_i/d_i$ for $i \leq r$ and zero free components. The implementation checks both $UAV = D$ and $Ax = b$ using bounded 64-bit integer arithmetic. Rank decisions do not use floating-point thresholds. Growth beyond the arithmetic bound or exhaustion of the factorization work limit returns an unresolved status, not a negative membership decision.

For the nonnegative signature matrices used here, the stronger condition $Ax = b$, $x \in \mathbb{Z}_{\geq 0}^n$, is tested separately. At a search node with residual $b' \geq 0$, a nonzero column j has the finite bound

$$0 \leq x_j \leq \min_{i: A_{ij} > 0} \left\lfloor \frac{b'_i}{A_{ij}} \right\rfloor. \quad (6.30)$$

Row-wise gcds of the remaining columns and their zero supports prune the complete enumeration; zero columns can be set to zero. Every accepted witness is checked exactly. A caller-specified node

budget bounds the potentially exponential search. An exhausted budget is explicitly undecided and must not be interpreted as absence of an atomic decomposition.

These are decisions about the supplied columns and sampled rows. The Smith divisors of an incomplete reciprocal sample are not, by themselves, the space group's symmetry-indicator group. Nonnegative membership likewise does not prove triviality of a Bloch bundle. The integer routines do not add an automatic general-classification input mode or replace spectral isolation and compatibility checks.

6.13 Joint compatibility lattice and atomic quotient

The separate edge tests can be assembled into one integer system. Let $n \in \mathbb{Z}^N$ contain the point-specific irrep or full corepresentation multiplicities, and let P_a select the block at point a . For an edge $e : a \rightarrow b + G_e$, the unitary line restrictions give a block row

$$C_e = B_a R_a P_a - B_b R_b P_b, \quad Cn = 0, \quad (6.31)$$

where R_a is the endpoint corepresentation restriction, or the identity at a purely unitary point. The line corepresentation restriction is injective: its disjoint unitary supports and prescribed pairing or doubling uniquely determine valid line counts. Thus these unitary equations also enforce equality of the full line corepresentations. When $a = b$, both terms act on the same block, but their rephasing still uses G_e . A screw-loop relation must not be discarded as an edge connecting a point to itself.

Symmetry-equivalent points must not carry independent labels. Let s map k_a to $k_b + G$, and for a destination little-group operation h set $h_a = s^{-1}hs$. The actual coset representatives obey $hs = \{I|L\}sh_a$, with

$$L = \tau_h + W_h \tau_s - \tau_s - W_s \tau_{h_a} \in \mathbb{Z}^3. \quad (6.32)$$

Writing z for the spin factor at Gamma, including physical time-reversal squaring, the transported unitary character is

$$\chi_b(h) = \frac{z(h, s)}{z(s, h_a)} e^{-2\pi i k_b \cdot L} \begin{cases} \chi_a(h_a), & s \text{ unitary,} \\ \chi_a(h_a)^*, & s \text{ antiunitary.} \end{cases} \quad (6.33)$$

Independent integer changes of endpoint representatives therefore retain the necessary Bloch phase. We check conjugation of the complete product table and the twisted factor law, including antiunitary little-group operations. Character decomposition must give a bijective unitary-irrep map. Restriction to the unitary subgroup then determines the full corepresentation map M_s , which must preserve dimensions. A spanning set of star relations adds $M_s P_a - P_b$ to C . Edge winding and star identification remain distinct constraints.

For the supplied graph $\mathcal{G}$, define

$$\mathcal{B}_{\mathcal{G}} = \ker_{\mathbb{Z}} C, \quad \mathcal{X}_{\mathcal{G}} = \mathcal{B}_{\mathcal{G}} / \text{im}_{\mathbb{Z}} A. \quad (6.34)$$

This follows the compatible-representation/atomic-lattice construction of symmetry indicators [16], but explicitly retains the scope of the supplied graph. A Smith factorization $U_C C V_C = D_C$ of rank s supplies the saturated integer kernel $K = (V_C)_{:,s+1:N}$. Saturation matters: rescaling rational nullspace vectors separately need not produce the entire integer kernel. Every atomic column must admit an exact solution $A = KY$; failure invalidates the reference/compatibility construction.

Writing $q = N - s$, the Smith form $U_Y Y V_Y = D_Y$ of rank r gives

$$\mathcal{X}_{\mathcal{G}} \simeq \mathbb{Z}^{q-r} \oplus \bigoplus_{i=1}^r \mathbb{Z}_{d_i}. \quad (6.35)$$

Factors with $d_i = 1$ are trivial. For a compatible Gaussian signature $b = Kz$, its class is $U_Y z$ modulo the nonzero divisors, with the remaining integer coordinates retained. The class vanishes

precisely when a signed atomic decomposition exists. Nonnegative decomposition is a stronger, separate question. An incompatible signature has no quotient class; it is not assigned a nonzero topological index.

The native implementation checks $CK = 0$ and $KY = A$ and exports the matrices, Smith row map, factors and target residues for independent verification. It removes duplicate homogeneous equations without replacing the integer kernel by a floating-point nullspace. Resource or coefficient-growth limits remain unresolved outcomes. These calculations are replicated on MPI ranks, not distributed sparse integer factorizations.

Star relations between supplied points are now included. Missing points or global compatibility relations can leave free factors in Eq. (6.35). They are diagnostics of the supplied lattices, not additional physical topological invariants. Conversely, vanishing free rank only means that the atomic columns span the compatible space rationally at this sample. It does not certify omitted congruence constraints, complete reciprocal coverage, EBR completeness or spectral isolation throughout the Brillouin zone.

6.14 Connecting atomic references to Gaussian bands

The optional `ATOMIC_SIGNATURES` analysis connects the native atomic reference and integer kernels to the validated Gaussian-band representations. It uses explicit NNKP, Wilson or TRIM points, or the generated SYMMETRY points, and implies the little-group analysis. Each selected band space must be separated from computed excluded bands at the sampled point. Metric covariance, subspace closure, Hamiltonian commutation and the projective sewing law are checked before comparison. This supports scalar GPW/GAPW and the existing post-SCF GTH-SOC treatment; physical time reversal requires a restricted SCF. It does not establish a gap between sampled points.

The band and reference operation lists are matched bijectively by their fractional rotations, antiunitary flags and translations modulo the input lattice, not by their numerical list indices. If the reference representative is $g' = \{I|L_g\}g$, its character changes by

$$p_g = \exp(-2\pi i \mathbf{k} \cdot L_g), \quad \chi'(g') = p_g \chi(g). \quad (6.36)$$

The corresponding factor transformation must also agree,

$$\omega'(g, h) = \frac{p_g^{a_g} p_h}{p_{gh}} \omega(g, h), \quad (6.37)$$

with conjugation when g is antiunitary. After this exact convention conversion, characters are decomposed into the reference row basis. Reordered local irrep labels do not affect the result. Different band ranks, missing operations or inconsistent factors are rejected rather than repaired by fitted gauges.

The output retains the reference operations, characters and corepresentation restrictions, the integer matrix and target, and the site/local-representation identity of each column. Available signed and nonnegative witnesses accompany the membership decisions. The nonnegative search budget is controlled by `ATOMIC_SIGNATURE_MAX_NODES`; exhaustion is reported as unresolved. Invalid data or failed witness verification abort instead of producing a physical classification. Segment compatibility can be requested separately for NNKP/Wilson connections and is automatic for SYMMETRY paths. With atomic signatures enabled, these connections also define the joint lattice of Eq. (6.34). The comparison does not automatically identify complete reciprocal strata, standard EBR labels or the topology of an isolated Bloch bundle.

6.15 Band representations and limits of indicators

Elementary band representations (EBRs) provide the indecomposable building blocks of atomic band representations. Restricting them to little groups yields a vector of irrep multiplicities.

Compatibility relations constrain the vectors that can connect along high-symmetry lines. A candidate symmetry vector n can be compared with an integer matrix of EBR signatures,

$$Eq = n. \quad (6.38)$$

Integer solvability, nonnegative solvability, and compatibility constraints are different questions. A symmetry indicator is a quotient of compatible representation data by atomic data; it can miss topology invisible to those symmetry eigenvalues [15, 16]. An atomic-compatible signature is not proof of a trivial Bloch bundle. A signed-only decomposition is not, without additional information, a complete diagnosis of fragile topology.

The inversion-subgroup classifier retains its separate analytic signatures and named indicators. The general native little-group/corepresentation, explicit-segment, Wyckoff-family and site-induction machinery now also tests actual band signatures against the generated atomic columns, but does not supply a complete all-space-group classification. Conventional-to-primitive matching, standard labels, complete connectivity and a validated catalogue with elementarity checks are still required. Algebra tests over all ordinary space groups must not be presented as a completed classification of all their materials.

7 Parameter-space topology and phasons

A separate native postprocessor, `topology_phasons`, extends the overlap construction to closed two- or four-parameter families, including periodic atomic displacements (phasons) combined with reciprocal coordinates. It reads `STATE_EXPORT` snapshots and evaluates $M_{ab} = C_a^\dagger O_{ab} C_b$, where O_{ab} includes the Gaussian centers of both geometries, periodic images, and Bloch and boundary-sewing phases. Polar-unitary links give C_1 from determinant plaquette phases [23] and a four-parameter C_2 estimate from full non-Abelian plaquette logarithms [24]. Here C_2 denotes the second Chern character pairing, not generally the second Chern class. The postprocessor requires neither Python, Z2Pack, nor Wannier90 at runtime. It operates on fixed-cell, fixed-rank families with explicit physical endpoint identifications. The Supporting Information gives its separate development version, discretization, and model and Gaussian-snapshot tests. These tests complement the material examples below without changing their Hamiltonians, sampling, or results.

For a discrete parameter family, let $Q_\mu(x)$ denote the polar link from x to $x + \hat{\mu}$. An oriented plaquette is

$$P_{\mu\nu}(x) = Q_\mu(x) Q_\nu(x + \hat{\mu}) Q_\mu^\dagger(x + \hat{\nu}) Q_\nu^\dagger(x). \quad (7.1)$$

With the implemented directed-overlap convention, $C_1^{\text{mesh}} = -(2\pi)^{-1} \sum_x \arg \det P_{12}(x)$. For four parameters write $F_{\mu\nu}(x) = \log P_{\mu\nu}(x)$ using the principal matrix logarithm. The non-Abelian estimator is

$$C_2^{\text{mesh}} = -\frac{1}{4\pi^2} \sum_x \Re \text{Tr}[F_{12}F_{34} - F_{13}F_{24} + F_{14}F_{23}](x). \quad (7.2)$$

The logarithms already include the discrete plaquette area. Mesh refinement, physical endpoint identifications, and a fixed isolated rank are essential. Coefficient arrays from different geometries cannot be contracted without cross-geometry AO overlaps. The numerical value is retained before rounding, and a sampled gap does not certify isolation between sampled geometries.

8 Spectral localizers in finite and periodic systems

8.1 Energy-position tuples and Clifford structure

A spectral localizer combines operators associated with several observables into one self-adjoint matrix. For two positions and a Hamiltonian, Pauli matrices provide a Clifford representation. In

an orthonormal space define

$$\tilde{L} = (\tilde{H} - EI) \otimes \sigma_z + \kappa(\tilde{X} - xI) \otimes \sigma_x + \kappa(\tilde{Y} - yI) \otimes \sigma_y, \quad (8.1)$$

up to the explicitly fixed tensor ordering. The parameter $\kappa > 0$ has units of energy/length. It balances energy and position resolution rather than correcting the Hamiltonian. The connection between a localizer signature and an even index pairing holds under appropriate locality, gap, and finite-volume bounds [25]. Those hypotheses remain relevant when interpreting a finite Gaussian model.

In auxiliary-index-first ordering the covariant AO matrix is

$$L = \begin{pmatrix} H - ES & \kappa[(X - xS) - i(Y - yS)] \\ \kappa[(X - xS) + i(Y - yS)] & -(H - ES) \end{pmatrix}, \quad B = \begin{pmatrix} S & 0 \\ 0 & S \end{pmatrix}. \quad (8.2)$$

For SOC, H and S here denote the physical spinor matrices. Auxiliary doubling is separate from spin, so the localizer dimension is four times the scalar AO count. All AO directions are retained: there is no hidden occupied-space projection or automatic removal of nearly dependent basis functions.

8.2 Index, metric gap, and stability

For class A define

$$\nu_A = \frac{1}{2} \text{Sig}(L) = \frac{1}{2}(n_+ - n_-), \quad \mu = \min_j |\lambda_j|, \quad Lv_j = \lambda_j Bv_j. \quad (8.3)$$

Signature is invariant under nonsingular congruence, whereas ordinary eigenvalues of a covariant AO matrix are not. Inertia can therefore be evaluated directly from L , but the protection gap must retain B . Equivalently, μ is the smallest absolute eigenvalue of $B^{-1/2}LB^{-1/2}$; this identity does not require forming that dense matrix.

At fixed metric, the index cannot change under a perturbation satisfying

$$\|B^{-1/2}\delta LB^{-1/2}\| < \mu. \quad (8.4)$$

This finite-matrix statement gives local protection an operational meaning. If the metric changes, the corresponding orthonormalized problems must be compared, including the effect of δB . An unresolved near-zero gap yields an *unresolved* index, not evidence for trivial topology.

Moving the query and coordinate operators together, shifting the Hamiltonian and query energy together, or changing AO basis must leave the generalized gap unchanged. Reversing the oriented plane reverses the class A sign. Model comparisons establish the sign convention before applying it to a material.

8.3 Inertia and gap require different calculations

A pivoted Hermitian factorization has the form

$$P^\dagger LP = \mathcal{L}D\mathcal{L}^\dagger. \quad (8.5)$$

Here $\mathcal{L}$ is unit lower triangular and D contains 1×1 and 2×2 blocks. Sylvester's law transfers the inertia from L to D . For a Hermitian block,

$$D_b = \begin{pmatrix} a & b \\ \bar{b} & d \end{pmatrix}, \quad \lambda_{\pm} = \frac{a+d}{2} \pm \sqrt{\left(\frac{a-d}{2}\right)^2 + |b|^2}.$$

Stable evaluation must retain scaling and both signs. A zero-diagonal block with $b \neq 0$ has one positive and one negative direction; counting its diagonal entries is wrong. The dense reference

uses LAPACK ZHETRF, with DSYTRF for exactly real matrices [26]. A small imaginary part is not discarded to select real arithmetic.

Pivot magnitudes are not generalized gaps. For $\sigma > 0$, shifted inertia counts satisfy, away from singular endpoints,

$$\begin{aligned} n_-(L - \sigma B) - n_-(L) &= \#\{j : 0 < \lambda_j < \sigma\}, \\ n_-(L) - n_-(L + \sigma B) &= \#\{j : -\sigma < \lambda_j < 0\}. \end{aligned} \quad (8.6)$$

These counts bracket the nearest positive or negative generalized eigenvalue. Factorization failure or an ambiguous endpoint must not be treated as a successful certificate. The reported bounds are finite-precision numerical brackets, not interval-arithmetic guarantees.

8.4 Pfaffian orientation and physical time reversal

For a real skew-symmetric matrix A of even order, $\det A = \text{Pf}(A)^2$: the determinant loses the Pfaffian sign. Moreover, $\text{Pf}(P^T A P) = \det(P) \text{Pf}(A)$. An odd simultaneous row/column permutation reverses that sign without changing the determinant. Ordinary inertia cannot replace a skew factorization with explicit permutation bookkeeping.

For spinful time reversal with $\Theta^2 = -1$, the localizer has a real skew representation [27, 28]. In auxiliary/spin/AO ordering, let $C = J_{\text{aux}} \otimes J$ and $Q = (I - iC)/\sqrt{2}$. The symmetry relations imply that

$$A = iQ^\dagger L Q \quad (8.7)$$

is real and skew-symmetric. Both properties are checked numerically, with residuals small relative to the independently resolved metric gap. An arbitrary complex AO basis change also transforms time reversal; keeping J fixed would generally be inconsistent.

Let A_0 be the corresponding transform of $L_0 = \text{diag}(S, -S)$. The reference-calibrated index is

$$(-1)^{\nu_{\text{AH}}} = \text{sgn Pf}(A) \text{sgn Pf}(A_0). \quad (8.8)$$

Both factors use the same ordering and orientation. The dense algorithm accumulates signs rather than an overflowing Pfaffian product. This is a time-reversal local index, not an inversion-parity test.

The optional sparse skew-LDL backend implements a separate numerical task from MUMPS inertia [28]. Its present single-rank factorization must not be described as an MPI-distributed Pfaffian solver. The motivating photonic and aperiodic examples [9, 28] provide methodological context, not first-principles material validation for the CP2K path.

8.5 Periodic positions and the torus construction

A Cartesian coordinate truncated to a primitive cell is discontinuous at the periodic seam. Using it in the open-boundary localizer is not a periodic formulation. Instead, analytic periodic functions are used [29]. For analysis-torus reciprocal vectors $\mathbf{G}_j$ and $\phi_j = \mathbf{G}_j \cdot \mathbf{r}_0$, define

$$\begin{aligned} C_j &= \langle \phi_\mu | \cos(\mathbf{G}_j \cdot \mathbf{r} - \phi_j) | \phi_\nu \rangle, \\ T_j &= \langle \phi_\mu | \sin(\mathbf{G}_j \cdot \mathbf{r} - \phi_j) | \phi_\nu \rangle. \end{aligned} \quad (8.9)$$

In the implemented orientation,

$$L_{\text{per}} = \begin{pmatrix} H - ES - \eta(2S - C_1 - C_2) & \eta(T_1 - iT_2) \\ \eta(T_1 + iT_2) & -H + ES + \eta(2S - C_1 - C_2) \end{pmatrix}. \quad (8.10)$$

The energy scale η is distinct from κ , and the metric remains $\text{diag}(S, S)$. An overall sign and auxiliary-space conjugation relative to the cited periodic construction make its orientation consistent with the finite localizer used here.

The trigonometric operators are integrated directly, not evaluated as matrix functions of finite projected Cartesian positions. Projection need not preserve either exact unitarity of exponentials or their commutativity. The theorem for a gapped finite-range lattice Hamiltonian therefore does not automatically become a theorem for an arbitrary Gaussian projection. The output is an AO-torus diagnostic requiring independent basis-, torus-, and scale-converged material checks.

Explicit analysis k-points describe the enlarged torus through its complete mesh. Periodic exponentials shift between mesh sectors, so their cross-k links must be retained. A symmetry-reduced SCF can supply the potential, but scalar SCF weights cannot replace the full torus operator. The present periodic localizer covers two-dimensional periodic planes; a general three-dimensional AII implementation is not claimed.

8.6 Connection to the quadratic pseudospectrum

For Hermitian operators A_j and anticommuting Hermitian Clifford matrices γ_j , squaring $L = \sum_j A_j \otimes \gamma_j$ gives

$$L^2 = \left(\sum_j A_j^2 \right) \otimes I + \sum_{j < l} [A_j, A_l] \otimes \gamma_j \gamma_l. \quad (8.11)$$

The first term measures simultaneous approximate localization; the second retains noncommutativity. The quadratic gap and localizer gap are related, but not interchangeable. In an AO basis each projected product must include the metric, or be evaluated after orthonormalization.

9 Quadratic pseudospectrum and localized approximate states

The quadratic pseudospectrum asks for a normalized state that minimizes the combined residual of incompatible observables [30]. For a query $(E, \mathbf{x})$, set $A_0 = H - ES$, $A_j = X_j - x_j S$, $w_0 = 1$, and $w_j = \kappa$ for $j = 1, 2, 3$. The AO problem is

$$Qc = \lambda Sc, \quad Q = \sum_{j=0}^3 w_j^2 A_j^\dagger S^{-1} A_j, \quad c^\dagger Sc = 1. \quad (9.1)$$

The minimum $\sqrt{\lambda}$ is the quadratic gap. Its eigenvector is a joint approximate state, and several low eigenpairs can describe a localized subspace. The separated residuals are

$$\begin{aligned} \delta E^2 &= (A_0 c)^\dagger S^{-1} (A_0 c), \\ \delta x_j^2 &= (A_j c)^\dagger S^{-1} (A_j c), \\ \lambda &= \delta E^2 + \kappa^2 \sum_j \delta x_j^2. \end{aligned} \quad (9.2)$$

Expectations $c^\dagger H c$ and $c^\dagger X_j c$ distinguish bias from spread around the query. These states need not be eigenstates of H . Within degenerate minimizing spaces, projectors or subspace observables must be compared instead of individual coefficients.

Equation (9.1) is positive semidefinite and covariant under Eq. (2.3). Ordinary matrix products without metric solutions would destroy that invariance. The position residual concerns the projected position: $X_j S^{-1} X_j$ is not an independently integrated continuum $\langle r_j^2 \rangle$. Small bases can underestimate continuum localization errors even when the eigenproblem is solved accurately.

The dense reference uses a complete orthonormalized space. The iterative path applies Q without forming it: multiply by A_j , solve with S , multiply by $A_j^\dagger$, and accumulate. This avoids fill-in from explicitly squaring sparse matrices. A complex block Rayleigh–Ritz iteration, double metric reorthogonalization, and thick restarts address clustered low states. The residual

$$\|Qc - \lambda Sc\|_{S^{-1}} \leq \epsilon_{\text{eig}} \max(1, |\lambda|)$$

checks an eigenpair in atomic units; it does not independently prove that no lower state was missed.

Reusable MUMPS factors avoid forming S^{-1} , but the present trial vectors and multiple-right-hand-side communication are not fully distributed. Output includes separated residuals, expectations, and optional complex AO coefficients. The current analysis covers isolated GPW/GAPW systems and post-SCF GTH SOC. Periodic quadratic formulations and band unfolding remain separate extensions. A small quadratic gap is a localization diagnostic, not a Chern or $\mathbb{Z}_2$ index.

10 Materials-discovery applications and demonstrative tests

10.1 Hybrid-functional topological materials databases

TopoHSE-DB provides an application example of electronic-structure-based topological classification at database scale [31]. Mirhosseini, Elcoro, Knüpfer and Kühne combine consistently relaxed nonmagnetic crystal structures, PBE and screened-hybrid HSE06 electronic structures, and topological quantum chemistry. Comparing the resulting labels exposes the sensitivity of a symmetry-based classification to the underlying exchange–correlation approximation. A converged integer or symmetry label is therefore not, by itself, evidence that the electronic Hamiltonian has been described accurately.

The original database workflow uses VASP, VASP2Trace, and CheckTopologicalMat. It is cited here as a materials-discovery application and a source of future comparison cases, not as a calculation performed with the new CP2K topology routines. Connecting such a workflow to CP2K requires equivalent electronic structure and symmetry conventions as well as the general little-group, compatibility-relation, and EBR machinery discussed in Section 6. The currently implemented inversion analysis does not yet reproduce the database’s full classification workflow.

10.2 Generative–predictive discovery of candidate materials

The submitted study by Brocai, Kühne, Knüpfer and Mirhosseini, *Generative–predictive models enable the discovery of topological materials*, provides a complementary application [32]. It couples topology-conditioned crystal generation to a predictive filter using a shared pretrained representation, followed by hybrid-functional electronic-structure and TQC analysis. The submitted manuscript reports 1,033 successfully calculated hybrid-functional band structures; 629 receive an unambiguous classification and 523 are classified as topological. The corresponding 83.15% rate applies to the classified subset, whereas the confirmed yield across all 1,033 calculated cases is 50.6%. The 404 unresolved classifications must not be silently counted as either trivial or topological.

These results, obtained with the study’s VASP-based workflow, illustrate the demand for reliable post-processing and explicit failure diagnostics. They also motivate subsequent, distinct questions: Wilson spectra can probe isolated band subspaces, Kubo calculations can address transport at specified temperature and dissipation, and localizers can investigate finite, defective, or disordered realizations of selected candidates. These are prospective CP2K follow-up applications, not calculations already completed for the reported candidate set. A TQC label alone does not supply a conductivity or a localizer protection gap.

10.3 Small calculations as tests and demonstrations

The smaller calculations in this paper and the companion k-point study serve a different purpose [33]. They provide transparent tests and demonstrations of individual algorithmic steps, not a second materials-discovery survey. Table 1 states the role of each set.

For Si and Al, the largest fixed-mesh cell-energy or electronic-free-energy difference among the compared representations is 1.4×10^{-12} hartree. This tests representation equivalence, not complete basis or k-point convergence. In particular, the Al free energy still changes by 10.721

Calculation set	Demonstration and test objective
Si and Al, companion k-point study	Quadrature dependence and equivalent full-grid, time-reversal, K290, and SPGLIB representations. At fixed 4^3 sampling, 64 eigensystems reduce to four with atomic symmetry.
Periodic neon and buckled stanene	Trivial and nontrivial Wilson-loop examples, with $\nu = 0$ and $\nu = 1$, respectively; comparison of native and external analyses.
Ne/SOC, Bi ₂ /SOC, and diamond/SOC	Primitive-mesh versus supercell and reduced versus full-property-grid checks of the metric, Hamiltonian, and current operators.
Analytic localizer models and small GPW/GAPW systems	Known indices, gap closings, basis covariance, and dense/sparse solver consistency; quadratic-state residual and subspace checks.

Table 1: Test and demonstration calculations, distinct from the materials-discovery applications cited above. The Si/Al results are retained from the companion study, not rerun or reassigned to the new topology implementation.

meV per atom between the 6^3 and 8^3 meshes. Likewise, agreement between Wilson analyses using the same overlap matrices tests the analysis stage, while independent operator identities test the AO construction. The quantitative checks in Section 11 and the Supporting Information retain these distinctions and their original calculation provenance.

11 Implementation, validation, and development boundaries

11.1 Shared operators and optional numerical backends

The finite-system property paths share full AO Hamiltonians, overlap, analytic positions, and optional GTH-SOC blocks. DBCSR supplies distributed block storage and matrix products [34]. Localizer assembly retains the union of the operator sparsity patterns without additional dropping. DBCSR is not itself a general pivoted indefinite factorization or Pfaffian solver.

The dense LAPACK path is the small-system reference. MUMPS supplies distributed symmetric-indefinite factorizations and reusable metric solutions [35]. Complex Hermitian matrices are realified:

$$\mathcal{R}(A) = \begin{pmatrix} \Re A & -\Im A \\ \Im A & \Re A \end{pmatrix}. \quad (11.1)$$

Each eigenvalue is duplicated, so inertia-based formulas compensate for that duplication. A positive metric is checked separately from the indefinite localizer. Static pivot perturbations and low-rank compression are not silently accepted as exact sign counts. Symbolic analysis is reused across appropriate shift families.

The optional Tacho skew-LDL extension is accessed through a small C interface and Fortran interoperability [39]. No MATLAB dependency is introduced. Permutation parity, skew-pivot conventions, numerical residuals, and the trivial-reference orientation are explicit. Its sparse factorization currently resides on one rank, while DBCSR assembly and MUMPS metric/gap checks can be distributed. PARDISO and PEXSI are possible alternative backends, not prerequisites and not part of the tested backend set reported here.

MUMPS factor storage and process-summed solver memory are not the whole application’s memory consumption. Sparse fill-in, source-rank coordinates, dense reference arrays, trial vectors, and MPI buffers must be accounted for separately. Similarly, single-node timings cannot establish multi-node scaling.

11.2 Three levels of validation

The first level tests algebra independently of an SCF calculation: metric congruences, origins, energy shifts, degeneracies, time reversal, permutation orientation, and analytic model indices. The second compares numerical algorithms on identical matrices: dense eigenvalues against LDL inertia, dense against sparse gap brackets, dense against skew-LDL signs, and full against iterative quadratic eigenproblems. The third exercises operator construction from actual converged CP2K GPW/GAPW calculations.

Retained checks include open and periodic Chern models, spin-mixed time-reversed model pairs, gap closings, singular and indefinite metrics, and deliberately broken symmetry. They establish nontrivial topology in controlled models. Small molecular and bulk integration fixtures instead establish that the AO data reach the algorithms consistently; most are not nontrivial first-principles topological materials.

The retained neon and stanene Wilson results in Section 5 are small band-topology test and demonstration calculations, with matching inversion indices. They complement the Si/Al k-point demonstrations summarized in Section 10, rather than constituting a materials-discovery application study. They were produced with their recorded executable and sampling. They are not relabelled as localizer or Kubo calculations merely because the underlying operators are now shared.

11.3 Quantitative test and demonstration calculations

Table 2 records representative comparisons from the retained development runs. Agreement is measured between calculations at identical numerical settings unless noted otherwise. These are consistency checks, not a claim of basis- and material-converged conductivity.

Comparison	Retained result and meaning
Ne/SOC primitive mesh versus Gamma supercell	A $3 \times 1 \times 2$ primitive mesh and its explicit Gamma supercell give $6.21105320 \times 10^{-9}$ S at the converged comparison settings.
Bi ₂ /SOC property symmetry	Six of 27 eigensystems suffice; the maximum printed full-tensor relative difference from the unreduced calculation is 2.71×10^{-15} .
Primitive diamond/SOC	Both K290 and SPGLIB use three of eight representatives; the full-tensor relative difference is 2.21×10^{-11} .
Bi ₂ quadratic states	Dense and iterative gaps agree within 6×10^{-15} hartree; four metric-subspace singular values differ from unity by less than 10^{-10} .

Table 2: Operator-level checks retained on 24 September 2026. Raw settings and limitations are recorded in the Supporting Information. The transport numbers are numerical fixtures, not material predictions.

For the diamond property reconstruction, metric, Hamiltonian, and current residuals are checked separately rather than inferred from the final scalar conductivity. Degenerate subspaces retain their transformed gauge. The representative count measures avoided diagonalizations, not total speedup.

The crystalline representation tests cover all 530 Hall settings of the 230 ordinary space groups, scalar and spinful factors, and their grey extensions. They validate 6,360 unitary projective character tables, 5,324 antiunitary corepresentation tables, and 12,720 directed segment comparisons; 4,462 segments retain antiunitary symmetry. Release and bounds-checked debug runs agree. The Gaussian little-group directory passes 69 numerical assertions in both serial and two-rank MPI runs, including GPW, GAPW, post-SCF SOC, screw monodromy and refined Wilson links. These counts describe algebra and small integration tests, not 230 separate converged material studies.

The Supporting Information separates this evidence from a supplied magnetic-group algebra sweep and its independent reference comparison.

The site-induction kernel adds 38,160 site/k-point cases over the same Hall settings, using three positions per cell. Their 67,824 induced local-representation columns have valid little-group decompositions and satisfy the checked directed compatibility relations. Independent Cartesian s/p/d orbital traces, with and without physical spin, give 114,480 matching character comparisons. This tests atomic reference construction rather than a Wyckoff catalogue or an EBR classification of first-principles bands.

The separate geometric enumeration generates 3,467 Wyckoff families over all 530 Hall settings. Every family matches a tabulated affine coordinate family bijectively; the largest matching residual is 2.22×10^{-16} . The independent comparison also verifies 5,648 directed specialization relations. Each generated site is accepted by the spinful induction kernel with the correct multiplicity and stabilizer. This supplies geometric reference positions, not an elementary-band classification of electronic-structure results.

Combining all generated sites into matrices at six explicit reciprocal points gives 2,120 catalogues across ordinary/grey and scalar/spinful variants. Their 31,079 atomic columns contain 707,038 integer entries. All column ranks, antiunitary partner conditions and 217,553 directed column/segment comparisons agree in serial release, bounds-checked debug and two-rank MPI runs. These are sampled atomic reference matrices, not 2,120 separate material calculations or complete EBR catalogues.

The integer layer factors all 2,120 generated matrices and recovers signed test vectors and 33,199 nonnegative witnesses. Independent small-matrix tests compare the Smith invariants with gcds of all minors and check unimodular transforms for 384 rectangular or rank-deficient matrices. All 6,642 two-Kramers-pair inversion patterns at four/eight TRIM agree with the separate analytic Fourier criterion; 18,225 additional nonnegative systems agree with exhaustive bounded coefficient enumeration. Arithmetic and search limits are tested as unresolved outcomes. This evidence concerns exact signature equations, not material topology or general EBR elementarity.

The Gaussian-band integration supplies signed and nonnegative atomic witnesses for the small He/Ne GPW, GAPW and SOC examples, including sheared cells and screw sectors. The single screw branch fails integer membership, consistently with its independent reciprocal-loop monodromy obstruction. A zero-budget TRIM test reports signed membership and unresolved nonnegative membership rather than mistaking an unfinished search for an obstruction. On all 2,120 generated reference matrices, reordered operations and irreps and integer coset shifts preserve the recovered signature after checking the independently regenerated factors. Corrupted factors and unmatched operations are rejected. The focused serial and two-rank MPI drivers each pass 135 assertions, combining six native unit programs with 129 Gaussian assertions, including automatic reciprocal sampling and the joint compatibility quotient. Another 36 export/k-point assertions pass in each configuration.

The reciprocal-geometry backend independently checks all 530 Hall settings and their grey extensions. It generates 18,700 fixed-family closures, 57,357 specialization incidences and 18,730 primitive torus cycles. Stabilizers agree with the full spinful little-group kernel; 333,900 rational-grid points have an unambiguous matching family. Coordinate changes preserve the families and incidence graph. The induced scalar/spinful atomic columns satisfy 2,481,572 directed incidence/cycle comparisons, including antiunitary restrictions when present. Serial, two-rank MPI and instrumented-kernel runs agree. These are geometric and representation tests, not additional DFT materials or a proof of complete global band connectivity.

The automatic Gaussian path is tested for He/GPW, all-electron He/GAPW, Ne/GTH-SOC and an equivalent sheared He cell. Each generates 15 fixed families, 56 points and 82 compatible directed segments. Replaying these points and shifts through explicit NNKP input gives selected eigenvalues within 2.1×10^{-13} eV and band characters within 2.2×10^{-14} ; all original segment restrictions and atomic integer witnesses agree exactly. Refining to 138 points and 164 segments retains compatibility. A deliberately selected single screw branch instead fails the band-isolation

guard at a generated crossing. Invalid sample budgets and Wilson-mode combinations are also rejected.

The joint integer construction additionally certifies the compatibility kernel and atomic quotient for every ordinary/grey and scalar/spinful Hall-setting graph. Exact tests recover the four/eight-point inversion quotients $\mathbb{Z}_2$ and $\mathbb{Z}_2^3 \times \mathbb{Z}_4$ and compare quotient classes with direct signed membership for all 6,642 two-pair signatures. Independent arbitrary-precision Smith calculations validate the exported Gaussian certificates. The automatic scalar GPW/GAPW and sheared fixtures have compatible rank six and a single $\mathbb{Z}_2$ factor; the SOC fixture has rank five and factors $\mathbb{Z}_2^2 \times \mathbb{Z}_4$. Their occupied signatures have zero class. The single screw branch is incompatible and receives no class. Star identifications additionally include nonsymmorphic phases and antiunitary conjugation. In all 2,120 Hall-setting/symmetry combinations, both the compatible rank and finite quotient factors agree with the published 230-space-group tables [16]. This is an independent algebraic benchmark, not a proof of complete EBR labels or material topology. Partial explicit graphs can still retain free factors: the refined He Wilson example changes from 23 to four after star identification and remains incomplete.

Real-space specialization tests now independently check 22,592 induction relations in all 2,120 ordinary/grey scalar/spinful Hall variants. Eight reciprocal points per relation give 180,736 character identities; recursive composite expansions preserve the complete tested signatures and band dimensions. Analytic cases exercise lifted intermediate paths, origin/cell/coset conventions, antiunitary pairing, and malformed-witness rejection. The affected-space-group coverage agrees with the exception tables of Ref. [37]. An independent comparison with the Bilbao EBR data archived by Crystalline.jl [38] also matches the maximal-site generator count for every one of the 2,120 variants. These counts are not canonical-irrep-by-irrep matches. Optimized SSMP, each of two MPI ranks and an instrumented build give identical sweep records. Arbitrary-precision checks independently validate the exported integer expansions, lifted geometry and provenance in 15 Gaussian fixtures per build. The official drivers pass 171/171 assertions in both SSMP and MPI without changing physical reference values.

An independent affine-reference comparison goes beyond total counts: all 1,731 Wyckoff families of the 230 conventional space groups match, giving 6,924 correspondences across the four symmetry variants. Simultaneous basis shear, origin shift, integer coset changes and operation reversal preserve the named specialization graphs. At each individual Wyckoff position, the primitive-cell band-dimension multiset of the unreduced maximal-site generators agrees with the archived EBR table, covering 10,398 columns. Equal-dimensional onsite irreps are not distinguished by this dimension check; canonical identification of the complete onsite EBR columns requires additional character and convention information.

A separate character-level test compares all 2,700 scalar and spinful tables at the tabulated reciprocal points of the 230 space groups in `irreptables` 3.1.0 [21, 22]. All 8,907 irreps match bijectively, with largest residual 1.21×10^{-5} against the finite-precision reference data. The reference adapter explicitly uses the opposite tabulated Bloch wave-vector convention and the actual native spin lifts; it does not selectively conjugate characters or fit a phase independently for each irrep. The native algebra is checked at 10^{-9} , independently of the 2×10^{-4} tolerance for the rounded external tables. Optimized serial, both processes of a two-rank MPI run, and a focused instrumented executable give identical 11,607 normalized match and label records. This establishes named reference correspondences, not automatic standard labels in the Gaussian analysis or complete EBR identities and connectivity.

A Gaussian post-SCF GTH-SOC test independently checks that the spin lifts used for the factor algebra and for the actual band sewing agree. For a Ne Kramers doublet in a primitive body-centered cubic cell, independent quaternion reconstruction gives a product-law residual of 2 despite passing the metric, subspace and energy tests. Reusing the stored lifts reduces this residual to 8.5×10^{-14} without altering the input or validation tolerances; the SCF energies agree within 3×10^{-14} hartree. The property point and the SCF point are both Gamma; the primitive-basis $2 \times 2 \times 2$ mesh is not a full-cubic-symmetry reference. Details and the separate

quadrature convergence checks are given in the Supporting Information.

11.4 Convergence and material validation

Before submission, the material set should include matched crystalline band and real-space indices, finite-size SOC flakes, disordered configurations, and controlled transitions. Transport studies need independent convergence of basis, mesh or size, temperature, and dissipation. Localizer studies need a stable scale interval and a spatial map of both the index and its protection gap. The paper should report failed or unresolved parameter regions rather than select only points with the desired integer.

12 Conclusions

A common nonorthogonal operator representation connects several otherwise distinct ways of analysing electronic states in CP2K. Metric-aware current operators support finite-temperature transport; directed Gaussian overlaps support Wilson and parameter-space invariants; and Hamiltonian-position tuples support local indices and joint approximate states. Sparse assembly and factorization address their numerical cost without replacing the physical assumptions of each construction.

The present evidence establishes operator and solver consistency for the tested models and small GPW/GAPW/SOC calculations, alongside the retained neon and stanene band-topology demonstrations and the Si/Al k-point tests of the companion study. The TopoHSE-DB and generative-predictive discovery studies illustrate the materials-science applications motivating this infrastructure; their published or submitted results are not new CP2K validation data. The present tests do not establish all-space-group TQC, a general Hall-response implementation, or size-converged first-principles localizer classifications. Those boundaries are part of the method specification.

The crystalline framework now resolves projective unitary irreps, antiunitary pairing and doubling, and their restrictions along explicit reciprocal-space segments. Its algebraic coverage across conventional cell settings and supplied magnetic groups is distinct from a magnetic electronic-structure implementation or a complete EBR catalogue. The next material studies must converge the electronic structure, geometry or torus size, analysis sampling, and numerical scales independently, rather than interpreting a stable integer from a single small matrix as a bulk-material result.

Code and Data Availability

The manuscript sources, Supporting Information, and retained validation records are maintained in the development repository <https://github.com/DCM-Uni-Paderborn/TopologicalCP2K>. The repository currently has restricted access during manuscript preparation; public archival availability will be established before publication. CP2K is distributed under the GNU General Public License, version 2 or later, at <https://github.com/cp2k/cp2k>. The working implementation and retained calculation records are described in the Supporting Information. Not all developments described here are present in an upstream release. A final reproducible source snapshot, external-library versions, and public data archive must accompany submission. The present draft does not assign an archival DOI or imply that the development branch has already been released.

Authorship and Declarations

The author list and corresponding author follow the agreed manuscript scope. Affiliations, CRediT roles, funding, competing-interest declarations, and any required disclosure concerning writing

tools are to be confirmed by all authors before submission.

References

- [1] T. D. Kühne, M. Iannuzzi, M. Del Ben, V. V. Rybkin, P. Seewald, F. Stein, T. Laino, R. Z. Khaliullin, O. Schütt, F. Schiffmann, D. Golze, J. Wilhelm, S. Chulkov, M. H. Bani-Hashemian, V. Weber, U. Borstnik, M. TAILLEFUMIER, A. S. Jakobovits, A. Lazzaro, H. Pabst, T. Müller, R. Schade, M. Guidon, S. Andermatt, N. Holmberg, G. K. Schenter, A. Hehn, A. Bussy, F. Belleflamme, G. Tabacchi, A. Glöß, M. Lass, I. Bethune, C. J. Mundy, C. Plessl, M. Watkins, J. VandeVondele, M. Krack and J. Hutter, CP2K: An electronic structure and molecular dynamics software package – Quickstep: Efficient and accurate electronic structure calculations, *J. Chem. Phys.* **152**, 194103 (2020), doi:10.1063/5.0007045.
- [2] G. Lippert, J. Hutter and M. Parrinello, A hybrid Gaussian and plane wave density functional scheme, *Mol. Phys.* **92**, 477–487 (1997), doi:10.1080/002689797170220.
- [3] G. Lippert, J. Hutter and M. Parrinello, The Gaussian and augmented-plane-wave density functional method for *ab initio* molecular dynamics simulations, *Theor. Chem. Acc.* **103**, 124–140 (1999), doi:10.1007/s002140050523.
- [4] J. VandeVondele, M. Krack, F. Mohamed, M. Parrinello, T. Chassaing and J. Hutter, QUICKSTEP: Fast and accurate density functional calculations using a mixed Gaussian and plane waves approach, *Comput. Phys. Commun.* **167**, 103–128 (2005), doi:10.1016/j.cpc.2004.12.014.
- [5] T. D. Kühne, J. Heske, and E. Prodan, Disordered crystals from first principles. II: Transport coefficients, *Ann. Phys.* **421**, 168290 (2020), doi:10.1016/j.aop.2020.168290.
- [6] J. Bellissard, A. van Elst, and H. Schulz-Baldes, The noncommutative geometry of the quantum Hall effect, *J. Math. Phys.* **35**, 5373–5451 (1994), doi:10.1063/1.530758.
- [7] E. Prodan and H. Schulz-Baldes, *Bulk and Boundary Invariants for Complex Topological Insulators: From K-Theory to Physics* (Springer, Cham, 2016), doi:10.1007/978-3-319-29351-6.
- [8] G. Pizzi et al., Wannier90 as a community code: new features and applications, *J. Phys.: Condens. Matter* **32**, 165902 (2020). doi:10.1088/1361-648X/ab51ff.
- [9] K. Y. Dixon, T. A. Loring, and A. Cerjan, Classifying topology in photonic heterostructures with gapless environments, *Phys. Rev. Lett.* **131**, 213801 (2023), doi:10.1103/PhysRevLett.131.213801.
- [10] J. J. Esteve-Paredes and J. J. Palacios, A comprehensive study of the velocity, momentum and position matrix elements for Bloch states using a local orbital basis, *SciPost Phys. Core* **6**, 002 (2023), doi:10.21468/SciPostPhysCore.6.1.002.
- [11] A. A. Soluyanov and D. Vanderbilt, Computing topological invariants without inversion symmetry, *Phys. Rev. B* **83**, 235401 (2011). doi:10.1103/PhysRevB.83.235401.
- [12] D. Gresch, G. Autès, O. V. Yazyev, M. Troyer, D. Vanderbilt, B. A. Bernevig, and A. A. Soluyanov, Z2Pack: Numerical implementation of hybrid Wannier centers for identifying topological materials, *Phys. Rev. B* **95**, 075146 (2017). doi:10.1103/PhysRevB.95.075146.
- [13] Y. Xu, B. Yan, H.-J. Zhang, J. Wang, G. Xu, P. Tang, W. Duan, and S.-C. Zhang, Large-gap quantum spin Hall insulators in tin films, *Phys. Rev. Lett.* **111**, 136804 (2013). doi:10.1103/PhysRevLett.111.136804.
- [14] L. Fu and C. L. Kane, Topological insulators with inversion symmetry, *Phys. Rev. B* **76**, 045302 (2007). doi:10.1103/PhysRevB.76.045302.

- [15] B. Bradlyn et al., Topological quantum chemistry, *Nature* **547**, 298–305 (2017). doi:10.1038/nature23268.
- [16] H. C. Po, A. Vishwanath, and H. Watanabe, Symmetry-based indicators of band topology in the 230 space groups, *Nat. Commun.* **8**, 50 (2017). doi:10.1038/s41467-017-00133-2.
- [17] SPGLIB developers, Definitions and conventions, SPGLIB documentation, <https://spglib.readthedocs.io/en/stable/definition.html> (accessed 24 September 2026).
- [18] Bilbao Crystallographic Server, Brillouin-zone databases: definitions of reciprocal-space groups and wave-vector types, https://cryst.ehu.es/cryst/help/definitions_kvec.html (accessed 24 September 2026).
- [19] K. Shinohara, Space-group irreducible representations and spin representations, SPGREP formulation documentation, https://spglib.github.io/spgrep/formulation/irreps/spacegroup_irreps.html (accessed 24 September 2026).
- [20] K. Shinohara, Co-representation of magnetic point and space group, SPGREP formulation documentation, <https://spglib.github.io/spgrep/formulation/irreps/corep.html> (accessed 24 September 2026).
- [21] M. Iraola, J. L. Mañes, B. Bradlyn, T. Neupert, M. G. Vergniory, and S. S. Tsirkin, IrRep: Symmetry eigenvalues and irreducible representations of ab initio band structures, *Comput. Phys. Commun.* **272**, 108226 (2022). doi:10.1016/j.cpc.2021.108226.
- [22] IrRep developers, irreptables 3.1.0: tables of characters for single and double space groups, <https://pypi.org/project/irreptables/3.1.0/> (accessed 24 September 2026).
- [23] T. Fukui, Y. Hatsugai, and H. Suzuki, Chern numbers in discretized Brillouin zone: Efficient method of computing (spin) Hall conductances, *J. Phys. Soc. Jpn.* **74**, 1674–1677 (2005). doi:10.1143/JPSJ.74.1674.
- [24] M. Mochol-Grzelak, A. Dauphin, A. Celi, and M. Lewenstein, Efficient algorithm to compute the second Chern number in four dimensional systems, *Quantum Sci. Technol.* **4**, 014009 (2019). doi:10.1088/2058-9565/aae93b.
- [25] T. Loring and H. Schulz-Baldes, The spectral localizer for even index pairings, *J. Noncommut. Geom.* **14**, 1–23 (2020); arXiv:1802.04517.
- [26] LAPACK developers, ZHETRF and DSYTRF: pivoted Hermitian and symmetric factorizations, <https://www.netlib.org/lapack/complex16/zhetrf.f>, <https://www.netlib.org/lapack/double/dsytrf.f> (accessed 24 September 2026).
- [27] N. Doll and H. Schulz-Baldes, Skew localizer and $\mathbb{Z}_2$ -flows for real index pairings, *Adv. Math.* **392**, 108038 (2021), doi:10.1016/j.aim.2021.108038.
- [28] S. Wong, I. Yamazaki, C. Siefert, I. Duff, T. A. Loring, and A. Cerjan, Efficient local classification of parity-based material topology, *Phys. Rev. B* **114**, 084201 (2026), doi:10.1103/6hj9-tgct.
- [29] N. Doll, T. Loring, and H. Schulz-Baldes, Topological indices for periodic gapped Hamiltonians and fuzzy tori, *Math. Phys. Anal. Geom.* **28**, 13 (2025), doi:10.1007/s11040-025-09508-0.
- [30] A. Cerjan, T. A. Loring, and F. Vides, Quadratic pseudospectrum for identifying localized states, *J. Math. Phys.* **64**, 023501 (2023), doi:10.1063/5.0098336.
- [31] H. Mirhosseini, L. Elcoro, A. Knüpfer, and T. D. Kühne, TopoHSE-DB: a topological materials database for generative and predictive models, *Mach. Learn.: Sci. Technol.* **7**, 040601 (2026), doi:10.1088/2632-2153/ae97d8.

- [32] L. Brocai, T. D. Kühne, A. Knüpfer, and H. Mirhosseini, Generative–predictive models enable the discovery of topological materials, submitted manuscript (2026).
- [33] J. V. Pototschnig, J. Kloppenburg, S. He, C. Zhou, S. Wang, H. Mirhosseini, S. Azadi, J. Hutter, and T. D. Kühne, Symmetry-reduced k-point sampling and Wannier analysis with localized Gaussian orbitals, companion manuscript in preparation (2026).
- [34] U. Borstnik, J. VandeVondele, V. Weber and J. Hutter, Sparse matrix multiplication: The distributed block-compressed sparse row library, *Parallel Comput.* **40**, 47–58 (2014), doi:10.1016/j.parco.2014.03.012.
- [35] MUMPS developers, MUMPS user documentation, <https://www.mumps-solver.org/index.php?page=doc> (accessed 24 September 2026).
- [36] J. Cano and B. Bradlyn, Band Representations and Topological Quantum Chemistry, *Annu. Rev. Condens. Matter Phys.* **12**, 225–246 (2021), doi:10.1146/annurev-conmatphys-041720-124134.
- [37] J. Cano, B. Bradlyn, Z. Wang, L. Elcoro, M. G. Vergniory, C. Felser, M. I. Aroyo and B. A. Bernevig, Building blocks of topological quantum chemistry: Elementary band representations, *Phys. Rev. B* **97**, 035139 (2018), doi:10.1103/PhysRevB.97.035139.
- [38] T. Christensen and contributors, Crystalline.jl: three-dimensional elementary-band-representation data archived from the Bilbao Crystallographic Server, [versioned reference dataset](#) (accessed 24 September 2026).
- [39] I. Yamazaki and collaborators, Tacho skew-LDL extension, https://github.com/iyamazaki/Trilinos/tree/tacho-sk/packages/shylu/shylu_node/tacho (accessed 24 September 2026).