

Supporting Information

Topological Systems with CP2K

Thomas D. Kühne, Alexander Cerjan, Vladislav Efremkin,
Hermann Schulz-Baldes, and Emil Prodan

Corresponding author: tkuehne@cp2k.org

S1 Scope and Reproduction Boundaries

This working document combines the retained band-topology calculations transferred from the k-point manuscript with separately validated transport, localizer, and quadratic-pseudospectrum developments. The original results are not recomputed or relabelled by moving the manuscript text. The development status is recorded as of 24 September 2026. A source freeze and complete public reproduction archive remain submission requirements.

Applications versus test calculations

TopoHSE-DB and the submitted generative-predictive discovery study are cited in the main text as materials-science applications. Their VASP-based calculations are not part of the CP2K validation records collected here. In particular, their database labels and candidate counts do not validate the new CP2K localizer or transport implementations.

The smaller calculations below are tests and demonstrations. The companion k-point paper additionally reports 18 converged Si/Al calculations in conventional cubic cells with eight Si atoms ($a = 5.43$ Å) and four Al atoms ($a = 4.05$ Å). These use PBE, DZVP-MOLOPT-SR-GTH bases, GTH-PBE pseudopotentials, 600/60 Ry grid cutoffs, and an SCF threshold of 10^{-9} . Al uses Fermi-Dirac occupations at 1000 K, so its compared quantity is the electronic free energy. The same-mesh comparison uses $4 \times 4 \times 4$ sampling: 64 full-grid points, 32 with time reversal, and four with either K290 or SPGLIB atomic symmetry. The maximum cell-energy/free-energy difference is 1.4×10^{-12} hartree; the Al atomic-symmetry stress difference is 0.098 bar. These tests demonstrate representation consistency, not independently converged material properties or general force accuracy. Their source and raw records remain those of the companion manuscript; no new Si/Al calculations were performed for the present discussion.

S2 Native Wilson-Loop Test and Demonstration Calculations

The topology examples form a separate calculation set from the Si/Al mainline tests. The native extension supplies directed Gaussian-basis overlaps, second-variational spinors, and the Wilson-loop analysis described in the main text. The retained calculations were completed on 17 September 2026 with one MPI process and one OpenMP thread. Source provenance and artifact checksums are recorded in `validation/topology/README.md` and the figure manifest.

Both examples use PBE, the DZVP-MOLOPT-GGA-GTH basis from `BASIS_MOLOPT_UZH`, SOC pseudopotentials from `GTH_SOC_POTENTIALS`, a 200 Ry density cutoff and a 40 Ry relative cutoff. The potential labels are `GTH-PBE-q8` for neon and `GTH-PBE-q4` for tin. Complex orbitals and explicit full SCF meshes are used. The neon reference contains one atom at fractional coordinate $(1/2, 1/2, 1/2)$ in a cubic cell of side 5 Å, with a $2 \times 2 \times 2$ SCF mesh and convergence threshold 10^{-9} . The stanene cell has lattice vectors $(4.674, 0, 0)$, $(2.337, 4.047802737288466, 0)$, and $(0, 0, 20)$ Å, and two tin atoms at fractional positions $(1/3, 1/3, 0.5213)$ and $(2/3, 2/3, 0.4787)$. It therefore

has a buckling height of 0.852 Å. These fixed geometric parameters follow the pristine-stanene reference reported by Zhang et al.¹ An $8 \times 8 \times 1$ SCF mesh and convergence threshold 10^{-8} are used for stanene. The examples test the topology implementation at these fixed settings, rather than constitute a new structural or band-gap prediction.

Eight occupied spinors are retained in each case. The stanene second-variational calculation includes all 26 scalar basis states, giving 52 spinor states before occupied-band selection. For neon, five scalar states produce ten spinors. The selected half-plane is $\mathbf{k}(s, t) = (t, s/2, 0)$ in fractional reciprocal coordinates, with $0 \leq s, t \leq 1$. The loop closes with reciprocal shift $(1, 0, 0)$. Starting meshes contain four points on each of three loops for neon and 12 points on each of 13 loops for stanene. Refinement doubles the points per loop and inserts a midpoint between neighboring transverse loops. The accepted surfaces have 8×5 and 192×193 points, respectively. The last center changes are 1.28×10^{-5} and 2.73×10^{-4} in reduced units. The smallest link singular values are 0.99826 and 0.95248.

The native invariants are zero for neon and one for stanene. Z2Pack 2.2.1 gives the same stanene parity using either the raw overlaps or their unitary polar factors. For the latter, the maximum Wilson-eigenvalue difference from the native calculation is 7.95×10^{-14} . This comparison tests the loop algebra with identical input matrices. Separate operator checks enforce zero-step normalization, reverse-link adjoint relations, loop reversal, and atomic-position phases. A bonded two-atom full-spinor check gives maximum adjoint and zero-step residuals of 2.79×10^{-14} and 2.00×10^{-13} , respectively.

An additional adaptive Z2Pack-driven stanene surface uses 42 CP2K line calculations and ends with 15 accepted transverse lines, all satisfying the line-position, movement, and gap checks, and again gives $\nu = 1$. This calculation chooses its sampling independently but uses the same CP2K Hamiltonian and overlap implementation. The main-text figure shows the native samples without interpolation or artificial separation of degenerate centers. It is regenerated by `python3 validation/topology/plot_wilson.py`, which requires NumPy and Matplotlib. Raw `.wilson` files, inputs, logs, comparison reports, generated comma-separated figure data, and checksums accompany the figure.

S3 Native Chern and Cross-Geometry Extension

S3.1 Source versions and independent analysis routes

The Chern and cross-geometry checks were completed on 18 September 2026 with a separate development executable. They are distinct from the retained 17 September neon/stanene calculations and do not reassign those earlier results to a newer implementation. Source patches, executable records, library versions, and independent comparison scripts accompany the validation archive. The numerical comparison tools require NumPy and SciPy; Z2Pack 2.2.1 is an optional independent analysis dependency.

Table S1: Topology analysis routes and runtime requirements. All native routes act on Gaussian-basis electronic states without constructing maximally localized Wannier orbitals.

Route	Quantity and sampling	Dependencies and scope
Native Wilson/ $\mathbb{Z}_2$	Wilson spectra, Berry phases, and time-reversal half-plane $\mathbb{Z}_2$.	CP2K native kernels, including second-variational SOC. No Z2Pack or Wannier90 runtime dependency.
Native Wilson/ C_1	Determinant-phase winding on a full closed reciprocal surface, selected by <code>CHERN</code> .	CP2K native kernels. Fixed-rank isolated subspace, closure and refinement checks.

¹R.-w. Zhang et al., Room temperature quantum spin Hall insulator in ethynyl-derivative functionalized stanene films, *Sci. Rep.* **6**, 18879 (2016), <https://doi.org/10.1038/srep18879>.

Route	Quantity and sampling	Dependencies and scope
Native inversion analysis	Even/odd irreps, Fu–Kane indices and inversion-subgroup EBR signatures at TRIM.	Separate native extension, Section S4. No general space-group catalogue.
Native parameter-mesh analysis	C_1 on two parameters and the second Chern character pairing C_2 on four parameters, using physical cross-geometry overlaps.	<code>topology_phasons</code> reads <code>STATE_EXPORT</code> files. One process with optional OpenMP. No Python, Z2Pack, or Wannier90 runtime dependency.
External Z2Pack route	Independent loop/surface analysis of CP2K overlaps, with optional adaptive CP2K sampling.	Python and Z2Pack. Optional analysis and cross-validation, not the implementation of the native invariants.

S3.2 Parameter-mesh discretization and physical closure

Let $Q_\mu(x)$ be the polar-unitary selected-state overlap directed from mesh vertex x to $x + \hat{\mu}$. All boundary links include the specified physical sewing. The oriented plaquette is

$$P_{\mu\nu}(x) = Q_\mu(x)Q_\nu(x + \hat{\mu})Q_\mu^\dagger(x + \hat{\nu})Q_\nu^\dagger(x). \quad (\text{S1})$$

With this link orientation, the two-parameter estimator is

$$C_1^{\text{mesh}} = -\frac{1}{2\pi} \sum_x \text{Arg det } P_{12}(x). \quad (\text{S2})$$

Its sign matches the determinant Wilson-phase winding convention of the main text. For four parameters, define the full principal matrix logarithm $F_{\mu\nu}(x) = \log P_{\mu\nu}(x)$. Each $F_{\mu\nu}$ is anti-Hermitian and already includes the plaquette area. The estimator is

$$C_2^{\text{mesh}} = -\frac{1}{4\pi^2} \sum_x \text{Re Tr } [F_{12}F_{34} - F_{13}F_{24} + F_{14}F_{23}](x). \quad (\text{S3})$$

The full matrices are needed for a non-Abelian occupied subspace. Equation (S3) approximates the second Chern character pairing and is not generally the second Chern class. Its finite-mesh value is reported without integer rounding.

Physical links use $C_a^\dagger O_{ab} C_b$, with both spinor components summed when present. The cross-geometry AO matrix O_{ab} retains the moving contracted Gaussian centers, periodic images, and reciprocal boundary phases. The cell, atom count and selected rank remain fixed throughout the family. Endpoint checks use explicit atom permutations, lattice translations, basis matching, and selected-subspace sewing. The code reports AO-metric errors, sampled selected/excluded-band gaps, minimum link singular values, and principal plaquette phases. A refined mesh must preserve the same physical family. Sampled isolation and invariant convergence do not by themselves establish a bulk spectral gap between samples. An indirect gap is interpreted only for a lowest-band prefix with a justified common energy reference across geometries.

S3.3 Retained validation results

The extension was rebuilt with GNU Fortran 16 and OpenBLAS. Its curvature and snapshot unit tests and the existing Wilson/ $C_1/\mathbb{Z}_2$ unit test passed with two OpenMP threads. All four inputs in `QS/regtest-topology` passed with two MPI ranks and two OpenMP threads per rank. The auxiliary parameter-mesh executable itself was tested as one process, not as an MPI-distributed postprocessor.

The native tests cover singular or underresolved links, metric and seam failures, determinant phases, non-Abelian gauge covariance, and model refinement. For the occupied doublet of the

four-dimensional Dirac model at mass -3 , the 8^4 estimate is -0.7424128463 and is closer to the continuum value -1 than the 6^4 result. This is a finite-resolution regression check, not a claim of a converged integer at 8^4 . Independent Python/native comparisons additionally check gauge changes and reversal of parameter orientation.

For archived CP2K state exports covering skew cells, multiple centers, SOC spinors and reciprocal-boundary links, the largest difference between native and independently evaluated Gaussian overlaps is 1.03×10^{-15} with either one or two OpenMP threads. A cross-geometry overlap agrees to 5.56×10^{-16} . Table S2 reports physical atomic-translation pumps. The two-parameter case varies one reciprocal coordinate and one periodic atomic displacement. The separable four-parameter case uses the ordered coordinates $(k_x, k_y, \phi_x, \phi_y)$.

Table S2: Native physical Gaussian-snapshot tests. The minimum singular value is measured before polar unitarization. The phase column reports the largest diagnostic plaquette phase in radians. The 4^4 separable case tests the four-parameter physical workflow, not generic C_2 convergence.

Mesh	Unrounded pairing	Minimum singular value	Maximum phase
4×4	$C_1 = -1.0000000000$	0.234278	0.402344
8×8	$C_1 = -1.0000000000$	0.609933	0.099083
12×12	$C_1 = -1.0000000000$	0.780474	0.044040
4^4	$C_2 = -1.0000000000$	0.233877	0.402347

The native 4×4 to 8×8 physical refinement check passed at an invariant tolerance of 10^{-8} . The independent comparison-tool suite passed 34 tests without Z2Pack, with five optional-adaptor tests skipped, and all 39 tests with Z2Pack installed. The accompanying `validation/topology/native-phasons/` directory contains the machine-readable overlap and pump results, test logs, and source-version record. These are implementation tests and introduce no new material prediction or change to the neon/stanene results.

S4 Native Inversion Representations and Symmetry Signatures

The inversion extension was validated separately from the phason extension and from the earlier Si/Al and Wilson calculations. Its source patch, base snapshot, inputs and outputs remain in `validation/topology/inversion-tqc/`. These records identify the original tests without retrospectively assigning them to another executable.

The WANNIER90 print section accepts KPOINTS_SOURCE TRIM, SOC T, TIME_REVERSAL T and INVERSION_TQC T. TQC_DIMENSION selects the four $k_z = 0$ TRIM or all eight three-dimensional TRIM. PARITY_ORIGIN specifies the inversion center in fractional direct coordinates. The occupied spinor bands must be a complete lowest-band prefix, with excluded states available for checking spectral separation. A PARITY-only mode also accepts isolated scalar subspaces. Neither route requires Z2Pack, Wannier90 execution or a Wannier fit. The TRIM calculation holds the converged SCF potential fixed and does not require the SCF integration mesh to contain those points. The two-dimensional option samples the fractional $k_z = 0$ plane of a three-dimensionally periodic cell, not a separate two-dimensional Poisson boundary condition. The inversion center is specified explicitly rather than inferred from a space-group search.

For an orbital of angular momentum l on atom a , inversion about $\mathbf{c}$ maps it to an identical Gaussian orbital on atom b when $2\mathbf{c} - \mathbf{r}_a = \mathbf{r}_b + \mathbf{L}$ with integer $\mathbf{L}$. The coefficient multiplier is $(-1)^l \exp(2\pi i \mathbf{k} \cdot \mathbf{L})$. The code verifies atom-kind matching, involution, metric covariance, $C^\dagger S C = I$, unitarity and Hermiticity of the projected inversion matrix, and its commutator with the selected eigenvalues. Spinful time reversal is checked through the sewing matrix of $\Theta = i\sigma_y K$. Parity multiplicities must be even before division into Kramers pairs. The dimensionless and geometry tolerances are 10^{-6} and 10^{-6} bohr, respectively. The energy-commutator tolerance is

10^{-6} hartree.

For N occupied Kramers pairs and odd counts n_{κ}^{-} at $\mathbf{k} = \kappa/2$, $\kappa \in \{0, 1\}^d$, the parity index is $\nu = \sum_{\kappa} n_{\kappa}^{-} \bmod 2$. In three dimensions the weak indices restrict this sum to $\kappa_i = 1$, and the convention for the inversion indicator is $z_4 = \sum_{\kappa} n_{\kappa}^{-} \bmod 4$. The analytic inversion-subgroup EBRs are local even or odd Kramers pairs at centers $\mathbf{c} + \mathbf{a}/2$. Define

$$d_{\mathbf{a}} = 2^{-d} \sum_{\kappa} (-1)^{\mathbf{a} \cdot \kappa} (N - 2n_{\kappa}^{-}). \quad (\text{S4})$$

An integer atomic-signature decomposition exists when all $d_{\mathbf{a}}$ are integers and $N - \sum_{\mathbf{a}} d_{\mathbf{a}}$ is even. A nonnegative decomposition additionally requires $N \geq \sum_{\mathbf{a}} |d_{\mathbf{a}}|$. The implementation makes these decisions with integer arithmetic. Atomic compatibility concerns the symmetry signature, not a proof of trivial topology. Likewise, a signed-only decomposition does not by itself establish fragile topology. This inversion-only EBR test is distinct from the subsequent native little-group and explicit-segment analysis described below; it does not become a general EBR classifier when that additional analysis is used.

Table S3: Additional native parity checks with four occupied Kramers pairs. The four columns refer to $(k_x, k_y) = (0, 0), (1/2, 0), (0, 1/2), (1/2, 1/2)$ at $k_z = 0$. Coordinates of the chosen inversion center are fractional.

System	$\mathbf{c}$	n_{κ}^{-}				ν	Atomic signature
Neon	$(1/2, 1/2, 1/2)$	3	3	3	3	0	Compatible
Stanene	$(0, 0, 0)$	1	2	2	2	1	Obstructed

These tests retain the basis, cutoff, geometry and SCF meshes of the earlier material examples, replacing the Wilson paths by TRIM sampling. The three-dimensional neon test, with inversion center at the origin, also gives zero strong and weak indices and $z_4 = 0$. All seven inputs in the extended `QS/regtest-topology` set passed with two MPI ranks and two OpenMP threads per rank. The stanene parity result was independently reproduced with one rank and one thread. A single-rank, multiple-thread run exposed inconsistent state normalization in the existing local k-point diagonalization path for this build. The new metric checks reject that case. This extension does not resolve or validate that threaded path.

The lightweight CI fixture subsequently reduces the stanene SCF mesh from $8 \times 8 \times 1$ to $4 \times 4 \times 1$ and suppresses unused restart files. It still evaluates all four TRIM and reproduces the odd-pair counts $(1, 2, 2, 2)$ and $\nu = 1$. This fixture is distinct from the archived $8 \times 8 \times 1$ manuscript calculation and does not replace it or establish k-mesh convergence.

The native unit test covers multidimensional-character decomposition, nonorthogonal metrics, degenerate-state gauge changes, broken-symmetry rejection, all inversion-center EBRs and strong, $z_4 = 2$ and signed-only examples. Independent enumeration of all two-Kramers-pair atomic sums matches the solver for all 81 two-dimensional and 6,561 three-dimensional candidate parity signatures. A deliberately incorrect inversion center is rejected in an end-to-end calculation. TRIM sampling does not prove a gap over the full Brillouin zone, and the SOC subspace still requires its usual convergence checks.

S5 Native Little-Group and Compatibility Validation

The native algebra constructs unitary projective character tables, Wigner corepresentation restrictions, and endpoint-to-segment branching without a runtime dependency on Spgrep or a pretabulated irrep catalogue. The electronic-structure path uses SPGLIB to detect spatial operations and, when requested, adds their nonmagnetic time-reversal extension. It retains centering translations and tests the metric, band-subspace closure, energy commutators and the complete semilinear multiplication law before accepting band multiplicities.

The all-setting sweep covers all 530 Hall settings of the 230 ordinary space groups. At each setting, it samples $(0, 0, 0)$, $(1/2, 1/2, 1/2)$, $(1/2, 0, 0)$, $(1/3, 1/3, 0)$, $(0, 0, 1/4)$ and $(0.173, 0.217, 0.319)$ with scalar and spinful factors. The invariant cell metric is obtained by averaging over the supplied rotations. Both the ordinary group and its grey extension are checked. Generic-site orbits with physical spin multiplicity provide fixed-rank compatible band data. Segments run from Gamma to the other sampled points and from Gamma to its reciprocal image $(0, 0, 1)$. This tests directed-image monodromy as well as ordinary endpoint restrictions; it does not sample every possible line or material.

Checked object	Successful cases
Ordinary/grey scalar/spinful factor systems	12,720
Complete unitary projective character tables	6,360
Endpoint tables with an antiunitary stabilizer	5,324
Ordinary/grey directed segment comparisons	12,720
Segments retaining an antiunitary stabilizer	4,462
Gaussian directory assertions, serial / two MPI ranks	47 / 47

Table S4: Native representation validation on 24 September 2026. The counts refer to algebraic cases or numerical assertions, not distinct material calculations.

Analytic cases separately test magnetic half translations, spin quaternion doublets, complex conjugate C_3 pairs, twisted changes of factor gauge, and invalid pairing/doubling counts. Segment cases include displaced-origin PT pairs, endpoint-only time reversal, magnetic-glide Kramers planes, and spinful glide partner switching: a two-band hourglass connection is obstructed, whereas the complete four-band space is compatible. Unitary screw and mirror branching, reciprocal-image monodromy and changed coset representatives are also checked. The complete Hall-setting suite passes in release and with bounds checking and floating-point traps enabled.

The Gaussian directory uses UZH MOLOPT bases for its GPW fixtures and a contracted all-electron basis for GAPW. Ne tests exercise scalar and post-SCF GTH-SOC multiplets at Gamma, a generic PT-symmetric point and nonsymmorphic screw sectors. Scalar He/GPW, He/GAPW and Ne/SOC close a four-segment reciprocal loop; a separately selected single screw branch must report its monodromy obstruction. The generated Wilson-link test also exercises repeated refinement. Serial runs use two OpenMP threads; MPI runs use two ranks with two threads each and the thread-safe BLAS settings stated below. An earlier wider corepresentation integration pass additionally gave 110/110 serial and 111/111 MPI assertions across topology, Kubo, localizer and quadratic directories. These are separate test scopes, not numbers to add into a material benchmark total.

A separate local algebra sweep uses SPGLIB 2.7 magnetic data in the supplied setting for every one of the 1,651 magnetic UNI numbers. At the same six points and two spin choices, all 19,812 factor cases pass; 14,088 contain an antiunitary stabilizer, and 5,724 do not. For 10,154 primitive-cell tables supported by an independent Spgrep 0.7.0 comparison, dimensions, Wigner types and unitary restriction characters agree within 7.11×10^{-15} . Both calculations use the same supplied factor system, so this is not an independent validation of symmetry detection. Moreover, a separate semilinear-product check did not validate every explicit antiunitary matrix produced by that reference calculation. Those matrices are not imported into CP2K; the comparison is limited to types, dimensions and unitary restrictions. CP2K’s actual Gaussian sewing matrices pass their own full product check.

The separate native site-induction test samples fractional positions $(0, 0, 0)$, $(1/2, 0, 1/4)$ and $(0.173, 0.217, 0.319)$ at the same six reciprocal points. Ordinary and grey groups, each with scalar and spinful factors, give 38,160 site/k-point cases over all 530 Hall settings. There are 67,824 local-representation columns evaluated and decomposed, and the same number of directed compatibility comparisons. Their atomic band rank equals the orbit multiplicity times the local irrep or corepresentation dimension. This construction preserves conventional-cell translation

sectors rather than silently converting them to a primitive-cell catalogue.

An independent check constructs Cartesian s/p/d orbital traces and, for spinors, multiplies them by the physical $SU(2)$ trace. After matching the onsite orbital content, its explicitly translated orbit trace agrees with the corresponding induced character sum in 114,480 comparisons. The numerical acceptance tolerance is 10^{-7} for these summed traces and decomposition checks; group and orbit geometry use 10^{-9} . Analytic checks additionally cover inversion-center parities, screw orbits, spin-quaternion doublets, and complex local orbitals related by an antiunitary half translation. These are algorithmic reference tests, not new material calculations or a new production input mode. Additional scalar and spinful checks preserve the characters under passive origin shifts and unimodular changes to a sheared cell, and verify the expected Bloch rephasing under integer changes of operation representatives. The full database sweeps pass in serial, two-rank MPI, and a separate bounds-checked build with floating-point traps.

These algebra sweeps do not implement magnetic-space-group detection, a magnetic non-collinear ground state, canonical irrep labels, a complete Brillouin-zone connectivity graph or a full elementary-band catalogue. Explicit-segment compatibility is necessary, not sufficient, for a gapped band subspace. The retained summaries and test logs are in `validation/development/`; no production material classification is inferred solely from a successful database sweep.

S6 Bloch Kubo and Property-Symmetry Validation

The retained development report is `validation/development/bloch-response-summary.md`. It records distinct finite-cutoff regression fixtures and the higher-cutoff primitive/supercell comparison. For Ne with post-SCF GTH SOC and PBE at 2000 Ry, the $3 \times 1 \times 2$ primitive mesh gives -34.923548219306689 hartree, compared with -209.541289276241002 hartree for six cells sampled explicitly at Gamma. Both print $6.21105320 \times 10^{-9}$ S. At the inexpensive 200 Ry PADE regression settings, the respective conductivities are $6.42389197 \times 10^{-9}$ and $6.35401974 \times 10^{-9}$ S. Those distinct references are retained: the mismatch is not removed by changing a reference value.

Ne/SOC on a 2^3 property mesh uses four representatives. Bi_2 /SOC on 3^3 uses six; its printed nine-component tensor differs from full-grid by at most 2.71×10^{-15} relative to its largest entry. Primitive diamond/SOC on 2^3 uses three representatives with either K290 or SPGLIB. Its tensor difference is 2.21×10^{-11} in the same normalization. Metric, Hamiltonian and current reconstruction residuals are approximately 2.1×10^{-13} , 6×10^{-11} and 1.3×10^{-11} , respectively. These are operator-covariance fixtures, not converged material transport predictions.

The final recorded directory drivers passed 132 serial and 162 MPI assertions. The MPI layout was four ranks with two OpenMP threads, and the serial build also used two OpenMP threads. Coverage includes moments, k-point/Wannier export, topology, Kubo, localizer, and quadratic directories; the MPI build additionally tests MUMPS and Tacho. Mathematical unit tests cover the constituent operator identities separately.

The Apple ARM runtime required one BLAS thread, a consistent thread-safe OpenBLAS preload, and an OpenMP stack of 128 MiB. Without these settings, failures occurred before the property code. An implicit-Gamma high-cutoff conversion also exposed a slow Miller-index lookup; the completed comparison uses an explicit Gamma input. These runtime qualifications must accompany the results, rather than be presented as production scalability.

S7 Localizer and Quadratic Validation Records

The transferred developer notes in `validation/development/` document finite and periodic formulations, dense and sparse factorization, and their limitations. They are implementation records, not independent material benchmarks. The matrix tests include nontrivial and trivial Chern models, spin-mixed Kramers-pair models, origin and energy shifts, nonorthogonal congruences, invalid metrics, and gap closings.

Dense LDL tests cover both 1×1 and 2×2 blocks. Pfaffian tests independently cover odd permutations and recursive small-matrix reference polynomials. Sparse gap brackets are compared with generalized dense spectra, not with pivot sizes. Realification duplicates the spectrum and is accounted for in every inertia comparison.

For a larger molecular quadratic check, Bi_2 uses TZVP-MOLOPT-GGA-GTH-q5, GTH SOC, 68 spinor coefficients, four requested states and a 32-dimensional trial space. GPW and GAPW dense/iterative comparisons on four MPI ranks agree in gaps within 6×10^{-15} hartree, with all four metric-subspace singular values within 10^{-10} of unity. This is separate from the smaller mandatory regression fixtures.

S8 Submission Validation Still Required

The current record does not establish a nontrivial, size-converged DFT localizer classification for a large SOC flake, multi-node sparse scaling, or general 230-space-group TQC. The final archive should contain complete material inputs and potentials, basis and cutoff studies, spatial/energy/scale scans, independent band references, and timed runs with whole-process memory. Development diagnostics and planned work must not be merged into tables of completed material results.