

# Electronic Supplementary Information

Native implementation of the machine-learned Skala exchange–correlation functional in CP2K:  
Unified one-centre reconstruction for molecular and condensed-phase calculations

Johann Pototschnig, Franz Pöschel,  
Jürg Hutter, and Thomas D. Kühne

## Contents

|                                                                             |            |
|-----------------------------------------------------------------------------|------------|
| <b>S1 Scope and Notation</b>                                                | <b>S2</b>  |
| <b>S2 Reconstruction and Derivative Consistency</b>                         | <b>S2</b>  |
| S2.1 Spin-gradient cross terms . . . . .                                    | S2         |
| S2.2 The discrete adjoint and moving grids . . . . .                        | S2         |
| S2.3 Finite-difference protocol . . . . .                                   | S3         |
| <b>S3 Molecular Numerical Settings and Convergence</b>                      | <b>S3</b>  |
| S3.1 Basis, grid, and boundary conditions . . . . .                         | S3         |
| S3.2 Regular-grid cutoff . . . . .                                          | S4         |
| S3.3 Radial and angular quadrature . . . . .                                | S4         |
| S3.4 Molecular cell padding . . . . .                                       | S4         |
| <b>S4 Molecular Reaction Data and Cross-Code Comparisons</b>                | <b>S5</b>  |
| S4.1 Direct comparison with the previous molecular implementation . . . . . | S6         |
| <b>S5 K-Point and Symmetry Checks</b>                                       | <b>S9</b>  |
| <b>S6 Molecular-Crystal Energies and Numerical Controls</b>                 | <b>S9</b>  |
| S6.1 Geometries and electronic protocol . . . . .                           | S9         |
| S6.2 Paired energies and controls . . . . .                                 | S10        |
| S6.3 Other density-functional approximations . . . . .                      | S10        |
| <b>S7 Ten-Solid Equations of State</b>                                      | <b>S11</b> |
| S7.1 Structures and common fit windows . . . . .                            | S11        |
| S7.2 Computational details and fit stability . . . . .                      | S11        |
| S7.3 Representation and material trends . . . . .                           | S12        |
| S7.4 Matched literature comparisons . . . . .                               | S12        |
| <b>S8 Pressure Derivatives and Compressibilities</b>                        | <b>S16</b> |
| <b>S9 Reproducibility and Numerical Limits</b>                              | <b>S17</b> |

## S1 Scope and Notation

This Electronic Supplementary Information (ESI) documents the numerical definition and validation of the molecular and condensed-phase comparisons. Self-consistent-field (SCF) convergence, basis resolution, and grid resolution are assessed separately. The SCF solvers are orbital transformation (OT) and diagonalization. Electronic occupations and the overlap-matrix trace define the explicitly treated particle number. Integration on the model quadrature provides an additional discretization diagnostic. A nonzero quadrature error is therefore distinct from an incorrect occupation count. The calculations do not renormalize the model density.

The Gaussian and plane-wave (GPW) and Gaussian and augmented-plane-wave (GAPW) methods provide the native density representations. Their exchange–correlation (XC) fields enter Skala. GAPW-XC denotes the XC-specific augmented density decomposition, and not merely the name of an input switch. All-electron (AE) GAPW is denoted GAPW-AE. Goedecker–Teter–Hutter (GTH) effective-core potentials (ECPs) define the pseudopotential Hamiltonian. GTH-direct and GTH-1C denote GAPW-GTH with direct valence and one-centre (1C) reconstructed fields, respectively. The mixed molecular and solid protocols combine AE and GTH treatments. They are not hybrid XC functionals. AE and GAPW-XC already contain their intrinsic one-centre terms. The independent with/without comparison applies to the two ordinary pseudopotential GAPW routes. Compact tables use GX for GAPW-XC/GTH, AE for GAPW-AE, and HD/HOC for the mixed AE/GTH protocols with direct/1C GTH fields, respectively.

Machine-readable reaction values, selected equation-of-state (EOS) curves, molecular-crystal total energies, and exact inputs are organized separately in the accompanying benchmark repository. The numerical tables below are generated from those datasets rather than transcribed from working logs.

## S2 Reconstruction and Derivative Consistency

### S2.1 Spin-gradient cross terms

Write the two reconstructed gradient fields as  $\mathbf{g}_\alpha = \tilde{\mathbf{g}}_\alpha + \mathbf{g}_\alpha^h - \mathbf{g}_\alpha^s$  and analogously for  $\beta$ . Their three spin invariants are  $\gamma_{\alpha\alpha} = \mathbf{g}_\alpha^2$ ,  $\gamma_{\beta\beta} = \mathbf{g}_\beta^2$ , and  $\gamma_{\alpha\beta} = \mathbf{g}_\alpha \cdot \mathbf{g}_\beta$ . For example,

$$\begin{aligned} \gamma_{\alpha\beta} = & \tilde{\mathbf{g}}_\alpha \cdot \tilde{\mathbf{g}}_\beta + \mathbf{g}_\alpha^h \cdot \mathbf{g}_\beta^h + \mathbf{g}_\alpha^s \cdot \mathbf{g}_\beta^s \\ & + \tilde{\mathbf{g}}_\alpha \cdot \mathbf{g}_\beta^h + \mathbf{g}_\alpha^h \cdot \tilde{\mathbf{g}}_\beta - \tilde{\mathbf{g}}_\alpha \cdot \mathbf{g}_\beta^s - \mathbf{g}_\alpha^s \cdot \tilde{\mathbf{g}}_\beta \\ & - \mathbf{g}_\alpha^h \cdot \mathbf{g}_\beta^s - \mathbf{g}_\alpha^s \cdot \mathbf{g}_\beta^h. \end{aligned} \quad (\text{S1})$$

The six component vectors admit  $\binom{6}{2} = 15$  unordered off-diagonal dot products. Three are opposite-spin products within the same smooth, hard, or soft component, while twelve couple different components. Thus “15 cross terms” counts all off-diagonal pairs, not fifteen terms all absent from three independent spin-resolved evaluations. Separate smooth/hard/soft evaluations also combine diagonal terms with the wrong nonlinear structure. Reconstructing the fields first avoids this ambiguity and retains the nonlocal coupling of the model, which cannot be recovered by adding local algebraic corrections afterwards.

### S2.2 The discrete adjoint and moving grids

For regular-grid values  $u_g$  and atom-grid interpolation coefficients  $I_{ig}$ ,

$$u_i = \sum_g I_{ig} u_g + u_i^h - u_i^s, \quad \frac{\partial E}{\partial u_g} = \sum_i I_{ig} \frac{\partial E}{\partial u_i}. \quad (\text{S2})$$

The reverse mapping uses the transpose of the forward interpolation, with the energy quadrature weights included consistently in the derivative. Hard and soft adjoints carry plus and minus signs. This applies separately to density, its Cartesian gradient components, and kinetic-energy density. At fixed orbital density matrix  $P$ , the explicit XC derivative for a nuclear coordinate or strain parameter  $\lambda$  includes both field and geometric terms,

$$\left. \frac{\partial E_{\text{xc}}}{\partial \lambda} \right|_P = \sum_i \left. \frac{\partial E_{\text{xc}}}{\partial u_i} \frac{\partial u_i}{\partial \lambda} \right|_P + \left. \frac{\partial E_{\text{xc}}}{\partial \lambda} \right|_u. \quad (\text{S3})$$

The last term encompasses explicit quadrature, partition, coordinate, and model-environment dependence. The total-energy force and stress additionally use electronic stationarity and the Gaussian-basis/overlap Pulay contributions, including cell and reciprocal-lattice dependence. The XC energy alone is not stationary in  $P$ . An adjoint test at fixed fields is necessary but does not replace a self-consistent total-energy finite-difference (FD) test.

### S2.3 Finite-difference protocol

For a displaced coordinate  $R_{A,x}$ , the central FD estimate is

$$F_{A,x}^{\text{FD}}(h) = -\frac{E(R_{A,x} + h) - E(R_{A,x} - h)}{2h}. \quad (\text{S4})$$

For a homogeneous  $xx$  strain, both cell and Cartesian coordinates are transformed while fractional coordinates remain fixed. In the positive-pressure stress convention,

$$\sigma_{xx}^{\text{FD}}(\eta) = -\frac{E(+\eta) - E(-\eta)}{2\eta\Omega}. \quad (\text{S5})$$

At least two step sizes distinguish truncation error from SCF noise. An SCF threshold tightened for these checks does not change the production thresholds. A molecular coordinate-scaling virial is not a periodic stress-tensor validation, and energy agreement under symmetry reduction does not establish force equivalence after a symmetry-breaking displacement.

Self-consistent checks use one water molecule in a periodically repeated 8 Å cube at Gamma, without D3, to isolate the native electronic derivatives. The grid settings are 400/60 Ry, five multigrids, and 100 radial/434 angular points. The basis/core and reconstruction definitions match the corresponding production representations. The first hydrogen is displaced along  $x$  by  $h = 10^{-3}$  and  $5 \times 10^{-4}$  bohr. Homogeneous  $xx$  strains are  $\eta = 10^{-4}$  and  $5 \times 10^{-5}$ . Every perturbed calculation starts from the exact central wavefunction and converges independently to an OT residual below  $10^{-9}$ . All 45 executions have normal termination and zero execution/time exits. The regular-grid dimensions remain unchanged under these perturbations.

Table S1: Signed FD minus analytic derivatives. Force errors are in  $\mu E_h/\text{bohr}$  and stress errors in MPa. Columns give the two displacement/strain steps.

| Representation      | $\Delta F(10^{-3})$ | $\Delta F(5 \times 10^{-4})$ | $\Delta \sigma(10^{-4})$ | $\Delta \sigma(5 \times 10^{-5})$ |
|---------------------|---------------------|------------------------------|--------------------------|-----------------------------------|
| GPW/GTH             | +0.228              | +2.239                       | +0.038                   | -0.208                            |
| GAPW-XC/GTH         | -0.377              | +2.380                       | +0.012                   | -0.332                            |
| GAPW-AE             | +2.855              | -2.404                       | +0.007                   | -0.472                            |
| GAPW-GTH direct     | +0.212              | +1.414                       | +0.044                   | -0.185                            |
| GAPW-GTH one centre | -2.726              | -1.977                       | -0.032                   | -0.017                            |

These checks test the self-consistent total-energy derivative, including the reconstruction and moving-grid terms, rather than only model differentiation at fixed fields. The smaller steps do not uniformly reduce the error: subtraction of converged energies amplifies finite SCF and numerical noise. Consequently, no ideal quadratic step-size scaling is claimed. The test covers one Cartesian force component and one diagonal strain component in this periodic Gamma-point system. It is not a validation of every force component, shear strain, finite-k-point derivative, or symmetry-breaking displacement under point-group reduction. Exact inputs, outputs, timing records, and file/model/binary hashes accompany the machine-readable analysis.

## S3 Molecular Numerical Settings and Convergence

### S3.1 Basis, grid, and boundary conditions

The 70-reaction comparison uses the University of Zurich (UZH) 2026.2 AE TZVPP-MOLOPT-PBE-ae basis and corresponding explicit-electron Hamiltonian where available. The GTH part uses TZV2P-MOLOPT-PBE-GTH bases and GTH-PBE potentials.<sup>1,2</sup> The all-GTH protocol uses GAPW-XC throughout. The mixed protocols share the light-element AE energies and use GTH for the retained heavy-element species, including Bi, Te, and I. This differs from the def2-based AE/ECP protocol in the molecular GauXC paper.

The molecule-centred cells add 25 Å to each molecular Cartesian extent. Cell and Poisson periodicity are disabled and the analytic isolated-system Poisson solver is used. The production atom-composite quadrature has 100 radial points and the 434-point Lebedev rule. Plane-wave/relative cutoffs are 400/60 Ry for GAPW-XC/GTH and 640/60 Ry for the shared AE route. The executed inputs document the corresponding GTH one-centre settings. OT and diagonalization thresholds are  $5 \times 10^{-7}$  and  $5 \times 10^{-6}$ , respectively. The threshold controls the solver residual, not an energy error in  $\text{kcal mol}^{-1}$ .

Basis/core changes must be kept distinct from quadrature refinement. The reported AE/GTH differences are not a basis-only convergence series, and no complete-basis limit is inferred from them.

### S3.2 Regular-grid cutoff

The paired conformational test uses the same two ACONF geometries and evaluates the relative energy within each representation. The 800 Ry result of that representation is the numerical reference. The small difference between the two 800 Ry relative energies is not set to zero: convergence within a representation and agreement between representations are separate questions.

Table S2: Conformational energy of the paired ACONF geometries in kcal mol<sup>-1</sup>. Each error is relative to its own 800 Ry result. The relative cutoff is 60 Ry.

| Cutoff/Ry | GPW      | GAPW-XC  | $\Delta$ GPW | $\Delta$ GAPW-XC |
|-----------|----------|----------|--------------|------------------|
| 240       | 0.774857 | 0.726730 | 0.027401     | -0.018666        |
| 280       | 0.704008 | 0.739410 | -0.043448    | -0.005985        |
| 320       | 0.747292 | 0.745360 | -0.000165    | -0.000035        |
| 360       | 0.746783 | 0.745610 | -0.000673    | 0.000215         |
| 400       | 0.749608 | 0.745215 | 0.002152     | -0.000181        |
| 480       | 0.750496 | 0.745307 | 0.003039     | -0.000089        |
| 560       | 0.748038 | 0.745300 | 0.000581     | -0.000095        |
| 640       | 0.747593 | 0.745297 | 0.000136     | -0.000099        |
| 800       | 0.747457 | 0.745395 | 0.000000     | 0.000000         |

At 400 Ry, the deviations from the respective 800 Ry conformational energies are +0.002152 kcal mol<sup>-1</sup> for GPW and -0.000181 kcal mol<sup>-1</sup> for GAPW-XC. The faster GAPW-XC convergence supports separating hard local XC structure from the smooth auxiliary grid. This is an observable-specific cutoff test, distinct from comparison with an AE Hamiltonian. The complete inputs and outputs accompany the table.

### S3.3 Radial and angular quadrature

Table S3: Self-consistent GPW water quadrature test at 800/60 Ry. Total-energy shifts are relative to 200/974 and are in microhartree. Electron-integral errors refer to eight explicit electrons.

| Radial | Angular | $\Delta E/\mu E_h$ | $\Delta N/e$ |
|--------|---------|--------------------|--------------|
| 50     | 50      | -369.201           | +4.486e-04   |
| 50     | 100     | +237.729           | -4.318e-05   |
| 50     | 194     | +263.762           | -8.268e-06   |
| 50     | 302     | +307.424           | -1.074e-06   |
| 50     | 434     | +245.349           | -1.215e-07   |
| 100    | 50      | -592.561           | +4.491e-04   |
| 100    | 100     | -10.739            | -4.333e-05   |
| 100    | 302     | +72.144            | -8.919e-07   |
| 100    | 434     | +8.379             | -3.031e-07   |
| 100    | 590     | +6.084             | -2.206e-07   |
| 150    | 434     | +12.627            | -1.352e-07   |
| 150    | 590     | +10.143            | -2.079e-07   |
| 150    | 770     | +14.433            | -1.869e-07   |
| 200    | 590     | +11.319            | -2.275e-07   |
| 200    | 770     | +15.822            | -1.715e-07   |
| 200    | 974     | +0.000             | -1.347e-07   |

The water test refines radial and angular integration independently of the plane-wave representation. The electron integral is retained as a diagnostic, without rescaling the density. Neither the small water total-energy differences nor the conformational cutoff differences define a universal reaction-energy error bound. Frozen-density calculations that did not satisfy the stated SCF acceptance are not used as converged-energy evidence.

### S3.4 Molecular cell padding

Selected matched isolated-system calculations tested added extents of 22, 24, and 25 Å. The maximum species-energy changes are 0.159, 0.189, and 0.348  $\mu E_h$  for 22–24, 24–25, and 22–25 Å, respectively. The corresponding maximum reaction-energy changes are 0.000138, 0.000163, and 0.000298 kcal mol<sup>-1</sup>. Maximum changes of the atom-grid electron integral are  $7.94 \times 10^{-8}$ ,  $1.43 \times 10^{-7}$ , and  $9.72 \times 10^{-8}$  electrons. These comparisons retain the nonperiodic analytic electrostatic treatment. Old periodic or differently centred cells are not pooled with the final molecular data.

## S4 Molecular Reaction Data and Cross-Code Comparisons

The common set contains 70 reactions from 36 parent subsets of dietGMTKN55.<sup>3,4</sup> Each reaction has complete accepted species energies in every compared protocol. The remaining cases required longer SCF convergence. The final common set is fixed across methods. No additional reaction is removed on the basis of its reference error. The reaction index and source records in the repository define the selection explicitly.

The same D3 contribution is used in the native reaction comparison. Signed errors are calculation minus the official reference, and mean absolute errors (MAEs) are unweighted averages over the fixed common set. They are not full-100 dietGMTKN55 statistics or type-2 weighted total mean absolute deviations (WTMAD-2). The ozone-containing W4-11/138 reaction remains included, despite its large GTH error. It contributes 58.8% of the all-GTH squared error. Removing it would lower the 69-reaction GAPW-XC MAE to 2.563 kcal mol<sup>-1</sup>. This is an influence diagnostic, not the headline result. The direct/one-centre mixed comparison shares all light-element AE energies, and only one retained reaction changes by more than 10<sup>-9</sup> kcal mol<sup>-1</sup>.

Table S4: Complete common-set reaction energies in kcal mol<sup>-1</sup>. Reference values are official benchmark energies. Protocol labels follow Sec. S1. No reference-error outlier is omitted.

| Subset    | Reaction | Ref.       | GX         | HD         | HOC        |
|-----------|----------|------------|------------|------------|------------|
| ACONF     | 2        | 0.61400    | 0.30663    | 0.68210    | 0.68210    |
| ACONF     | 5        | 0.59500    | 0.30757    | 0.58761    | 0.58761    |
| ACONF     | 8        | 1.17800    | 0.77590    | 1.22152    | 1.22152    |
| ADIM6     | 2        | 1.99000    | 2.72718    | 2.08787    | 2.08787    |
| AHB21     | 1        | -17.79000  | -22.18751  | -20.29619  | -20.29619  |
| AHB21     | 15       | -8.62000   | -11.59294  | -10.86032  | -10.86032  |
| Amino20x4 | 23       | 4.04500    | 3.68879    | 4.42456    | 4.42456    |
| Amino20x4 | 48       | 0.54000    | 0.27631    | 0.90052    | 0.90052    |
| Amino20x4 | 56       | 1.88900    | 1.93478    | 1.58118    | 1.58118    |
| BH76      | 6        | 17.80000   | 18.56504   | 17.86395   | 17.86395   |
| BH76      | 18       | 2.50000    | -8.63844   | -1.27459   | -1.27459   |
| BH76RC    | 16       | 13.40000   | 10.89326   | 13.59788   | 13.59788   |
| BHPERI    | 1        | 35.30000   | 33.09940   | 39.25014   | 39.25014   |
| BHPERI    | 17       | 13.10000   | 10.81645   | 12.70129   | 12.70129   |
| BHPERI    | 23       | 27.80000   | 28.29192   | 26.47568   | 26.47568   |
| BHPERI    | 26       | 31.30000   | 31.40095   | 30.85649   | 30.85649   |
| BHROT27   | 1        | 2.73000    | 2.69043    | 2.83649    | 2.83649    |
| BHROT27   | 10       | 8.03000    | 8.43958    | 8.00707    | 8.00707    |
| BSR36     | 4        | 8.85000    | 8.44034    | 8.81562    | 8.81562    |
| BSR36     | 21       | 9.78000    | 8.34085    | 8.82335    | 8.82335    |
| BUT14DIOL | 28       | 2.83000    | 2.69982    | 4.33282    | 4.33282    |
| BUT14DIOL | 39       | 3.10000    | 3.11306    | 4.93629    | 4.93629    |
| BUT14DIOL | 63       | 4.31000    | 4.09728    | 5.97195    | 5.97195    |
| CARBHB12  | 3        | 2.42100    | 2.69672    | 3.68798    | 3.68798    |
| CARBHB12  | 4        | 9.96700    | 9.97430    | 15.57723   | 15.57723   |
| CARBHB12  | 6        | 3.02100    | 2.95159    | 5.16892    | 5.16892    |
| CARBHB12  | 11       | 3.24100    | 3.92467    | 3.06786    | 3.06786    |
| CDIE20    | 2        | 4.40000    | 4.48598    | 4.99901    | 4.99901    |
| CDIE20    | 14       | 4.40000    | 4.56044    | 4.54726    | 4.54726    |
| CHB6      | 1        | -34.43000  | -33.49431  | -34.18723  | -34.18723  |
| DC13      | 6        | -25.70000  | -32.03193  | -31.81763  | -31.81763  |
| DC13      | 12       | -58.70000  | -60.73559  | -67.15589  | -67.15589  |
| FH51      | 25       | -20.27000  | -21.62745  | -19.42107  | -19.42107  |
| G21EA     | 5        | 17.30000   | -1.42565   | 10.29928   | 10.29928   |
| G21EA     | 25       | 54.70000   | 50.95114   | 51.72694   | 51.72694   |
| G21IP     | 28       | 243.70900  | 239.85434  | 243.88740  | 243.88740  |
| G2RC      | 15       | -39.43000  | -40.17337  | -44.40802  | -44.40802  |
| HAL59     | 21       | 10.54000   | 12.10722   | 9.89950    | 9.89950    |
| HEAVY28   | 5        | 0.98000    | 2.32113    | 1.35292    | 1.35483    |
| ICONF     | 16       | 3.66000    | 4.07657    | 3.99415    | 3.99415    |
| IL16      | 3        | -116.91000 | -116.64846 | -119.94685 | -119.94685 |
| INV24     | 12       | 10.30000   | 11.08966   | 10.79061   | 10.79061   |
| INV24     | 19       | 6.20000    | 5.46217    | 6.10591    | 6.10591    |
| INV24     | 22       | 27.20000   | 27.35357   | 27.74964   | 27.74964   |
| ISO34     | 12       | 45.65000   | 48.13511   | 45.85998   | 45.85998   |

| Subset  | Reaction | Ref.      | GX        | HD        | HOC       |
|---------|----------|-----------|-----------|-----------|-----------|
| ISOL24  | 13       | 33.02000  | 34.47575  | 34.65021  | 34.65021  |
| MB16-43 | 4        | 356.84810 | 320.43581 | 358.96453 | 358.96453 |
| MCONF   | 7        | 2.92000   | 3.11166   | 2.61484   | 2.61484   |
| MCONF   | 22       | 5.31000   | 5.83140   | 4.73929   | 4.73929   |
| MCONF   | 30       | 4.86000   | 5.03309   | 5.16786   | 5.16786   |
| MCONF   | 42       | 7.32000   | 8.16255   | 7.33699   | 7.33699   |
| PArel   | 3        | 2.94000   | 2.90267   | 3.05754   | 3.05754   |
| PArel   | 5        | 7.19000   | 8.68203   | 7.75014   | 7.75014   |
| PArel   | 10       | 2.98000   | 2.64094   | 2.92259   | 2.92259   |
| PArel   | 20       | 2.15000   | 1.17708   | 1.12222   | 1.12222   |
| PNICO23 | 13       | 3.98000   | 4.16981   | 4.46224   | 4.46224   |
| PNICO23 | 17       | 7.10000   | 7.39778   | 7.31551   | 7.31551   |
| RG18    | 1        | 0.08000   | 0.17488   | 0.13391   | 0.13391   |
| RG18    | 15       | 0.24000   | 0.48594   | 0.28332   | 0.28332   |
| S66     | 11       | 5.42000   | 5.85282   | 5.96275   | 5.96275   |
| S66     | 17       | 17.18000  | 17.67587  | 19.17660  | 19.17660  |
| S66     | 24       | 2.82000   | 3.50720   | 3.46501   | 3.46501   |
| S66     | 45       | 1.75000   | 2.59070   | 2.03582   | 2.03582   |
| SCONF   | 10       | 5.65000   | 6.09695   | 8.67396   | 8.67396   |
| SIE4x4  | 7        | 31.30000  | 45.77575  | 51.75030  | 51.75030  |
| W4-11   | 136      | 126.38500 | 92.90307  | 129.75705 | 129.75705 |
| W4-11   | 138      | 147.42800 | 78.94035  | 156.69640 | 156.69640 |
| WATER27 | 17       | 76.50000  | 76.32827  | 82.77137  | 82.77137  |
| WATER27 | 20       | 26.60000  | 29.31104  | 33.76403  | 33.76403  |
| YBDE18  | 1        | 57.17000  | 55.99377  | 60.22366  | 60.22366  |

#### S4.1 Direct comparison with the previous molecular implementation

All PySCF/Skala reference values use Becke partitioning without an atomic-radius adjustment.

The source is the reaction-resolved collaborator dataset archived with the earlier molecular GauXC study, at repository revision 25e80ba.<sup>5,6</sup> Each native reaction is matched by its species geometries after translation, charge, multiplicity, and signed stoichiometric coefficients. Official references and weights are checked independently. All 70 retained reactions have complete GauXC AE/ECP and PySCF energies. Converged GauXC GPW-GTH energies cover 65 of them. Source reaction numbers differ from the native benchmark identifiers, and the deposited mapping resolves those aliases explicitly.

The comparisons are separated by core treatment. Native GAPW-XC/GTH is compared with GauXC GPW-GTH on the 65-reaction intersection. The mixed native AE/GTH protocols are compared with GauXC AE/def2-ECP and PySCF on the full 70-reaction common set. The earlier AE/ECP calculations use def2-TZVP/ma-def2-TZVP bases, with def2 ECPs for the heavier elements, rather than the native UZH basis/core protocol specified in Sec. S3.1. This is the principal protocol distinction: the finite Gaussian space changes and, where ECPs differ, so does the core Hamiltonian. The remaining reaction-energy differences are consistent with these basis/core and numerical integration choices. The earlier GPW-GTH calculations instead use a SCAN-optimized basis/pseudopotential combination, whereas the native GTH protocol uses its PBE-optimized counterpart. The comparison complements the numerical consistency tests in Secs. S2 and S3, which validate the native implementation.

The following tables abbreviate GauXC AE/def2-ECP as GauXC AE and GauXC GPW-GTH as GauXC GPW where space is limited. Mean absolute differences (MADs) refer to direct differences between calculated reaction energies, not differences of reference MAEs. Root-mean-square (RMS) errors and differences are reported separately.

Table S5: Reference errors on reaction-matched populations, in kcal mol<sup>-1</sup>. The first five rows use the fixed native 70-reaction set. The last two use its 65-reaction intersection with converged GauXC GPW-GTH data. Each source retains its recorded dispersion contribution.

| Method        | <i>N</i> | MAE    | Median absolute | RMS     | Maximum absolute |
|---------------|----------|--------|-----------------|---------|------------------|
| Native GX     | 70       | 3.5051 | 0.6025          | 10.6749 | 68.4877          |
| Native HD     | 70       | 1.8905 | 0.5549          | 3.6502  | 20.4503          |
| Native HOC    | 70       | 1.8905 | 0.5549          | 3.6502  | 20.4503          |
| GauXC AE/ECP  | 70       | 1.2378 | 0.2777          | 2.7842  | 14.9074          |
| PySCF         | 70       | 1.2314 | 0.3048          | 2.7662  | 14.9606          |
| Native GX     | 65       | 2.2193 | 0.4919          | 8.8172  | 68.4877          |
| GauXC GPW-GTH | 65       | 2.1811 | 0.5820          | 8.6673  | 67.4910          |

On the 65-reaction pseudopotential intersection, native GAPW-XC/GTH and GauXC GPW-GTH have reference MAEs of 2.2193 and 2.1811 kcal mol<sup>-1</sup>. Their direct MAD is 0.4631 kcal mol<sup>-1</sup>, with signed mean -0.2250, RMS difference 0.7799, and maximum absolute difference 3.9442 for DC13/6. By contrast, on all 70 reactions the mixed native protocols have reference MAEs of 1.8905, compared with 1.2378 for GauXC AE/ECP and 1.2314 for PySCF. The Mixed D-GauXC AE/ECP MAD is 1.8889 kcal mol<sup>-1</sup>, with signed mean +0.6494 and median absolute difference 0.7151. The largest difference, +18.6055 kcal mol<sup>-1</sup> for W4-11/138, remains included. No outlier is removed from either intersection.

Table S6: Direct reaction-energy differences, in kcal mol<sup>-1</sup>. The first three difference statistics use the recorded total energies. The last column removes each source's own D3 contribution to isolate its influence on the comparison.

| Method pair               | <i>N</i> | MAD    | RMS    | Maximum | Electronic MAD |
|---------------------------|----------|--------|--------|---------|----------------|
| Native GX / GauXC GPW-GTH | 65       | 0.4631 | 0.7799 | 3.9442  | 0.4624         |
| Native HD / GauXC AE/ECP  | 70       | 1.8889 | 3.5718 | 18.6055 | 1.8880         |
| Native HOC / GauXC AE/ECP | 70       | 1.8889 | 3.5718 | 18.6055 | 1.8880         |
| Native HD / PySCF         | 70       | 1.8925 | 3.5585 | 18.3927 | 1.8857         |
| GauXC AE/ECP / PySCF      | 70       | 0.0456 | 0.0787 | 0.4059  | 0.0400         |

All headline comparisons retain the total energies as recorded, including each source's separately evaluated D3(BJ) correction. The source calculations use the B3LYP5 dispersion convention, but the recorded corrections are not numerically identical across every protocol. For native Mixed D versus GauXC AE/ECP, the D3 reaction-contribution differences have a MAD of only 0.001396 kcal mol<sup>-1</sup> and a maximum of 0.014549. Removing each source's own correction changes the direct MAD from 1.8889 to 1.8880 kcal mol<sup>-1</sup>. Dispersion bookkeeping therefore does not explain the larger native-GauXC differences. For native GAPW-XC/GTH versus GauXC GPW-GTH, the corresponding MAD changes from 0.4631 to 0.4624. The latter reproduces the earlier comparison in which the same native D3 term was added to both electronic reaction energies.

There is one non-negligible recorded GauXC AE/ECP-PySCF dispersion mismatch within the retained set: HEAVY28/5 differs by +0.419041 kcal mol<sup>-1</sup> in its D3 reaction contribution. Its origin is not established, and neither source value is overwritten. Across the 70 reactions, the GauXC-PySCF direct MAD is 0.04560 for the recorded totals and 0.03999 for the electronic contributions alone. Their total-energy differences should therefore not all be attributed to quadrature. The comparison files preserve electronic, dispersion, and total energies separately, together with input/output hashes and the pinned source revision.

Table S7: Reaction-resolved collaborator energies on the native common set, in kcal mol<sup>-1</sup>. Reaction identifiers and official references match Table S4. A dash denotes an unavailable converged GPW-GTH value, not a zero. Recorded dispersion contributions are included.

| Subset    | Reaction | Ref.      | GauXC GPW | GauXC AE  | PySCF     |
|-----------|----------|-----------|-----------|-----------|-----------|
| ACONF     | 2        | 0.61400   | 0.36237   | 0.64648   | 0.68837   |
| ACONF     | 5        | 0.59500   | 0.31848   | 0.55307   | 0.52619   |
| ACONF     | 8        | 1.17800   | 0.77859   | 1.18813   | 1.15144   |
| ADIM6     | 2        | 1.99000   | 2.30458   | 1.85184   | 1.85984   |
| AHB21     | 1        | -17.79000 | -20.45402 | -17.39888 | -17.40971 |
| AHB21     | 15       | -8.62000  | -11.71918 | -8.79444  | -8.79323  |
| Amino20x4 | 23       | 4.04500   | 3.85348   | 3.71127   | 3.63208   |
| Amino20x4 | 48       | 0.54000   | 0.01575   | 0.60110   | 0.61964   |
| Amino20x4 | 56       | 1.88900   | 2.20238   | 2.03964   | 2.03641   |
| BH76      | 6        | 17.80000  | 19.11753  | 17.03316  | 17.15220  |
| BH76      | 18       | 2.50000   | —         | -0.21392  | -0.22005  |
| BH76RC    | 16       | 13.40000  | 10.99734  | 14.26280  | 14.25226  |
| BHPERI    | 1        | 35.30000  | 34.24032  | 36.03270  | 36.07884  |
| BHPERI    | 17       | 13.10000  | 11.23120  | 12.99901  | 13.04625  |
| BHPERI    | 23       | 27.80000  | 28.41555  | 28.73250  | 28.79864  |
| BHPERI    | 26       | 31.30000  | 31.41251  | 31.55939  | 31.44178  |
| BHROT27   | 1        | 2.73000   | 2.83023   | 2.88882   | 2.89208   |
| BHROT27   | 10       | 8.03000   | 8.43419   | 8.32717   | 8.32723   |
| BSR36     | 4        | 8.85000   | 8.73777   | 8.15235   | 8.24610   |
| BSR36     | 21       | 9.78000   | 9.14667   | 7.99739   | 8.10612   |
| BUT14DIOL | 28       | 2.83000   | 2.76824   | 2.95418   | 2.96389   |
| BUT14DIOL | 39       | 3.10000   | 3.33710   | 3.62532   | 3.65624   |

| Subset    | Reaction | Ref.       | GauXC GPW  | GauXC AE   | PySCF      |
|-----------|----------|------------|------------|------------|------------|
| BUT14DIOL | 63       | 4.31000    | 4.08891    | 4.48423    | 4.52609    |
| CARBHB12  | 3        | 2.42100    | 2.64009    | 2.55364    | 2.56030    |
| CARBHB12  | 4        | 9.96700    | 9.78532    | 10.38773   | 10.36877   |
| CARBHB12  | 6        | 3.02100    | 2.80568    | 2.36925    | 2.38662    |
| CARBHB12  | 11       | 3.24100    | 3.96770    | 3.39157    | 3.39480    |
| CDIE20    | 2        | 4.40000    | 4.52498    | 4.36188    | 4.36983    |
| CDIE20    | 14       | 4.40000    | 4.80957    | 4.23473    | 4.22647    |
| CHB6      | 1        | -34.43000  | -32.06178  | -35.71422  | -35.72727  |
| DC13      | 6        | -25.70000  | -28.08768  | -32.02494  | -32.00898  |
| DC13      | 12       | -58.70000  | -62.03843  | -63.80513  | -63.59949  |
| FH51      | 25       | -20.27000  | -22.33482  | -21.14074  | -21.08624  |
| G21EA     | 5        | 17.30000   | –          | 17.43069   | 17.44739   |
| G21EA     | 25       | 54.70000   | 52.40220   | 56.66247   | 56.67548   |
| G21IP     | 28       | 243.70900  | 240.91373  | 246.10544  | 246.06756  |
| G2RC      | 15       | -39.43000  | -40.58825  | -36.07029  | -35.92457  |
| HAL59     | 21       | 10.54000   | 13.16196   | 10.18052   | 10.22753   |
| HEAVY28   | 5        | 0.98000    | –          | 1.01418    | 0.60831    |
| ICONF     | 16       | 3.66000    | 4.21525    | 3.91693    | 3.89953    |
| IL16      | 3        | -116.91000 | -116.14007 | -117.17430 | -117.26645 |
| INV24     | 12       | 10.30000   | 11.29756   | 10.42813   | 10.40776   |
| INV24     | 19       | 6.20000    | 6.15822    | 5.90506    | 5.91207    |
| INV24     | 22       | 27.20000   | 28.71502   | 27.03272   | 27.02526   |
| ISO34     | 12       | 45.65000   | 48.12960   | 44.81214   | 44.83039   |
| ISOL24    | 13       | 33.02000   | 34.15054   | 32.82246   | 32.75595   |
| MB16-43   | 4        | 356.84810  | –          | 349.30785  | 349.27767  |
| MCONF     | 7        | 2.92000    | 3.40584    | 2.76081    | 2.75618    |
| MCONF     | 22       | 5.31000    | 5.98878    | 5.70352    | 5.65919    |
| MCONF     | 30       | 4.86000    | 5.02760    | 5.10589    | 5.07451    |
| MCONF     | 42       | 7.32000    | 8.08468    | 7.86694    | 7.84842    |
| PArel     | 3        | 2.94000    | 3.04128    | 2.97323    | 2.95625    |
| PArel     | 5        | 7.19000    | 9.07152    | 7.36215    | 7.35467    |
| PArel     | 10       | 2.98000    | 2.86582    | 3.35368    | 3.35918    |
| PArel     | 20       | 2.15000    | 1.55954    | 1.95014    | 2.02341    |
| PNICO23   | 13       | 3.98000    | 4.24282    | 4.57238    | 4.57621    |
| PNICO23   | 17       | 7.10000    | 7.50137    | 7.35457    | 7.39182    |
| RG18      | 1        | 0.08000    | 0.12640    | 0.13584    | 0.13272    |
| RG18      | 15       | 0.24000    | 0.34321    | 0.48691    | 0.49133    |
| S66       | 11       | 5.42000    | 5.52785    | 5.65511    | 5.66394    |
| S66       | 17       | 17.18000   | 16.58982   | 17.11875   | 17.10879   |
| S66       | 24       | 2.82000    | 2.91409    | 3.02703    | 3.01553    |
| S66       | 45       | 1.75000    | 2.48973    | 1.89424    | 1.90961    |
| SCONF     | 10       | 5.65000    | 6.23204    | 6.22279    | 6.19781    |
| SIE4x4    | 7        | 31.30000   | 45.74439   | 46.20737   | 46.26060   |
| W4-11     | 136      | 126.38500  | –          | 119.95421  | 120.14384  |
| W4-11     | 138      | 147.42800  | 79.93703   | 138.09093  | 138.30367  |
| WATER27   | 17       | 76.50000   | 76.34800   | 81.45738   | 81.39384   |
| WATER27   | 20       | 26.60000   | 30.45634   | 26.30882   | 26.24992   |
| YBDE18    | 1        | 57.17000   | 54.63500   | 58.85771   | 58.76963   |

For context, the prior GauXC manuscript’s updated primary AE/ECP–PySCF comparison comprises all 100 reactions, not these 65 or 70.<sup>5</sup> The values in Table S8 are from that updated analysis and are consistent with the reaction-resolved source supplied on 11 September 2026. They supersede earlier draft aggregates and differ from the first arXiv version. They must not be compared with the native aggregates as though they used the same reaction population.

Table S8: Context from the updated 100-reaction molecular GauXC manuscript. These aggregates are quoted for their original set, not rescored or compared as if they belonged to the native 70-reaction set. Energies are in kcal mol<sup>-1</sup>.

| Metric                | CP2K/GauXC AE/ECP | Microsoft PySCF |
|-----------------------|-------------------|-----------------|
| MAE against reference | 1.255             | 1.243           |
| WTMAD-2               | 3.486             | 3.459           |

Direct cross-code MAD: 0.058; signed-error  $R^2$ : 0.998; all absolute cross-code residuals below 0.5.

The difference of the 100-reaction aggregate MAEs,  $0.012 \text{ kcal mol}^{-1}$ , is not the direct cross-code MAD of  $0.058 \text{ kcal mol}^{-1}$ . Nor can either quantify native/GauXC equivalence under a changed basis and core treatment. Tables S5–S7 instead provide the explicit common-set comparison directly from the deposited species energies.

## S5 K-Point and Symmetry Checks

Symmetry reduction is a change in the quadrature representation of a fixed Brillouin-zone integral. Correct weights, rotations of orbital components, and Bloch-phase conventions are required. For FD forces, the perturbed geometry may have lower symmetry than the reference geometry. Consistent unreduced sampling or correctly regenerated symmetry is needed. Inversion/time-reversal reduction and full point-group reduction are not interchangeable validation labels.

Seven archived controls compare complete and reduced  $4^3$  meshes for C and MgO at unchanged geometry and scientific input. The full point-group reduction uses 10 irreducible points from 64, at symmetry tolerance  $10^{-8}$ . Table S9 lists the absolute energy difference per reported cell. These controls establish agreement for their specific geometries, meshes, and executables. They are not a proof for all future restarts or all symmetry operations.

Table S9: Selected full/reduced-grid energy checks. Residuals are the final printed SCF values.  $|\Delta E|$  is in  $\text{kJ mol}^{-1}$  of simulation cells.

| System | Representation             | Full SCF residual    | Reduced SCF residual | $ \Delta E $ |
|--------|----------------------------|----------------------|----------------------|--------------|
| C      | GAPW-XC/GTH                | $3.0 \times 10^{-7}$ | $2.1 \times 10^{-7}$ | 0.0200       |
| C      | GAPW-AE                    | $8.0 \times 10^{-8}$ | $9.0 \times 10^{-8}$ | 0.0229       |
| C      | Mixed direct (AE here)     | $8.0 \times 10^{-8}$ | $9.0 \times 10^{-8}$ | 0.0229       |
| C      | Mixed one centre (AE here) | $8.0 \times 10^{-8}$ | $9.0 \times 10^{-8}$ | 0.0229       |
| MgO    | GAPW-XC/GTH                | $4.7 \times 10^{-7}$ | $3.6 \times 10^{-7}$ | 0.0291       |
| MgO    | Mixed direct               | $2.7 \times 10^{-7}$ | $4.9 \times 10^{-7}$ | 0.0493       |
| MgO    | Mixed one centre           | $1.4 \times 10^{-7}$ | $9.0 \times 10^{-8}$ | 0.0378       |

The C inputs of both mixed protocols use an AE carbon basis and potential, so those rows do not constitute additional GTH validations. Mixed MgO uses GTH Mg and AE O. The molecular-crystal production uses full point-group reduction of  $3^3$  meshes. The independently reported  $\text{CO}_2$  <sup>53</sup> and  $\text{NH}_3$  tolerance checks below constrain sampling and approximate-symmetry sensitivity for those cases. They should not be relabelled as full-versus-reduced tests.

## S6 Molecular-Crystal Energies and Numerical Controls

### S6.1 Geometries and electronic protocol

The crystal and gas-phase geometries are taken from Sec. 5.6 of the versioned supporting dataset arXiv:2402.13059v1 of Della Pia *et al.*<sup>7</sup> Crystal coordinates are wrapped into the supplied cells without relaxation or symmetrization.  $\text{CO}_2$  and  $\text{NH}_3$  have cubic cells with  $a = 5.624$  and  $5.1305 \text{ \AA}$  and four molecules each. Urea has a tetragonal cell with  $a = b = 5.565$ ,  $c = 4.684 \text{ \AA}$ , and two molecules. The separately supplied gas molecules are rigidly translated into centred cells whose dimensions are the molecular extents plus  $25 \text{ \AA}$ . They use nonperiodic cell/Poisson settings and the analytic isolated-system solver.

All four representations use Skala-1.1 Rev1 on a smoothly partitioned atom-composite quadrature with 150 radial points and the 770-point Lebedev rule. Five multigrids use cutoffs of 800/60 Ry in both phases. AE calculations use UZH 2026.2 TZVPP-MOLOPT-PBE-ae bases and explicit nuclei. GTH calculations use TZV2P-MOLOPT-PBE-GTH and GTH-PBE potentials. One-centre calculations employ the small extended expansion and accurate augmented-XC integration. The AE hard-minus-soft fields are reconstructed before the joint model evaluation. Direct and reconstructed GTH calculations share their Hamiltonian and basis.

D3(BJ) is included once, with  $s_6 = 1$ ,  $s_8 = 1.9889$ ,  $a_1 = 0.3981$ ,  $a_2 = 4.4211$ , and interaction cutoff 20 in the executed input convention.<sup>8,9</sup> The B3LYP parameter alias selects these damping constants, not the electronic functional. All systems are neutral restricted singlets. OT convergence is  $5 \times 10^{-7}$ . Complex k-point orbitals use iterative refinement of the approximative congruence transformation with the full AE preconditioner. No density renormalization, energy alignment, or fitting to the reference lattice energies is applied.

Every crystal uses a Gamma-centred  $3^3$  mesh with full point-group reduction. With symmetry tolerance  $10^{-6}$ , AE and GAPW-XC give 4, 10, and 10 irreducible points for  $\text{CO}_2$ ,  $\text{NH}_3$ , and urea. The GTH-direct and GTH-1C calculations use  $10^{-4}$  and give 4, 4, and 6. These finite tolerances recognize near-symmetries in the decimal coordinates without moving the atoms. Their numerical sensitivity is distinct from a full-versus-reduced-grid comparison.

Table S10: Accepted total energies including D3(BJ), in hartree, and paired lattice energies in  $\text{kJ mol}^{-1}$ . Crystal energies refer to the complete cell. Extra digits support reproducibility and are not an accuracy claim.

| System          | Representation       | Z | $E_{\text{solid}}$ | $E_{\text{molecule}}$ | $E_{\text{latt}}$ |
|-----------------|----------------------|---|--------------------|-----------------------|-------------------|
| CO <sub>2</sub> | GAPW-XC-GTH          | 4 | -151.045238668728  | -37.747061778360      | -37.407827        |
| CO <sub>2</sub> | GAPW-AE              | 4 | -754.204172576782  | -188.537114495983     | -36.569661        |
| CO <sub>2</sub> | GAPW-GTH, direct     | 4 | -151.045104012149  | -37.747040234695      | -37.376005        |
| CO <sub>2</sub> | GAPW-GTH, one center | 4 | -151.045343896582  | -37.747094149196      | -37.391906        |
| NH <sub>3</sub> | GAPW-XC-GTH          | 4 | -47.028180200438   | -11.741781495152      | -40.074458        |
| NH <sub>3</sub> | GAPW-AE              | 4 | -226.289314085879  | -56.557571602094      | -38.744287        |
| NH <sub>3</sub> | GAPW-GTH, direct     | 4 | -47.028284463885   | -11.741799977614      | -40.094368        |
| NH <sub>3</sub> | GAPW-GTH, one center | 4 | -47.028166197256   | -11.741777335794      | -40.076187        |
| Urea            | GAPW-XC-GTH          | 2 | -88.087642387844   | -44.000505611404      | -113.725046       |
| Urea            | GAPW-AE              | 2 | -450.589911407247  | -225.247483909549     | -124.637178       |
| Urea            | GAPW-GTH, direct     | 2 | -88.087842193572   | -44.000575209028      | -113.804613       |
| Urea            | GAPW-GTH, one center | 2 | -88.087646982008   | -44.000510082308      | -113.719339       |

Table S11: Bounded numerical sensitivities in  $\text{kJ mol}^{-1}$  per molecule. All unlisted scientific settings are held fixed. Seven accepted control executions produce six comparisons because the atom-grid refinement requires both phases.

| System and representation     | Change                                    | Observable                             | Shift     |
|-------------------------------|-------------------------------------------|----------------------------------------|-----------|
| CO <sub>2</sub> , GAPW-XC-GTH | 150/770 to 200/974                        | $\Delta E_{\text{latt}}$ , both phases | +0.044554 |
| CO <sub>2</sub> , GAPW-XC-GTH | 800 to 1200 Ry                            | $\Delta(E_{\text{solid}}/Z)$ only      | -0.015499 |
| CO <sub>2</sub> , GAPW-XC-GTH | $3^3$ to $5^3$                            | $\Delta E_{\text{latt}}$               | +0.000391 |
| NH <sub>3</sub> , GAPW-XC-GTH | Symmetry tolerance $10^{-6}$ to $10^{-4}$ | $\Delta E_{\text{latt}}$               | -0.000429 |
| CO <sub>2</sub> , GAPW-XC-GTH | CPU minus CUDA                            | $\Delta E_{\text{latt}}$               | -0.000127 |
| Urea, GAPW-AE                 | CPU minus CUDA                            | $\Delta E_{\text{latt}}$               | -0.000379 |

## S6.2 Paired energies and controls

Table S10 gives all paired total energies. We use  $E_{\text{latt}} = E_{\text{solid}}/Z - E_{\text{molecule}}$  and  $1 E_h = 2625.4996394799 \text{ kJ mol}^{-1}$ . The electronic diffusion Monte Carlo (DMC) references have statistical uncertainties of 0.2, 0.1, and 0.3  $\text{kJ mol}^{-1}$  for CO<sub>2</sub>, NH<sub>3</sub>, and urea. Their estimated overall uncertainty is about 2  $\text{kJ mol}^{-1}$ , rather than the smaller statistical bars alone.<sup>7</sup> Thermal and zero-point corrections to experimental sublimation enthalpies are not added to the DMC energies.

Device comparisons distinguish central processing unit (CPU) inference from accelerator inference through CUDA. The CPU/CUDA labels refer only to this device choice.

For NH<sub>3</sub>, CO<sub>2</sub>, and urea, respectively, the absolute AE errors are 0.544, 7.170, and 16.137  $\text{kJ mol}^{-1}$ , or 1.4%, 24.4%, and 14.9% of the DMC binding magnitudes. GAPW-XC/GTH gives 1.874, 8.008, and 5.225  $\text{kJ mol}^{-1}$ , or 4.9%, 27.2%, and 4.8%. Thus NH<sub>3</sub> has the smallest absolute error in each representation, whereas the CO<sub>2</sub>/urea ordering depends on both core treatment and the error measure. These errors describe the combined Skala-D3(BJ) protocol at the stated basis and geometry, not separate XC, dispersion, or deformation-energy components.

Refining the paired CO<sub>2</sub> quadrature from 150/770 to 200/974 changes  $E_{\text{solid}}/Z$  by -0.110687 and  $E_{\text{molecule}}$  by -0.155241  $\text{kJ mol}^{-1}$ . Their difference, +0.044554  $\text{kJ mol}^{-1}$ , measures the lattice-energy sensitivity. The  $5^3$  crystal check uses 11 of 125 irreducible points, compared with four of 27 at  $3^3$ . For k-point, symmetry-tolerance, and device checks the molecule is unchanged, so the crystal shift per molecule is also the lattice-energy shift.

The accepted 1200 Ry cutoff result concerns only the CO<sub>2</sub> crystal and is not a paired binding-energy convergence test. The separate gas-phase calculation lacks a complete runtime-provenance record and is not used. CO<sub>2</sub> CPU/CUDA and urea-AE CPU/CUDA controls retain the respective executable and scientific input apart from model device selection. Their sub-0.001  $\text{kJ mol}^{-1}$  changes constrain device effects for these cases. No complete-basis limit, basis-set superposition error (BSSE) correction, equilibrium volume, elastic response, or vibrational thermodynamics is inferred from the fixed-geometry energies.

## S6.3 Other density-functional approximations

Table S12 compares the native results with the DMC references and published electronic lattice energies collected by Della Pia *et al.*<sup>7;10</sup> Their plane-wave calculations use hard projector augmented-wave (PAW) datasets, a 1000 eV cutoff, and dense system-specific k-point meshes. The PBE0+MBD values use FHI-aims. These are literature protocols, not same-basis native-code controls or Skala reruns. The MAE is the unweighted mean of  $|E_{\text{latt}} - E_{\text{latt}}^{\text{DMC}}|$  over CO<sub>2</sub>, NH<sub>3</sub>,

and urea, using unrounded native energies and the reported literature values.

Table S12: Electronic lattice energies and three-crystal MAEs, all in  $\text{kJ mol}^{-1}$ . DMC references are from Ref. 7, with statistical uncertainties in parentheses. Native Skala-D3(BJ) values are from Table S10. Other density-functional values are transcribed from Tables S5, S8, and S22 of Ref. 10. Each dispersion model is part of the stated method. MAEs summarize only these three crystals and are not full-X23 statistics. Different geometrical and numerical protocols preclude interpreting the differences as implementation errors or a definitive method ranking.

| Method                            | CO <sub>2</sub> | NH <sub>3</sub> | Urea      | MAE   |
|-----------------------------------|-----------------|-----------------|-----------|-------|
| DMC reference                     | -29.4(2)        | -38.2(1)        | -108.5(3) | –     |
| Skala-D3(BJ), GAPW-XC/GTH         | -37.41          | -40.07          | -113.73   | 5.04  |
| Skala-D3(BJ), GAPW-AE             | -36.57          | -38.74          | -124.64   | 7.95  |
| Skala-D3(BJ), GAPW-GTH direct     | -37.38          | -40.09          | -113.80   | 5.06  |
| Skala-D3(BJ), GAPW-GTH one-centre | -37.39          | -40.08          | -113.72   | 5.03  |
| PBE                               | -6.51           | -28.24          | -73.24    | 22.70 |
| PBE+D3                            | -25.07          | -43.76          | -109.90   | 3.76  |
| PBE+MBD                           | -22.74          | -42.10          | -109.11   | 3.72  |
| revPBE+D3                         | -22.60          | -38.32          | -99.88    | 5.18  |
| SCAN+rVV10                        | -31.47          | -41.69          | -113.55   | 3.54  |
| vdW-DF2                           | -32.75          | -39.70          | -103.85   | 3.17  |
| PBE0+MBD (FHI-aims)               | -23.74          | -38.99          | -109.40   | 2.45  |

The native pseudopotential MAEs of 5.03–5.06  $\text{kJ mol}^{-1}$  are close to revPBE+D3 (5.18), although several other dispersion-inclusive approximations have smaller three-crystal MAEs. The AE MAE of 7.95  $\text{kJ mol}^{-1}$  is dominated by urea. These aggregates conceal useful material dependence: NH<sub>3</sub> is accurate in both native core treatments, and pseudopotential urea errors are close to SCAN+rVV10 and vdW-DF2. CO<sub>2</sub> and AE urea remain the main limitations. The approximately 2  $\text{kJ mol}^{-1}$  overall DMC uncertainty and the differing computational protocols caution against ranking methods by small residual differences.<sup>7</sup> The results support partially successful condensed-phase transfer with a consistent overbinding tendency, not a general conclusion about all molecular crystals.

The accompanying benchmark directory contains six source geometries, 24 base inputs, accepted executed inputs, and compact energy and provenance records. Calculations use CP2K revision 37aedacbb3, with full source and host-specific executable hashes in the archive. The crystal energies and numerical-control results are retained separately.

## S7 Ten-Solid Equations of State

### S7.1 Structures and common fit windows

The selected solids are AlN, AlP, BN, BP, C, LiCl, MgO, MgS, Si, and SiC from the Goldzak set.<sup>11</sup> All use conventional cubic eight-atom cells. Ten initial volumes are evenly spaced from 0.90 to 1.10 times the static-lattice experimental volume. The common fit window contains the first nine volumes for AlN and AlP, ending at 1.0777778, and all ten for the other solids. The 40 method–solid comparisons derive from 36 independent curves and 352 unique energies: the AE curves for BN and C are shared by both mixed protocols. Third-order Birch–Murnaghan fits give the equilibrium lattice constant and elastic response under isotropic compression.

The comparison uses the ten solids for which consistent EOS data were available in every representation. LiF and LiH are not included. The selection is identical across protocols and was not based on the magnitude of the reference error.

### S7.2 Computational details and fit stability

All selected points use 800/60 Ry cutoffs, 150 radial points, the 770-point Lebedev rule, and the small extended one-centre basis. All are neutral, spin-restricted calculations with UZH 2026.2 basis and potential definitions. The mixed protocols retain AE H, Li, B, C, N, O, and F and use GTH Mg, Al, Si, P, S, and Cl. Diagonalization and OT use residual thresholds of  $5 \times 10^{-6}$  and  $5 \times 10^{-7}$ , respectively. Exact inputs specify the dispersion and integration conventions.

The actual Gamma-centred meshes are listed in Table S22. The mesh is identical across representations at each volume but can change with volume within an EOS. Consequently, a smooth fit is not by itself evidence of k-point convergence. The selected full/reduced-mesh checks in the preceding section constrain symmetry reduction for their tested cases, not all possible geometries and restarts.

Every full-window minimum is bracketed. The largest fit RMS residual is 0.674 meV per atom. Omitting one endpoint changes the fitted lattice constant by at most 0.002444 Å (mixed-direct MgS, smallest-volume point omitted)

Table S13: Selected  $a_0$  values (Å). Experimental references follow Ref. 11. BN and C reuse the AE curve for HD and HOC. These entries are not independent calculations.

| Solid | Experiment | GX     | AE     | HD     | HOC    |
|-------|------------|--------|--------|--------|--------|
| AlN   | 4.3680     | 4.2811 | 4.3025 | 4.2747 | 4.2746 |
| AlP   | 5.4480     | 5.3622 | 5.3781 | 5.3616 | 5.3621 |
| BN    | 3.5930     | 3.5777 | 3.5675 | 3.5675 | 3.5675 |
| BP    | 4.5250     | 4.4439 | 4.4696 | 4.4527 | 4.4527 |
| C     | 3.5530     | 3.5345 | 3.5292 | 3.5292 | 3.5292 |
| LiCl  | 5.0720     | 4.9081 | 4.9155 | 4.9178 | 4.9179 |
| MgO   | 4.1890     | 4.1059 | 4.0935 | 4.0923 | 4.0919 |
| MgS   | 5.1880     | 5.0689 | 5.0536 | 5.0700 | 5.0697 |
| Si    | 5.4210     | 5.3386 | 5.3740 | 5.3383 | 5.3383 |
| SiC   | 4.3470     | 4.2517 | 4.2920 | 4.2556 | 4.2556 |

and the bulk modulus by at most 9.997 GPa (GAPW-XC AlN, largest-volume point omitted). The smallest-volume point is essential to bracketing the LiCl minimum. These tests assess fit-window sensitivity. They do not measure basis or quadrature errors and are not transferable error bars for other curves.

### S7.3 Representation and material trends

All native lattice constants are below the static-lattice experimental reference. LiCl contracts most, by approximately 0.154–0.164 Å, while C and BN contract by about 0.015–0.026 Å. MgS is also strongly contracted. The common sign spans diamond, zincblende, and rocksalt structures rather than a single bonding class.

GAPW-XC and AE differ on average by 0.01893 Å in  $a_0$  and 10.55 GPa in  $B_0$ , but this comparison changes basis and core treatment as well as density representation. At fixed mixed-core Hamiltonian, the one-centre correction changes  $a_0$  by 0.000135 Å on average and at most 0.000542 Å (AlP). The corresponding changes in  $B_0$  average 1.11 GPa and reach 5.75 GPa (MgO). Excluding the two shared AE curves gives mean changes of 0.000169 Å and 1.39 GPa over eight independent pairs. The main contraction trend is therefore insensitive to this representation choice.

Curvature errors are less uniform. GAPW-XC underestimates the Si bulk modulus, 92.76 versus 100.3 GPa, despite a contracted lattice, whereas AE gives 100.56 GPa. For C, GAPW-XC is closer to the reference modulus than AE. A shorter equilibrium lattice does not necessarily imply a larger bulk modulus, and AE does not improve every property of every material. Neither  $a_0$  nor  $B_0$  determines a cohesive energy without atomic references.

### S7.4 Matched literature comparisons

All methods are compared against the same primary static-lattice experimental values from Goldzak *et al.*<sup>11</sup> Their Hartree–Fock (HF), second-order Møller–Plesset perturbation theory (MP2), spin-component-scaled MP2 (SCS-MP2), and scaled opposite-spin MP2 (SOS-MP2) results cover all ten selected solids. Zhang *et al.*<sup>12</sup> supply local-density approximation (LDA), PBE, PBEsol, M06-L, SCAN, and HSE06 values for nine, excluding AlN. We use their uncorrected static electronic values, not predictions with zero-point corrections. Their SCAN calculations were non-self-consistent on PBE densities. Schimka *et al.*<sup>13</sup> provide PBE, PBEsol, HSE06, and HSEsol lattice constants for nine, excluding MgS. No missing bulk modulus is inferred. Self-consistent Tao–Mo results<sup>14</sup> cover eight, excluding AlN and BN, and are rescored against the same Goldzak reference.

Our periodic GFN manuscript<sup>15</sup> supplies GFN1-xTB and GFN2-xTB lattice constants for nine shared solids, but no bulk moduli in the curated comparison table. Its ten-material set excludes MgO and LiH, whereas the present set excludes LiF and LiH. The two ten-material selections must therefore not be treated as identical.

SCS-MP2 and SOS-MP2 give matched ten-solid lattice MAEs of 0.0125 and 0.0149 Å, and bulk-modulus MAEs of 4.10 and 3.88 GPa. The native ranges are 0.0728–0.0844 Å and 16.61–21.77 GPa. Established semilocal and screened-hybrid methods also have smaller lattice errors on the corresponding intersections. These results identify a limitation of the present structural predictions. The comparisons do not uniquely separate functional and numerical contributions.

Alternative experimental moduli listed by Goldzak *et al.* for BN, BP, and MgO are 410.2, 168.0, and 169.8 GPa, instead of the primary 388.5, 176.5, and 173.0 GPa. Simultaneously replacing all three lowers each native ten-solid modulus MAE by 1.00 GPa and leaves the qualitative conclusion intact. Broader 44-solid statistics, such as those of Tran *et al.*,<sup>16</sup> are not relabelled as matched ten-solid errors.

Table S14: Selected  $B_0$  values (GPa). Experimental references follow Ref. 11. BN and C reuse the AE curve for HD and HOC. These entries are not independent calculations.

| Solid | Experiment | GX     | AE     | HD     | HOC    |
|-------|------------|--------|--------|--------|--------|
| AlN   | 206.00     | 239.07 | 225.89 | 240.42 | 240.40 |
| AlP   | 87.00      | 94.34  | 101.35 | 96.36  | 93.12  |
| BN    | 388.50     | 423.39 | 420.87 | 420.87 | 420.87 |
| BP    | 176.50     | 187.93 | 183.64 | 185.86 | 185.86 |
| C     | 453.30     | 457.33 | 491.35 | 491.35 | 491.35 |
| LiCl  | 37.30      | 47.98  | 46.37  | 47.13  | 47.16  |
| MgO   | 173.00     | 188.28 | 198.71 | 213.13 | 207.37 |
| MgS   | 81.00      | 87.90  | 90.27  | 84.17  | 86.22  |
| Si    | 100.30     | 92.76  | 100.56 | 94.13  | 94.13  |
| SiC   | 228.90     | 263.89 | 241.68 | 263.73 | 263.71 |

Table S15: EOS fit diagnostics, part 1. All full-window minima are bracketed. RMS residuals are in meV per atom.

| Method | Solid | $N_V$ | $B'_0$ | RMS    |
|--------|-------|-------|--------|--------|
| AE     | AlN   | 9     | 3.707  | 0.0418 |
| AE     | AlP   | 9     | 4.179  | 0.0880 |
| AE     | BN    | 10    | 3.772  | 0.1250 |
| AE     | BP    | 10    | 3.701  | 0.0311 |
| AE     | C     | 10    | 3.756  | 0.0542 |
| AE     | LiCl  | 10    | 4.029  | 0.0205 |
| AE     | MgO   | 10    | 4.092  | 0.0190 |
| AE     | MgS   | 10    | 3.682  | 0.0117 |
| AE     | Si    | 10    | 4.544  | 0.3496 |
| AE     | SiC   | 10    | 3.854  | 0.0449 |
| GX     | AlN   | 9     | 2.802  | 0.6739 |
| GX     | AlP   | 9     | 3.959  | 0.1431 |
| GX     | BN    | 10    | 3.713  | 0.1306 |
| GX     | BP    | 10    | 4.081  | 0.0723 |
| GX     | C     | 10    | 3.611  | 0.1734 |
| GX     | LiCl  | 10    | 4.027  | 0.0435 |
| GX     | MgO   | 10    | 3.864  | 0.2921 |
| GX     | MgS   | 10    | 4.349  | 0.1597 |

Table S16: EOS fit diagnostics, part 2. All full-window minima are bracketed. RMS residuals are in meV per atom.

| Method | Solid | $N_V$ | $B'_0$ | RMS    |
|--------|-------|-------|--------|--------|
| GX     | Si    | 10    | 5.096  | 0.5702 |
| GX     | SiC   | 10    | 4.107  | 0.1309 |
| HD     | AlN   | 9     | 3.339  | 0.3028 |
| HD     | AlP   | 9     | 4.778  | 0.3362 |
| HD     | BP    | 10    | 3.918  | 0.0532 |
| HD     | LiCl  | 10    | 3.905  | 0.0330 |
| HD     | MgO   | 10    | 4.208  | 0.3562 |
| HD     | MgS   | 10    | 3.167  | 0.2585 |
| HD     | Si    | 10    | 5.257  | 0.3634 |
| HD     | SiC   | 10    | 4.168  | 0.0561 |
| HOC    | AlN   | 9     | 3.338  | 0.3027 |
| HOC    | AlP   | 9     | 3.787  | 0.2878 |
| HOC    | BP    | 10    | 3.918  | 0.0532 |
| HOC    | LiCl  | 10    | 3.915  | 0.0328 |
| HOC    | MgO   | 10    | 4.181  | 0.0393 |
| HOC    | MgS   | 10    | 3.865  | 0.0993 |
| HOC    | Si    | 10    | 5.257  | 0.3634 |
| HOC    | SiC   | 10    | 4.167  | 0.0557 |

Table S17: Matched comparison with Ref. 11: ten-solid set. Every row uses the identical intersection and the Goldzak static-lattice experimental reference.  $\text{MAE}(a_0)$  is in Å and  $\text{MAE}(B_0)$  in GPa. A dash denotes unavailable data, not zero error. Native rows retain the numerical qualifications discussed in the text.

| Method  | $N$ | $\text{MAE}(a_0)$ | $\text{MAE}(B_0)$ |
|---------|-----|-------------------|-------------------|
| GX      | 10  | 0.0831            | 16.61             |
| AE      | 10  | 0.0728            | 16.89             |
| HD      | 10  | 0.0844            | 21.77             |
| HOC     | 10  | 0.0844            | 21.07             |
| HF      | 10  | 0.0563            | 15.23             |
| MP2     | 10  | 0.0201            | 5.71              |
| SCS-MP2 | 10  | 0.0125            | 4.10              |
| SOS-MP2 | 10  | 0.0149            | 3.88              |

Table S18: Matched comparison with Ref. 12: nine solids, excluding AlN. Every row uses the identical intersection and the Goldzak static-lattice experimental reference.  $\text{MAE}(a_0)$  is in Å and  $\text{MAE}(B_0)$  in GPa. A dash denotes unavailable data, not zero error. Native rows retain the numerical qualifications discussed in the text.

| Method | $N$ | $\text{MAE}(a_0)$ | $\text{MAE}(B_0)$ |
|--------|-----|-------------------|-------------------|
| GX     | 9   | 0.0827            | 14.78             |
| AE     | 9   | 0.0737            | 16.56             |
| HD     | 9   | 0.0834            | 20.36             |
| HOC    | 9   | 0.0835            | 19.59             |
| LDA    | 9   | 0.0324            | 4.66              |
| PBE    | 9   | 0.0451            | 12.96             |
| PBEsol | 9   | 0.0113            | 4.72              |
| M06-L  | 9   | 0.0128            | 3.46              |
| SCAN   | 9   | 0.0123            | 3.08              |
| HSE06  | 9   | 0.0170            | 5.86              |

Table S19: Matched comparison with Ref. 13: nine solids, excluding MgS. Every row uses the identical intersection and the Goldzak static-lattice experimental reference.  $\text{MAE}(a_0)$  is in Å and  $\text{MAE}(B_0)$  in GPa. A dash denotes unavailable data, not zero error. Native rows retain the numerical qualifications discussed in the text.

| Method | $N$ | $\text{MAE}(a_0)$ | $\text{MAE}(B_0)$ |
|--------|-----|-------------------|-------------------|
| GX     | 9   | 0.0791            | –                 |
| AE     | 9   | 0.0660            | –                 |
| HD     | 9   | 0.0807            | –                 |
| HOC    | 9   | 0.0807            | –                 |
| PBE    | 9   | 0.0438            | –                 |
| PBEsol | 9   | 0.0136            | –                 |
| HSE06  | 9   | 0.0132            | –                 |
| HSEsol | 9   | 0.0117            | –                 |

Table S20: Matched comparison with Ref. 14: eight solids, excluding AlN and BN. Every row uses the identical intersection and the Goldzak static-lattice experimental reference.  $\text{MAE}(a_0)$  is in Å and  $\text{MAE}(B_0)$  in GPa. A dash denotes unavailable data, not zero error. Native rows retain the numerical qualifications discussed in the text.

| Method | $N$ | $\text{MAE}(a_0)$ | $\text{MAE}(B_0)$ |
|--------|-----|-------------------|-------------------|
| GX     | 8   | 0.0911            | 12.27             |
| AE     | 8   | 0.0797            | 14.58             |
| HD     | 8   | 0.0907            | 18.86             |
| HOC    | 8   | 0.0907            | 18.00             |
| Tao-Mo | 8   | 0.0194            | 4.26              |

Table S21: Matched comparison with Ref. 15: nine solids, excluding MgO. Every row uses the identical intersection and the Goldzak static-lattice experimental reference.  $\text{MAE}(a_0)$  is in Å and  $\text{MAE}(B_0)$  in GPa. A dash denotes unavailable data, not zero error. Native rows retain the numerical qualifications discussed in the text.

| Method   | $N$ | $\text{MAE}(a_0)$ | $\text{MAE}(B_0)$ |
|----------|-----|-------------------|-------------------|
| GX       | 9   | 0.0831            | —                 |
| AE       | 9   | 0.0703            | —                 |
| HD       | 9   | 0.0831            | —                 |
| HOC      | 9   | 0.0830            | —                 |
| GFN1-xTB | 9   | 0.1360            | —                 |
| GFN2-xTB | 9   | 0.0659            | —                 |

Table S22: Actual Gamma-centered k meshes by volume index. Indices 01–10 refer to the original evenly spaced volume ratios 0.90–1.10. Each selected volume uses the same mesh in all representations, but the mesh can change along an EOS.

| Solid | Mesh and volume indices                        |
|-------|------------------------------------------------|
| AlN   | $5^3$ : 01,02,03,04,05,06,07,08,09             |
| AlP   | $4^3$ : 01,02,03,04,05,06,07,08,09             |
| BN    | $6^3$ : 02,03,04,05,06,07,08,09,10; $7^3$ : 01 |
| BP    | $5^3$ : 01,02,03,04,05,06,07,08,09,10          |
| C     | $6^3$ : 04,05,06,07,08,09,10; $7^3$ : 01,02,03 |
| LiCl  | $5^3$ : 01,02,03,04,05,06,07,08,09,10          |
| MgO   | $5^3$ : 06,07,08,09,10; $6^3$ : 01,02,03,04,05 |
| MgS   | $4^3$ : 07,08,09,10; $5^3$ : 01,02,03,04,05,06 |
| Si    | $4^3$ : 02,03,04,05,06,07,08,09,10; $5^3$ : 01 |
| SiC   | $5^3$ : 01,02,03,04,05,06,07,08,09,10          |




Figure S1: Signed lattice-constant and bulk-modulus errors against the common Goldzak static-lattice reference. SCS-MP2 provides matched correlated-method context. Systematic native lattice contraction persists across representations. Its sign alone does not identify the source of the discrepancy.

## S8 Pressure Derivatives and Compressibilities

The third-order Birch–Murnaghan fit supplies  $V_0$ ,  $B_0$ , and  $B'_0 = (\partial B/\partial P)_{P=0}$  in addition to  $E_0$ . Its energy expression is

$$E(V) = E_0 + \frac{9V_0B_0}{16} [B'_0(x-1)^3 + (6-4x)(x-1)^2], \quad x = (V_0/V)^{2/3}. \quad (\text{S6})$$

The zero-pressure compressibility is  $\kappa_0 = 1/B_0$ . These quantities require no new SCF calculation.  $B'_0$  depends on the third derivative of the fitted energy and is more sensitive to the fit interval and numerical noise than the minimum location. We therefore supplement the full-window values with endpoint-omission sensitivity rather than assigning artificial high precision from the least-squares fit.

Table S23: Additional predictions from the existing EOS data.  $\kappa_0$  is in  $10^{-3} \text{ GPa}^{-1}$ .  $V_0$  is the eight-atom cell volume in  $\text{\AA}^3$ . The endpoint column is the largest absolute change of  $B'_0$  after removing either endpoint. A dagger means at least one omitted-window minimum is no longer bracketed.

| Method | Solid | $V_0$    | $B_0/\text{GPa}$ | $B'_0$ | $10^3\kappa_0$ | max. $ \Delta B'_0 $ |
|--------|-------|----------|------------------|--------|----------------|----------------------|
| AE     | AlN   | 79.6434  | 225.89           | 3.707  | 4.427          | 0.061                |
| AE     | AlP   | 155.5561 | 101.35           | 4.179  | 9.867          | 0.292                |
| AE     | BN    | 45.4045  | 420.87           | 3.772  | 2.376          | 0.106                |
| AE     | BP    | 89.2889  | 183.64           | 3.701  | 5.445          | 0.050                |
| AE     | C     | 43.9583  | 491.35           | 3.756  | 2.035          | 0.092                |
| AE     | LiCl  | 118.7693 | 46.37            | 4.029  | 21.568         | 0.073 <sup>†</sup>   |
| AE     | MgO   | 68.5923  | 198.71           | 4.092  | 5.032          | 0.049                |
| AE     | MgS   | 129.0645 | 90.27            | 3.682  | 11.077         | 0.031                |
| AE     | Si    | 155.2040 | 100.56           | 4.544  | 9.944          | 0.921                |
| AE     | SiC   | 79.0664  | 241.68           | 3.854  | 4.138          | 0.032                |
| GX     | AlN   | 78.4652  | 239.07           | 2.802  | 4.183          | 1.895                |
| GX     | AlP   | 154.1793 | 94.34            | 3.959  | 10.600         | 0.466                |
| GX     | BN    | 45.7944  | 423.39           | 3.713  | 2.362          | 0.129                |
| GX     | BP    | 87.7590  | 187.93           | 4.081  | 5.321          | 0.038                |
| GX     | C     | 44.1546  | 457.33           | 3.611  | 2.187          | 0.350                |
| GX     | LiCl  | 118.2330 | 47.98            | 4.027  | 20.844         | 0.230 <sup>†</sup>   |
| GX     | MgO   | 69.2214  | 188.28           | 3.864  | 5.311          | 0.625                |
| GX     | MgS   | 130.2384 | 87.90            | 4.349  | 11.377         | 0.264                |
| GX     | Si    | 152.1509 | 92.76            | 5.096  | 10.780         | 0.810                |
| GX     | SiC   | 76.8604  | 263.89           | 4.107  | 3.789          | 0.230                |
| HD     | AlN   | 78.1094  | 240.42           | 3.339  | 4.159          | 0.634                |
| HD     | AlP   | 154.1285 | 96.36            | 4.778  | 10.378         | 0.750                |
| HD     | BN    | 45.4045  | 420.87           | 3.772  | 2.376          | 0.106                |
| HD     | BP    | 88.2826  | 185.86           | 3.918  | 5.380          | 0.091                |
| HD     | C     | 43.9583  | 491.35           | 3.756  | 2.035          | 0.092                |
| HD     | LiCl  | 118.9380 | 47.13            | 3.905  | 21.217         | 0.212 <sup>†</sup>   |
| HD     | MgO   | 68.5343  | 213.13           | 4.208  | 4.692          | 0.337                |
| HD     | MgS   | 130.3228 | 84.17            | 3.167  | 11.880         | 0.631                |
| HD     | Si    | 152.1244 | 94.13            | 5.257  | 10.624         | 1.074                |
| HD     | SiC   | 77.0688  | 263.73           | 4.168  | 3.792          | 0.099                |
| HOC    | AlN   | 78.1082  | 240.40           | 3.338  | 4.160          | 0.634                |
| HOC    | AlP   | 154.1753 | 93.12            | 3.787  | 10.739         | 0.766                |
| HOC    | BN    | 45.4045  | 420.87           | 3.772  | 2.376          | 0.106                |
| HOC    | BP    | 88.2820  | 185.86           | 3.918  | 5.381          | 0.091                |
| HOC    | C     | 43.9583  | 491.35           | 3.756  | 2.035          | 0.092                |
| HOC    | LiCl  | 118.9405 | 47.16            | 3.915  | 21.203         | 0.240 <sup>†</sup>   |
| HOC    | MgO   | 68.5140  | 207.37           | 4.181  | 4.822          | 0.070                |
| HOC    | MgS   | 130.2985 | 86.22            | 3.865  | 11.598         | 0.224                |
| HOC    | Si    | 152.1238 | 94.13            | 5.257  | 10.624         | 1.074                |
| HOC    | SiC   | 77.0679  | 263.71           | 4.167  | 3.792          | 0.099                |

The full-window  $B'_0$  values span 2.80–5.26. The largest endpoint-induced change, 1.90, occurs for GAPW-XC AlN, whose full-window value is 2.80. This substantial sensitivity cautions against assigning physical significance to small differences between fitted pressure derivatives. The accompanying machine-readable analysis retains both endpoint tests and identifies cases in which an omitted endpoint no longer brackets the minimum.

The matched primary literature tables provide  $a_0$  and  $B_0$  but no consistently curated common  $B'_0$  reference set. We report  $B'_0$  as a derived model prediction with a fit-window diagnostic, not an accuracy ranking against an invented reference. Compressibility comparisons follow directly from the same matched  $B_0$  values. They contain no independent evidence beyond that observable. The three molecular crystals have only paired single-point energies, so no bulk modulus, pressure derivative, elastic tensor, or equilibrium volume is inferred for them.

## S9 Reproducibility and Numerical Limits

The molecular, molecular-crystal, and solid subsets use distinct documented protocols and are analysed separately. Each reported energy is linked to its executed input, model and executable versions, SCF convergence, and normal termination record. Replicate calculations count once. The archive distinguishes production energies from the independent numerical controls and preserves the input and source hashes needed to reproduce the reported quantities.

The ESI reports available convergence and sensitivity data rather than claiming a universal bound on basis, integration, or k-point error. In particular, the present work does not provide complete-basis extrapolation, BSSE correction for the molecular crystals, converged phonons, finite-temperature thermodynamics, or cohesive energies for the ten solids. Such quantities would require additional calculations and reference choices. The reported lattice-energy and structural trends remain useful, provided those distinctions are maintained.

## References

- [1] H. Mirhosseini, T. M. A. Müller, M. Krack, T. D. Kühne and J. Hutter, *J. Chem. Phys.*, 2026, **165**, 104103.
- [2] S. Goedecker, M. Teter and J. Hutter, *Phys. Rev. B*, 1996, **54**, 1703–1710.
- [3] T. Gould, *Phys. Chem. Chem. Phys.*, 2018, **20**, 27735–27739.
- [4] L. Goerigk, A. Hansen, C. Bauer, S. Ehrlich, A. Najibi and S. Grimme, *Phys. Chem. Chem. Phys.*, 2017, **19**, 32184–32215.
- [5] F. Pöschel, J. Pototschnig, F. Stein, A. Knüpfer, T. Vogels, S. Battaglia, S. Ehlert, J. Hutter and T. D. Kühne, arXiv:2608.19033; updated author manuscript, September 2026.
- [6] S. Battaglia, *Reaction-resolved molecular Skala validation data*, 2026, <https://github.com/DCM-Uni-Paderborn/Molecular-Skala-in-CP2K>, DietGMTKN55 validation dataset, repository revision 25e80baf68bee449d1dc2fd11cc359fa503272b8, accessed 11 September 2026.
- [7] F. Della Pia, A. Zen, D. Alfè and A. Michaelides, *Phys. Rev. Lett.*, 2024, **133**, 046401.
- [8] S. Grimme, J. Antony, S. Ehrlich and H. Krieg, *J. Chem. Phys.*, 2010, **132**, 154104.
- [9] S. Grimme, S. Ehrlich and L. Goerigk, *J. Comput. Chem.*, 2011, **32**, 1456–1465.
- [10] F. Della Pia, B. X. Shi, V. Kapil, A. Zen, D. Alfè and A. Michaelides, *Chemical Science*, 2025, **16**, 11419–11433.
- [11] T. Goldzak, X. Wang, H.-Z. Ye and T. C. Berkelbach, *J. Chem. Phys.*, 2022, **157**, 174112.
- [12] G.-X. Zhang, A. M. Reilly, A. Tkatchenko and M. Scheffler, *New J. Phys.*, 2018, **20**, 063020.
- [13] L. Schimka, J. Harl and G. Kresse, *J. Chem. Phys.*, 2011, **134**, 024116.
- [14] Y. Mo, R. Car, V. N. Staroverov, G. E. Scuseria and J. Tao, *Phys. Rev. B*, 2017, **95**, 035118.
- [15] V. Alizadeh, J. Pototschnig, L. M. Seidler, S. Ehlert and T. D. Kühne, Accepted in *J. Chem. Theory Comput.*, manuscript ct-2026-010347.
- [16] F. Tran, J. Stelzl and P. Blaha, *J. Chem. Phys.*, 2016, **144**, 204120.