




Regression & Mixed-Effects Modelling with R

Fang Jackson-Yang

13 May 2024

Course outline


- Session 1. Simple Regression; Individual Difference; Intro to Linear mixed-effect models (LMMs)
- Session 2. LMMs (lmer); Introduction to Generalised LMMs (glmer)
- Session 3. Generalised LMMs (glmer); Model assumptions and diagnostics



Session 3

Roadmap (today)

- Model Comparison & Selection
- Generalised LMMs
- Model Assumptions
- Exercise and Q&A





Model Comparison and Selection

Recall: Structure of LMMS

- *R* code:

```
lmer (y ~  $x_1 * x_2$  +  
      (  $1 + x_1 * x_2$  | Grouping1 ) +  
      (  $1 + x_1$  | Grouping2 ) ,  
      data = datafilename )
```

Recall: Model Fit

- No R^2
- Use Maximum likelihood Ratio test to compare models
- R code: `anova(model1, model2)`

```
anova(mMixed1_pval, mMixed_reduced)

## Data: vocabdata
## Models:
## mMixed1_pval: vocab_test_score ~ week * proficiency + (1 | participant)
## mMixed_reduced: vocab_test_score ~ week * proficiency + (1 + week | participant)
##           npar    AIC    BIC  logLik deviance  Chisq Df Pr(>Chisq)
## mMixed1_pval      6 2451.4 2474.8 -1219.7   2439.4
## mMixed_reduced    8 2416.2 2447.5 -1200.1   2400.2 39.151  2 3.151e-09 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Model Comparison & Selection

- In what order should you build your models?
- Which model should you select as final model?
 - **Barr, Levy, Scheepers, & Tily, (2013)** : keep it maximal
 - **Matuschek, Kliegl, Vasishth, Baayen & Bates (2017)** : make it parsimonious to balance Type1 error and power

Common Issues

- Convergence

```
Warning message:  
In checkConv(attr(opt, "derivs"), opt$par, ctrl = control$checkConv, :  
  Model failed to converge with max|grad| = 0.0206805 (tol = 0.002, component 1)
```

- Overfitting

```
boundary (singular) fit: see ?isSingular
```


Deal with Convergence Issue

Warning message:

```
In checkConv(attr(opt, "derivs"), opt$par, ctrl = control$checkConv, :  
  Model failed to converge with max|grad| = 0.0206805 (tol = 0.002, component 1)
```

Solutions:

- Adjust stopping (convergence) tolerances for the nonlinear optimizer, using the `optCtrl()` argument to `lmerControl`.
- Centre and standardise continuous predictor variables
 - the `scale()` function

Deal with Overfitting Issue

boundary (singular) fit: see `?isSingular`

Solutions:

- Remove the most complex part of the random effects structure (i.e. random slopes)
- Maybe acceptable to remove a specific random effect term when its variance estimates are very low



Generalised Linear Mixed-effects Models - `glmer()`

Binary Outcome

- Pass or Fail
- Invest or Not
- Vote 'Yes' or 'No'
- 'Correct' or 'incorrect' answer
- Fixation on the target image or not
- Align with partner or not
 - Lexical choice
 - Grammatical choice

'potato'
or
'tattie'



(a) *Gromit gave Wallace some cheese.*

(b) *Gromit gave some cheese to Wallace.*

Generalised Linear Mixed-effects Models

- `glmer()` for binary outcome



- *R* code:

```
glmer( y ~ x1 * x2 +  
        ( 1 + x1 * x2 | Grouping1 ) +  
        ( 1 + x1 | Grouping2 ) ,  
        data = datafilename,  
        family = 'binomial'  
        )
```

Example: Lexical Choice Study

‘Noob’ or ‘Beginner’ ?

- Internet slang data (simulated data)
- Hypothesis: People were more likely to use **internet slang words** (as **relative to alternative expressions**) after seeing their conversational partner uses a slang rather than a stand word.



Example: Lexical Choice Study

- Repeated measure (36 items/participant)
- One predictor (3 conditions):
‘prime’ = slang / standard words / other
- Two sources of random variance

	Participant	Item	List	Trial	Prime	Response
15	1	11	1	81	standard	0
16	1	11	1	109	other	0
17	1	12	1	8	slang	0
18	1	13	1	106	standard	0
19	1	14	1	25	other	0
20	1	15	1	1	slang	1
21	1	16	1	103	standard	0
22	1	17	1	38	slang	1
23	1	18	1	49	standard	0
24	1	18	1	70	other	0
25	1	19	1	42	other	0
26	1	21	1	46	slang	1
27	1	22	1	87	standard	0
28	1	23	1	18	other	0
29	1	24	1	120	slang	0
30	1	25	1	116	standard	0
31	1	26	1	78	slang	0

Showing 15 to 32 of 5,434 entries, 6 total columns

```
> summary(data_ISD)
```

Participant	Item	List	Trial	Prime	Response
1 : 36	18 : 248	6 : 457	Min. : 1.00	other :1787	0 :2874
2 : 36	14 : 234	1 : 410	1st Qu.: 31.00	standard:1833	1 :2512
5 : 36	28 : 215	5 : 401	Median : 63.00	slang :1814	NA's: 48
6 : 36	2 : 208	10 : 386	Mean : 62.45		
8 : 36	19 : 208	14 : 377	3rd Qu.: 95.00		
10 : 36	12 : 201	11 : 374	Max. :123.00		
(Other):5218	(Other):4120	(Other):3029			

Build a Glmer Model

```
Model <- glmer( response ~ prime +  
                ( 1 | participant ) +  
                ( 1 | item ) ,  
                data = dataISD,  
                family = 'binomial'  
                )
```


Interpret Coefficients

```
Generalized linear mixed model fit by maximum likelihood (Laplace Approximation) ['glmerMod']
Family: binomial (logit)
Formula: Response ~ Prime + (1 | Participant) + (1 | Item)
Data: data_ISD
Control: glmerControl(optimizer = "bobyqa", optCtrl = list(maxfun = 2e+05))

      AIC      BIC   logLik deviance df.resid
 5625.4   5658.3  -2807.7   5615.4     5381

Scaled residuals:
    Min       1Q   Median       3Q      Max
-4.9925 -0.5758 -0.2008  0.6024  4.8213

Random effects:
 Groups      Name      Variance Std.Dev.
Participant (Intercept) 2.4345   1.5603
Item        (Intercept) 0.1262   0.3552
Number of obs: 5386, groups: Participant, 160; Item, 30

Fixed effects:
              Estimate Std. Error z value Pr(>|z|)
(Intercept)  -0.27907    0.15230   -1.832   0.0669 .
Primestandar -0.93888    0.08594  -10.925 <2e-16 ***
Primeslang   1.12819    0.08583   13.145 <2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Correlation of Fixed Effects:
      (Intr) Prmstn
Primestndrd -0.266
Primeslang  -0.277  0.441
```

β_0 Probability of the intercept:

$$\frac{e^{-0.28}}{1 + e^{-0.28}} = 0.43$$

β_1 Probability of slope 1:

$$\frac{e^{(-0.28 + (-0.94))}}{1 + e^{(-0.28 + (-0.94))}} = 0.23$$

β_2 Probability of slope 2:

$$\frac{e^{(-0.28 + 1.13)}}{1 + e^{(-0.28 + 1.13)}} = 0.70$$

- Coefficients are in logit units (log-odds)
- Can be transformed to probabilities: $prob(x) = \frac{e^x}{1 + e^x}$




Model Assumptions

Model Assumptions: Linear regression

- Nature of the model
 - the relation between predictor and outcome has to be linear
- Nature of the errors (i.e., residuals)
 - normal and independent of each other

Model Assumptions: Simple regression `lm()`


- Required(LINE):
 - Linearity of relationships
 - Independence of residuals
 - Normality of residuals
 - Equal variances for residuals
- Desirable:
 - uncorrelated predictors
(no collinearity)
 - no outliers



Model Assumptions: Simple regression glm()

For binomial DVs, (logistic regression)

- Required:
 - LINEAR relationships between IVs and log-odds
 - ~~Normality of residuals~~
 - ~~homogeneity of variance~~
 - **Independence of residuals**
- Desirable:
 - uncorrelated predictors (no collinearity)
 - no “bad” (overly influential) observations
 - large samples (due to maximum likelihood fitting)







Model Assumptions: Mixed-effects Models

Similar to simple linear regressions model

- Error is random
- Residuals at multiple levels
 - Level1 residuals: mean = 0, variance constant (R code: `residual()`)
 - Level2+ residuals: mean = 0, variance constant (R code: `ranef()`)



Model Assumptions: Mixed Models – lmer()

```
plot_model(mMixed_reduced , type = "diag")
```



Model Assumptions: Mixed Models – glmer()

```
plot_model(m3_dataISD, type = "diag")
```






Exercise & Q&A

Exercise

The 'cheese data': Simulated data based on real psycholinguistic findings on structural priming.

- Two Predictors :
 - 2-level factor "Prime"
 - (a): Tom gave Spike some flowers.
 - (b): Tom gave some flowers to Spike.
 - 2-level factor "communication" (video- vs audio-call)
- Binary outcome: (a) or (b)?
 - (a) Gromit gave ... (Wallace some cheese)
 - (b) Gromit gave ... (some cheese to Wallace).



Exercise



Can you investigate the priming effect in the 'cheese data'?

- *R* code:


```
glmer( y ~ x1 * x2 +  
        ( 1 + x1 * x2 | Grouping1 ) +  
        ( 1 + x1 | Grouping2 ) ,  
        data = datafilename,  
        family = 'binomial'  
        )
```



Further Reading

- Paper


Brown, VA. (2021). An Introduction to Linear Mixed-Effects Modeling in R. *Advances in Methods and Practices in Psychological Science*. 4(1). doi:10.1177/2515245920960351

Advances in Methods and Practices in Psychological Science

 ASSOCIATION FOR PSYCHOLOGICAL SCIENCE

 Open access |  | Research article | First published online March 25, 2021

An Introduction to Linear Mixed-Effects Modeling in R

[Violet A. Brown](#)   [View all authors and affiliations](#)

[All Articles](#) | <https://doi.org/10.1177/2515245920960351>

- E-Book

https://vasishth.github.io/Freq_CogSci/

Shravan Vasishth, Daniel Schad, Audrey Bürki, Reinhold Kliegl

***Linear Mixed Models in
Linguistics and Psychology:
A Comprehensive
Introduction***

THANK YOU



THE UNIVERSITY of EDINBURGH
Centre for Data, Culture & Society