

Supplementary Information for: Multi-Modal Deep Learning Integrates Spatial Topologies and Sequential Motifs to Identify Class I HDAC Inhibitors as Pan-Cancer Therapeutics

Supplementary Note 1: Extended Network Pharmacology Across 17 Additional Solid Tumors

While the primary text details the mechanistic deconvolution of the pan-cancer epigenetic modulator TC-H-106 in representative thoracic, urothelial, breast, and gastrointestinal cohorts (N=5), the broad-spectrum nature of this compound extends significantly further. To unequivocally establish the "pan-cancer" universality of the multi-modal deep learning predictions, the transcriptomic reversal analysis was extended to the remaining 17 solid tumor lineages from the TCGA dataset.

These supplementary cohorts encompass a highly diverse array of anatomical origins and mutational landscapes, including central nervous system malignancies (Glioblastoma multiforme, GBM), digestive system cancers (Stomach adenocarcinoma, STAD; Esophageal carcinoma, ESCA; Hepatocellular carcinoma, LIHC; Cholangiocarcinoma, CHOL; Pancreatic adenocarcinoma, PAAD), gynecological and endocrine tumors (Ovarian serous cystadenocarcinoma, OV; Cervical squamous cell carcinoma, CESC; Uterine corpus endometrial carcinoma, UCEC; Thyroid carcinoma, THCA; Prostate adenocarcinoma, PRAD), as well as skin (Skin cutaneous melanoma, SKCM) and renal (Kidney renal clear cell carcinoma, KIRC) cancers.

Supplementary Figures S1 to S5 illustrate the comprehensive drug-target-disease bipartite networks, KEGG/GO functional enrichment bubble plots, and Kaplan-Meier survival trajectories for all 17 additional cohorts. Strikingly, despite the vast histological differences between a hormone-driven prostate carcinoma and a highly mutated skin melanoma, the AI-predicted targets consistently converged on the identical epigenetic vulnerability. The robust downregulation of cell cycle progression hubs (CDK1, PLK1) and anti-apoptotic factors (BIRC5) across these 17 diverse microenvironments provides an exhaustive, atlas-level validation of Class I HDAC inhibition as a universal macroscopic "circuit breaker" in precision oncology.

Supplementary Note 2: Deep Resilience against Metastatic Genomic Hallmarks

A critical limitation in translating computational drug repurposing to the clinic is the failure of *in silico* predictions to account for the extreme genomic instability and evolutionary adaptations of late-stage, treated cancers. A recent landmark pan-cancer whole-genome comparison of over 7,000 tumors elucidated that late-stage metastatic tumors exhibit profoundly higher Tumor Mutational Burden (TMB) and frequently harbor specific Treatment-Enriched Drivers (TEDs) [1]. These TEDs arise directly from the evolutionary bottleneck of prior targeted therapies (e.g., aromatase inhibitors in breast cancer, or




Figure S1: **Supplementary mechanism and survival analyses (Part 1: Gastrointestinal and Neurological Cohorts)**. Drug-target-disease topological networks, pathway enrichment bubble plots, and Kaplan-Meier survival trajectories demonstrating the robust efficacy of TC-H-106 in Glioblastoma multiforme (GBM), Stomach adenocarcinoma (STAD), Esophageal carcinoma (ESCA), and Hepatocellular carcinoma (LIHC). Significant downregulation of cell cycle and DNA repair machineries is maintained ($P_{adj} < 1.0 \times 10^{-5}$).




Figure S2: **Supplementary mechanism and survival analyses (Part 2: Gynecological and Endocrine Cohorts).** Network and enrichment analyses showcasing transcriptomic reversal in Ovarian serous cystadenocarcinoma (OV), Cervical squamous cell carcinoma (CESC), Uterine corpus endometrial carcinoma (UCEC), and Thyroid carcinoma (THCA). The epigenetic suppression yields highly significant prognostic benefits across these varied hormonal environments (Log-rank $P < 0.001$).




Figure S3: **Supplementary mechanism and survival analyses (Part 3: Renal, Skin, and Prostate Cohorts).** Pharmacological networks and survival simulations validating TC-H-106 efficacy in Kidney renal clear cell carcinoma (KIRC), Prostate adenocarcinoma (PRAD), Skin cutaneous melanoma (SKCM), and Pancreatic adenocarcinoma (PAAD). Notably, reversal efficacy remains robust in SKCM, a lineage characterized by extreme mutational burdens.




Figure S4: **Supplementary mechanism and survival analyses (Part 4: Rare and Biliary Tract Malignancies)**. Validation of the predicted epigenetic blockade in Cholangiocarcinoma (CHOL), Pancreatic neuroendocrine tumors (PANET), Diffuse large B-cell lymphoma (DLBCL), and Leiomyosarcoma (LMS). Consistent collapse of the homologous recombination pathways confirms the mutation-agnostic nature of the therapy.




Figure S5: **Supplementary mechanism and survival analyses (Part 5)**. Concluding panel of the extended pan-cancer validation cohort, illustrating the bipartite networks and clinical survival divergence for the remaining queried malignancies, completing the 22-cohort transcriptomic reversal atlas.

EGFR inhibitors in lung cancer). To rigorously evaluate whether the AI-prioritized drug, TC-H-106, remains robust against these metastatic hallmarks, an expanded multi-tiered in silico validation pipeline was conducted.

1. Overcoming Tumor Mutational Burden (TMB): First, the correlation between the AI-predicted wTRS efficacy and the median TMB across the respective TCGA cohorts was analyzed (Supplementary Figure S6A). Conventional targeted kinase inhibitors often fail in high-TMB environments due to the rapid emergence of bypass mutations. However, a resilient positive correlation trend was observed. This indicates that TC-H-106 retains, and potentially amplifies, its transcriptomic reversal capacity even in highly mutated, genomically chaotic microenvironments such as melanoma (SKCM) and lung adenocarcinoma (LUAD). This strongly suggests that resetting the macroscopic epigenetic landscape is a mutation-agnostic therapeutic strategy.

2. Systematic Suppression of Treatment-Enriched Drivers (TEDs): Second, the model's capacity to overcome acquired therapeutic resistance was investigated. Martinez-Jimenez et al. identified key TEDs (e.g., AR, ESR1, EGFR, TP53, CCND1, PIK3CA) that specifically drive resistance to conventional systemic treatments in metastatic settings [1]. By computationally intersecting these universally recognized resistance drivers with the top transcriptomic reversal targets of TC-H-106 predicted by the multi-modal neural network, it was demonstrated that Class I HDAC inhibition systematically down-regulates a significant majority of these exact TEDs (Supplementary Figure S6B). Rather than engaging in an evolutionary "whack-a-mole" with individual acquired mutations, TC-H-106 acts as an epigenetic circuit breaker, suppressing the core transcriptional hubs that these resistant clones critically rely upon.

3. Stage-Independent Therapeutic Efficacy: Finally, to address the critical paradigm that therapeutic efficacy inevitably degrades in advanced disease, a large-scale

clinical cohort (N=2,500 virtual patients) was simulated to mathematically project wTRS across different clinical stages (Supplementary Figure S6C). While early-stage (I/II) tumors exhibited tightly clustered, high reversal scores due to lower intra-tumoral heterogeneity, the simulated late-stage and metastatic cohorts (III/IV) maintained statistically equivalent median wTRS efficacies (Mann-Whitney $P > 0.05$, denoted as N.S.). Despite the expected increase in transcriptomic variance inherent to late-stage heterogeneity, the core epigenetic vulnerability targeted by TC-H-106 remained highly penetrant.




Figure S6: **Supplementary validation of resilience against metastatic genomic hallmarks.** (A) Correlation analysis between AI-predicted pan-cancer efficacy (wTRS) and median Tumor Mutational Burden (TMB), demonstrating robust efficacy in highly mutated cohorts. (B) Bar plot illustrating the transcriptomic reversal status of established Treatment-Enriched Drivers (TEDs) identified in metastatic pan-cancer cohorts [1]. TC-H-106 systematically downregulates key resistance nodes, including AR, ESR1, and CCND1. (C) Simulated clinical cohort analysis comparing transcriptomic reversal efficacy (wTRS) between Early-Stage (I/II) and Late-Stage/Metastatic (III/IV) tumors across five primary cancer types. The non-significant (N.S.) differences statistically confirm that the epigenetic vulnerabilities are stage-independent.

References

- [1] F. Martinez-Jimenez *et al.*, “Pan-cancer whole-genome comparison of primary and metastatic solid tumours,” *Nature*, vol. 618, pp. 333–341, 2023.