



MENTICATION

ELITE INTERVENTION LIBRARY

17 approved flagship pathways - evidence, experience systems and production direction

PERSONALISATION, MEMORY AND PRIVACY CONTRACT

1

Remember only with a clear benefit

Session text stays temporary unless the user deliberately saves it.

2

Four memory tiers

Session state, user-saved artefacts, confirmed preferences and optional pattern summaries.

3

No hidden psychological inference

No diagnoses, nervous-system claims, risk scores, emotion labels or personality conclusions from behaviour.

4

Visible and correctable

Every suggestion can explain its source; saved items and patterns can be edited, rejected or deleted.

5

Consent at handoff

The user sees what will carry over, why it helps and can remove fields before transfer.

6

Evidence thresholds

One observation is a fact, three similar outcomes may form a tentative pattern, five may support a stronger descriptive pattern.

BOX BREATHING

CALM / RHYTHMIC PACING

1 PURPOSE

A paced breathing experience for moments of physical activation, tension or scattered attention. It gives the user a predictable rhythm and a steady attentional anchor without requiring them to force calm.

PSYCHOEDUCATION

Slow, structured breathing can support autonomic regulation and increase heart-rate variability. Evidence is stronger for slow-paced breathing generally than for the exact 4-4-4-4 pattern.

2 USER FLOW

- 1 Arrive and check breathing comfort
- 2 Choose standard or no-hold route
- 3 Follow the expanding field
- 4 Adapt pace, guidance or holds
- 5 Notice breathing comfort
- 6 Finish or hand off to grounding

4 VISUAL AND AUDIO DIRECTION

Deep emerald volumetric glass square; expansion travels around four edges with no sudden flashes. Quiet low-frequency bed, soft directional cues and warm optional narration; static edge highlighting for reduced motion.

3 MENTICATION UPGRADE

The breathing object is both pacer and progress cue. Holds can be removed without restarting, guidance changes in place, and success is breathing comfort or usable attention rather than a required feeling of calm.

ELITE STANDARD

One mechanism. One memorable interaction. Honest outcomes, accessible alternatives and a consented route onwards.

SIGNATURE VISUAL



EVIDENCE REGISTER

A for slow-paced breathing; C for the exact branded pattern

Do not imply that the exact four-part ratio is superior. Stop or remove holds if dizziness, air hunger or respiratory discomfort occurs.

Reviewed 6 Sep 2026 | Next review Sep 2027

PROGRESSIVE MUSCLE RELAXATION

CALM / INTERACTIVE BODY MAPPING

1 PURPOSE

A body-led experience for prominent muscular tension when safe tensing or release-only attention is possible. It helps the user notice the contrast between tension and release and choose what their body can comfortably soften.

PSYCHOEDUCATION

Progressive muscle relaxation has evidence for reducing stress and anxiety in adults. The useful mechanism is learning the contrast between tension and release, not forcing every muscle to relax.

2 USER FLOW

- 1 Check pain, injury and movement comfort
- 2 Choose Immediate, Standard, Supported or release-only
- 3 Select or follow body region
- 4 Tense gently if suitable
- 5 Release and compare
- 6 Whole-body soften
- 7 Notice remaining tension and finish

4 VISUAL AND AUDIO DIRECTION

Neutral inclusive body silhouette with user-selectable regions; warm points gather during tension and dissolve on release. Slow low pulse, breath-safe cues and precise narration; list and tap alternatives replace the body map when needed.


3 MENTICATION UPGRADE

Suitability is checked before tensing, every region can be skipped, release-only remains a full pathway, and the outcome records where tension changed rather than demanding total relaxation.

ELITE STANDARD

One mechanism. One memorable interaction. Honest outcomes, accessible alternatives and a consented route onwards.

SIGNATURE VISUAL



EVIDENCE REGISTER

A for PMR generally; C for this interaction system

Evidence covers varied PMR protocols and populations. Avoid painful or injured areas and use release-only when tensing is unsuitable.

Reviewed 6 Sep 2026 | Next review Sep 2027

5-4-3-2-1 GROUNDING

GROUND / SEQUENTIAL DISCOVERY

1 PURPOSE

An eyes-open sensory orientation experience for overwhelm, disconnection or narrowed attention. It rebuilds contact with the immediate environment one sense at a time without asking the user to analyse why they feel this way.

PSYCHOEDUCATION

Grounding uses present sensory information to redirect attention toward the here and now. The exact 5-4-3-2-1 sequence is widely used, but direct efficacy research on that specific sequence remains limited.

2 USER FLOW

- 1 Check immediate safety and orient eyes open
- 2 See five
- 3 Feel four
- 4 Hear three
- 5 Smell two
- 6 Taste one or use a neutral alternative
- 7 Notice reconnection
- 8 Finish or reroute attention

4 VISUAL AND AUDIO DIRECTION

A quiet world progressively rebuilds through light, ripple, wave, wisp and droplet layers. Each sense has one tactile sound identity; narration is optional and every gesture has a large tap target.

3 MENTICATION UPGRADE

Discovery replaces a checklist: the environment visibly returns as senses are named. Counts can be shortened, inaccessible senses skipped, and success is orientation rather than symptom disappearance.


ELITE STANDARD

One mechanism. One memorable interaction. Honest outcomes, accessible alternatives and a consented route onwards.

SIGNATURE VISUAL

ROUNDING — PRODUCTION VISUAL SYSTEM

FIGURE • FIVE SENSORY TRANSFORMATIONS • GROUND



EVIDENCE REGISTER

C: mechanism-informed; exact sequence under-studied

The review notes that grounding has lacked agreed definitions and efficacy studies. Never present this as proven treatment for dissociation or trauma.

Reviewed 6 Sep 2026 | Next review Sep 2027

THOUGHT OR FACT

CALM / COGNITIVE SORTING

1 PURPOSE

A private evidence review for distressing thoughts that have begun to feel certain. It helps the user separate observable facts from interpretation, prediction and catastrophe while keeping the feeling itself valid.

PSYCHOEDUCATION

Cognitive approaches can reduce anxiety and depression by helping people examine interpretations and generate more balanced alternatives. This is a brief sorting tool, not a diagnostic test or a verdict on whether a person is right.

2 USER FLOW

- 1 Check that reflective work fits
- 2 Capture one thought
- 3 Separate fact, interpretation, prediction or catastrophe
- 4 Review evidence and uncertainty
- 5 Write a balanced statement
- 6 Choose next action or safe experiment
- 7 Finish without a verdict

4 VISUAL AND AUDIO DIRECTION

A softened private evidence chamber: dark amber glass, movable evidence cards and one illuminated clarity line. Sparse paper movement and optional measured narration; no courtroom verdict imagery.


3 MENTICATION UPGRADE

The experience makes uncertainty visible, prevents forced positive thinking and offers Test the Prediction only when a real-world question is safely testable.

ELITE STANDARD

One mechanism. One memorable interaction. Honest outcomes, accessible alternatives and a consented route onwards.

SIGNATURE VISUAL



EVIDENCE REGISTER

B for CBT/cognitive change; **C** for this brief standalone sorter

Do not use to challenge trauma memories, abuse disclosures, delusional or paranoid beliefs, or immediate danger. Evidence is for broader CBT packages, not this screen sequence alone.

URGE SURFING

CALM / EMOTIONAL WAVE TRACKING

1 PURPOSE

A choice-window experience for a strong urge that can be observed safely without immediate action. It helps the user notice that urge intensity and behaviour are separate, creating room for a deliberate next choice.

PSYCHOEDUCATION

Mindfulness practice may weaken the link between craving and action even when the urge does not immediately disappear. The goal is a more available choice, not a promise that every urge will pass within a fixed time.

2 USER FLOW

- 1 Check danger and leave-trigger options
- 2 Name the urge without storing detail
- 3 Choose a 60-second, 90-second or supported window
- 4 Locate and trace the wave
- 5 Pause action while sensations change
- 6 Notice available choice
- 7 Finish, leave the trigger or seek support

4 VISUAL AND AUDIO DIRECTION

A dark oceanic wave responds to a simple intensity trace; the line remains honest when flat or rising. Low tide texture and optional sparse voice; static trace and automatic pacing are equivalent routes.


3 MENTICATION UPGRADE

Safety precedes immersion, duration is a user-selected pause rather than biology, and unchanged or stronger urges remain valid outcomes. The standout measure is choice availability.

ELITE STANDARD

One mechanism. One memorable interaction. Honest outcomes, accessible alternatives and a consented route onwards.

SIGNATURE VISUAL



EVIDENCE REGISTER

B for mindfulness and craving-response decoupling; **C** for this brief wave interaction

Results for brief app-based mindfulness are mixed. This is not suitable when an urge involves immediate danger, overdose risk, self-harm or harm to others.

Reviewed 6 Sep 2026 | Next review Sep 2027

THEN WHAT?

CALM / COGNITIVE SORTING

1 PURPOSE

A bounded coping-appraisal pathway for a feared future frozen at its worst frame. It continues the timeline far enough to identify coping, support and what remains intact without promising that the feared event is harmless or unlikely.

PSYCHOEDUCATION

Cognitive change and perspective-taking can reduce emotional impact by changing how a situation is understood. This pathway tests coping beyond the worst frame; it is not reassurance and does not establish what will happen.

2 USER FLOW

- 1 Check danger, activation and reassurance looping
- 2 Name the frozen frame briefly
- 3 Name what it has come to mean
- 4 Map first move
- 5 Map first person and first resource
- 6 Name what remains intact
- 7 Decide whether action is needed now
- 8 Save the coping route and stop

4 VISUAL AND AUDIO DIRECTION

A cinematic strip begins as one frozen frame and opens into five bounded illuminated nodes. Each node appears once with a soft click; no endless zoom, branching catastrophe or dramatic crescendo.

3 MENTICATION UPGRADE

The system detects when analysis is becoming reassurance-seeking and routes to Grounding or Urge Surfing. It stores a coping route, never a catastrophe rehearsal.

ELITE STANDARD

One mechanism. One memorable interaction. Honest outcomes, accessible alternatives and a consented route onwards.

SIGNATURE VISUAL



EVIDENCE REGISTER

B for cognitive change/perspective-taking; **C** for this bounded standalone sequence

Not for immediate danger, acute panic, trauma re-experiencing, impaired reality testing or repetitive certainty-seeking. Broader CBT evidence does not prove the exact sequence.

Reviewed 6 Sep 2026 | Next review Sep 2027

IGNITION POINT

LIFT / PERSONALISED CHOICE PATHWAY

1 PURPOSE

A low-pressure activation pathway for low reward expectancy, flatness or reduced motivation. It helps the user choose one reachable source of Pleasure, Mastery, Connection or Dream and make contact with it before waiting to feel motivated.

PSYCHOEDUCATION

Behavioural activation works by increasing meaningful contact between a person and their environment. A small action can be worthwhile before mood changes, so success is engagement with value or reward rather than instant happiness.

2 USER FLOW

1 Check whether rest or safety is the real need

2 Choose Pleasure, Mastery, Connection or Dream

3 Select a reachable possibility

4 Scale it to current capacity

5 Begin the first move

6 Return when ready

7 Notice whether it felt worthwhile

8 Save only with consent

4 VISUAL AND AUDIO DIRECTION

Four premium energy constellations orbit a small ignition core, each with a distinct material and gentle pulse. Warm rise without triumphal music; optional narration uses invitational language.

3 MENTICATION UPGRADE

The permanent Dream pathway protects future-directed meaning alongside immediate reward. The system learns only user-confirmed reachability and worthwhileness, not hidden mood labels.

ELITE STANDARD

One mechanism. One memorable interaction. Honest outcomes, accessible alternatives and a consented route onwards.

SIGNATURE VISUAL



EVIDENCE REGISTER

A for behavioural activation packages; **C** for the four-pathway branded selector

Evidence is mainly from depression treatment protocols delivered over time. One micro-action should not be described as treatment or guaranteed mood improvement.

Reviewed 6 Sep 2026 | Next review Sep 2027

CHANGE THE SCENE

LIFT / SPATIAL ATTENTION MAPPING

1 PURPOSE

An environmental interruption for flatness or inertia that is being reinforced by the current setting. It helps the user make one safe change of position, view or place so a different next behaviour becomes available.

PSYCHOEDUCATION

Stable contexts can cue automatic behaviour, while a context change can interrupt those cues and create a window for a different choice. Evidence supports the broader context-discontinuity principle, not this exact short intervention.

2 USER FLOW

- 1 Check that movement and the destination are safe
- 2 Choose turn, doorway, room or outside
- 3 Preview the smallest transition
- 4 Cross into the new scene
- 5 Orient to one difference
- 6 Notice what behaviour is now available
- 7 Finish or continue with an action pathway

4 VISUAL AND AUDIO DIRECTION

One physical portal cuts through a muted scene; parallax and light temperature change only after the transition. A single threshold sound and environmental bed replace motivational music.


3 MENTICATION UPGRADE

The app checks whether a scene change is realistic and does not claim the new place must improve mood. The system remembers useful scene types only after repeated user confirmation.

ELITE STANDARD

One mechanism. One memorable interaction. Honest outcomes, accessible alternatives and a consented route onwards.

SIGNATURE VISUAL



EVIDENCE REGISTER

C: mechanism-adjacent context research

The cited context-discontinuity evidence concerns longer-term habit contexts, not momentary mood shifts. Keep claims modest and never direct users into unsafe environments.

Reviewed 6 Sep 2026 | Next review Sep 2027

TEST THE PREDICTION

LIFT / COGNITIVE SORTING

1 PURPOSE

A real-world behavioural experiment for a specific negative prediction that can be tested safely, ethically and proportionately. It replaces imagined certainty with direct observation while allowing confirming, mixed and unanswered results.

PSYCHOEDUCATION

Behavioural experiments can help people compare a prediction with observable experience. Evidence suggests benefit within anxiety treatment, but results must be read fairly and should never be designed to prove the user wrong.

2 USER FLOW

- 1 Capture a specific prediction
- 2 Run the eligibility and safety screen
- 3 Define a small fair action
- 4 Define an observable outcome
- 5 Preview and edit the experiment
- 6 Leave the app and test
- 7 Record confirming, disconfirming, mixed or unanswered evidence
- 8 Update the prediction or stop

4 VISUAL AND AUDIO DIRECTION

Two balanced glass panels hold Prediction and Observation with a neutral bridge between them. Evidence particles move without winning colours, fanfare or scales that imply disconfirmation is the correct result.


3 MENTICATION UPGRADE

Eligibility blocks medical, legal, dangerous, coercive and reality-testing use. All result types have equal visual dignity and can hand back to Thought or Fact for fair interpretation.

ELITE STANDARD

One mechanism. One memorable interaction. Honest outcomes, accessible alternatives and a consented route onwards.

SIGNATURE VISUAL



EVIDENCE REGISTER

B for behavioural experiments in anxiety treatment; **C** for this standalone digital path

The review found methodological limitations and studied behavioural experiments inside treatment. Never create high-stakes, unsafe or boundary-violating tests.

Reviewed 6 Sep 2026 | Next review Sep 2027

OPEN CHANNEL

LIFT / LANGUAGE TRANSFORMATION

1 PURPOSE

A relational re-entry experience after safe connection has been interrupted by silence, withdrawal, cancellation or shame. It helps the user choose and review one honest bridge without requiring a reply or accessing contacts.

PSYCHOEDUCATION

Interventions that increase access to people or address barriers to connection can improve social contact and perceived connection. One message is only a bridge, not a guaranteed repair or cure for loneliness.

2 USER FLOW

- 1 Name where the channel closed
- 2 Name the barrier
- 3 Check safety, consent and no-contact boundaries
- 4 Choose Signal, Reopen, Repair, Ask or Presence
- 5 Draft in the user's own voice
- 6 Review every word
- 7 Complete outside the app or deliberately decline
- 8 Record the decision, not the other person's response

4 VISUAL AND AUDIO DIRECTION

Two quiet points separated by fractured space; a bridge forms segment by segment only after the user approves language. Restrained connection tone, no message-send sound unless the app actually sends something.

3 MENTICATION UPGRADE

The system separates connection from contact-at-any-cost, makes not contacting a valid outcome, and never reads contacts, sends messages or fabricates a response.

ELITE STANDARD

One mechanism. One memorable interaction. Honest outcomes, accessible alternatives and a consented route onwards.

SIGNATURE VISUAL



EVIDENCE REGISTER

B for broader social-connection interventions; **C** for a single digital re-entry prompt

Intervention effects vary and much evidence concerns multi-session programmes. Do not use for abuse, coercion, unsafe former partners or when no contact has been requested.

Reviewed 6 Sep 2026 | Next review Sep 2027

COUNTERMOVE

LIFT / ACTION INITIATION AND TIMING

1 PURPOSE

A graded opposite-direction experience for withdrawal or emotion-driven avoidance when the pull is neither protective nor restorative. It creates the smallest behaviour that widens life while allowing the feeling to remain present.

PSYCHOEDUCATION

Behavioural activation increases meaningful interaction with the environment, and specific action plans can help close the gap between intention and behaviour. A small directional move can count before mood changes.

2 USER FLOW

- 1 Name the pull
- 2 Check whether it protects, restores or avoids
- 3 Choose a 5-degree, 20-degree or 45-degree trajectory
- 4 Generate and edit one countermove
- 5 Make it lighter or change direction
- 6 Act outside the app
- 7 Notice whether the situation widened
- 8 Finish or hand off

4 VISUAL AND AUDIO DIRECTION

A matte sphere sits in a soft gravitational field; one selected trajectory bends by the chosen degree. Low hum, directional tone and subtle haptic confirmation; reduced motion shows a static before-and-after line.


3 MENTICATION UPGRADE

Function is checked before action so safety and genuine rest are never mislabeled as avoidance. The trajectory engine scales one mechanism without ranking stronger moves as better.

ELITE STANDARD

One mechanism. One memorable interaction. Honest outcomes, accessible alternatives and a consented route onwards.

SIGNATURE VISUAL



EVIDENCE REGISTER

B for behavioural activation and implementation intentions; **C** for the graded trajectory interface

Evidence comes from broader interventions, not the 5/20/45-degree metaphor. Do not oppose behaviour that is protective, restorative or required for safety.

Reviewed 6 Sep 2026 | Next review Sep 2027

PULSE SHIFT

LIFT / INTERACTIVE BODY MAPPING

1 PURPOSE

A capacity-matched movement experience for flatness, heaviness, fogginess or physical inertia. It creates a small amount of usable activation and then directs it toward a real-life action without becoming exercise coaching.

PSYCHOEDUCATION

A single bout of exercise is associated on average with modest improvements in mood and anxiety, but responses vary widely. Here the aim is simply to make a little more movement available, not to prescribe intensity or promise a mood lift.

2 USER FLOW

- 1 Check pain, dizziness, instability and genuine need for rest
- 2 Choose lying, seated, standing or already moving
- 3 Choose Flicker, Pulse or Surge
- 4 Unlock movement
- 5 Build a short rhythm
- 6 Direct it toward one real action
- 7 Notice available movement
- 8 Stop, rest or hand off

4 VISUAL AND AUDIO DIRECTION

A near-static body field receives one travelling pulse that grows only to the selected capacity. Percussive texture stays soft and tempo-stable; captions, seated alternatives and tap pacing are native.

3 MENTICATION UPGRADE

The pathway starts from position and capacity rather than a fitness norm. Rest, no change and deliberate stopping are valid outcomes, and concerning symptoms route away immediately.

ELITE STANDARD

One mechanism. One memorable interaction. Honest outcomes, accessible alternatives and a consented route onwards.

SIGNATURE VISUAL



EVIDENCE REGISTER

B for acute exercise affect; C for micro-dose graded activation

The meta-analysis found substantial heterogeneity and typically studied exercise bouts, not tiny movements. This is not medical or exercise advice.

Reviewed 6 Sep 2026 | Next review Sep 2027

1 PURPOSE

A brief orientation and attentional-redirection experience for genuine distress when a longer grounding exercise is unlikely to be usable. It gives attention one safe, absorbing destination while preserving the option to return to the problem later.

PSYCHOEDUCATION

Distraction can reduce emotional impact in some contexts, but its usefulness depends on the emotion, timing and purpose. Reroute is a deliberate short-term shift of attention, not suppression or a claim that the problem no longer matters.

2 USER FLOW

- 1 Check immediate danger and urgent practical action
- 2 Orient to one stable point
- 3 Choose Social, Sensory, Familiar, Active or Absorbing destination
- 4 Select a reachable route
- 5 Take the first action
- 6 Leave the app
- 7 Notice whether attention entered
- 8 Return, repeat once or choose another mechanism

4 VISUAL AND AUDIO DIRECTION

Visual interference clears around one stable point, then a single route opens through the field. Directional sound follows the route; reduced-motion mode swaps travelling lines for progressive contrast.

3 MENTICATION UPGRADE

The tool distinguishes deliberate redirection from avoidance, asks whether attention actually entered, and hands to Next Easiest Step when the destination fits but the first action is unclear.

ELITE STANDARD

One mechanism. One memorable interaction. Honest outcomes, accessible alternatives and a consented route onwards.

SIGNATURE VISUAL



EVIDENCE REGISTER

C: mechanism-informed attentional deployment

The meta-analysis found strategy and context differences; attentional deployment was not uniformly effective. Do not use when urgent safety or practical action is required.

NEXT EASIEST STEP

FOCUS / ACTION INITIATION AND TIMING

1 PURPOSE

An executive-friction reducer for tasks that feel unclear, effortful, distracting or too large. It converts a general intention into one observable physical action that can begin now or be taken outside the app.

PSYCHOEDUCATION

Specific action plans can improve goal attainment by linking intention to a concrete response. The useful target is a visible next behaviour, not a perfect plan or a judgment about motivation.

2 USER FLOW

- 1 Name the task
- 2 Identify the barrier
- 3 Compress to one observable action
- 4 Check that it can begin quickly
- 5 Edit or make easier
- 6 Take the step outside the app
- 7 Record complete, partial or could not start
- 8 Hand off to Ignition Point or Signal Lock if useful

4 VISUAL AND AUDIO DIRECTION

A dense cloud of task fragments compresses into one bright physical tile. Soft mechanical clicks mark simplification; no ticking clock unless the user chooses a timed handoff.


3 MENTICATION UPGRADE

Barrier-specific compression avoids generic advice. Could not start becomes information that either reduces the step again or changes mechanism rather than a failure state.

ELITE STANDARD

One mechanism. One memorable interaction. Honest outcomes, accessible alternatives and a consented route onwards.

SIGNATURE VISUAL



EVIDENCE REGISTER

B for implementation intentions; C for task-compression micro-intervention

If-then planning evidence does not establish an optimal 20-second start rule. Keep the action user-editable and do not score productivity.

Reviewed 6 Sep 2026 | Next review Sep 2027

SIGNAL LOCK

FOCUS / ACTION INITIATION AND TIMING

1 PURPOSE

A task-scaffolding and bounded sprint system for a chosen task that is startable but vulnerable to interruption. It clarifies the target, contains predictable distractions and supports deliberate focus without pretending to control the device.

PSYCHOEDUCATION

Reducing interruption costs and using deliberate time-management strategies can support task accuracy, resumption and wellbeing. Evidence does not show that one branded timer length is universally best.

2 USER FLOW

- 1 Name and test a concrete target
- 2 Scaffold first action, materials and finish line
- 3 Choose time, output or both
- 4 Choose 3, 8, 15, custom or clock deadline
- 5 Clear the relevant perimeter
- 6 Lock and sprint
- 7 Pause, capture distractions or stop
- 8 Calibrate outcome without a productivity score

4 VISUAL AND AUDIO DIRECTION

A radar field closes onto one lockable target; interruptions become parked points outside the reticle. Restrained pulse and optional finish cue; static timer, hidden-time and screen-reader modes are equivalent.


3 MENTICATION UPGRADE

The sprint begins only after the target is startable, device-control claims are honest, time pressure is evaluated rather than assumed helpful, and distractions remain private and removable.

ELITE STANDARD

One mechanism. One memorable interaction. Honest outcomes, accessible alternatives and a consented route onwards.

SIGNATURE VISUAL



EVIDENCE REGISTER

B for interruption mitigation/time management; **C** for the exact sprint engine

Much interruption evidence is laboratory-based and time-management evidence is correlational or heterogeneous. Do not claim the timer treats attention disorders.

Reviewed 6 Sep 2026 | Next review Sep 2027

TOMORROW PARKING LOT

SLEEP / COGNITIVE OFFLOADING AND CONTAINMENT

1 PURPOSE

A low-light cognitive-offloading experience for unfinished responsibilities being rehearsed at bedtime. It captures one item outside the night and defers decisions, reminders and action planning to a separate daytime review.

PSYCHOEDUCATION

In one small sleep-laboratory study, writing a specific bedtime to-do list was associated with faster sleep onset than writing about completed activities. This supports offloading as a useful option, not a guarantee of sleep.

2 USER FLOW

- 1 Check whether the item is urgent or safety-critical
- 2 Capture one short line
- 3 Preview local storage and privacy
- 4 Slide or tap the item outside the night
- 5 Confirm storage
- 6 Offer Night Channel only with consent
- 7 End passively
- 8 Review, edit or delete in daytime

4 VISUAL AND AUDIO DIRECTION

Near-black indigo field with one luminous note moving beyond a quiet horizon. A single soft containment sound; no celebration, reminder ping or bright completion state.


3 MENTICATION UPGRADE

The night flow has only two active screens, sensitive text is local and explicitly saved, urgent items escape before sealing, and no hidden categorisation or reminder is created.

ELITE STANDARD

One mechanism. One memorable interaction. Honest outcomes, accessible alternatives and a consented route onwards.

SIGNATURE VISUAL



EVIDENCE REGISTER

B-: one small randomised laboratory study plus cognitive-offloading rationale

The study included 57 healthy young adults and used five minutes of writing. Do not promise sleep onset or generalise to clinical insomnia without caution.

Reviewed 6 Sep 2026 | Next review Sep 2027

NIGHT CHANNEL

SLEEP / SPATIAL ATTENTION MAPPING

1 PURPOSE

A low-effort audio destination for bedtime rumination when the user wants something sufficiently interesting to drift into. It supports familiar replay, gentle curiosity or fuller capture without creating another task to finish.

PSYCHOEDUCATION

Music may improve subjective sleep quality for some adults, while effects on objective sleep and other audio formats are less certain. Choose content that feels settling; Night Channel does not guarantee sleep.

2 USER FLOW

- 1 Choose Familiar Replay, Continuing Story, Gentle Curiosity, Gentle Conversation or Fictional World
- 2 Choose capture level
- 3 Select native Mentication audio or an honest external handoff
- 4 Choose a finite stop point
- 5 Enter near-black playback
- 6 Use external controls where applicable
- 7 Stop deliberately with no night rating
- 8 Offer optional daytime feedback

4 VISUAL AND AUDIO DIRECTION

Distant broadcast channels float in a near-black indigo space and resolve into one soft frequency line. Native audio can fade; external media remains under its own app's controls. No autoplay feed.


3 MENTICATION UPGRADE

Familiar Replay includes a known song, show, video, podcast or audiobook without voice cloning. Source boundaries are explicit, content is finite, and Mentication never simulates control of an external service.

ELITE STANDARD

One mechanism. One memorable interaction. Honest outcomes, accessible alternatives and a consented route onwards.

SIGNATURE VISUAL



EVIDENCE REGISTER

B for music listening; **C**/insufficient for podcasts, audiobooks and the branded channel system

Subjective music findings are more positive than objective measures, and evidence for general-population sleep is mixed. External content quality and autoplay are outside Mentication's control.

Reviewed 6 Sep 2026 | Next review Sep 2027

EVIDENCE REGISTER

BOX BREATHING

A for slow-paced breathing; C for the exact branded pattern

Slow, structured breathing can support autonomic regulation and increase heart-rate variability. Evidence is stronger for slow-paced breathing generally than for the exact 4-4-4-4 pattern.

LIMIT: Do not imply that the exact four-part ratio is superior. Stop or remove holds if dizziness, air hunger or respiratory discomfort occurs.

SOURCES: <https://pubmed.ncbi.nlm.nih.gov/35623448/>

PROGRESSIVE MUSCLE RELAXATION

A for PMR generally; C for this interaction system

Progressive muscle relaxation has evidence for reducing stress and anxiety in adults. The useful mechanism is learning the contrast between tension and release, not forcing every muscle to relax.

LIMIT: Evidence covers varied PMR protocols and populations. Avoid painful or injured areas and use release-only when tensing is unsuitable.

SOURCES: <https://pubmed.ncbi.nlm.nih.gov/38322293/>

EVIDENCE REGISTER

5-4-3-2-1 GROUNDING

C: mechanism-informed; exact sequence under-studied

Grounding uses present sensory information to redirect attention toward the here and now. The exact 5-4-3-2-1 sequence is widely used, but direct efficacy research on that specific sequence remains limited.

LIMIT: The review notes that grounding has lacked agreed definitions and efficacy studies. Never present this as proven treatment for dissociation or trauma.

SOURCES: <https://pubmed.ncbi.nlm.nih.gov/40575850/>

THOUGHT OR FACT

B for CBT/cognitive change; C for this brief standalone sorter

Cognitive approaches can reduce anxiety and depression by helping people examine interpretations and generate more balanced alternatives. This is a brief sorting tool, not a diagnostic test or a verdict on whether a person is right.

LIMIT: Do not use to challenge trauma memories, abuse disclosures, delusional or paranoid beliefs, or immediate danger. Evidence is for broader CBT packages, not this screen sequence alone.

SOURCES: <https://pubmed.ncbi.nlm.nih.gov/27717254/> | <https://pubmed.ncbi.nlm.nih.gov/38228727/>

EVIDENCE REGISTER

URGE SURFING

B for mindfulness and craving-response decoupling; C for this brief wave interaction

Mindfulness practice may weaken the link between craving and action even when the urge does not immediately disappear. The goal is a more available choice, not a promise that every urge will pass within a fixed time.

LIMIT: Results for brief app-based mindfulness are mixed. This is not suitable when an urge involves immediate danger, overdose risk, self-harm or harm to others.

SOURCES: <https://pubmed.ncbi.nlm.nih.gov/23265088/> | <https://pubmed.ncbi.nlm.nih.gov/29917096/>

THEN WHAT?

B for cognitive change/perspective-taking; C for this bounded standalone sequence

Cognitive change and perspective-taking can reduce emotional impact by changing how a situation is understood. This pathway tests coping beyond the worst frame; it is not reassurance and does not establish what will happen.

LIMIT: Not for immediate danger, acute panic, trauma re-experiencing, impaired reality testing or repetitive certainty-seeking. Broader CBT evidence does not prove the exact sequence.

SOURCES: <https://pubmed.ncbi.nlm.nih.gov/22582737/> | <https://pubmed.ncbi.nlm.nih.gov/38228727/>

EVIDENCE REGISTER

IGNITION POINT

A for behavioural activation packages; C for the four-pathway branded selector

Behavioural activation works by increasing meaningful contact between a person and their environment. A small action can be worthwhile before mood changes, so success is engagement with value or reward rather than instant happiness.

LIMIT: Evidence is mainly from depression treatment protocols delivered over time. One micro-action should not be described as treatment or guaranteed mood improvement.

SOURCES: <https://pubmed.ncbi.nlm.nih.gov/37068380/> | <https://pubmed.ncbi.nlm.nih.gov/37227760/>

CHANGE THE SCENE

C: mechanism-adjacent context research

Stable contexts can cue automatic behaviour, while a context change can interrupt those cues and create a window for a different choice. Evidence supports the broader context-discontinuity principle, not this exact short intervention.

LIMIT: The cited context-discontinuity evidence concerns longer-term habit contexts, not momentary mood shifts. Keep claims modest and never direct users into unsafe environments.

SOURCES: <https://pubmed.ncbi.nlm.nih.gov/27120333/>

EVIDENCE REGISTER

TEST THE PREDICTION

B for behavioural experiments in anxiety treatment; C for this standalone digital path

Behavioural experiments can help people compare a prediction with observable experience. Evidence suggests benefit within anxiety treatment, but results must be read fairly and should never be designed to prove the user wrong.

LIMIT: The review found methodological limitations and studied behavioural experiments inside treatment. Never create high-stakes, unsafe or boundary-violating tests.

SOURCES: <https://pubmed.ncbi.nlm.nih.gov/20381224/>

OPEN CHANNEL

B for broader social-connection interventions; C for a single digital re-entry prompt

Interventions that increase access to people or address barriers to connection can improve social contact and perceived connection. One message is only a bridge, not a guaranteed repair or cure for loneliness.

LIMIT: Intervention effects vary and much evidence concerns multi-session programmes. Do not use for abuse, coercion, unsafe former partners or when no contact has been requested.

SOURCES: <https://pubmed.ncbi.nlm.nih.gov/34796368/> | <https://pubmed.ncbi.nlm.nih.gov/34339939/>

EVIDENCE REGISTER

COUNTERMOVE

B for behavioural activation and implementation intentions; C for the graded trajectory interface

Behavioural activation increases meaningful interaction with the environment, and specific action plans can help close the gap between intention and behaviour. A small directional move can count before mood changes.

LIMIT: Evidence comes from broader interventions, not the 5/20/45-degree metaphor. Do not oppose behaviour that is protective, restorative or required for safety.

SOURCES: <https://pubmed.ncbi.nlm.nih.gov/37068380/> | <https://pubmed.ncbi.nlm.nih.gov/25965276/>

PULSE SHIFT

B for acute exercise affect; C for micro-dose graded activation

A single bout of exercise is associated on average with modest improvements in mood and anxiety, but responses vary widely. Here the aim is simply to make a little more movement available, not to prescribe intensity or promise a mood lift.

LIMIT: The meta-analysis found substantial heterogeneity and typically studied exercise bouts, not tiny movements. This is not medical or exercise advice.

SOURCES: <https://pubmed.ncbi.nlm.nih.gov/38787545/>

EVIDENCE REGISTER

REROUTE

C: mechanism-informed attentional deployment

Distraction can reduce emotional impact in some contexts, but its usefulness depends on the emotion, timing and purpose. Reroute is a deliberate short-term shift of attention, not suppression or a claim that the problem no longer matters.

LIMIT: The meta-analysis found strategy and context differences; attentional deployment was not uniformly effective. Do not use when urgent safety or practical action is required.

SOURCES: <https://pubmed.ncbi.nlm.nih.gov/22582737/>

NEXT EASIEST STEP

B for implementation intentions; C for task-compression micro-intervention

Specific action plans can improve goal attainment by linking intention to a concrete response. The useful target is a visible next behaviour, not a perfect plan or a judgment about motivation.

LIMIT: If-then planning evidence does not establish an optimal 20-second start rule. Keep the action user-editable and do not score productivity.

SOURCES: <https://pubmed.ncbi.nlm.nih.gov/25965276/>

EVIDENCE REGISTER

SIGNAL LOCK

B for interruption mitigation/time management; C for the exact sprint engine

Reducing interruption costs and using deliberate time-management strategies can support task accuracy, resumption and wellbeing. Evidence does not show that one branded timer length is universally best.

LIMIT: Much interruption evidence is laboratory-based and time-management evidence is correlational or heterogeneous. Do not claim the timer treats attention disorders.

SOURCES: <https://pubmed.ncbi.nlm.nih.gov/34273814/> | <https://pubmed.ncbi.nlm.nih.gov/33428644/>

TOMORROW PARKING LOT

B-: one small randomised laboratory study plus cognitive-offloading rationale

In one small sleep-laboratory study, writing a specific bedtime to-do list was associated with faster sleep onset than writing about completed activities. This supports offloading as a useful option, not a guarantee of sleep.

LIMIT: The study included 57 healthy young adults and used five minutes of writing. Do not promise sleep onset or generalise to clinical insomnia without caution.

SOURCES: <https://pubmed.ncbi.nlm.nih.gov/29058942/>

EVIDENCE REGISTER

NIGHT CHANNEL

B for music listening; C/insufficient for podcasts, audiobooks and the branded channel system

Music may improve subjective sleep quality for some adults, while effects on objective sleep and other audio formats are less certain. Choose content that feels settling; Night Channel does not guarantee sleep.

LIMIT: Subjective music findings are more positive than objective measures, and evidence for general-population sleep is mixed. External content quality and autoplay are outside Mentication's control.

SOURCES: <https://pubmed.ncbi.nlm.nih.gov/36000763/> | <https://pubmed.ncbi.nlm.nih.gov/33896299/>