


# Graph Theory and Algorithms

An Introduction to the world of Graphs



**Devansh Jain,**  
Indian Institute of Technology, Bombay

## Contents

1 Preface

ii

## 1 Preface

Preface

### About the project

adedsd

### About the report

sdsd

### References and Acknowledgement

sdfsd

(23)

CLASSMATE


Date \_\_\_\_\_

Page \_\_\_\_\_

L-6

Graphs are incredibly very useful structures in computer science. They come up in all sorts of applications, scheduling, optimization, communications, the design and analysis of algorithms.

⇒ Informally, a graph is just a bunch of dots and lines connecting the dots



⇒ A graph  $G$  is a pair of sets  $(V, E)$  where

- $V$  is a non-empty set of items called vertices or nodes
- $E$  is a set of 2-item subsets of  $V$  called edges (can be empty set).


$$V = \{x_1, x_2, x_3, x_4, x_5, x_6, x_7\}$$

$$E = \{ (x_1, x_2), (x_2, x_4), (x_4, x_3), (x_3, x_1), (x_5, x_6), (x_3, x_5), (x_5, x_7) \}$$

equivalent

to  $x_1 - x_2$


24




$\Rightarrow$  Two nodes  $x_i$  &  $x_j$  are adjacent if they're connected by an edge i.e.  $x_i - x_j \in E$

$\Rightarrow$  An edge  $e = (x_i, x_j)$  is said to be incident to its end-points  $x_i$  &  $x_j$ .

$\Rightarrow$  The number of edges incident to a node is called the degree of the node.



$\Rightarrow$  Graph is simple if it has no loops or multiples edges (multi-edge).



$|V| \rightarrow$  cardinality notation  
i.e. no. of elements in set V.

Men-Women partners ratio

25

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

6.041

6.002

6.042

6.003

6.034

Slots

5-7

7-9

9-11

11-1

1-3

### Graph-coloring problem


Given a graph  $G$  and  $K$  colors, assign a color to each node so adjacent nodes get different colors.

Minimum value of  $K$  for which such a coloring exists is the Chromatic number of  $G$   
 $\chi(G)$


→ Color  $\Rightarrow$  Slots

Nodes  $\Rightarrow$  Courses

Option 1



Option 2



$$\chi(G) = 3$$

2 colors can't work because of triangle formation

(26)


classmate

Date \_\_\_\_\_


Page \_\_\_\_\_

Basic graph coloring algorithm

1. Order the nodes  $v_1, \dots, v_n$
2. Order the colors  $c_1, \dots$
3. For  $i = 1, 2, \dots, n$
4. Assign the lowest legal color to  $v_i$



We know this is not best case.



So, if you change the ordering, you may get a better answer.

Now this algorithm is an example of what's known as a greedy algorithm.

You just go one step after the next, taking the best you can do at each step. You never go back and try to make things better.

(27)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

Thm.

If every node in  $G$  has degree  $d$ , then this basic algorithm uses atmost  $d+1$  colors for  $G$ .

Proof

By induction,

Induction hypothesis:

//Never induction on  $d$ , induction on  $n$  or  $e$   
↑  
node

Predicate

↓ "If every node in  $n$ -node graph  $G$  with max. degree  $d$   
 $P(n) =$  then this basic algorithm uses atmost  $d+1$  colors  
 for  $G$ ."

Base case:

 $n=1 \Rightarrow |E|=0 \Rightarrow d=0 \Rightarrow$  one color∴  $P(1)$  is trueInd. step: Assume  $P(n)$  is trueLet  $G$  be any  $n+1$  node graph.Let  $d$  be the max. degree of nodes in  $G$ Order the nodes:  $v_1, \dots, v_n, v_{n+1}$ Let  $G'$  be the graph after removing  $v_{n+1}$   
max. degree of nodes in  $G' \leq d$ ∴ Basic algorithm uses atmost  $d+1$  colors  
for  $G'$ .

(28)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

In  $G$ , color nodes  $v_1, \dots, v_n$  same as  $G'$ .

$v_{n+1}$  has atmost  $d$  neighbours therefore  
atleast one color is not present in  
 $v_{n+1}$ 's neighbours.

∴ Give  $v_{n+1}$  that color.

∴  $P(n+1)$  is true

$K_n \rightarrow$  n-node complete graph (or clique)



$$\text{Degree} = n-1$$

$$\chi(K_n) = n$$

(n)




$$d = n/2$$

$$\chi(G) = 2$$

Bipartite graph

(n) star graph




$$d = n-1$$

$$\chi(G) = 2$$

The Algorithm may be much better than the theorem

(29)

classmate

Date \_\_\_\_\_  
Page \_\_\_\_\_

Good ordering (2 colors)

(C <sub>1</sub> ) v <sub>1</sub>	v <sub>5</sub> (C <sub>2</sub> )
(C <sub>1</sub> ) v <sub>2</sub>	v <sub>6</sub> (C <sub>2</sub> )
(C <sub>1</sub> ) v <sub>3</sub>	v <sub>7</sub> (C <sub>2</sub> )
(C <sub>1</sub> ) v <sub>4</sub>	v <sub>8</sub> (C <sub>2</sub> )

Bad ordering ( $\lceil \frac{n}{2} \rceil$  colors)



(C <sub>1</sub> ) v <sub>1</sub>	v <sub>3</sub> (C <sub>2</sub> )
(C <sub>2</sub> ) v <sub>3</sub>	v <sub>4</sub> (C <sub>2</sub> )
(C <sub>3</sub> ) v <sub>5</sub>	v <sub>6</sub> (C <sub>3</sub> )
(C <sub>4</sub> ) v <sub>7</sub>	v <sub>8</sub> (C <sub>5</sub> )

A graph  $G = (V, E)$  is said to be bipartite if  $V$  can be split into  $V_L, V_R$  so that all the edges connect  $V_L$  to  $V_R$ .

resource allocation problems (load balancing traffic on internet)


CLASSMATE

(30)



Matching algorithm - usage: online dating agencies, assignments problems (matching interns to hospitals on match day)


L-7  $\Rightarrow$  Given a graph  $G = (V, E)$ , a matching is a subgraph of  $G$  where every node has degree 1.



{x1-x3, x2-x5, x3-x6} is a matching of size 3.

Size 4 matching is not possible in this graph.

$\Rightarrow$  A matching is perfect if it has size  $\frac{|V|}{2}$



(31)

classmate


Date \_\_\_\_\_

Page \_\_\_\_\_


Weights?

$\Rightarrow$  The weight of a matching ( $M$ ) is the sum of the weights on the edge of  $M$ .

$\Rightarrow$  A minimum weight matching for graph  $G$  is a perfect matching for  $G$  with the minimum weight.



Preference list - similar to weights



If  $b_1-g_1, b_2-g_2$ , then  $b_1-g_2$  will be rogue couple.

Given a matching  $M$ ,  $x \& y$  form rogue couple if they prefer each other to their mates in  $M$ .


A matching is stable if there aren't any rogue couple.

(32)

classmate

Date \_\_\_\_\_


Page \_\_\_\_\_



$b_1-g_2$  &  $b_2-g_1$  is stable matching

If  $b-g$  matching is only allowed then there always exist a stable perfect matching.

If  $b-b$  &  $g-g$  is allowed, it may not always possible to find stable perfect matching.



$g_2$  (Preference doesn't matter)

Theorem: There is no stable perfect matching

Proof: By contradiction,

If there is a stable matching,  $g_2$  would be paired to someone without loss of generality (by symmetry) (the triangle is symmetric)

$\{g_2-b_2, g_1-b_1\} \Rightarrow b_1, b_2$  form rogue couple.

(33)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

1962

Stable Marriage Algorithm (<sup>David Lloyd</sup> Gale-Shapley Algo)

- N boys & N girls

- each boy has his own ranked list of all the girls. Same for each girl

- to find stable perfect matching.

Eg:- 5 boys & 5 girls

1 ♂	CBEAD	A 35214
2 ♂	ABECD	B 52143
3 ♂	DCBAE	C 43512
4 ♂	ACDBE	D 12345
5 ♂	ABDEC	E 23415

Using greedy algorithm,

1-C, 2-A, 3-D, 4-B, 5-E

Rogue Couple : 4-C

Using mating algorithm,  
Serendaders

Girls	Day 1	Day 2	Day 3	Day 4	
1 ABEAD	A 2 4(5)	5	5	5	A-5
2 ABEDC	B	2	21	2	B-2
3 DCBAE	C 1	14	4	4	C-4
4 ACDBE	D 3	3	3	3	D-3
5 ABDEC	E			1	E-1

34

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

Need to show:

- ① Algorithm terminates (Matching returned)
- ② Everyone gets matched (Perfect matching)
- ③ No rogue couple (Stable matching)
- ④ It runs quickly
- ⑤ Fairness

Theorem 1 Algo terminates in less than  $N^2+1$  days

Proof By contradiction.

Suppose, algo doesn't terminate in  $N^2+1$  days.

Claim: If we don't terminate on a day,  
then atleast one girl had more than  
one boy and so she rejected atleast  
one boy who crosses her name.

If the algo doesn't terminate in  $N^2+1$  days  
then atleast  $N^2+1$  cross-outs have happened.

As there are  $N^2$  names on  $N$  lists, there  
are atmost  $N^2$  cross-outs. ( $< N^2+1$ )

Thus, not possible.

(35)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

Predicate  $P = \text{"If a girl } G \text{ ever rejected a boy } B, \text{ then } G \text{ has a suitor who she prefers to } B."$

Lemma 1 :  $P$  is an invariant for the algo

Proof: Base case : By induction on number of days.

Base case: Day 0 (Vacuously true)

(No one rejected yet)

Ind step:  $P$  holds at the end of Day  $d$ .

Case 1  $G$  rejects  $B$  on day  $d+1$ .

Then there was a better boy

$\Rightarrow P$  holds on Day  $d+1$

Case 2  $G$  rejected  $B$  before day  $d+1$

$P$  on day  $d \Rightarrow G$  had a better boy on day  $d$ .

$\Rightarrow P$  holds on Day  $d+1$ .

Theorem 2 Everyone is married in Algo

Proof By contradiction

Assume that some boy  $B$  is not married,

then he was rejected by every girl which

$\Rightarrow$  every girl by Lemma 1 has a better suitor

$\Rightarrow$  every girl is married  $\Rightarrow$  every boy is married

$\Rightarrow B$  is married.

(36)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

Theorem 3 Algo produces a stable matching (No rogue couple)

Proof

let B & G be any pair that are not married.

To prove: B & G aren't rogue.

Case 1 G rejected B

⇒ G has a better suitor than B (Lemma 1)

⇒ G married someone whom she prefers over B.

⇒ G is not rogue with B.

Case 2 G didn't reject B

⇒ B never serenaded G

⇒ G is lower in order than B's spouse.

⇒ B is not rogue with G

∴ No rogue couples.

37

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

Let  $S$  be the set of all stable matching.  
 $S \neq \emptyset$  as algo produces one stable matching.

For each person  $P$ , we define the realm of possibility for  $P$  to be  $\{Q \mid \exists M \in S \mid \{P, Q\} \in M\}$

A person's optimal mate is his/her favorite from the realm of possibility.

A person's pessimal mate is his/her least favorite from the realm of possibility.

Theorem 4 Algo marries every boy with his optimal mate.

Proof —

Theorem 5 Algo marries every girl with her pessimal mate.

Proof —

TMA  $\rightarrow$  The Marriage Algorithm

38

## classmate

Date \_\_\_\_\_  
Page \_\_\_\_\_


- L-8

  - Walks & Paths
  - Connectivity
  - Cycles & Closed walk
  - Spanning Tree (ST)
  - Min-weight spanning tree (MST)

## Walks & Paths

⇒ A walk is a sequence of vertices that are connected by edges

It has  $k$ -edges of length =  $k$



$\Rightarrow$  A path is a walk where all  $v$ 's are distinct

(39)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

Lemma 1 If I walk from  $u$  to  $v$ , then I path from  $u$  to  $v$ .

Proof I walk  $u$  to  $v$ .

By well-ordering principle: Walk of minimal length.

$$u = v_0 - v_1 - \dots - v_k = v$$

We prove that this walk (of minimal length) is a path.

Case  $k=0$  No edge length = 0

Case  $k=1$  Just one edge length = 1

Case  $k \geq 2$

Suppose this walk is not a path

then  $i \neq j$   $v_i = v_j$

$$u = v_0 - \dots - v_i = v_j - \dots - v_k$$

Now this a shorter walk

thus contradicting our assumption.

(40)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_


Connectivity

- $\Rightarrow$   $u$  and  $v$  are connected if there is a path from  $u$  to  $v$ .
- $\Rightarrow$  A graph  $G = (V, E)$  is connected if every pair of vertices  $(v_1, v_2) \in V \times V$  are connected.

Not connected graph



Connected graph

Cycles & closed walks

(also known as loops)

- $\Rightarrow$  A closed walk is a walk starts and ends at exactly same vertex.

$$v_0 - v_1 - \dots - v_k = v_0$$

a closed walk with and

- $\Rightarrow$  If  $k \geq 3$ , if all  $v_0, \dots, v_{k-1}$  are distinct then it is called a cycle

(41)

classmate


Date \_\_\_\_\_

Page \_\_\_\_\_

Trees

⇒ A connected and acyclic graph is called a tree.

⇒ A leaf is a node with degree = 1 in a tree.



Lemma 2 Any connected subgraph of a tree is a tree.

Proof By contradiction, if the connected subgraph is not a tree then it must have a cycle.

As the subgraph is a part of a graph, the graph must have a cycle.

But the graph is a tree.

We get a contradiction. ■

(42)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

Lemma 3 A tree with  $n$  vertices has  $n-1$  edges.

Proof By Induction

P( $n$ ) Ind. Hypo. :  $P(n)$  = "There are  $n-1$  edges in an  $n$ -vertex tree"

Base case:  $P(1)$  is true. ①

Ind. step: Assume that  $P(n)$  is true

Let  $T$  be an  $(n+1)$ -vertex tree,

Let  $v$  be a leaf of the tree.

Delete  $v$ : this creates a connected subgraph  
(which is also a tree (Lemma))

By  $P(n)$ : this subgraph has  $n-1$  edges.

Re-attach  $v$ :  $T$  has  $(n-1)+1=n$  edges  
(degree of  $v=1$ )

$P(n+1)$  is true



43

classmate

Date \_\_\_\_\_  
Page \_\_\_\_\_

## Spanning Trees

→ A spanning tree (ST) of a connected graph  $G = (V, E)$  is a subgraph that is a tree with same vertices as the graph.


Theorem Every connected graph has a ST.

Proof By contradiction.

Assume a connected graph  $G$  with no ST.

Let  $T$  be a connected subgraph of  $G$  and has same vertices as  $G$  and has minimum number of edges.

As  $T$  is not a spanning tree, it must have atleast one cycle.



(44)


classmate

Date \_\_\_\_\_  
Page \_\_\_\_\_

Subgraph we obtain is smaller than T  
(w.r.t. number of edges)

thus contradicting our minimum condition on T.

Minimum-weighted spanning tree



$$ST_1 = \{v_1-v_4, v_4-v_5, v_5-v_2, v_5-v_6, v_6-v_3, v_6-v_7, v_5-v_8\}$$

$$\text{Weight of } ST_1 = 2 + 2 + 3 + 1 + 3 + 7 + 1 = 19$$

$$ST_2 = \{v_1-v_2, v_1-v_4, v_4-v_5, v_5-v_6, v_6-v_3, v_6-v_7, v_5-v_8\}$$

$$\text{Weight of } ST_2 = 1 + 2 + 2 + 1 + 3 + 7 + 1 = 17$$

(45)

classmate


Date \_\_\_\_\_

Page \_\_\_\_\_

⇒ The minimum spanning tree (MST) of an edge-weighted graph is defined as the ST of  $G$  with the smallest possible sum of edge-weights.

Algo

- Grow a subgraph one edge at a time, such that at each step:
- Add the minimum weighted edge keeps the subgraph acyclic.



$$+1 \quad (1) \quad v_6 - v_8$$

$$+1 \quad (2) \quad v_1 - v_2$$

$$+1 \quad (3) \quad v_5 - v_6 \Rightarrow v_5 - v_8 \text{ } \times$$

$$+2 \quad (4) \quad v_1 - v_4$$

$$+2 \quad (5) \quad v_4 - v_5 \Rightarrow v_4 - v_8 \text{ } \times, v_2 - v_5 \text{ } \textcircled{R}$$

$$+3 \quad (6) \quad v_2 - v_3 \Rightarrow v_3 - v_8 \text{ } \times$$

$$+2 \quad (7) \quad v_6 - v_7$$

46

classmate

Date \_\_\_\_\_  
Page \_\_\_\_\_

Theorem For any connected weighted graph  $G$ , the algo produces a MST.

Lemma Let  $S$  consists of first  $k$ -edges, by the algo, then  $\exists$  MST  $T = (V, E)$  for  $G$  such that  $S \subseteq E$

Proof

(Assumed lemma is true)

Graph  $G = (V, E)$  $|V| = n$ 

1) Suppose  $< n-1$  edges are picked, then  $\exists$  edge in  $E - S$  that can be added without creating a cycle.

2) Once  $n-1$  edges are <sup>picked</sup>, we know  $S$  is an MST.

Proof

By Induction:


$P(m) = \# G \# S$  consisting of first  $m$  selected edges  $\exists$  MST of  $G$  such that  $S \subseteq E$ .  
 $T = (V, E)$

Base Case:  $m = 0 \Rightarrow S = \emptyset \subseteq E$  for any MST.  
 $T = (V, E)$

Assume


Ind. Step:  $P(m)$  holdslet  $e$  denote the  $(m+1)$ th selected edge.Let  $S$  denote the first  $m$  selected edges.

47



Let  $T^* = (V, E^*)$  be MST of  $G$  such that  
 $S \subseteq E^*$

Case 1:  $e \in E^* \Rightarrow S \cup \{e\} \subseteq E^* \Rightarrow P(m+1)$  holds



From the algo,

$\{ \text{algo} \rightarrow S \cup \{e\} \}$  has no cycle

(This is how we choose  $(m+1)^{\text{th}}$  edge)

$T^*$  is a tree  $\Rightarrow (V, E^* \cup \{e\})$  must contain  
 $= (V, E^*)$  a cycle. (Proof in book)

→ this cycle has edge  $e' \in E^* - S$

(48)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

Algo could have selected  $e$  or  $e'$   
 $\Rightarrow$  weight of  $e \leq$  weight of  $e'$ .

Swap  $e$  and  $e'$  in  $T^*$ :

$$\text{Let } T^{**} = (V, E^{**}) : E^{**} = (E^* - \{e'\}) \cup \{e\}$$

We need to prove that  $T^{**}$  is an MST.

- $T^{**}$  is acyclic because we removed  $e'$  from the only cycle in  $E^* \cup \{e\}$ .
- $T^{**}$  is connected since  $e'$  was on a cycle.
- $T^{**}$  contains all vertices of  $G$

$\Rightarrow T^{**}$  is ST of  $G$ .

$$\begin{aligned} \text{weight of } T^{**} &= \text{weighted sum of } E^{**} \\ &= \text{Weighted sum of } E^* - \text{weight of } e' \\ &\quad + \text{weight of } e \\ &= \text{Weight } \cancel{\text{sum}} \text{ of } T^* - \text{weight of } e' \\ &\quad + \text{weight of } e \end{aligned}$$

As weight of  $e \leq$  weight of  $e'$

Weight of  $T^{**} \leq$  weight of  $T^*$


And as  $T^*$  is MST, equality holds &  $T^{**}$  is also MST.  $\square$

49


classmate

Date \_\_\_\_\_  
Page \_\_\_\_\_L-9L-10Communication Networks

- great application of graph theory
- how do you route packets through networks

 $N \times N$  Networks ( $N = 2^k$ )Complete Binary Tree $\circ$  = switch (direct packets through a network) $\square$  = terminal (source and destination of data)4x4 Network $N \times N$  NetworkDiameter =  $2(1 + \log_2 N)$ Max. Switch size =  $3 \times 3$ Switch count =  $2N - 1$ Max Congestion =  $N$

(50)



**Latency:** Time that is required for a packet to travel from an input to an output.

**Diameter:** Length of the shortest path of a network b/w Input and Output that are farthest apart  
(Maximum length of shortest path)

A permutation is a function  $\pi: \{0, \dots, n-1\} \rightarrow \{0, \dots, n-1\}$  such that no two numbers are mapped to the same value  
( $\pi(i) = \pi(j) \text{ iff } i=j$ )


$$\text{Eg: } \begin{aligned} \pi(i) &= n-1-i \\ \pi(i) &= i \end{aligned}$$


Permutation Routing Problem for  $\pi$

- For each  $i$  direct the packet at  $I_i$  to  $O_{\pi(i)}$   
path taken is denoted by  $P_{i, \pi(i)}$

The congestion of paths  $P_{0, \pi(0)}, \dots, P_{n-1, \pi(n-1)}$  is equal to the largest number of paths that pass through a single switch.

Max congestion = max possible value of congestion. (over permutations with best path)






53

## classmate

Date

Page

## Butterfly Network Structure



Switch is uniquely identified by its row & col

binary rep. of input

$$\left( \underbrace{(b_1 \dots b_{l+1} \dots b_{\log N})}_{\uparrow}, l+1 \right) \quad \left( (b_1 \dots b_{l+1} \dots b_{\log N}), l+1 \right)$$

Complement  
of  $(1+)^n$  bit.

## Routing

$$(x_1 \dots x_{\log N}, 0) \rightarrow (y_1 x_2 \dots x_{\log N}, 1)$$

$$(x_1, x_2, x_3, \dots, x_{\log N}), 2)$$

$$((y_1, \dots, y_{\log N}), \log N) \leftarrow ((y_1, \dots, y_{\frac{N}{2}}, x_{\frac{N}{2}+1}, \dots, x_{\log N}), l)$$

(54)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

 $\text{In}_3 \rightarrow \text{Out}_5$ Eg:-  $011 \rightarrow 101$  $\text{In}_3 \rightarrow (011, 0)$  $(111, 1)$  $(101, 2)$  $\text{Out}_5 \leftarrow (101, 3)$ NxN Network

$$\text{Diameter} = 2 + \log N$$

$$\text{Max. Switch size} = 2 \times 2$$

$$\text{No. of Switches} = N(1 + \log N)$$

$$\text{Congestion} = \sqrt{N} (N \cdot 2^{2n})$$

$$= \sqrt{N/2} (N \cdot 2^{2n+1})$$

classmate  
Date \_\_\_\_\_  
Page \_\_\_\_\_

Benes

1960s - Benes, a Bell Labs researcher

$N \times N$  Network

Diameter =  $1 + 2 \log N$

Max switch size =  $2 \times 2$

No of switches =  $2N \log N$

Congestion = 1


(56)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

Theorem | Congestion of  $N$ -input Benes network  
is 1, when  $N = 2^a$  for some  $a \geq 1$

ProofBy induction on  $a$ , $P(a) =$  "This theorem is true for  $a$ ".Base case:  $a=1 N=2$ 

$$\pi(0) = 0$$

$$\pi(1) = 1$$

Congestion = 1  
(Just straight)

$$\pi(0) = 1$$

$$\pi(1) = 0$$

Congestion = 1  
(Just cross-over)

Ind. Step: Assume  $P(a)$  is true

Example:	$\pi(0) = 1$	$\pi(4) = 3$
	$\pi(1) = 5$	$\pi(5) = 6$
	$\pi(2) = 4$	$\pi(6) = 0$
	$\pi(3) = 7$	$\pi(7) = 2$

Constraint graph

If two packets must pass through different subnetworks then there is an edge b/w them.

(57)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

L-10

## Euler tours



## Directed graphs

- Definitions
- No. of walks
- Strong connectivity
- DAGs

## Tournament Graphs

Euler tour


- He lived in Konigsberg
- Seven bridges (birth of graph theory)

An Euler tour is a walk that traverses every edge exactly once, and starts and finishes at same vertex.

Theorem A connected graph has an Euler tour iff every vertex has even degree.

Proof ( $\Rightarrow$ )

Assume  $G = (V, E)$  has an Euler tour




(58)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

Since every edge in  $E$  is traversed once;  
 clearly  $\deg(u) = (\# \text{ times } u \text{ appears in this tour } v_0 \dots v_k) \times 2$



(≤)

For  $G = (V, E)$ , assume  $\deg(v)$  is even  $\forall v \in V$

Let  $N: v_0 - v_1 - \dots - v_k$  be the longest walk that traverses no edge more than once.

①  $v_k - u$  not in  $W \Rightarrow v_0 - v_1 - \dots - v_k - u$  is longer than  $W \Rightarrow$  Contradicts  
 $\therefore$  All the edges  $\overset{\text{int}}{\in}$  incident of  $v_k$  are traversed in  $W$ .


②  $v_k \neq v_0 \Rightarrow v_k$  has odd degree in walk  $W$   
 As we have showed that all edges  $\overset{\text{int}}{\in}$  of  $v_k$  are in  $W$ , so  $\deg(v_k)$  is odd in  $G$   
 $\Rightarrow$  Contradicts as  $\deg(v)$  is even  $\forall v \in V$ .

(59)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

Directed graph (digraphs)

$$\text{indegree}(v_2) = 2$$

$$\text{outdegree}(v_2) = 1$$

Theorem Let  $G = (V, E)$  be an  $n$ -node graph with  $V = \{v_1, \dots, v_n\}$ . Let  $A = \{a_{ij}\}$  denote the adjacency matrix for  $G$  that is

$$a_{ij} = \begin{cases} 1 & \text{if } v_i \rightarrow v_j \text{ is an edge.} \\ 0 & \text{otherwise} \end{cases}$$


Let  $p_{ij}^{(k)} = \#$  directed walks of length  $k$  from  $v_i$  to  $v_j$ .

$$\text{then } A^k = \{p_{ij}^{(k)}\}$$

$$A = \begin{bmatrix} v_1 & v_2 & v_3 \\ v_1 & 1 & 1 & 1 \\ v_2 & 0 & 0 & 1 \\ v_3 & 0 & 1 & 0 \end{bmatrix} \quad A^k = \begin{bmatrix} 1 & 2 & 2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$A^3 = \begin{bmatrix} 1 & 3 & 3 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

(60)

Proof

$a_{ij}^{(k)}$  denote the  $(i, j)^{\text{th}}$  entry in  $A^k$

By induction:

Predicate  $\rightarrow P(k) = \text{"Theorem is true for } k\text{"}$   
 $= \text{"}\forall i, j \quad a_{ij}^{(k)} = p_{ij}^{(k)}\text{"}$

Base case  $k=1$

Edge  $v_i \rightarrow v_j : p_{ij}^{(1)} = 1 = a_{ij}^{(1)}$

No edge  $v_i \rightarrow v_j : p_{ij}^{(1)} = 0 = a_{ij}^{(1)}$

$\therefore \forall i, j \quad a_{ij}^{(1)} = p_{ij}^{(1)}$

Ind. step

Assume  $P(k)$  holds.

$$p_{ij}^{(k+1)} = \sum_{h: v_h \rightarrow v_j \text{ is an edge in } E} p_{ih}^{(k)} = \sum_{h=1}^n p_{ih}^{(k)} \cdot a_{hj}^{(k)}$$

$v_i \xrightarrow{k} v_n \xrightarrow{?} v_j$

Ind. step  $\rightarrow$

$$= \sum_{h=1}^n a_{ih}^{(k)} \cdot a_{hj}^{(k)}$$

$$= a_{ij}^{(k+1)} \Rightarrow P(k+1)$$

Matrix multiplication holds

(6)

classmate


Date \_\_\_\_\_

Page \_\_\_\_\_

$\Rightarrow$  A digraph  $G = (V, E)$  is called strongly connected if  $\forall u, v \in V \exists$  directed path from  $u$  to  $v$  in  $G$ .

$\Rightarrow$  A digraph is called directed acyclic graph (DAG) if it does not contain any directed cycles.

### Tournament Graph



$\Rightarrow$  A directed Hamiltonian path is a directed walk that visits every vertex exactly once.

Theorem Every tournament graph contains a directed Hamiltonian path.

(62)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

Proof By induction (on number of nodes)

$P(n)$  = "Every tournament graph on  $n$ -nodes contains a directed Ham-pair."

Base case :  $P(1)$  holds

Ind. step : Assume  $P(n)$  holds

Consider a tournament graph on  $(n+1)$ -nodes

Take out a node  $v$ . We are left with a tournament graph on  $n$ -nodes.

By  $P(n)$ ,  $v_1 \rightarrow v_2 \rightarrow \dots \rightarrow v_n$

Applying  $\text{Q} \rightarrow$  If  $v \rightarrow v_i$  is an edge, then  $P(n+1)$  holds.

(if  $v_i \rightarrow v$  and  $v \rightarrow v_n$ ) then

Claim:  $\exists i$  such that  $v_i \rightarrow v$  &  $v \rightarrow v_{i+1}$

Proof by contradiction

If no such  $i$  exists and we know

$v \rightarrow v \not\in$  then  $v_i \rightarrow v \nexists i \in \{1, 2, \dots, n\}$  (by induction)

But  $v \rightarrow v_n$  contradicts this.

There is also proof by strong induction.  
See it.

63

CLASSMATE  
Date \_\_\_\_\_  
Page \_\_\_\_\_

Approach

Case 1  $v \rightarrow v_i$

Case 2  $v_i \rightarrow v$

Smallest  $i$  such that  $v \rightarrow v_i$   
 $\Rightarrow v_k \rightarrow v$  &  $k < i$   
 $\Rightarrow v_{i-1} \rightarrow v$

$(v_1 \rightarrow \dots \rightarrow v_{i-1} \rightarrow v_i \rightarrow v_{i+1} \rightarrow \dots \rightarrow v_n)$

directed  
A new Hamiltonian path  
is formed

$\therefore P(n+1)$  holds

(64)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_


King Chicken Problem

either chicken  $u$  pecks chicken  $v$ :  $u \rightarrow v$   
 or chicken  $v$  pecks chicken  $u$ :  $v \rightarrow u$

$u$  "virtually" pecks  $v$  if

- $u \rightarrow v$ , or
- $\exists w \ u \rightarrow w \rightarrow v$

A chicken that virtually pecks every other chicken is called king chicken.




Theorem The chicken with highest outdegree is definitely a king.

Proof By contradiction,

Let  $u$  have highest outdegree.  
 Suppose  $u$  is not king

(65)



$\Rightarrow \exists_v : v \rightarrow u$ , and  $\nexists_w : u \rightarrow w \Rightarrow v \rightarrow w$

(Outdegree of  $v$ )  $\geq$  (Outdegree of  $u$ ) + 1

So, our assumption is false. ■

L-11

- Relations
  - Properties
  - Equivalent Relations
  - Partial Orders
    - Hasse diagram
    - Total Order
    - Topological Sort
  - Parallel Task Scheduling
    - Dilworth's Lemma

### Relations

$\rightarrow$  A relation from a set  $A$  to a set  $B$   
is a subset  $R \subseteq A \times B$

Ex :-  $R = \{(a, b) : \text{student } a \text{ is taking class } b\}$

$(a, b) \in R : a R b, a \underset{\substack{\text{Relational} \\ \uparrow \\ \text{Symbol}}}{\sim} b$

(66)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_


→ Relation on A is a subset  $R \subseteq A \times A$

Eg:-  $A = \mathbb{Z} : xRy \text{ iff } x \equiv y \pmod{5}$

$A = \mathbb{N} : xRy \text{ iff } x|y$

$A = \mathbb{N} : xRy \text{ iff } x \leq y$

Set A together with R is a directed graph  $G = (V, E)$  with  $V = A, E = R$ .



### Properties

A relation R on A is

\* reflexive if  $xRx \quad \forall x \in A$

\* symmetric if  $xRy \Rightarrow yRx \quad \forall x, y \in R$

\* anti-symmetric if  $xRy \wedge yRx \Rightarrow x=y$

\* transitive if  $xRy \wedge yRz \Rightarrow xRz$

(67)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

Ex

Reflexive Symmetric Anti-Symmetric Transitive

 $x \equiv y \pmod{5}$  ✓ ✓ ✗ (2, 7) → Equivalence

 $x \mid y$  ✓ ✗ (1, 2) → Divisibility

 $x \leq y$  ✓ ✗ (1, 2) → Order

⇒ An equivalence relation is reflexive, symmetric, transitive.

Ex!  $A = \mathbb{Z} : x R y \text{ iff } x = y \in \mathbb{Z}$   
 $A = \mathbb{Z} : x R y \text{ iff } x \equiv y \pmod{5}$

⇒ The equivalence class of  $x \in A$  is the set of all elements in A related to  $x$  by  $R$ : denoted by  $[x]$   
 $[x] = \{y : x R y\}$

Ex:  $A = \mathbb{Z} : x R y \text{ iff } x \equiv y \pmod{5}$

$[7] = \{ \dots, -3, 2, 7, 12, 17, \dots \}$

$[2] = [7] = [12] = [17] = \dots$

(68)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

A partition of  $A$  is a collection of disjoint nonempty sets  $A_1, \dots, A_n \subseteq A$  whose union is  $A$ .

Ex:  $A = \mathbb{Z} : xRy \text{ iff } x \equiv y \pmod{5}$

$$A_0 = \{ \dots -5, 0, 5, \dots \}$$

$$A_1 = \{ \dots -4, 1, 6, \dots \}$$

$$A_2 = \{ \dots -3, 2, 7, \dots \}$$

$$A_3 = \{ \dots -2, 3, 8, \dots \}$$

$$A_4 = \{ \dots -1, 4, 9, \dots \}$$

Theorem The equivalence classes of equivalence relation on a set  $A$  form a partition of  $A$ .

### Partial Orders

A relation is a (weak) partial order if it is reflexive, antisymmetric and transitive.

A relation is a (strong) partial order if it is irreflexive, antisymmetric and transitive  
 (See book).

(69)


classmate

Date \_\_\_\_\_  
Page \_\_\_\_\_

A partial order relation is denoted by  $\leq$  instead of  $R$ .


$(A, \leq)$  is actually called partial ordered set. or poset

A poset is a directed graph with vertex set  $A$  and edge set  $\leq$ .



Transitive  $\Rightarrow$  No directed cycles

Antisymmetric  $\Rightarrow$  Only one directed edges b/w two vertices



(70)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

A Hasse diagram for a poset  $(A, \leq)$  is a directed graph with vertex set  $A$  and edge set  $\leq$  minus:

- \* all self-loops, and
- \* all edges implied by transitivity

Theorem A poset has no directed cycles other than self loops.

Proof By contradiction

Suppose  $\exists n (\geq 2)$  distinct elements

$a_1, \dots, a_n$  such that

$a_1 \leq a_2 \leq a_3 \leq \dots \leq a_n \leq a_1 \Rightarrow a_1 \leq a_1$

(induction)

By transitivity  $\rightarrow a_1 \leq a_2 \leq \dots \leq a_n \rightarrow a_1 \leq a_n$

( $\neg$ )

$a_1 = a_n$

Contradicts distinct element assumption.  $\blacksquare$

(IV)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

Conclusion of the theorem

So, deleting self-loops from a poset, makes a directed acyclic graph (DAG)

⇒ a and b are incomparable if neither  $a \leq b$  nor  $b \leq a$

⇒ a and b are comparable if ~~neither~~  $a \leq b$  or  $b \leq a$ .

⇒ A total order is a partial order in which every pair of elements is comparable.

Hesse diagram of a total order is a straight line.

A total order consistent with a partial order is called a Topological sort.

A topological of a poset  $(A, \leq)$  is a total order  $(A, \leq_T)$  such that

$$\leq \subseteq \leq_T$$

$$(x \leq y \Rightarrow x \leq_T y)$$

(72)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

Theorem Every finite poset has a topological sort.

Proof  $\Rightarrow x \in A$  is minimal if there doesn't exist  $y \in A : y \neq x$  such that  $y \leq x$ .

$\Rightarrow x \in A$  is maximal if  $\forall y \in A, y \neq x$  s.t.  $x \leq y$ .  
 ↓  
 not there exists

Lemma Every finite poset has a minimum element.

$\Rightarrow$  A chain is a sequence of elements such that  $a_1 \leq a_2 \leq \dots \leq a_t$  (length of chain is  $t$ ).  
 distinct

Proof Let  $c: (a_1, \leq a_2, \dots, a_n)$  be the max length chain.

Case 1:  $a \notin \{a_1, \dots, a_n\}$   
 if  $a \leq a_1$ , then  $c$  is not the longest chain.

So,  $a \nleq a_1$

(73)

classmate

Date \_\_\_\_\_  
Page 61Case 2:  $a \in \{a_1, \dots, a_n\}$ 

if  $a \leq a_i$ , then we have  
 a directed cycle which  
 contradicts Theorem 1.

So,  $\nexists a \leq a_i$ , $\therefore \nexists a \in A : a \leq a_i$ ,So, by definition,  $a_i$  is the  
 minimum element.

Add Parallel Task Scheduling

&amp; Dilworth's Lemma.

L-13

(45:30)

Asymptotic Notationtilde  $f(x) \sim g(x)$  if  $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = 1$ oh, big-oh  $f(x) = O(g(x))$  if  $\lim_{x \rightarrow \infty} \left| \frac{f(x)}{g(x)} \right| < \infty$ 

(finite)

Multiple usage:

 $f(x) \leq O(g(x))$ ;  $f(x)$  is  $O(g(x))$ ;  $f(x) \in O(g(x))$  $O(g(x))$  is aset of all func.  
 that grow slowly than  $g(x)$

(74)

classmate

Date \_\_\_\_\_  
Page 62Theorem

Let  $f(x) = x$ ,  $g(x) = x^2$   
 Then  $f(x) \in O(g(x))$

Proof

$\lim_{x \rightarrow \infty} \frac{x}{x^2} = 0$  which is finite

Theorem

$x^2 \notin O(x)$

Proof

$\lim_{x \rightarrow \infty} \frac{x^2}{x} = \lim_{x \rightarrow \infty} x$  is infinite.

Q. Is  $x^2 \in O(10^6 x)$  true?

A. No.

Q. Is  $10^6 x^2 \in O(x^2)$  true?

A. Yes

Q. Is  $x^2 + 100x + 10^7 = O(x^2)$  true?

A. Yes

Theorem  $x^{10} \in O(e^x)$

Proof

$\lim_{x \rightarrow \infty} \frac{x^{10}}{e^x} = 0$  which is finite.  
 (L'Hopital's rule)

75

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

63

Theorem

$$4^x \notin O(2^x)$$

Proof

$$\lim_{x \rightarrow \infty} \frac{4^x}{2^x} = \lim_{x \rightarrow \infty} 2^x \rightarrow \infty$$

Q.  
A:Is  $10 \in O(1)$  true?

Yes

 $f(x) \geq O(g(x))$  is meaningless.

omega

$$f(x) = \Omega(g(x)) \text{ if } \lim_{x \rightarrow \infty} \left| \frac{f(x)}{g(x)} \right| > 0$$

Theorem

$$f(x) = O(g(x)) \text{ iff } g(x) = \Omega(f(x))$$

$$f(x) \leq O(g(x)) \text{ iff } g(x) \geq \Omega(f(x))$$

$$x^2 = \Omega(x)$$

$$2^x = \Omega(x^2)$$

$$\frac{x}{100} = \Omega(100x + 25)$$

(36)

classmate

Date \_\_\_\_\_  
Page 64

theta  $f(x) = \Theta(g(x))$  if  $\lim_{x \rightarrow \infty} \left| \frac{f(x)}{g(x)} \right| < \infty$

Theorem  $f(x) = \Theta(g(x))$  iff  $f(x) = O(g(x))$  &  
 $f(x) = \Omega(g(x))$

$$10x^3 - 20x + 1 \in \Theta(x^3)$$

$$\frac{x}{\ln(x)} \notin \Theta(x) \quad \left| \frac{x}{\ln(x)} \in O(x) \right.$$

$T(n) = \Theta(n^2)$  means  $T$  grows quadratically in  $n$ .

$$O \Rightarrow \leq$$

$$L \Rightarrow \geq$$

$$\Theta \Rightarrow =$$

$$o \Rightarrow < \quad (\leq \text{ not } =)$$

$$w \Rightarrow > \quad (\geq \text{ not } =)$$

(77)

classmate

Date \_\_\_\_\_

Page \_\_\_\_\_

65

little oh  $f(x) = o(g(x))$  iff  $\lim_{x \rightarrow \infty} \left| \frac{f(x)}{g(x)} \right| = 0$

little omega  $f(x) = \omega(g(x))$  iff  $\lim_{x \rightarrow \infty} \left| \frac{f(x)}{g(x)} \right| = \infty$

$$\frac{x}{\ln(x)} \in o(x)$$

$$\frac{x}{100} \notin o(x) \quad \frac{x}{100} \in \Theta(x)$$

$$x^2 \in \omega(o)$$

what not do  
example.