# Foundations of Full House Methodology

Daniel J. Eck[1], Shen Yan[1*]

1. Department of Statistics, University of Illinois Urbana-Champaign

**Abstract**

The Full House Methodology is an approach for computing era-adjusted or system-adjusted statistics that balances the size of the eligible population in each system with the component's performance via "versus their peers" in a specific system. We assume that each component has a latent aptitude that can be calculated from this balancing act provided the system inclusion mechanism for components of the system is known. Without making any assumptions about the distribution of the system's components, the components' distribution is well-established from the nonparametric probability distribution.

## 1 Introduction

Several academics have attempted to develop methods that can correct for era or system biases. The starting place for the answer to the problem of comparing players across eras can be found in the work of Stephen Jay Gould. Gould [1996] suggested that the entire distribution of achievement, or "full house of variation," is relevant when considering the achievements of baseball players across eras. This idea was also recognized as the starting plac e for an answer by Schell [2013] and Schell [2016].

Berry et al. [1999], Petersen et al. [2011], Schell [2013, 2016], Petersen and Penner [2020] provide several techniques to compare baseball players across eras, but they have fallen short in their objective and their results remain biased, as discussed in Eck [2020a] and Eck [2020b]. In short, these previous attempts do not incorporate the evolving talent pool in calculations, which is necessary for understanding the changing distributions of achievement over time. The result is that existing methodologies produce a disproportionate number of players who began their career before baseball was integrated players in the rankings of baseball's greatest of all time. This era of baseball did not include a crucial population of eligible players. Yu et al. [2016], Meyer [2021], Campbell [1990] also develop methods for comparing components in historical popularity, educational attainment, and evaluation of employees. But they have similar problems as stated before and fail to account for the evolving talent pool.

In some circumstances, such as test equating, it is not required to take into account an evolving talent pool. Andersson et al. [2013] offers a collaborative framework for test equating and builds the cumulative density function using the kernel method. Nevertheless,

*Daniel J. Eck (dje13@illinois.edu); email addresses: Shen Yan (shenyan3@illinois.edu)

they are unable to resolve the extrapolation issue in non-parametric estimation, and the distribution's tail is not given special consideration.

Our Full House methodology directly incorporates the quality of the talent pool as a central modeling component. It offers metrics for assessing the performance of components in various eras or systems. With the application of educational attainment simulation, our Full House methodology reveals that the most talented students could perform worse than the other students who attend schools with lower mean scores.

The next section establishes the model setting and specifies the assumptions in our Full House methodology. Section 3 considers an intriguing paradox that the component with more talent underperforms the component with less talent. We compute the probability of this paradox under different scenarios. Section 4 discusses the application of Full House methodology for the educational attainment simulation.

# 2 Model Setting

We now motivate the structure of the Full House Model. Let $N_i$ denote the size of population $i$. We will suppose the every individual $k = 1, \ldots, N_i$ has an underlying latent aptitude $X_{i,1}, \ldots, X_{i,N_i} \overset{\text{iid}}{\sim} F_X$. We denote $X_{i,(j)}$ as the $j$th ordered latent aptitude in population $i$. We define $g_i(\cdot, \cdot)$ as the system inclusion function, where $g_i(X_{i,j}, \mathbf{X}_{i,-j}) = 1$ indicates that individual $j$ is included in the $i$th system, and $g_i(X_{i,j}, \mathbf{X}_{i,-j}) = 0$ indicates that individual $j$ is not included in the $i$th system. Let $\mathbf{X}_{i,-j}$ be the vector of all latent aptitudes not including the component $j$.

We now suppose that an individual $j$ included in system $i$ has an observed statistic $Y_{i,j}$, $j = 1, \ldots, n_i$ and we suppose that $Y_{i,j}$ is a function of $X_{i,j}$ given the $X_{i,j}$ is included in the ith system, which is

$$Y_{i,j} = p_i(X_{i,j} | g_i(X_{i,j}, \mathbf{X}_{i,-j}) = 1)$$

where $p_i(\cdot)$ is an unknown continuous function.

Apparently $Y_{i,1}, \ldots, Y_{i,n_i} \overset{iid}{\sim} F_{Y_i}$, $F_{Y_i}$ is continuous, and $n_i < N_i$ is the number of individuals included in system $i$. We denote $Y_{i,(j)}$ as the $j$th ordered statistic for system $i$.

The key to the Full House Model is connecting latent aptitudes $X$ with observed statistics $Y$. To achieve this we form pairs $(Y_{i,(j)}, X_{i,(r(N_i,n_i,j))})$, $j = 1, \ldots, n_i$, where $r(N_i, n_i, k)$ is the index function that connects this latent aptitude to the $k$th components in the system $i$. The function $r(N_i, n_i, k)$ determines the relationship between the components in the system and the selected latent aptitudes in the population. As an example, $r(N_i, n_i, k) = N_i - n_i + k$, then

We now demonstrate how the Full House Model works in a parametric setting. Consider the pair $(Y_{i,(j)}, X_{i,(r(N_i,n_i,j))})$ and suppose that the distribution corresponding to $Y_{i,j}$ from the $i$th system is known to belong to a continuous distribution family.

In order to connect the $Y_{i,(j)}$ and $X_{i,(r(N_i,n_i,j))}$ and obtain an estimate of the underlying talents, we will make use of the following classical order statistics properties,

$$F_{Y_i}\left(Y_{i,(j)}\right) \sim U_{i,(j)}, F_{Y_{i,(j)}}\left(Y_{i,(j)}\right) \sim U_{i,j},$$

where $\sim$ means distributed as, $U_{i,j} \sim U(0,1)$, and $U_{i,(j)} \sim \text{Beta}(j, n_i + 1 - j)$.

We now connect the order statistics to the underlying talent distribution that comes from a population with $N_i \geq n_i$ observations when $F_X$ is known. This connection is established with

$$F^{-1}_{X_{i,(r(N_i,n_i,j))}}\left(F_{Y_{i,(j)}}\left(Y_{i,(j)}\right)\right) \sim F^{-1}_{X_{i,(r(N_i,n_i,j))}}\left(U_{i,j}\right) = X_{i,(r(N_i,n_i,j))}. \tag{1}$$

Similarly, we can obtain $Y$ from $X$,

$$F^{-1}_{Y_{i,(j)}}\left(F_{X_{i,(r(N_i,n_i,j))}}\left(X_{i,(r(N_i,n_i,j))}\right)\right) = Y_{i,(j)}, \tag{2}$$

# 3 Paradoxes

Consider two populations $i = 1, 2$ with sizes $N_1$ and $N_2$, and systems of sizes $n_1$ and $n_2$. We define a paradox (or a Gould paradox) when

$$X_{1,(N_1-k)} > X_{2,(N_2-j)} \quad \text{and} \quad Y_{1,(n_1-k)} < Y_{2,(n_2-j)}$$

both occur, where k, j $\in \mathbb{Z}$,. A central interest is evaluating the probability of such a paradox,

$$\text{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right). \tag{3}$$

As a simple illustrative example of some mechanics concerning paradoxes, suppose that $Y_{1,j} < Y_{2,k}$ a.s. Then (3) is equivalent to $\text{P}\left(X_{1,(N_1)} > X_{2,(N_2)}\right)$, and we have $\text{P}\left(X_{1,(N_1)} > X_{2,(N_2)}\right) \to 1$ as $N_1 \to \infty$. We motivate a convenient representation of (3) in the following proposition.

**Proposition 3.1.** *Consider two populations $i = 1, 2$ with sizes $N_1$ and $N_2$, and systems of sizes $n_1$ and $n_2$. Suppose that the system inclusion mechanism has the property that $g(X_{i,(N_i)}) = 1$, and that the Full House Model is of the form*

$$\begin{aligned} Y_{i,j} &\sim F_Y(y), \qquad i = 1, 2, \qquad j = 1, \ldots, n_1, \\ X_{i,j} &\sim F_X(x), \qquad i = 1, 2, \qquad j = 1, \ldots, N_i, \end{aligned}$$

*where $F_Y^{-1}(y)$ and $F_X^{-1}(x)$ exist. Then,*

$$\text{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \text{P}\left(X_{2,(N_2)} < X_{1,(N_1)} < h_{1,2}\left(X_{2,(N_2)}\right)\right), \tag{4}$$

*where $h_{i,j}(\bullet) = (f_i^{-1} \circ f_j)(\bullet)$, $f_i(\bullet) = F^{-1}_{Y_{i,(n_i)}}\left(F_{X_{i,(N_i)}}(\bullet)\right)$, and both $f$ and $h$ exist.*

*Proof.* If $f$ and $h$ exist, then we have

$$\begin{aligned} &\text{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\ &\quad = \text{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, f_1(X_{1,(N_1)}) < f_2(X_{2,(N_2)})\right) \\ &\quad = \text{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, f_1(X_{1,(N_1)}) < f_2(X_{2,(N_2)})\right) \\ &\quad = \text{P}\left(X_{2,(N_2)} < X_{1,(N_1)} < h_{1,2}\left(X_{2,(N_2)}\right)\right). \end{aligned}$$

Existance of $f_i$ follows directly from the existance of $F_Y^{-1}(y)$, and existance of $h_{i,j}$ follows directly from the existance of $f_i$ and the existance of $F_X^{-1}(x)$. □

**Lemma 3.2.** *Suppose that the system inclusion mechanism has the property that* $g(X_{i,(N_i)}) = 1$*, and that the Full House Model is of the form*

$$\begin{aligned} Y_{i,j} &\sim F_Y(y), \qquad i = 1, 2, \qquad j = 1, \ldots, n_1, \\ X_{i,j} &\sim F_X(x), \qquad i = 1, 2, \qquad j = 1, \ldots, N_i, \end{aligned}$$

*where* $F_Y^{-1}(y)$ *and* $F_X^{-1}(x)$ *exist. Then,*

$$\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \begin{cases} \frac{n_2}{n_1+n_2} - \frac{N_2}{N_1+N_2}, & \text{if } \frac{n_1}{N_1} \le \frac{n_2}{N_2} \\ 0, & \text{otherwise} \end{cases}.$$

*Proof.* We can use the convenient representation of (4) since the conditions of Proposition 3.1 hold. Thus, we have

$$\begin{aligned} &\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \mathrm{P}\left(X_{2,(N_2)} < X_{1,(N_1)} < h_{1,2}\left(X_{2,(N_2)}\right)\right) \\ &= \int_{-\infty}^{+\infty} \left[ \int_x^{h_{1,2}(x)} f_{X_{1,(N_1)}}(y)\, dy \right] f_{X_{2,(N_2)}}(x)\, dx \\ &= \int_{-\infty}^{+\infty} \left[ \int_x^{h_{1,2}(x)} N_1 (F_X(y))^{N_1-1} f_X(y) dy \right] N_2 (F_X(x))^{N_2-1} f_X(x)\, dx \\ &= \int_{-\infty}^{+\infty} \left[ \int_x^{F_X^{-1}(F_X(x)^{\frac{N_2}{N_1}\frac{n_1}{n_2}})} N_1 (F_X(y))^{N_1-1} f_X(y) dy \right] N_2 (F_X(x))^{N_2-1} f_X(x)\, dx \\ &= \int_{-\infty}^{+\infty} (F_X(x)^{N_2\frac{n_1}{n_2}} - F_X(x)^{N_1})\, N_2 (F_X(x))^{N_2-1} f_X(x)\, dx \\ &= \int_0^1 t^{\frac{n_1}{n_2}} - t^{\frac{N_1}{N_2}}\, dt \\ &= \frac{n_2}{n_1+n_2} - \frac{N_2}{N_1+N_2}, \end{aligned}$$

where

$$h_{1,2}(x) = F_X^{-1}((F_Y(F_Y^{-1}(F_X^{\frac{N_2}{n_2}})))^{\frac{n_1}{N_1}})(x) = F_X^{-1}(F_X^{\frac{N_2}{N_1}\frac{n_1}{n_2}})(x).$$

In order to ensure a positive probability, we assume $h_{1,2}(x) > x$, which indicates $\frac{n_1}{N_1} < \frac{n_2}{N_2}$ □

It is interesting to note that the paradox probability $\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \frac{n_2}{n_1+n_2}$ as $N_1 \to \infty$ under the setup of Lemma 3.2.

We verify Lemma 3.2 with a Monte Carlo simulation. In each simulation, we first randomly generate two groups of observation: $Y_1$ and $Y_2$ from the standard normal distribution, and the number of each group is given by $n_1$ and $n_2$. Based on our setting, we assume the

latent aptitudes follow the Pareto distribution with $\alpha = 1.16$. Then we obtain the latent aptitudes for each group and check whether the paradox happens. We repeat this process 1000 times and record the frequency of the paradox. 1000 Monte Carlo iterations of our simulation are performed and we compute the mean of the frequency.

| | Prob from Lemma 3.2 | Prob from simulation |
|---|---|---|
| $n_1 = 400, n_2 = 600, N_1 = 1e6, N_2 = 1e6$ | 0.1 | 0.10084 |
| $n_1 = 300, n_2 = 700, N_1 = 1e6, N_2 = 1e6$ | 0.2 | 0.200309 |
| $n_1 = 600, n_2 = 400, N_1 = 1e6, N_2 = 1e6$ | 0 | 0 |
| $n_1 = 700, n_2 = 300, N_1 = 1e6, N_2 = 1e6$ | 0 | 0 |

Table 1: Simulation result to validate the lemma about the probability of paradox

**Lemma 3.3.** *Suppose that the system inclusion mechanism has the property that* $g(X_{i,(N_i)}) = 1$*, and that the Full House Model is of the form*

$$\begin{aligned} Y_{i,j} &\sim F_Y(y), \qquad i = 1, 2, \qquad j = 1, \ldots, n_1, \\ X_{i,j} &\sim F_X(x), \qquad i = 1, 2, \qquad j = 1, \ldots, N_i, \end{aligned}$$

*where* $F_Y^{-1}(y)$ *and* $F_X^{-1}(x)$ *exist. Then,*

$$\mathrm{P}\left(X_{1,(1)} > X_{2,(1)}, Y_{1,(1)} < Y_{2,(1)}\right) = \begin{cases} \frac{N_2}{N_1+N_2} - \frac{n_2}{n_1+n_2}, & \text{if } \frac{n_1}{N_1} \geq \frac{n_2}{N_2} \\ 0, & \text{otherwise} \end{cases}.$$

*Proof.* We can use the convenient representation of (4) since the conditions of Proposition 3.1 hold. Thus, we have

$$\begin{aligned} &\mathrm{P}\left(X_{1,(1)} > X_{2,(1)}, Y_{1,(1)} < Y_{2,(1)}\right) = \mathrm{P}\left(X_{2,(1)} < X_{1,(1)} < h_{1,2}\left(X_{2,(1)}\right)\right) \\ &= \int_{-\infty}^{+\infty} \left[\int_x^{h_{1,2}(x)} f_{X_{1,(1)}}(y)\, dy\right] f_{X_{2,(1)}}(x)\, dx \\ &= \int_{-\infty}^{+\infty} \left[\int_x^{h_{1,2}(x)} N_1(1 - F_X(y))^{N_1-1} f_X(y) dy\right] N_2(1 - F_X(x))^{N_2-1} f_X(x)\, dx \\ &= \int_{-\infty}^{+\infty} \left[\int_x^{F_X^{-1}(1-(1-F_X(x))^{\frac{N_2}{N_1}\frac{n_1}{n_2}})} N_1(1 - F_X(y))^{N_1-1} f_X(y) dy\right] N_2(1 - F_X(x))^{N_2-1} f_X(x)\, dx \\ &= \int_{-\infty}^{+\infty} -((1 - F_X(x))^{N_2\frac{n_1}{n_2}} + (1 - F_X(x))^{N_1})\, N_2(1 - F_X(x))^{N_2-1} f_X(x)\, dx \\ &= \int_0^1 -(1-t)^{\frac{n_1}{n_2}} + (1-t)^{\frac{N_1}{N_2}}\, dt \\ &= \frac{N_2}{N_1+N_2} - \frac{n_2}{n_1+n_2}, \end{aligned}$$

where

$$h_{1,2}(x) = F_X^{-1}(1 - (F_Y(F_Y^{-1}((1 - F_X)^{\frac{N_2}{n_2}})))^{\frac{n_1}{N_1}})(x) = F_X^{-1}(1 - (1 - F_X)^{\frac{N_2}{N_1}\frac{n_1}{n_2}})(x).$$

□

We verify Lemma 3.3 with a Monte Carlo simulation. In each simulation, we first randomly generate two groups of observation: $Y_1$ and $Y_2$ from the standard normal distribution, and the number of each group is given by $n_1$ and $n_2$. Based on our setting, we assume the latent aptitudes follow the Pareto distribution with $\alpha = 1.16$. Then we obtain the latent aptitudes for each group and check whether the paradox happens. We repeat this process 1000 times and record the frequency of the paradox. 1000 Monte Carlo iterations of our simulation are performed and we compute the mean of the frequency.

| | Prob from Lemma 3.3 | Prob from simulation |
|---|---|---|
| $n_1 = 400, n_2 = 600, N_1 = 1e6, N_2 = 1e6$ | 0 | 0 |
| $n_1 = 300, n_2 = 700, N_1 = 1e6, N_2 = 1e6$ | 0 | 0 |
| $n_1 = 600, n_2 = 400, N_1 = 1e6, N_2 = 1e6$ | 0.1 | 0.1012 |
| $n_1 = 700, n_2 = 300, N_1 = 1e6, N_2 = 1e6$ | 0.2 | 0.2 |

Table 2: Simulation result to validate the lemma about the probability of paradox

**Lemma 3.4.** *If we assume*

$$\begin{aligned} Y_{i,j} &\sim F_Y(y), \qquad i = 1, 2, \qquad j = 1, \ldots, n_1, \\ X_{i,j} &\sim F_X(x), \qquad i = 1, 2, \qquad j = 1, \ldots, N_i, \end{aligned}$$

*where $F_Y^{-1}(y)$ and $F_X^{-1}(x)$ are assumed to exist. Then,*

$$\mathrm{P}\left(X_{1,(N_1-k)} > X_{2,(N_2-j)}, Y_{1,(n_1-k)} < Y_{2,(n_2-j)}\right) = .$$

*Proof.*

$$\begin{aligned} &\mathrm{P}\left(X_{1,(N_1-k)} > X_{2,(N_2-j)}, Y_{1,(n_1-k)} < Y_{2,(n_2-j)}\right) = \mathrm{P}\left(X_{2,(N_2-j)} < X_{1,(N_1-k)} < h_{1,2}\left(X_{2,(N_2-j)}\right)\right) \\ &= \int_{-\infty}^{+\infty} \left[ \int_x^{F_X^{-1} g_{N_1,k}^{-1} g_{n_1,k} g_{n_2,j}^{-1} g_{N_2,j} F_X(x)} f_{X_{1,(N_1-k)}}(y)\, dy \right] f_{X_{2,(N_2-j)}}(x)\, dx \\ &= \int_{-\infty}^{+\infty} \left[ \int_x^{F_X^{-1} g_{N_1,k}^{-1} g_{n_1,k} g_{n_2,j}^{-1} g_{N_2,j} F_X(x)} d\, g_{N_1,k}(F_X(y)) \right] f_{X_{2,(N_2-j)}}(x)\, dx \\ &= \int_{-\infty}^{+\infty} \left[ g_{n_1,k} g_{n_2,j}^{-1} g_{N_2,j} F_X(x) - g_{N_1,k} F_X(x) \right] f_{X_{2,(N_2-j)}}(x)\, dx \\ &= \int_{-\infty}^{+\infty} \left[ g_{n_1,k} g_{n_2,j}^{-1} g_{N_2,j} F_X(x) - g_{N_1,k} F_X(x) \right] d\, g_{N_2,j}(F_X(x)) \\ &= \int_0^1 g_{n_1,k} g_{n_2,j}^{-1}(t) - g_{N_1,k} g_{N_2,j}^{-1}(t)\, dt \end{aligned}$$

where

$$\begin{aligned} h_{1,2}(x) &= (g_1 \circ f_2)(x) \\ &= F_X^{-1} g_{N_1,k}^{-1} g_{n_1,k} g_{n_2,j}^{-1} g_{N_2,j} F_X(x) \\ g_{n,i}(x) &= I_x(n - i, i + 1) \end{aligned}$$

and $I_x(\alpha, \beta)$ is the regularized incompleted beta function. □

**Lemma 3.5.** *If we assume*

$$\begin{aligned} Y_{i,j} &\sim F_Y(y), \qquad i = 1, 2, \qquad j = 1, \ldots, n_1, \\ X_{i,j} &\sim F_X(x), \qquad i = 1, 2, \qquad j = 1, \ldots, N_i, \end{aligned}$$

*where $F_Y^{-1}(y)$ and $F_X^{-1}(x)$ are assumed to exist. Then,*

$$\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2-k)}, Y_{1,(n_1)} < Y_{2,(n_2-k)}\right) =$$

*Proof.*

$$\begin{aligned} &\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2-k)}, Y_{1,(n_1)} < Y_{2,(n_2-k)}\right) = \mathrm{P}\left(X_{2,(N_2-k)} < X_{1,(N_1)} < h_{1,2}\left(X_{2,(N_2-k)}\right)\right) \\ &= \int_0^1 g_{n_1,k} g_{n_2,0}^{-1}(t) - g_{N_1,k} g_{N_2,0}^{-1}(t)\, dt \\ &= \int_0^1 g_{n_1,k}(t^{\frac{1}{n_2}}) - g_{N_1,k}(t^{\frac{1}{N_2}})\, dt \\ &= \int_0^1 \int_0^{t^{\frac{1}{n_2}}} \frac{n_1!}{(n_1-k-1)!} x^{n_1-k-1}(1-x)^k dx - \int_0^{t^{\frac{1}{N_2}}} \frac{N_1!}{(N_1-k-1)!} x^{N_1-k-1}(1-x)^k dx\, dt \\ &= \int_0^1 \frac{n_1!}{(n_1-k-1)!} \sum_{i=0}^{k} \frac{\binom{k}{i}}{n_1-k+i} t^{\frac{n_1-k+i}{n_2}} - \frac{N_1!}{(N_1-k-1)!} \sum_{i=0}^{k} \frac{\binom{k}{i}}{N_1-k+i} t^{\frac{N_1-k+i}{N_2}} dt \\ &= \sum_{i=0}^{k} \frac{n_1!}{(n_1-k-1)!} \frac{\binom{k}{i}}{(n_1-k+i)} \frac{n_2}{(n_1-k+i+n_2)} - \frac{N_1!}{(N_1-k-1)!} \frac{\binom{k}{i}}{(N_1-k+i)} \frac{N_2}{(N_1-k+i+N_2)} \end{aligned}$$

□

**Definition 3.6.** *We will say that $Y_2$ is first order $\varepsilon$-weakly stochastically dominant over $Y_1$ if, for $\varepsilon > 0$, $\mathrm{P}(Y_2 \leq y) \leq \mathrm{P}(Y_1 \leq y) \leq \mathrm{P}(Y_2 \leq y) + \varepsilon$ for all $y$ and $\mathrm{P}(Y_2 \leq y) < \mathrm{P}(Y_1 \leq y) \leq \mathrm{P}(Y_2 \leq y) + \varepsilon$ for some $y$.*

**Lemma 3.7.** *Suppose that the system inclusion mechanism has the property that $g(X_{i,(N_i)}, \boldsymbol{X}_{i,-(N_i)}) = 1$, and that the Full House Model is of the form*

$$\begin{aligned} Y_{i,j} &\sim F_{Y_i}(y), \qquad i = 1, 2, \qquad j = 1, \ldots, n_i, \\ X_{i,j} &\sim F_X(x), \qquad i = 1, 2, \qquad j = 1, \ldots, N_i, \end{aligned}$$

*where $F_{Y_i}^{-1}(y)$ and $F_X^{-1}(x)$ exist. Further suppose that $Y_2$ is first order $\varepsilon$-weakly stochastically dominant over $Y_1$ for some $\varepsilon > 0$. Then,*

$$\begin{aligned} \frac{n_2}{n_1+n_2} - \frac{N_2}{N_1+N_2} &\leq \mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\ &\leq n_2 \sum_{i=0}^{n_1} \frac{\binom{n_1}{i}\varepsilon^i}{n_1+n_2-i} - \frac{N_2}{N_1+N_2} \end{aligned} \tag{5}$$

*Proof.* First, observe that the assumption of first order $\varepsilon$-weakly stochastically dominant gives

$$\left(F_{Y_{2,(n_2)}}(y)\right)^{n_1/n_2} \leq (F_{Y_1}(y))^{n_1} \leq (F_{Y_2}(y)+\varepsilon)^{n_1} = \left(\left(F_{Y_{2,(n_2)}}(y)\right)^{1/n_2}+\varepsilon\right)^{n_1},$$

and this implies that

$$F_X^{-1}\left(F_X^{\frac{N_2}{N_1}\frac{n_1}{n_2}}(X_{2,(N_2)})\right) \leq h_{1,2}(X_{2,(N_2)}) \leq F_{X_{1,(N_1)}}^{-1}\left(\left(\left(F_{X_{2,(N_2)}}(X_{2,(N_2)})\right)^{1/n_2}+\varepsilon\right)^{n_1}\right).$$

In combination with Lemma 3.2 and the proof of Lemma 3.2, the first inequality yields

$$\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) \geq \frac{n_2}{n_1+n_2} - \frac{N_2}{N_1+N_2}.$$

This verifies the first inequality of (5). We now verify the second inequality of (5). Now,

$$\begin{aligned}
&\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\
&\quad\leq \int_{-\infty}^{\infty}\left[\int_x^{F_{X_{1,(N_1)}}^{-1}\left(\left(\left(F_{X_{2,(N_2)}}(x)\right)^{1/n_2}+\varepsilon\right)^{n_1}\right)} f_{X_{1,(N_1)}}(y)dy\right] f_{X_{2,(N_2)}}(x)dx \\
&\quad= \int_{-\infty}^{\infty}\left[\left((F_X(x))^{N_2/n_2}+\varepsilon\right)^{n_1} - F_X(x)^{N_1}\right] N_2 F_X(x)^{N_2-1} f_X(x)dx \\
&\quad= \int_{-\infty}^{\infty}\left((F_X(x))^{N_2/n_2}+\varepsilon\right)^{n_1} N_2 F_X(x)^{N_2-1} f_X(x)dx - \frac{N_2}{N_1+N_2} \\
&\quad= \int_0^1 \left(t^{1/n_2}+\varepsilon\right)^{n_1} dt - \frac{N_2}{N_1+N_2} \\
&\quad= n_2 \sum_{i=0}^{n_1} \frac{\binom{n_1}{i}\varepsilon^i}{n_1+n_2-i} - \frac{N_2}{N_1+N_2}
\end{aligned}$$

□

**Example: Uniform distributions**. In this example we demonstrate a case where Lemma 3.7 applies, but we can explicitly calculate the paradox probability. Suppose that the system inclusion mechanism has the property that $g(X_{i,(N_i)}, \mathbf{X}_{i,-(N_i)}) = 1$, and we will consider the following Full House Model for two systems,

$$\begin{aligned}
Y_{1,j} &\sim U(0,1), \qquad j = 1,\ldots,n_1, \\
Y_{2,j} &\sim U(0,1+\delta), \qquad j = 1,\ldots,n_2, \\
X_{i,j} &\sim F_X(x), \qquad j = 1,\ldots,N_i,
\end{aligned}$$

where $\delta > 0$. We can pick $\varepsilon = \delta/(1+\delta)$ so that $Y_2$ is first order $\varepsilon$-weakly stochastically dominant over $Y_1$ with this choice of $\varepsilon$, and the conditions of Lemma 3.7 are satisfied. However,

our setup allows for direct calculation of the paradox probability. From Proposition 3.1 and a bit of manipulation we have that

$$
\begin{aligned}
&\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \mathrm{P}\left(X_{2,(N_2)} < X_{1,(N_1)} < h_{1,2}\left(X_{2,(N_2)}\right)\right) \\
&= \int_{-\infty}^{+\infty} \int_{x}^{h_{1,2}(x)} f_{X_{1,(N_1)}}(y)\ dy\ f_{X_{2,(N_2)}}(x)\ dx \\
&= \int_{-\infty}^{+\infty} \int_{x}^{h_{1,2}(x)} N_1 F_X(y)^{N_1-1}\ dy\ N_2 F_X(x)^{N_2-1}\ dx \\
&= \int_{-\infty}^{+\infty} \int_{x}^{F_X^{-1}(((1+\delta)F_X^{\frac{N_2}{n_2}})^{\frac{n_1}{N_1}})(x)} N_1 F_X(y)^{N_1-1}\ dy\ N_2 F_X(x)^{N_2-1}\ dx \\
&= \int_{-\infty}^{+\infty} (((1+\delta)F_X(x)^{\frac{N_2}{n_2}})^{n_1} - F_X(x)^{N_1})\ N_2 F_X(x)^{N_2-1}\ dx \\
&= \int_{0}^{1} (1+\delta)^{n_1} t^{\frac{n_1}{n_2}} - t^{\frac{N_1}{N_2}}\ dt \\
&= \frac{n_2}{n_1+n_2}(1+\delta)^{n_1} - \frac{N_2}{N_1+N_2}.
\end{aligned}
$$

Here we see that the probability of paradox is 1 for all $\delta \geq [2(n_1/n_2+1)]^{1/n_1} - 1$. When $n_1 \gg n_2$, then $[2(n_1/n_2+1)]^{1/n_1} - 1 \approx n_1^{n_1} - 1$ and we see that the probability of paradox is 1 for extremely small values of $\delta$.

We verify the uniform example with a Monte Carlo simulation. In each simulation, we first randomly generate two groups of observation: $Y_1$ and $Y_2$ from the standard normal distribution, and the number of each group is given by $n_1$ and $n_2$. Based on our setting, we assume the latent aptitudes follow the Pareto distribution with $\alpha = 1.16$ and the $\delta = 0.001$. Then we obtain the latent aptitudes for each group and check whether the paradox happens. We repeat this process 1000 times and record the frequency of the Gould paradox happening. 1000 Monte Carlo iterations of our simulation are performed and we calculate the average of the frequency divided by 1000 as the probability of the Gould paradox happening.

| | Prob from the result | Prob from simulation |
|---|---|---|
| $n_1 = 40, n_2 = 60, N_1 = 1e5, N_2 = 1e5$ | 0.124 | 0.122 |
| $n_1 = 30, n_2 = 70, N_1 = 1e5, N_2 = 1e5$ | 0.221 | 0.221 |
| $n_1 = 60, n_2 = 40, N_1 = 1e5, N_2 = 1e5$ | 0 | 0 |
| $n_1 = 70, n_2 = 30, N_1 = 1e5, N_2 = 1e5$ | 0 | 0.002 |

Table 3: Simulation result to validate the lemma about the probability of paradox

**Example: Normal distributions**

In this example, we demonstrate a case where Lemma 3.7 applies. Suppose that the system inclusion mechanism has the property that $g(X_{i,(N_i)}, \mathbf{X}_{i,-(N_i)}) = 1$, and we will consider

the following Full House Model for two systems,

$$\begin{aligned} Y_{1,j} &\sim N(\mu_1, \sigma_1^2), \qquad j = 1, \ldots, n_1, \\ Y_{2,j} &\sim N(\mu_2, \sigma_2^2), \qquad j = 1, \ldots, n_2, \\ X_{i,j} &\sim F_X(x), \qquad j = 1, \ldots, N_i, \end{aligned}$$

where $\mu_1 \leq \mu_2, \sigma_1 > 0$ and $\sigma_2 > 0$.

Supposed $\sigma_1 = \sigma_2 = \sigma$, it is obvious to see $\mathrm{P}(Y_2 \leq y) \leq \mathrm{P}(Y_1 \leq y)$ and we can also find the $\varepsilon$ such that $\mathrm{P}(Y_1 \leq y) \leq \mathrm{P}(Y_2 \leq y) + \varepsilon$

$$\begin{aligned} F_{Y_1}(y) - F_{Y_2}(y) &= \int_{-\infty}^{y} \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu_1)^2}{2\sigma^2}} dx - \int_{-\infty}^{y} \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu_2)^2}{2\sigma^2}} dx \\ &= \int_{y-\frac{\mu_2-\mu_1}{2}}^{y+\frac{\mu_2-\mu_1}{2}} \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\frac{\mu_2+\mu_1}{2})^2}{2\sigma^2}} dx \end{aligned}$$

Since $\frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\frac{\mu_2+\mu_1}{2})^2}{2\sigma^2}}$ is the density function for $N(\frac{\mu_2+\mu_1}{2}, \sigma^2)$, then the maximum value of the difference between these two cumulative distribution function of the normal distribution is achieved when $y = \frac{\mu_2+\mu_1}{2}$.

We can pick $\varepsilon = \int_{\mu_1}^{\mu_2} \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\frac{\mu_2+\mu_1}{2})^2}{2\sigma^2}} dx$ so that $Y_2$ is first order $\varepsilon$-weakly stochastically dominant over $Y_1$ with this choice of $\varepsilon$, and the conditions of Lemma 3.7 are satisfied.

From Proposition 3.1 and a bit of manipulation, we have that

$$
\begin{aligned}
&\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \mathrm{P}\left(X_{2,(N_2)} < X_{1,(N_1)} < h_{1,2}\left(X_{2,(N_2)}\right)\right) \\
&= \int_{-\infty}^{+\infty} \int_{x}^{h_{1,2}(x)} f_{X_{1,(N_1)}}(y)\ dy\ f_{X_{2,(N_2)}}(x)\ dx \\
&= \int_{-\infty}^{+\infty} \int_{x}^{h_{1,2}(x)} N_1 (F_X(y))^{N_1-1}\ dy\ N_2 F_X(x)^{N_2-1}\ dx \\
&= \int_{-\infty}^{+\infty} \int_{x}^{F_X^{-1}((\Phi(\Phi^{-1}(F_X^{\frac{N_2}{n_2}})+\frac{\mu_2-\mu_1}{\sigma}))^{\frac{n_1}{N_1}})(x)} N_1 F_X(y)^{N_1-1}\ dy\ N_2 F_X(x)^{N_2-1}\ dx \\
&= \int_{-\infty}^{+\infty} ((\Phi(\Phi^{-1}(F_X(x)^{\frac{N_2}{n_2}}) + \frac{\mu_2-\mu_1}{\sigma}))^{n_1} - F_X(x)^{N_1})\ N_2 F_X(x)^{N_2-1}\ dx \\
&= \int_{0}^{1} (\Phi(\Phi^{-1}(t^{\frac{1}{n_2}}) + \frac{\mu_2-\mu_1}{\sigma}))^{n_1} - t^{\frac{N_1}{N_2}})\ dt \\
&= \int_{0}^{1} (\Phi(\Phi^{-1}(t^{\frac{1}{n_2}}) + \frac{\mu_2-\mu_1}{\sigma}))^{n_1}\ dt - \frac{N_2}{N_1+N_2} \\
&= \int_{-\infty}^{+\infty} \Phi(x + \frac{\mu_2-\mu_1}{\sigma})^{n_1} (n_2 \Phi(x)^{n_2-1}\phi(x))\ dx - \frac{N_2}{N_1+N_2} \\
&= n_2 \int_{-\infty}^{+\infty} \mathbb{P}(Z_{1,1} - X \le \frac{\mu_2-\mu_1}{\sigma}, \ldots, Z_{1,n_1} - X \le \frac{\mu_2-\mu_1}{\sigma}, Z_{2,1} - X \le 0, \ldots, Z_{2,n_2-1} - X \le 0 | X = x)\phi(x \\
&- \frac{N_2}{N_1+N_2} \\
&= n_2 \mathbb{P}(Z_{1,1} - X \le \frac{\mu_2-\mu_1}{\sigma}, \ldots, Z_{1,n_1} - X \le \frac{\mu_2-\mu_1}{\sigma}, Z_{2,1} - X \le 0, \ldots, Z_{2,n_2-1} - X \le 0) - \frac{N_2}{N_1+N_2} \\
&= n_2 F_{\mathbf{W}}(\mathbf{w} = (\frac{\mu_2-\mu_1}{\sigma}, \ldots, \frac{\mu_2-\mu_1}{\sigma}, 0, \ldots, 0)) - \frac{N_2}{N_1+N_2}
\end{aligned}
$$

where

$$
\begin{aligned}
h_{1,2}(x) &= (g_1 \circ f_2)(x) \\
&= F_X^{-1}((F_{Y_2}(F_{Y_1}^{-1}(F_X^{\frac{N_2}{n_2}})))^{\frac{n_1}{N_1}})(x) \\
&= F_X^{-1}((\Phi(\Phi^{-1}(F_X^{\frac{N_2}{n_2}}) - \frac{\mu}{\sigma}))^{\frac{n_1}{N_1}})(x)
\end{aligned}
$$

$\mathbf{W} \sim \mathcal{N}_{(n_1+n_2-1)}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ with $(n_1+n_2-1)$-dimensional mean vector

$$\boldsymbol{\mu} = (0, \ldots, 0)$$

and $(n_1+n_2-1)\times(n_1+n_2-1)$ covariance matrix

$$\mathbf{\Sigma}=\begin{bmatrix}2 & 1 & \dots & 1\\ 1 & 2 & \ddots & \vdots\\ \vdots & \ddots & \ddots & 1\\ 1 & \dots & 1 & 2\end{bmatrix}$$

The first $n_1$ elements of $\mathbf{w}$ is $\frac{\mu_2-\mu_1}{\sigma}$ and the last $n_2-1$ elements are 0.

Then when $\sigma_1\neq\sigma_2$, we do not have a closed form for $\epsilon$ but we can calculate the probability of the paradox.

$$\begin{aligned}
&\mathrm{P}\left(X_{1,(N_1)}>X_{2,(N_2)},Y_{1,(n_1)}<Y_{2,(n_2)}\right)=\mathrm{P}\left(X_{2,(N_2)}<X_{1,(N_1)}<h_{1,2}\left(X_{2,(N_2)}\right)\right)\\
&=\int_{-\infty}^{+\infty}\int_{x}^{h_{1,2}(x)} f_{X_{1,(N_1)}}(y)\ dy\ f_{X_{2,(N_2)}}(x)\ dx\\
&=\int_{-\infty}^{+\infty}\int_{x}^{h_{1,2}(x)} N_1(F_X(y))^{N_1-1}\ dy\ N_2F_X(x)^{N_2-1}\ dx\\
&=\int_{-\infty}^{+\infty}\int_{x}^{F_X^{-1}(\Phi(\frac{\Phi^{-1}(F_X^{\frac{N_2}{n_2}})\sigma_2+\mu_2-\mu_1}{\sigma_1})^{\frac{n_1}{N_1}})(x)} N_1F_X(y)^{N_1-1}\ dy\ N_2F_X(x)^{N_2-1}\ dx\\
&=\int_{-\infty}^{+\infty}((\Phi(\frac{(\Phi^{-1}(F_X(x)^{\frac{N_2}{n_2}})\sigma_2+\mu_2-\mu_1}{\sigma_1}))^{n_1}-F_X(x)^{N_1})\ N_2F_X(x)^{N_2-1}\ dx\\
&=\int_0^1(\Phi(\frac{\Phi^{-1}(t^{\frac{1}{n_2}})\sigma_2+\mu_2-\mu_1}{\sigma_1}))^{n_1}-t^{\frac{N_1}{N_2}})\ dt\\
&=\int_0^1(\Phi(\frac{\Phi^{-1}(t^{\frac{1}{n_2}})\sigma_2+\mu_2-\mu_1}{\sigma_1}))^{n_1}\ dt-\frac{N_2}{N_1+N_2}\\
&=\int_{-\infty}^{+\infty}\Phi(\frac{\sigma_2x+\mu_2-\mu_1}{\sigma_1})^{n_1}(n_2\Phi(x)^{n_2-1}\phi(x))\ dx-\frac{N_2}{N_1+N_2}\\
&=n_2\int_{-\infty}^{+\infty}\mathbb{P}(Z_{1,1}-\frac{\sigma_2}{\sigma_1}X\leq\frac{\mu_2-\mu_1}{\sigma_1},\dots,Z_{1,n_1}-\frac{\sigma_2}{\sigma_1}X\leq-\frac{\mu_2-\mu_1}{\sigma_1},\\
&Z_{2,1}-X\leq0,\dots,Z_{2,n_2-1}-X\leq0|X=x)\phi(x)\ dx-\frac{N_2}{N_1+N_2}\\
&=n_2\mathbb{P}(Z_{1,1}-\frac{\sigma_2}{\sigma_1}X\leq\frac{\mu_2-\mu_1}{\sigma_1},\dots,Z_{1,n_1}-\frac{\sigma_2}{\sigma_1}X\leq-\frac{\mu_2-\mu_1}{\sigma_1},Z_{2,1}-X\leq0,\dots,Z_{2,n_2-1}-X\leq0)\\
&-\frac{N_2}{N_1+N_2}\\
&=n_2F_{\mathbf{W}}(\mathbf{w}=(\frac{\mu_2-\mu_1}{\sigma_1},\dots,\frac{\mu_2-\mu_1}{\sigma_1},0,\dots,0))-\frac{N_2}{N_1+N_2}
\end{aligned}$$

where

$$
\begin{aligned}
h_{1,2}(x) &= (g_1 \circ f_2)(x) \\
&= F_X^{-1}((F_{Y_2}(F_{Y_1}^{-1}(F_X^{\frac{N_2}{n_2}})))^{\frac{n_1}{N_1}})(x) \\
&= F_X^{-1}((\Phi(\frac{\Phi^{-1}(F_X^{\frac{N_2}{n_2}})\sigma_2 + \mu_2 - \mu_1}{\sigma_1}))^{\frac{n_1}{N_1}})(x)
\end{aligned}
$$

$\mathbf{W} \sim \mathcal{N}_{(n_1+n_2-1)}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ with $(n_1 + n_2 - 1)$-dimensional mean vector

$$\boldsymbol{\mu} = (0, \ldots, 0)$$

and $(n_1 + n_2 - 1) \times (n_1 + n_2 - 1)$ covariance matrix

$$\boldsymbol{\Sigma} = \begin{bmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{bmatrix}$$

$$\Sigma_{11} = \begin{bmatrix} \frac{\sigma_2^2}{\sigma_1^2} + 1 & \frac{\sigma_2^2}{\sigma_1^2} & \ldots & \frac{\sigma_2^2}{\sigma_1^2} \\ \frac{\sigma_2^2}{\sigma_1^2} & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \frac{\sigma_2^2}{\sigma_1^2} \\ \frac{\sigma_2^2}{\sigma_1^2} & \ldots & \frac{\sigma_2^2}{\sigma_1^2} & \frac{\sigma_2^2}{\sigma_1^2} + 1 \end{bmatrix}_{n_1 \times n_1}, \Sigma_{12}^T = \Sigma_{21} = \begin{bmatrix} \frac{\sigma_2}{\sigma_1} & \ldots & \ldots & \frac{\sigma_2}{\sigma_1} \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ \frac{\sigma_2}{\sigma_1} & \ldots & \ldots & \frac{\sigma_2}{\sigma_1} \end{bmatrix}_{(n_2-1) \times n_1}$$

$$\Sigma_{22} = \begin{bmatrix} 2 & 1 & \ldots & 1 \\ 1 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & 1 \\ 1 & \ldots & 1 & 2 \end{bmatrix}_{(n_2-1) \times (n_2-1)}$$

The first $n_1$ elements of $\mathbf{w}$ is $\frac{\mu_2 - \mu_1}{\sigma_1}$ and the last $n_2 - 1$ elements are 0.

We verify this example with a Monte Carlo simulation. In each simulation, we first randomly generate two groups of observation: $Y_1$ and $Y_2$ from the normal distribution with $\mu_1 = -1$ and $\mu_2 = 1$, and the number of each group is given by $n_1$ and $n_2$. Based on our setting, we assume the latent aptitudes follow the Pareto distribution with $\alpha = 1.16$, $N_1 = 1e5$ and $N_2 = 2e5$. Then we obtain the latent aptitudes for each group and check whether the paradox happens. We repeat this process 1000 times and record the frequency of the Gould paradox happening. 1000 Monte Carlo iterations of our simulation are performed and we calculate the average of the frequency divided by 1000 as the probability of the Gould paradox happening. Next, using the procedure in Mi et al. [2009], we calculate the probability of our outcome, which is the probability of a multivariate normal distribution.

| | Prob from simulation | Prob from multivariate normal distribution |
|---|---|---|
| $\sigma_1 = 2, \sigma_2 = 4, n_1 = 40, n_2 = 60$ | 0.331 | 0.333 |
| $\sigma_1 = 2, \sigma_2 = 4, n_1 = 30, n_2 = 70$ | 0.334 | 0.333 |
| $\sigma_1 = 2, \sigma_2 = 4, n_1 = 10, n_2 = 100$ | 0.332 | 0.333 |
| $\sigma_1 = 2, \sigma_2 = 4, n_1 = 100, n_2 = 10$ | 0.240 | 0.238 |
| $\sigma_1 = 1, \sigma_2 = 1, n_1 = 40, n_2 = 60$ | 0.332 | 0.331 |
| $\sigma_1 = 1, \sigma_2 = 1, n_1 = 30, n_2 = 70$ | 0.332 | 0.332 |
| $\sigma_1 = 1, \sigma_2 = 1, n_1 = 10, n_2 = 100$ | 0.334 | 0.333 |
| $\sigma_1 = 1, \sigma_2 = 1, n_1 = 100, n_2 = 10$ | 0.259 | 0.259 |

Table 4: Simulation result to validate the lemma about the probability of paradox

# 4 Application

## 4.1 Educational Attainment Simulation

Our Gould paradox is validated by simulating the process of students moving on to the next level of education. First, we randomly generate 1 million talents from Pareto distribution, which corresponds to the 1 million talents of primary school students. Then we randomly generate 2 thousand school indices for these 1 million talents, which represent the schools that the students will attend. Finally, we transform ordered talents corresponding to the eligible population to percentiles from the ordered statistics and convert percentiles to ordered scores from a normal distribution. The score distribution for each school follows the normal distribution with different parameters. The mean of the Normal distribution ranges from 60 to 80, which indicates the different levels of schools. Better schools have higher mean scores. The standard deviation for Figure 1 is 2 and for Figure 2 is 5.

50 Monte Carlo iterations of our simulation are performed. The tendencies of the most talented students in each of the 50 simulations are displayed in Figure 1 and Figure 2. The primary_index0 in the x-axis indicates the initial schools for the students with the highest talent. The levels of schools are represented by the index on the y-axis, which goes from 1 to 2000. High-achieving schools are a sign of superiority. The following are the schools that students will attend: primary_index1 through primary_index3. It is evident that even if gifted children begin their education at a lesser school, they may advance to a top institution within two to three stages.




Figure 1: The tendencies of the most talented students when sd = 2




Figure 2: The tendencies of the most talented students when sd = 5

### 4.2 Major League Baseball Example

Ted Williams and Bob Feller were two of the many great baseball players who fought in the Army during World War II(WWII) [1]. As a result, in the absence of these talented players, the players who continued in the league produced high-standard performances, such as Hal Newhouser. Now we are interested in the performance of those who played in the league during WWII when the players who served in the war would still be playing if World War II hadn't occurred.

Table 5 displays the performance of Hal Newhouser when the top talented players served in the Army or not. We assume only two pitchers would outperform Hal Newhouser. Additionally, we exclude the WWII correction for the MLB-eligible population when the top talented players did not serve in the Army.

| Player | Top talented players served in the Army | Estimated bWAR |
|---|---|---|
| Hal Newhouser | Yes | 10.35 |
| Hal Newhouser | No | 8.10 |

Table 5: the performance of Hal Newhouser when the top talented players served in the Army or not

## 5 Notes

Not sure where cross-contextual comparisons are needed. We can use the Full House Model to obtain estimated statistics in another context. Consider the the pair $(Y_{i,(j)}, X_{i,(r(N_i,n_i,j))})$ in the $i$th population. We first put the aptitude value $X_{i,(r(N_i,n_i,j))}$ obtained by (**??**) in a new population $m$ and reverse the process to obtain the hypothetical statistics in system $m$, which we will denote as $Y_{i,j,m}$. More formally, $Y_{i,j,m}$ are estimated as follows in the parametric context,

$$\widehat{Y}_{i,j,m} = F^{-1}_{Y_m}\left(F_{X_{m,(r(N_m,n_m,l_{i,j,m}))}}\left(\widehat{X}_{i,(r(N_i,n_i,j))}\right)|\hat{\theta}_m\right), \tag{6}$$

where $l_{i,j,m}$ is the rank of $\widehat{X}_{i,(r(N_i,n_i,j))}$ among the values

$$\left\{\widehat{X}_{i,(r(N_i,n_i,j))}, \widehat{X}_{m,(r(N_m,n_m,t))} : t = 1,\ldots,n_m\right\}.$$

The estimates (6) correspond to

$$Y_{i,j,m} = F^{-1}_{Y_m}\left(F_{X_{m,(r(N_m,n_m,l_{i,j,m}))}}\left(X_{i,(r(N_i,n_i,j))}\right) \mid \theta_m\right), \tag{7}$$

where $l_{i,j,m}$ is the rank of $X_{i,(r(N_i,n_i,j))}$ among the values

$$\left\{X_{i,(r(N_i,n_i,j))}, X_{m,(r(N_m,n_m,t))} : t = 1,\ldots,n_m\right\},$$

and $X_{i,(r(N_i,n_i,j))}$ is computed via (1).

[1] https://www.baseballinwartime.com/timeline/timeline_1942.htm

# References


Björn Andersson, Kenny Bränberg, and Marie Wiberg. Performing the kernel method of test equating with the package kequate. *Journal of Statistical Software*, 55:1–25, 2013.

Scott M Berry, C Shane Reese, and Patrick D Larkey. Bridging different eras in sports. *Journal of the American Statistical Association*, 94(447):661–676, 1999.

John P Campbell. Modeling the performance prediction problem in industrial and organizational psychology. 1990.

Daniel J Eck. Challenging nostalgia and performance metrics in baseball. *Chance*, 33(1): 16–25, 2020a.

Daniel J Eck. Challenging WAR and other statistics as era-adjustment tools. *FanGraphs*, 2020b. https://community.fangraphs.com/challenging-war-and-other-statistics-as-era-adjustment-tools/.

Stephen Jay Gould. *Full House: The Spread of Excellence from Plato to Darwin*. Harvard University Press, 1996.

Kirsten Meyer. Talents, abilities and educational justice. *Educational Philosophy and Theory*, 53(8):799–809, 2021.

Xuefei Mi, Tetsuhisa Miwa, and Torsten Hothorn. mvtnorm: New numerical algorithm for multivariate normal probabilities. *R Journal 1 (2009), Nr. 1*, 1(1):37–39, 2009.

Alexander M Petersen and Orion Penner. Renormalizing individual performance metrics for cultural heritage management of sports records. *Chaos, Solitons & Fractals*, 136:109821, 2020.

Alexander M Petersen, Orion Penner, and H Eugene Stanley. Methods for detrending success metrics to account for inflationary and deflationary factors. *The European Physical Journal B*, 79(1):67–78, 2011.

Michael J Schell. *Baseball's all-time best hitters*. Princeton University Press, 2013.

Michael J Schell. *Baseball's all-time best sluggers*. Princeton University Press, 2016.

Amy Zhao Yu, Shahar Ronen, Kevin Hu, Tiffany Lu, and César A Hidalgo. Pantheon 1.0, a manually verified dataset of globally famous biographies. *Scientific data*, 3(1):1–16, 2016.