A Formal Investigation of Paradoxes Between Latent Aptitude and Observed Achievement via Full House Methodology

Daniel J. Eck[1], Shen Yan[1*]

1. Department of Statistics, University of Illinois Urbana-Champaign

Abstract

1 Introduction

Comparisons of human performance across differing contexts, schools, regions, eras, institutions, and tasks, are central to education, labor economics, psychometrics, medicine, and sports analytics [Chetty et al., 2014, Iezzoni et al., 1997, Bartholomew et al., 2011, Bond and Fox, 2013, Berry et al., 1999, Imbens and Menzel, 2018, Holland and Wainer, 2012, Yan et al., 2025a]. A recurring challenge is that an individual in one context may have higher latent aptitude yet lower observed achievement than a counterpart elsewhere. We term this the aptitude-achievement paradox and develop a statistical framework to quantify its probability across contexts. Formally, let individual j in context i have an observed $\mathbb{R}$-valued statistic $Y_{i,j}$ and an underlying latent $\mathbb{R}$-valued aptitude $X_{i,j}$, and let individual n in context m has an observed $\mathbb{R}$-valued statistic $Y_{m,n}$,and an underlying latent $\mathbb{R}$-valued aptitude $X_{m,n}$. The probability of aptitude-achievement paradox is

$$\mathrm{P}\left(X_{i,j} > X_{m,n}, Y_{i,j} < Y_{m,n}\right)$$

The paradox is structurally impossible if a single, strictly increasing, context-invariant measurement map aptitude to achievement and if selection into tasks is ignorable. In practice, however, both measurement heterogeneity (non-invariance, differential item functioning) and non-ignorable selection (who attempts which tasks, faces which opponents, receives which treatment) are pervasive. These features can reverse cross-context rankings even when the latent ranking is unambiguous, echoing classic reversal phenomena such as Simpson's [Simpson, 1951] and Lord's paradoxes [Lord, 1967] while implicating modern concerns about fairness, transportability, and domain shift. What practitioners lack is a principled way to quantify how often such reversals should occur under an explicit model of aptitude, measurement, and selection.

*Daniel J. Eck (dje13@illinois.edu); email addresses: Shen Yan (shenyan3@illinois.edu)

A large class of methods measures how often observed orderings disagree with a target ordering-precisely the phenomenon at the heart of our cross-context paradox. At the most elementary level, discordance statistics summarize the share of ordered pairs whose observed outcomes align (or misalign) with a reference score or latent ordering. Classic measures include Kendall's τ [Kendall, 1938] and Somers'D [Somers, 1962], which compute differences between the proportions of concordant and discordant pairs and offer adjustments for ties; Somers'D is asymmetric and treats one variable (e.g., a score) as the predictor of the other (e.g., an outcome), making it particularly natural for "does the score rank the outcomes correctly?" questions. These rank-based indices provide assumption-lean, effect-size-like summaries of reversal risk. However, because these indices operate on observed rankings, they neither place individuals on a common latent scale nor disentangle measurement non-invariance or selection, motivating models that make the measurement process explicit.

Traditional latent trait models [Bartholomew et al., 2011] focus on estimating underlying traits from observed categorical data. One of the example of traditional latent trait models, the Rasch model [Bond and Fox, 2013] ensures that comparisons of individuals' abilities are independent of the specific items administered, and the Rasch model expect no differential item functioning (DIF) [Holland and Wainer, 2012], which means an item behaves differently for two (or more) groups of examinees who have the same underlying ability on the target construct. But it does not explicitly leverage the source population structure. Likewise, the bayesian hierarchical modeling [Berry et al., 1999] with the application in comparison of individuals' latent aptitudes typically captures population-level variation through seasonal random effects. In parallel, the causal bootstrap [Imbens and Menzel, 2018] highlights that valid causal inference similarly requires situating the observed sample within large source population to properly account for sampling uncertainty.

Full House Modeling offers a novel approach for comparing individuals across diverse contexts, initially developed through the analysis of baseball players across different eras [Yan et al., 2025a]. This analytical framework gained further historical significance [Burgos Jr et al., 2025], which was subsequently explored and presented in a separate study. These analyses are transforming baseball analytics and are increasingly recognized within the broader baseball community. The applicability of Full House Modeling was further extended to investigate the influence of historical figures [Cohen et al., 2025].

In this article, we formalize the Full House Methodology and investigate the probabilities of observing paradoxes between latent aptitude and observed achievement in comparative analysis of individuals across contexts. The aptitude-achievement paradox arises when an individual from one context demonstrates higher latent aptitude but simultaneously lower observed achievement than an individual from another context. We provide explicit probabilities for these paradoxes in two scenarios: first, when observed achievement is independent and identically distributed (i.i.d.), and second, when observed statistics across contexts follow different normal distributions but remain independent. We also explore bounds for these paradoxes in more generalized settings and demonstrate that when observed achievement is normally distributed, paradox probabilities can be reliably estimated using a plug-in method. Based on the sensitivity analysis of the paradox probability, it depends on population size ratios not absolute population sizes.

2 Model Setting

We now motivate the structure of the Full House Model. Let N_i denote the size of population i. We will suppose the every individual $k = 1, \ldots, N_i$ has an underlying latent $\mathbb{R}$-valued aptitude $X_{i,1}, \ldots, X_{i,N_i} \overset{iid}{\sim} F_X$. Let $\mathbf{X}_{i,-j}$ be the vector of all latent aptitudes not including the component j. We then define an indicator function $1_i\left(X_{i,j}, \mathbf{X}_{i,-j}\right)$ as a determined and known system inclusion function for population i where a 1 indicates inclusion and a 0 indicates exclusion. We now suppose that an individual j included in system i has an observed $\mathbb{R}$-valued statistic $Y_{i,j} \overset{iid}{\sim} F_{Y_i}$, $j = 1, \ldots, n_i$, $n_i \leq N_i$. As we will see later, the observed $Y_{i,j}$ values are one-to-one realizations of $X_{i,j}$, provided that $1_i\left(X_{i,j}, \mathbf{X}_{i,-j}\right) = 1$. Therefore $Y_{i,1}, \ldots, Y_{i,n_i} \overset{iid}{\sim} F_{Y_i}$.

Let $X_{i,(j)}$ and $Y_{i,(j)}$ denote the jth order statistic for system i. Here, larger numbers will correspond to a notion of "better." Thus $X_{i,(j)} > X_{i,(k)}$ whenever $j > k$, for example. We now form (Y, X) pairs of random variables for each individual included in the system. These pairs are of the form $(Y_{i,(j)}, X_{i,(r_{i,j})})$ where $r_{i,j}$ indexes the aptitude of the jth highest value observed from the system. As an example, let $1_i\left(X_{i,j}, \mathbf{X}_{i,-j}\right) = 1(X_{i,j} \geq X_{i,(N_i - n_i + 1)})$, then the $(Y_{i,(j)}, X_{i,(r_{i,j})})$ pairs take the form $(Y_{i,(j)}, X_{i,(N_i - n_i + j)})$ for $j = 1, \ldots, n_i$.

The key Full House Modeling is the explicit connection between the elements forming the $(Y_{i,(j)}, X_{i,(r_{i,j})})$ pairs. Full House Modeling states this connection as

$$\begin{aligned} Y_{i,(j)} &= F^{-1}_{Y_{i,(j)}} \circ F_{X_{i,(r_{i,j})}}\left(X_{i,(r_{i,j})}\right), \\ X_{i,(r_{i,j})} &= F^{-1}_{X_{i,(r_{i,j})}} \circ F_{Y_{i,(j)}}\left(Y_{i,(j)}\right), \end{aligned} \tag{1}$$

where $1_i(X_{i,(r_{i,j})}, \mathbf{X}_{i,-(r_{i,j})}) = 1$, and both F_{Y_i} and F_X are continuous and invertible. We summarise Full House Modeling in the definition below.

Definition 2.1 (Full House Model)**.** *Consider populations with sizes N_i and system sizes n_i, $N_i \geq n_i$. Let a system inclusion function $1_i(\cdot, \cdot)$ be specified. Let $X_{i,j} \overset{iid}{\sim} F_X$, $j = 1, \ldots, N_i$. Let $Y_{i,j} \overset{iid}{\sim} F_{Y_i}$, $j = 1, \ldots, n_i$. Form pairs $(Y_{i,(j)}, X_{i,(r_{i,j})})$, $j = 1, \ldots, n_i$ for all individuals such that $1_i(X_{i,(r_{i,j})}, \boldsymbol{X}_{i,-(r_{i,j})}) = 1$. Then the Full House Model generates $Y_{i,(j)}$ values following (1) where F_{Y_i} and F_X are continuous and invertible.*

Note that Definition 2.1 fixes F_X but allows F_{Y_i} to vary across systems.

Remarks:

1. The definition of a Full House Model immediately implies structural missingness. Every $X_{i,j}$ such that $1_i(X_{i,j}, \mathbf{X}_{i,-j}) = 0$ will not have a corresponding Y value. In the following sections we will consider probabilities involving both X and Y quantities. To handle this structural missingness, we will suppose that $Y_{i,j}$ values that are not observed are not P-measurable.

3 Paradoxes Under Identical Distributions

In this section we will provide explicit aptitude-achievement paradox probability results in a Full House Modeling setting involving two populations with the additional specification

that $F_{Y_i} = F_Y$, $i = 1, 2$ in Definition 2.1. The following definition formally defines the notion of paradox used throughout this article.

Definition 3.1. *Consider the Full House Model in Definition 2.1 with two populations denoted by $i = 1, 2$. We say a aptitude-achievement paradox has occurred when both $X_{1,(r_{1,j})} > X_{2,(r_{2,k})}$ and $Y_{1,(j)} < Y_{2,(k)}$ have occurred. The aptitude-achievement paradox probability is defined as*

$$\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right). \tag{2}$$

As a simple illustrative example of the above, let $r_{i,t} = N_i - n_i + t$ and set $j = n_1$ and $k = n_2$, and further suppose that $Y_{1,j} < Y_{2,k}$ a.s. Then the aptitude-achievement paradox probability (2) is equivalent to $\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}\right)$, and we have $\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}\right) \to 1$ when $N_1 \to \infty$. We now introduce a convenient representation which allows the paradox probability to be expressed in terms of the latent aptitudes alone.

Lemma 3.2. *Consider the Full House Model in Definition 2.1 with two populations, where $F_{Y_i} = F_Y$ for $i = 1, 2$. Then,*

$$\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right) = \mathrm{P}\left(X_{2,(r_{2,k})} < X_{1,(r_{1,j})} < h_{j,k}\left(X_{2,(r_{2,k})}\right)\right), \tag{3}$$

$$\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right) = \mathrm{P}\left(g_{j,k}\left(Y_{2,(k)}\right) < Y_{1,(j)} < Y_{2,(k)}\right) \tag{4}$$

where $h_{j,k}(\cdot) = F^{-1}_{X_{1,(r_{1,j})}} \circ F_{Y_{1,(j)}} \circ F^{-1}_{Y_{2,(k)}} \circ F_{X_{2,(r_{2,k})}}(\cdot)$, $g_{j,k}(\cdot) = F^{-1}_{Y_{1,(j)}} \circ F_{X_{1,(r_{1,j})}} \circ F^{-1}_{X_{2,(r_{2,k})}} \circ F_{Y_{2,(k)}}(\cdot)$.

Proof. Define $f_{1,j}(\cdot) = F^{-1}_{Y_{1,(j)}} \circ F_{X_{1,(r_{1,j})}}(\cdot)$ and $f_{2,k}(\cdot) = F^{-1}_{Y_{2,(k)}} \circ F_{X_{2,(r_{2,k})}}(\cdot)$. Then we have

$$\begin{aligned}
&\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right) \\
&\quad = \mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}), f_{1,j}(X_{1,(r_{1,j})}) < f_{2,k}(X_{2,(r_{2,k})})\right) \\
&\quad = \mathrm{P}\left(X_{2,(r_{2,k})} < X_{1,(r_{1,j})} < h_{j,k}\left(X_{2,(r_{2,k})}\right)\right),
\end{aligned}$$

where $h_{j,k}(\cdot) = (f^{-1}_{1,j} \circ f_{2,k})(\cdot)$. A similar technique can be applied to the second equation to obtain the result. □

Our first formal result provides conditions that allow for an explicit aptitude-achievement paradox probability result. Those conditions require the following definition.

Definition 3.3. *Let $\alpha, \beta > 0$. Then the regularized incomplete beta function is defined as*

$$I_{\alpha,\beta}(x) = \frac{1}{B(\alpha,\beta)} \int_0^x u^{\alpha-1}(1-u)^{\beta-1} du,$$

where $B(\alpha,\beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$ is the beta coefficient.

Note that the regularized incomplete beta function is the cumulative distribution function of a Beta(α, β) distribution.

Theorem 3.4. *Consider the Full House Model in Definition 2.1 with two populations, where* $F_{Y_i} = F_Y$ *for* $i = 1, 2$. *Denote by* I *the regularized incomplete beta function. Further assume that*

$$I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1}(t) - I_{r_{1,j},N_1-r_{1,j}+1} \circ I^{-1}_{r_{2,k},N_2-r_{2,k}+1}(t) \geq 0, \qquad \text{for all } t \in (0,1). \tag{5}$$

Under these assumptions,

$$\begin{aligned}
&\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right) \\
&= \sum_{p=j}^{n_1} \binom{n_1}{p} \frac{B(k+p, n_2-n_1-p+k+1)}{B(k, n_2-k+1)} - \sum_{p=r_{1,j}}^{N_1} \binom{N_1}{p} \frac{B(r_{2,k}+p, N_2-N_1-p+r_{2,k}+1)}{B(r_{2,k}, N_2-r_{2,k}+1)}.
\end{aligned}$$

Remarks:

1. The result of Theorem 3.4 is distribution free. It only depends on the sample sizes n_i, N_i, $i = 1, 2$, the system inclusion function, and the indexes of the order statistics under comparison. The distribution on Y is eliminated by Lemma 3.2. To see this, consider the distribution $F_{Y_{(j)}}$ for some $j = 1, \ldots, n_1$. The random variable $Y_{(j)} = F_Y^{-1} \circ U_{(j)}$ where $U_{(j)} \sim \text{Beta}(j, n_1 - j + 1)$. Thus, $F_{Y_{(j)}}(y) = I_{j,n_1-j+1} \circ F_Y(y)$. With this in mind, from Lemma 3.2 we have that

$$\begin{aligned}
h_{j,k}(x) &= F^{-1}_{X_{1,(r_{1,j})}} \circ F_{Y_{1,(j)}} \circ F^{-1}_{Y_{2,(k)}} \circ F_{X_{2,(r_{2,k})}}(x) \\
&= F^{-1}_{X_{1,(r_{1,j})}} \circ I_{j,n_1-j+1} \circ \left(F_Y \circ F_Y^{-1}\right) \circ I^{-1}_{k,n_2-k+1} \circ F_{X_{2,(r_{2,k})}}(x) \\
&= F^{-1}_{X_{1,(r_{1,j})}} \circ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1} \circ F_{X_{2,(r_{2,k})}}(x).
\end{aligned}$$

   The above is the upper bound of an integral with respect to Xs in the proof of Theorem 3.4. The influence of the distributional specification on Y does not influence the calculation of the aptitude-achievement paradox probability result in Theorem 3.4.

2. The distribution on X is eliminated when calculating the aptitude-achievement paradox probability (2) in its Lemma 3.2 representation, $\mathrm{P}\left(X_{2,(r_{2,k})} < X_{1,(r_{1,j})} < h_{j,k}\left(X_{2,(r_{2,k})}\right)\right)$. These calculations involve double integrals with respect to densities for $F_{X_{1,(r_{1,j})}}$ and $F_{X_{2,(r_{2,k})}}$. One integral evaluation and a reparameterization remove the specification of F_X from the probability calculation as seen in the proof of Theorem 3.4.

3. The steps mentioned in the last sentence of the previous remark allow the probability calculation for $\mathrm{P}\left(X_{2,(r_{2,k})} < X_{1,(r_{1,j})} < h_{j,k}\left(X_{2,(r_{2,k})}\right)\right)$ to simplify to

$$\int_0^1 \left\{ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1}(t) - I_{r_{1,j},N_1-r_{1,j}+1} \circ I^{-1}_{r_{2,k},N_2-r_{2,k}+1}(t) \right\}^+ dt.$$

   The technical condition (5) ensures that this integral defines a non-negative probability in strong terms. We demonstrate with analytic and numeric arguments when (5) is satisfied and the results of Theorem 3.4 therefore hold.

Proof. First, Lemma 3.2 allows us to write

$$\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right) = \mathrm{P}\left(X_{2,(r_{2,k})} < X_{1,(r_{1,j})} < h_{j,k}\left(X_{2,(r_{2,k})}\right)\right),$$

where $h_{j,k}(\cdot) = F^{-1}_{X_{1,(r_{1,j})}} \circ F_{Y_{1,(j)}} \circ F^{-1}_{Y_{2,(k)}} \circ F_{X_{2,(r_{2,k})}}(\cdot)$. We can rewrite $h_{j,k}(\cdot)$ into a form that will allow for specific computation involving Beta distributions. To see this, consider the distribution $F_{Y_{(j)}}$ for some $j = 1, \ldots, n_1$. The random variable $Y_{(j)} = F_Y^{-1} \circ U_{(j)}$ where $U_{(j)} \sim \mathrm{Beta}(j, n_1 - j + 1)$.

$$\begin{aligned}
&\mathrm{P}\left(X_{2,(r_{2,k})} < X_{1,(r_{1,j})} < h_{j,k}\left(X_{2,(r_{2,k})}\right)\right) \\
&= \int_{-\infty}^{+\infty} \left\{ \int_x^{F_X^{-1} \circ I^{-1}_{r_{1,j},N_1-r_{1,j}+1} \circ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1} \circ I_{r_{2,k},N_2-r_{2,k}+1} \circ F_X(x)} f_{X_{1,(r_{1,j})}}(y)\, dy \right\} f_{X_{2,(r_{2,k})}}(x)\, dx \\
&= \int_{-\infty}^{+\infty} \left\{ \int_x^{F_X^{-1} \circ I^{-1}_{r_{1,j},N_1-r_{1,j}+1} \circ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1} \circ I_{r_{2,k},N_2-r_{2,k}+1} \circ F_X(x)} d\, I_{r_{1,j},N_1-r_{1,j}+1}(F_X(y)) \right\} f_{X_{2,(r_{2,k})}}(x)\, dx \\
&= \int_{-\infty}^{+\infty} \left\{ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1} \circ I_{r_{2,k},N_2-r_{2,k}+1} \circ F_X(x) - I_{r_{1,j},N_1-r_{1,j}+1} \circ F_X(x) \right\}^+ f_{X_{2,(r_{2,k})}}(x)\, dx \\
&= \int_{-\infty}^{+\infty} \left\{ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1} \circ I_{r_{2,k},N_2-r_{2,k}+1} \circ F_X(x) - I_{r_{1,j},N_1-r_{1,j}+1} \circ F_X(x) \right\}^+ d\, I_{r_{2,k},N_2-r_{2,k}+1}(F_X(x)) \\
&= \int_0^1 \left\{ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1}(t) - I_{r_{1,j},N_1-r_{1,j}+1} \circ I^{-1}_{r_{2,k},N_2-r_{2,k}+1}(t) \right\}^+ dt.
\end{aligned}$$

Condition (5) states that

$$\begin{aligned}
&\int_0^1 \left\{ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1}(t) - I_{r_{1,j},N_1-r_{1,j}+1} \circ I^{-1}_{r_{2,k},N_2-r_{2,k}+1}(t) \right\}^+ dt \\
&= \int_0^1 \left\{ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1}(t) - I_{r_{1,j},N_1-r_{1,j}+1} \circ I^{-1}_{r_{2,k},N_2-r_{2,k}+1}(t) \right\} dt.
\end{aligned}$$

Thus,

$$\begin{aligned}
&\int_0^1 \left\{ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1}(t) - I_{r_{1,j},N_1-r_{1,j}+1} \circ I^{-1}_{r_{2,k},N_2-r_{2,k}+1}(t) \right\}^+ dt \\
&\quad = \int_0^1 I_{j,n_1-j+1}(x)\, dI_{k,n_2-k+1}(x) - \int_0^1 I_{r_{1,j},N_1-r_{1,j}+1}(x)\, dI_{r_{2,k},N_2-r_{2,k}+1}(x) \\
&\quad = \int_0^1 I_x(j, n_1 - j + 1) \frac{x^{k-1}(1-x)^{n_2-k}}{B(k, n_2 - k + 1)}\, dx \\
&\qquad - \int_0^1 I_x(r_{1,j}, N_1 - r_{1,j} + 1) \frac{x^{r_{2,k}-1}(1-x)^{N_2-r_{2,k}}}{B(k, N_2 - k + 1)}\, dx \\
&\quad = \int_0^1 \sum_{p=j}^{n_1} \binom{n_1}{p} x^p (1-x)^{n_1-p} \frac{x^{k-1}(1-x)^{n_2-k}}{B(k, n_2 - k + 1)}\, dx \\
&\qquad - \int_0^1 \sum_{p=r_{1,j}}^{N_1} \binom{N_1}{p} x^p (1-x)^{N_1-p} \frac{x^{r_{2,k}-1}(1-x)^{N_2-r_{2,k}}}{B(r_{2,k}, N_2 - r_{2,k} + 1)}\, dx
\end{aligned}$$

$$= \sum_{p=j}^{n_1} \binom{n_1}{p} \frac{B(k+p, n_2+n_1-p-k+1)}{B(j, n_2-k+1)}$$
$$- \sum_{p=r_{1,j}}^{N_1} \binom{N_1}{p} \frac{B(r_{2,k}+p, N_2+N_1-p-r_{2,k}+1)}{B(r_{2,k}, N_2-r_{2,k}+1)}.$$

□

Often it is of interest to compare the individuals with the highest aptitudes in their respective populations. Below we state the aptitude-achievement paradox probability in such setting using a Full House Model with two populations with the specification that $F_{Y_1} = F_Y$ for $i = 1, 2$.

Corollary 3.4.1. *Suppose that the conditions of Theorem 3.4 hold. Suppose that* $1_1(X_{1,(N_1)}, \boldsymbol{X}_{1,-(N_1)}) = 1_2(X_{2,(N_2)}, \boldsymbol{X}_{2,-(N_2)}) = 1$. *Then,*

$$\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \begin{cases} \frac{n_2}{n_1+n_2} - \frac{N_2}{N_1+N_2}, & \text{if } \frac{n_1}{N_1} \leq \frac{n_2}{N_2} \\ 0, & \text{otherwise} \end{cases}.$$

A similar result holds for the individuals with the lowest latent aptitudes in the population.

Corollary 3.4.2. *Suppose that the conditions of Theorem 3.4 hold. Suppose that* $1_1(X_{1,(1)}, \boldsymbol{X}_{1,-(1)}) = 1_2(X_{2,(1)}, \boldsymbol{X}_{2,-(1)}) = 1$. *Then,*

$$\mathrm{P}\left(X_{1,(1)} > X_{2,(1)}, Y_{1,(1)} < Y_{2,(1)}\right) = \begin{cases} \frac{N_2}{N_1+N_2} - \frac{n_2}{n_1+n_2}, & \text{if } \frac{n_1}{N_1} \geq \frac{n_2}{N_2} \\ 0, & \text{otherwise} \end{cases}.$$

We now demonstrate a setting in which technical condition (5) holds. This setting is based on a configuration of n_j, N_j, $j = 1, 2$ under the assumption that the systems are populated with individuals from the source populations who have the highest aptitudes.

Proposition 3.5. *Consider the Full House Model in Definition 2.1 with two populations, where* $F_{Y_i} = F_Y$ *for* $i = 1, 2$, *and let the system inclusion function be* $1_i(X_{i,j} \geq X_{i,(N_i-n_i+1)})$. *Denote by* I *the regularized incomplete beta function. Suppose that* $\frac{n_1}{n_2} \leq 1$ *and* $\frac{N_2}{N_1} \leq 1$, *and set* $k = j$. *Under these assumptions, we conclude that technical condition (5) holds and the conclusion of Theorem 3.4 applies.*

Proof. Specification of $1_i(X_{i,j} \geq X_{i,(N_i-n_i+1)})$ implies that $r_{i,j} = N_i + 1 - j$. When either $n_1 = N_1$ and $n_2 = N_2$ or $n_1 = n_2$ and $N_1 = N_2$ we have, for all $x \in (0, 1)$, that

$$I_{N_2-j,j+1} \circ I^{-1}_{N_1-j,j+1} \circ I_{n_1-j,j+1} \circ I^{-1}_{n_2-j,j+1}(t) \geq t.$$

The left hand side of the above equation increases whenever n_1 or N_1 are increased [Askitis, 2021], or n_2 or N_1 are decreased. Therefore, $I_{N_2-j,j+1} \circ I^{-1}_{N_1-j,j+1} \circ I_{n_1-j,j+1} \circ I^{-1}_{n_2-j,j+1}(t) \geq t$ whenever $\frac{n_1}{n_2} \leq 1$ and $\frac{N_2}{N_1} \leq 1$. Observe that

$$\begin{aligned} & I_{N_2-j,j+1} \circ I^{-1}_{N_1-j,j+1} \circ I_{n_1-j,j+1} \circ I^{-1}_{n_2-j,j+1}(t) \geq t, \\ \implies & I_{n_1-j,j+1} \circ I^{-1}_{n_2-j,j+1}(t) \geq I_{N_1-j,j+1} \circ I^{-1}_{N_2-j,j+1}(t), \\ \implies & I_{n_1-j,j+1} \circ I^{-1}_{n_2-j,j+1}(x) - I_{N_1-j,j+1} \circ I^{-1}_{N_2-j,j+1}(t) \geq t. \end{aligned}$$

Technical condition (5) holds and the conclusion of Theorem 3.4 applies. □

4 Paradox Under Different Distribution

In this section we will provide aptitude-achievement paradox probability results in a Full House Modeling setting involving two populations when we relax the specification that $F_{Y_1} = F_{Y_2}$ in Definition 2.1. We first provide bounds for aptitude-achievement paradox probability in general settings. We then provide explicit aptitude-achievement paradox probability results when Y_1 and Y_2 are normally distributed with different means and variances.

4.1 General bounds

The general bounds we derive require a relaxed form of first-order stochastic dominance that constrains the distributions of Y_1 and Y_2 are 'close' to each other. More formally, we introduce first-order ε-weak stochastic dominance, which is necessary for our aptitude-achievement paradox probability results.

Definition 4.1. *We will say that Y_2 is first order ε-weakly stochastically dominant over Y_1 if, for $\varepsilon > 0$, $\mathrm{P}(Y_2 \leq y) \leq \mathrm{P}(Y_1 \leq y) \leq \mathrm{P}(Y_2 \leq y) + \varepsilon$ for all y and $\mathrm{P}(Y_2 \leq y) < \mathrm{P}(Y_1 \leq y) \leq \mathrm{P}(Y_2 \leq y) + \varepsilon$ for some y.*

Under first-order ε-weak stochastic dominance, we derive a general lower and upper bound on the aptitude-achievement paradox probability when $F_{Y_1} \neq F_{Y_2}$.

Proposition 4.2. *Suppose that the system inclusion mechanism has the property that $1_i\left(X_{i,(N_i)}, \boldsymbol{X}_{i,-(N_i)}\right) = 1$, for $i = 1, 2$, and that the Full House Model is of the form*

$$\begin{aligned} Y_{i,j} &\sim F_{Y_i}(y), \qquad i = 1, 2, \qquad j = 1, \ldots, n_i, \\ X_{i,j} &\sim F_X(x), \qquad i = 1, 2, \qquad j = 1, \ldots, N_i, \end{aligned}$$

where $F_{Y_i}^{-1}(y)$ and $F_X^{-1}(x)$ exist. Further suppose that Y_2 is first order ε-weakly stochastically dominant over Y_1 for some $\varepsilon > 0$ and $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$. Then,

$$\begin{aligned} \frac{n_2}{n_1 + n_2} - \frac{N_2}{N_1 + N_2} &\leq \mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\ &\leq n_2 \sum_{i=0}^{n_1} \frac{\binom{n_1}{i}\varepsilon^i}{n_1 + n_2 - i} - \frac{N_2}{N_1 + N_2} \end{aligned} \tag{6}$$

Proof. First, in combination with the Definition 4.1 and the assumption $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$, we have

$$F_{Y_1}(x)^{\frac{n_1}{N_1}} \geq F_{Y_2}(x)^{\frac{n_2}{N_2}} \implies x \geq F_{Y_1}^{-1}\left(F_{Y_2}(x)^{\frac{n_2 N_1}{N_2 n_1}}\right).$$

By applying the Equation 4 from the Lemma 3.2 we have

$$\begin{aligned} &\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \mathrm{P}\left(g_{n_1,n_2}\left(Y_{2,(n_2)}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\ &= \mathrm{P}\left(F_{Y_1}^{-1}\left(F_{Y_2}(Y_{2,(n_2)})^{\frac{n_2 N_1}{N_2 n_1}}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\ &= \int_{-\infty}^{+\infty} \int_{F_{Y_1}^{-1}\left(F_{Y_2}(x)^{\frac{n_2 N_1}{N_2 n_1}}\right)}^{x} dF_{Y_1,(n_1)}(y) dF_{Y_2,(n_2)}(x) \end{aligned}$$

$$= \int_{-\infty}^{+\infty} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2 N_1}{N_2}} dF_{Y_{2,(n_2)}}(x)$$
$$= \int_0^1 F_{Y_1}\left(F_{Y_2}^{-1}(t^{\frac{1}{n_2}})\right)^{n_1} - t^{\frac{N_1}{N_2}} dt.$$

Based on the Definition 4.1, we have

$$\begin{aligned}
\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) &= \int_0^1 F_{Y_1}\left(F_{Y_2}^{-1}(t^{\frac{1}{n_2}})\right)^{n_1} - t^{\frac{N_1}{N_2}} dt \\
&\geq \int_0^1 F_{Y_2}\left(F_{Y_2}^{-1}(t^{\frac{1}{n_2}})\right)^{n_1} - t^{\frac{N_1}{N_2}} dt \\
&= \frac{n_2}{n_1+n_2} - \frac{N_2}{N_1+N_2},
\end{aligned}$$

and

$$\begin{aligned}
\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) &= \int_0^1 F_{Y_1}\left(F_{Y_2}^{-1}(t^{\frac{1}{n_2}})\right)^{n_1} - t^{\frac{N_1}{N_2}} dt \\
&\leq \int_0^1 \left(t^{1/n_2} + \varepsilon\right)^{n_1} - t^{\frac{N_1}{N_2}} dt \\
&= n_2 \sum_{i=0}^{n_1} \frac{\binom{n_1}{i}\varepsilon^i}{n_1+n_2-i} - \frac{N_2}{N_1+N_2}.
\end{aligned}$$

□

These bounds are a prescription under general distributions for achievement in Full House Modeling. However, we can still obtain explicit aptitude-achievement paradox probabilities when we specify $Y_i \sim N(\mu_i, \sigma_i^2)$, $i = 1, 2$.

4.2 Normal distributions

This section will develop explicit aptitude-achievement paradox probabilities (2) when a Full House Model in Definition 2.1 additionally specifies that $Y_i \sim N(\mu_i, \sigma_i^2)$, $i = 1, 2$. We will focus on the case where the system inclusion mechanism has the property that $1_i(X_{i,(N_i)}, \mathbf{X}_{i,-(N_i)}) = 1$ for $i = 1, 2$.

The outline for our results is similar to that in Section 3 where a technical condition is needed to prove the main result – like condition (5) in Theorem 3.4 –and further results provide configurations of the sample sizes that satisfy the technical condition. However, in this setting with $Y_i \sim N(\mu_i, \sigma_i^2)$, $i = 1, 2$, the technical condition leaves a remainder term in excess of a clean probability result, and further results will show that this remainder term is negligible.

Similar to the proof of Theorem 3.4, Lemma 3.2 will provide a representation more conducive to formal integration needed to prove our main result. Only this time we will use (4) in Lemma 3.2 to represent the aptitude-achievement paradox probability as

$$\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \mathrm{P}\left(g_{n_1,n_2}\left(Y_{2,(n_2)}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right),$$

where, from the definition of g in Lemma 3.2 applied to our current setting, we have

$$g_{n_1,n_2}(x) = F_{Y_1}^{-1}\left(F_{Y_2}(x)^{\frac{n_2N_1}{N_2n_1}}\right) = \Phi^{-1}\left(\Phi\left(\frac{x-\mu_2}{\sigma_2}\right)^{\frac{n_2N_1}{N_2n_1}}\right)\sigma_1 + \mu_1, \tag{7}$$

and Φ defines the cumulative distribution function of the standard normal distribution. From here, we have that

$$\begin{aligned}
&\mathrm{P}\left(g_{n_1,n_2}\left(Y_{2,(n_2)}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \mathrm{P}\left(F_{Y_1}^{-1}\left(F_{Y_2}(Y_{2,(n_2)})^{\frac{n_2N_1}{N_2n_1}}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\
&\quad = \int_A \int_{F_{Y_1}^{-1}\left(F_{Y_2}(x)^{\frac{n_2N_1}{N_2n_1}}\right)}^{x} dF_{Y_1,(n_1)}(y)dF_{Y_2,(n_2)}(x),
\end{aligned}$$

where $A := A(\mu_1, \mu_2, \sigma_1, \sigma_2, n_1, n_2, N_1, N_2)$ is the range of integration so that the inner integral above is properly specified. More formally,

$$\begin{aligned}
A &= \left\{\omega : F_{Y_1}^{-1}\left(F_{Y_2}(Y_{2,(n_2)}(\omega))^{\frac{n_2N_1}{N_2n_1}}\right) \le Y_{2,(n_2)}(\omega)\right\} \\
&= \left\{\omega : \Phi\left(\frac{Y_{2,(n_2)}(\omega)-\mu_2}{\sigma_2}\right)^{\frac{n_2}{N_2}} \le \Phi\left(\frac{Y_{2,(n_2)}(\omega)-\mu_1}{\sigma_1}\right)^{\frac{n_1}{N_1}}\right\},
\end{aligned} \tag{8}$$

where the second equality in (8) follows from (7). Continuing with the probability calculation above

$$\begin{aligned}
&\int_A \int_{F_{Y_1}^{-1}\left(F_{Y_2}(x)^{\frac{n_2N_1}{N_2n_1}}\right)}^{x} dF_{Y_1,(n_1)}(y)dF_{Y_2,(n_2)}(x) = \int_A F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}} dF_{Y_2,(n_2)}(x) \\
&\quad = \int_{-\infty}^{\infty} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}} dF_{Y_2,(n_2)}(x) - \int_{A^c} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}} dF_{Y_2,(n_2)}(x).
\end{aligned}$$

The left-hand side can (and will) be further evaluated and leads to the desired explicit aptitude-achievement paradox probability. The right-hand side represents a remainder term $R := R(\mu_1, \mu_2, \sigma_1, \sigma_2, n_1, n_2, N_1, N_2)$,

$$R = -\int_{A^c} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}} dF_{Y_2,(n_2)}(x), \tag{9}$$

where A in (9) is defined in (8). The remainder term may seem unintuitive. However, we will now motivate a simple setting that allows for R to be negligible when n_2 is moderately large.

From (8) and (9), we have

$$\begin{aligned}
|R| &= \left|\int_{A^c} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}} dF_{Y_2,(n_2)}(x)\right| \\
&\le \int_{A^c} \left|F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}}\right| dF_{Y_2,(n_2)}(x)
\end{aligned}$$

$$\leq \mathrm{P}\left(\Phi\left(\frac{Y_{2,(n_2)}-\mu_2}{\sigma_2}\right)^{\frac{n_2}{N_2}} \geq \Phi\left(\frac{Y_{2,(n_2)}-\mu_1}{\sigma_1}\right)^{\frac{n_1}{N_1}}\right).$$

Now, when we specify that $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$ as in Proposition 4.2, we have

$$\begin{aligned}
&\mathrm{P}\left(\Phi\left(\frac{Y_{2,(n_2)}-\mu_2}{\sigma_2}\right)^{\frac{n_2}{N_2}} \geq \Phi\left(\frac{Y_{2,(n_2)}-\mu_1}{\sigma_1}\right)^{\frac{n_1}{N_1}}\right) \\
&\quad \leq \mathrm{P}\left(\Phi\left(\frac{Y_{2,(n_2)}-\mu_2}{\sigma_2}\right) \geq \Phi\left(\frac{Y_{2,(n_2)}-\mu_1}{\sigma_1}\right)\right) \\
&\quad = \mathrm{P}\left(\frac{Y_{2,(n_2)}-\mu_2}{\sigma_2} \geq \frac{Y_{2,(n_2)}-\mu_1}{\sigma_1}\right).
\end{aligned}$$

A bit of algebra under the specification that $\sigma_1 \leq \sigma_2$ yields

$$\mathrm{P}\left(\frac{Y_{2,(n_2)}-\mu_2}{\sigma_2} \geq \frac{Y_{2,(n_2)}-\mu_1}{\sigma_1}\right) = \mathrm{P}\left(Y_{2,(n_2)} \leq \left(\frac{\mu_2}{\sigma_2}-\frac{\mu_1}{\sigma_1}\right)\left(\frac{1}{\sigma_2}-\frac{1}{\sigma_1}\right)^{-1}\right).$$

A bit of algebra under the specification that $\mu_1 \leq \mu_2$ implies that

$$\left(\frac{\mu_2}{\sigma_2}-\frac{\mu_1}{\sigma_1}\right)\left(\frac{1}{\sigma_2}-\frac{1}{\sigma_1}\right)^{-1} \leq \mu_2.$$

Putting it all together, we have

$$|R| \leq \mathrm{P}\left(Y_{2,(n_2)} \leq \mu_2\right) \leq \left(\frac{1}{2}\right)^{n_2},$$

when $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$, $\sigma_1 \leq \sigma_2$, and $\mu_1 \leq \mu_2$. Therefore, we have a trivial bound on the remainder term which shows that it is negligible when n_2 is moderately large under a familiar sample size specification (see Proposition 4.2) and a simple parametric configuration, despite the remainder term (9) being unintuitive. This argument will be necessary both for the statement and proof of our main result in this section. First some additional preliminaries.

Let $\mathbf{W} \sim \mathcal{N}_{(n_1+n_2-1)}(\mathbf{0}, \mathbf{\Sigma})$ and covariance matrix

$$\mathbf{\Sigma} = \begin{bmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{bmatrix}, \tag{10}$$

defined with the following structure

$$\begin{aligned}
\Sigma_{11} &= \begin{bmatrix} \frac{\sigma_2^2}{\sigma_1^2}+1 & \frac{\sigma_2^2}{\sigma_1^2} & \cdots & \frac{\sigma_2^2}{\sigma_1^2} \\ \frac{\sigma_2^2}{\sigma_1^2} & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \frac{\sigma_2^2}{\sigma_1^2} \\ \frac{\sigma_2^2}{\sigma_1^2} & \cdots & \frac{\sigma_2^2}{\sigma_1^2} & \frac{\sigma_2^2}{\sigma_1^2}+1 \end{bmatrix}_{n_1 \times n_1} \qquad \Sigma_{12}^T = \Sigma_{21} = \begin{bmatrix} \frac{\sigma_2}{\sigma_1} & \cdots & \cdots & \frac{\sigma_2}{\sigma_1} \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ \frac{\sigma_2}{\sigma_1} & \cdots & \cdots & \frac{\sigma_2}{\sigma_1} \end{bmatrix}_{(n_2-1)\times n_1} \\
\Sigma_{22} &= \begin{bmatrix} 2 & 1 & \ldots & 1 \\ 1 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & 1 \\ 1 & \ldots & 1 & 2 \end{bmatrix}_{(n_2-1)\times(n_2-1)}.
\end{aligned} \tag{11}$$

With this structure we now provide an explicit aptitude-achievement paradox probability, the main result of this section.

Theorem 4.3. *Assume a Full House Model in Definition 2.1 and additionally specify that* $Y_i \sim N(\mu_i, \sigma_i^2)$*, and suppose that the system inclusion mechanism has the property that* $1_i(X_{i,(N_i)}, \boldsymbol{X}_{i,-(N_i)}) = 1$ *for* $1 = 1, 2$*. Let* $\mathbf{W} \sim \mathcal{N}_{(n_1+n_2-1)}(\mathbf{0}, \boldsymbol{\Sigma})$ *where* $\boldsymbol{\Sigma}$ *is as defined in (10) and (11). Then,*

$$\begin{aligned} &\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\ &\quad = n_2 F_{\mathbf{W}}(\mathbf{w} = (\frac{\mu_2 - \mu_1}{\sigma_1}, \ldots, \frac{\mu_2 - \mu_1}{\sigma_1}, 0, \ldots, 0)) - \frac{N_2}{N_1 + N_2} + R, \end{aligned} \tag{12}$$

where R *is defined in (9),* $F_{\mathbf{W}}$ *is the distribution function of* $\mathbf{W}$*, and the first* n_1 *elements of* $\mathbf{w}$ *are* $\frac{\mu_2-\mu_1}{\sigma_1}$ *and the last* $n_2 - 1$ *elements are 0.*

Moreover, we have that

$$|R| \leq \left(\frac{1}{2}\right)^{n_2}, \tag{13}$$

when $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$*,* $\sigma_1 \leq \sigma_2$*, and* $\mu_1 \leq \mu_2$*.*

Proof. Equation 4 from Lemma 3.2 yields

$$g_{n_1,n_2}(x) = F_{Y_1}^{-1}\left(F_{Y_2}(x)^{\frac{n_2 N_1}{N_2 n_1}}\right) = \Phi^{-1}\left(\Phi\left(\frac{x-\mu_2}{\sigma_2}\right)^{\frac{n_2 N_1}{N_2 n_1}}\right)\sigma_1 + \mu_1.$$

Let A and R be as defined, respectively, in (8) and (9). We have,

$$\begin{aligned} &\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \mathrm{P}\left(g_{n_1,n_2}\left(Y_{2,(n_2)}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\ &= \mathrm{P}\left(F_{Y_1}^{-1}\left(F_{Y_2}(Y_{2,(n_2)})^{\frac{n_2 N_1}{N_2 n_1}}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\ &= \int_A \int_{F_{Y_1}^{-1}\left(F_{Y_2}(x)^{\frac{n_2 N_1}{N_2 n_1}}\right)}^{x} dF_{Y_1,(n_1)}(y) dF_{Y_2,(n_2)}(x) \\ &= \int_A F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2 N_1}{N_2}} dF_{Y_2,(n_2)}(x) \\ &= \int_{-\infty}^{\infty} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2 N_1}{N_2}} dF_{Y_2,(n_2)}(x) - \int_{A^c} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2 N_1}{N_2}} dF_{Y_2,(n_2)}(x) \\ &= \int_{-\infty}^{\infty} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2 N_1}{N_2}} dF_{Y_2,(n_2)}(x) + R. \end{aligned}$$

We continue with the left-hand side of the last equality,

$$\begin{aligned} &\int_{-\infty}^{\infty} \left[F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2 N_1}{N_2}}\right] dF_{Y_2,(n_2)}(x) = \int_0^1 \left[F_{Y_1}\left(F_{Y_2}^{-1}(t^{1/n_2})\right)^{n_1} - t^{N_1/N_2}\right] dt \\ &\quad = \int_0^1 \left(\Phi\left(\frac{\Phi^{-1}(t^{1/n_2})\sigma_2 + \mu_2 - \mu_1}{\sigma_1}\right)\right)^{n_1} dt \; - \; \frac{N_2}{N_1 + N_2} \end{aligned}$$

$$= \int_{-\infty}^{+\infty} \Phi\left(\frac{\sigma_2 x + \mu_2 - \mu_1}{\sigma_1}\right)^{n_1} \left(n_2 \Phi(x)^{n_2-1}\phi(x)\right)\, dx \;-\; \frac{N_2}{N_1+N_2}$$

$$= n_2 \int_{-\infty}^{+\infty} \mathbb{P}\left(\begin{array}{l} Z_{1,1} - \frac{\sigma_2}{\sigma_1}X \le \frac{\mu_2-\mu_1}{\sigma_1}, \ldots, Z_{1,n_1} - \frac{\sigma_2}{\sigma_1}X \le \frac{\mu_2-\mu_1}{\sigma_1}, \\ Z_{2,1} - X \le 0, \ldots, Z_{2,n_2-1} - X \le 0 \end{array} \;\middle|\; X = x \right) \phi(x)\, dx \;-\; \frac{N_2}{N_1+N_2}$$

$$= n_2\, \mathbb{P}\left(\begin{array}{l} Z_{1,1} - \frac{\sigma_2}{\sigma_1}X \le \frac{\mu_2-\mu_1}{\sigma_1}, \ldots, Z_{1,n_1} - \frac{\sigma_2}{\sigma_1}X \le \frac{\mu_2-\mu_1}{\sigma_1}, \\ Z_{2,1} - X \le 0, \ldots, Z_{2,n_2-1} - X \le 0 \end{array} \right) \;-\; \frac{N_2}{N_1+N_2}$$

$$= n_2\, F_{\mathbf{W}}\left(\mathbf{w} = (\frac{\mu_2-\mu_1}{\sigma_1}, \ldots, \frac{\mu_2-\mu_1}{\sigma_1}, 0, \ldots, 0)\right) \;-\; \frac{N_2}{N_1+N_2}.$$

The distribution function $F_{\mathbf{W}}$ corresponds to random variable $\mathbf{W} \sim \mathcal{N}_{(n_1+n_2-1)}(\mathbf{0}, \mathbf{\Sigma})$ where $\mathbf{\Sigma}$ is as defined in (10) and (11). The first n_1 elements of $\mathbf{w}$ is $\frac{\mu_2-\mu_1}{\sigma_1}$ and the last $n_2 - 1$ elements are 0.

The bound on the remainder (13) follows from the argument in Section 4.2. □

To validate the Theorem 4.3 in the numerical simulation, we conducted Monte Carlo simulations across a range of parameter settings. The simulated paradox probabilities closely matched the theoretical values. Because the simulations serve only as a verification of the closed-form result and do not provide additional methodological insight, we present them in Supplementary Materials for completeness.

We now show that plug-in estimates for normal distribution parameters μ_i, σ_i, $i = 1, 2$ allow for reliable estimate of the paradox probability (12). Our approach requires properties for matrices of the form

$$\Sigma = \begin{bmatrix} c+1 & c & \ldots & c \\ c & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & c \\ c & \ldots & c & c+1 \end{bmatrix}_{n\times n}.$$

These properties are

$$\det(\Sigma) = cn + 1, \tag{14}$$

and

$$\Sigma^{-1} = \frac{1}{c\cdot n + 1} \begin{bmatrix} c\cdot(n-1)+1 & -c & \ldots & -c \\ -c & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & -c \\ -c & \ldots & -c & c\cdot(n-1)+1 \end{bmatrix}_{n\times n}, \tag{15}$$

where (15) follows from an application of the Woodbury identity [Golub and Van Loan, 2013].

Before we state our result, we introduce the Lemma 4.4

Lemma 4.4. *Let* $\mathbf{W} \sim \mathcal{N}_{n_1+n_2-1}(\mathbf{0}, \mathbf{\Sigma})$ *be a multivariate normal random vector with mean vector* $\mathbf{0}$ *and covariance matrix* $\mathbf{\Sigma}$ *as defined in* (10) *and* (11). *We denote the cumulative*

distribution function (CDF) of $\mathbf{W}$ *by* $\mathbf{\Phi}(\mathbf{w};\mathbf{\Sigma})$*, where* $\mathbf{w} \in \mathbb{R}^n$ *is the quantile vector at which the CDF is evaluated.*

Define

$$
\begin{aligned}
c = \max \Big\{ & \Phi(W_{n_1+n_2-1} = w_{n_1+n_2-1} | W_1 = w_1, W_2 = w_2, \ldots, W_{n_1+n_2-2} = w_{n_1+n_2-2}), \\
& \Phi(W_{n_1+n_2-2} = w_{n_1+n_2-2} | W_1 = w_1, W_2 = w_2, \ldots, W_{n_1+n_2-3} = w_{n_1+n_2-3}), \ldots, \\
& \Phi(W_2 = w_2 | W_1 = w_1), \Phi(W_1 = w_1) \Big\},
\end{aligned} \tag{16}
$$

then c *is strictly less than 1 and*

$$
\mathbf{\Phi}(\mathbf{W}, \mathbf{\Sigma}) \leq c^{n_1+n_2-1} \tag{17}
$$

Proof. The proof can be found in the Supplementary Materials. □

Theorem 4.5. *Assume a Full House Model in Definition 2.1 and additionally specify that* $Y_i \sim N(\mu_i, \sigma_i^2)$*, and suppose that the system inclusion mechanism has the property that* $1_i(X_{i,(N_i)}, \boldsymbol{X}_{i,-(N_i)}) = 1$ *for* $1 = 1, 2$*. Let* $\mathbf{W} \sim \mathcal{N}_{(n_1+n_2-1)}(\mathbf{0}, \mathbf{\Sigma})$ *where* $\mathbf{\Sigma}$ *is as defined in (10) and (11). Suppose that* $\hat{\mu}_1$*,* $\hat{\mu}_2$*,* $\hat{\sigma}_1$ *and* $\hat{\sigma}_2$ *are, respectively, root sample-size consistent estimators of* μ_1*,* μ_2*,* σ_1 *and* σ_2*. Then*

$$
\begin{aligned}
& \mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)} | \hat{\mu_1}.\hat{\mu_2}, \hat{\sigma_1}, \hat{\sigma_2}\right) - \mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)} | \mu_1.\mu_2, \sigma_1, \sigma_2\right) \\
& = O_P(n_1^{\frac{1}{2}} n_2 c^{n_1+n_2-3} + n_1 n_2^{\frac{1}{2}} c^{n_1+n_2-3})
\end{aligned}
$$

where c *is defined in the Lemma 4.4*

Proof. The proof can be found in the Supplementary Materials. □

5 Examples

5.1 Sensitivity Analysis for Theorem 3.4

In this section, we conduct a sensitivity analysis to evaluate the influence of the parameters N_1, N_2, n_1, and n_2 in Theorem 3.4.

Under the setting of Theorem 3.4, we assume $X_1 = X_2 = X$, $Y_1 = Y_2 = Y$, where $X \sim$ Pareto(α) and $Y \sim \mathcal{N}(0, 1)$. Without loss of generality, we set $j = n_1 - 1$ and $k = n_2 - 1$, corresponding to the selection of the second-best performer in the paradox probability, as defined in Definition 3.1.

In each simulation, we randomly sample n_1 and n_2 observations from the distribution of Y, and compute their latent aptitudes using Equation (1). Based on these values, we evaluate the paradox probability according to Definition 3.1. We repeat this simulation 10000 times and report the average paradox probability.




Figure 1: Sensitivity Analysis for Paradox Probability

Figure 1 presents the results of the sensitivity analysis for the paradox probability described in Theorem 3.4. In Figure 1 (A), we fix $n_1 = 40$ and $n_2 = 60$, and vary N_1 and N_2 from 200 to 2000. In Figure 1 (B), we fix $N_1 = 10^3$ and $N_2 = 10^3$, and vary n_1 and n_2 from 20 to 200. In Figure 1 (C), we fix $n_1 = 40$ and $N_2 = 10^3$, and vary n_1 from 20 to 200 and N_2 from 200 to 2000. In Figure 1 (D), we fix $N_1 = 10^3$ and $n_2 = 60$, and vary N_1 from 200 to 2000 and n_2 from 20 to 200.

As shown in Figure 1 (A), a small N_2 and large N_1 increase the paradox probability by making $\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}\right)$ approach 1. Similarly, in Figure 1 (B), a small n_1 combined with a large n_2 leads to a higher paradox probability, since it drives $\mathrm{P}\left(Y_{1,(j)} < Y_{2,(k)}\right)$ towards 1.

We also investigate the effect of the ratio between N_2 and N_1 on the paradox probability. Let $c = \frac{N_2}{N_1}$, and vary c from 0.1 to 1 in increments of 0.1. Without loss of generality, we set $j = n_1 - 1$ and $k = n_2 - 1$, corresponding to the selection of the second-best performer in the paradox probability, as defined in Definition 3.1.

In each simulation, we fix $n_1 = 60$ and $n_2 = 40$, and vary N_1 from 100 to 2000. We repeat this simulation 10000 times and report the average paradox probability.




Figure 2: Effect of the ratio between N_2 and N_1 on the paradox probability

Figure 2 presents the results illustrating the effect of the ratio between N_2 and N_1 on the paradox probability under different settings. As shown in the figure, the paradox probability decreases as c increases, which is reasonable since $\mathbb{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}\right)$ decreases as N_2 increases.

Figure 2 also demonstrates that the paradox probability remains constant when the ratio between N_2 and N_1 is fixed. This implies that it is unnecessary to focus on the exact values of N_1 and N_2 when computing the paradox probability; rather, the ratio N_2/N_1 alone determines its value.

5.2 Recursive Full House Models

In this section we will investigate a simple recursive Full House Models in which Full House Models are repeatedly applied to individuals as they progress through iterations of some generic process. The general setup is that, in the first round, latent aptitude $X_j \overset{iid}{\sim} F_X$, $j = 1, \ldots, N$ will be generated and then a label will be randomly assigned to each individual. There will be m labels and each label will designate the outcome distribution. $F_{Y_{i,j}}$, $i = 1, \ldots, n$, $j = 1, \ldots, n_i$. Then outcomes will be realized as

$$Y_{i,(r_{i,(j)})} = F^{-1}_{Y_{i,j}} \circ I^{-1}_{r_{i,(j)}, n_i - r_{i,(j)}+1} \circ I_{j,N-j+1} \circ F_X(X_{(j)}), \tag{18}$$

where $r_{i,(j)}$ is the rank of $X_{(j)}$ among the n_i latent aptitudes also assigned label i.

In the next iteration, all observations will be combined and then sorted in descending order by $Y_{i,(j)}$. The labels will be assigned in descending order, the largest n_m values of $Y_{i,(j)}$ are assigned label m, the next largest n_{m-1} values of $Y_{i,(j)}$ are assigned label $m-1$, and so on. We will setup these simulation so that Y_i stochastically dominates Y_j whenever $i > j$ so that higher outcomes are associated with higher labels. Then new Y values are generated according to (18). The process will repeat until a pre-assigned stopping rule is satisfied. Note that this simulation setup assumes an immutable latent aptitude.

5.2.1 Stop when highest latent aptitude is included in the highest-performing group

Here we consider a recursive Full House Modeling setup described in Section 5.2 in which the recursion stops at the first iteration when $X_{(N)}$ is assigned label m. Here we specify that $N = 100,000$, $m = 200$, $n_i = 500$, $i = 1, \ldots, m$. We will assume that $X_j \overset{iid}{\sim}$ Pareto(1.16), $j = 1, \ldots, N$. And we will assume that $Y_{i,j} \overset{iid}{\sim} N(\mu_i, \sigma_i)$, $j = 1, \ldots, n_i$ and $i = 1, \ldots, m$. We will specify that the μ_is as a set of 200 equally spaced values between 60 and 80 with μ_i increasing in i, ie $\mu_i = 60 - 20/199 + 20i/199$, $i = 1, \ldots, m$. We will fix $\sigma_i = \sigma$ and consider $\sigma \in \{0.5, 1, 2\}$. These simulation will be repeated 500 times.

To illustrate the procedure, consider a simplified scenario in which all $X_{(j)}$ values are realized at their median value. With this simplification, $Y_{i,(r_{i,(j)})}$ in (18) is approximately $F^{-1}_{Y_{i,j}}((r_{i,(j)} - 1/3)/(n_i + 1/3))$ [Kerman, 2011]. And we have that $Y_{i,(n_i)} - Y_{i+1,(n_{i+1}-1)} > \mu_{i+1} - \mu_i$ for all considered values of σ. The individual with latent ability $X_{(N)}$ will consistently attain the value $Y_{i,(n_i)}$, regardless their assigned group. Consequently, this individual's position among the m groups will improve as the simulation advances.

Results can be seen in Figure 3. These Monte Carlo simulations reveal that it typically takes several rounds before the individual with the top aptitude belongs to the group that allows for the highest performances. On average, with σ set to 0.5, 1, and 2, it takes 2.61, 4.19, and 7.46 rounds, respectively, for the individual with the top aptitude to join the highest-performing group (respective standard errors: 0.05, 0.09, 0.18).

These findings resonate with several empirical studies examining bias and distortion in real-world evaluative systems. The GRADE algorithm developed by Waters and Miikkulainen [2014] demonstrates how admission systems can bypass nuanced information (e.g., personal statements) in favor of machine-learned proxies, potentially reinforcing existing filters. Similarly, Alvero et al. [2020] show that seemingly innocuous features (such as the word baseball) can serve as strong demographic signals. This raises concerns about indirect inference. A broader review by Baker and Hawn [2022] warns that algorithmic systems may harbor unknown unknowns, especially in global or under-examined subpopulations. Finally, the meta-analyses in Walton and Spencer [2009] provide direct evidence that outcome-based measures (e.g., grades, test scores) can systematically understate the aptitude of marginalized groups due to identity-based threat.

These phenomena of latent talent being misclassified, sorted, and repeatedly filtered through context-specific outcome layers is precisely what this presentation of recursive Full House Modeling aims to formalize and study. At any stage in the selection process, you cannot be sure that the highest marks correspond to individuals with the highest ability.




Figure 3: The number of rounds that it takes for $X_{(N)}$ to be assigned label m in a recursive Full House Modeling framework in Section 5.2 with specifications in Section 5.2.1.

While the setting we consider is stylized and assumes fixed latent aptitude, it serves to illustrate how structural features of evaluative systems, not just randomness, can delay or obscure the recognition of true potential.

5.3 Batting Averages in Major League Baseball

We now apply Theorem 4.3 and the reliability guarantees in Theorem 4.5 to evaluate the contextual extremeness of observed outcomes in a real-world system where the underlying distribution of aptitude is not directly observable and evolves over time. Major League Baseball (MLB) offers a canonical example of such a system. Here, player skill (latent batting aptitude) is inferred only indirectly through season-level observed outcomes (batting averages), which are shaped by changing rules, population structures, and environmental variance.

In particular, MLB introduced a series of rule changes in 2023 intended to increase in-game action, including a restriction on defensive positioning designed to raise batting averages on balls in play (BABIP) [Castrovince, 2023].

First, the 2023 mean batting average for full-time players increased by 0.0057 (corresponding to an additional hit about every four baseball games) with a reduction in variability. A two-sample tests for differences in batting averages among full-time players from the 2023 and 2022 seasons has a p-value of 0.039. Thus, the rule changes seemed to be effective in yielding increased batting averages[1] Moreover, the 2023 MLB batting leader Luis Arraez

[1]Technically, Castrovince [2023] referred to MLB's desire to increased batting averages on balls in play




Figure 4: Seasonal means and standard deviations of batting averages for full-time players from 1871-2023. Smoothing is applied to both trends.

had a 0.354 batting average which was the first batting average to exceed .350 since Josh Hamilton in 2010.

Now we turn to applying the results of Theorem 4.3 in this example to contextualize the effect of these rule changes. Here we assume that seasonal batting averages of full-time players are normally distributed. The true means and standard deviations of these distributions are unknown. Thus, we estimate them via plug-in sample mean and sample standard-deviation and invoke the reliability guarantees from Theorem 4.5. Figure 4 presents smoothed estimates of the mean and standard deviation of batting averages for full-time players from 1871 to 2023. Notably, both the average and variability of batting averages in 2023 are among the lowest observed in the dataset, indicating a historically suppressed environment for batting averages of full-time batters.

In this example we will investigate the paradox probability corresponding to maximum batting averages when comparing the 2023 season to all prior seasons. We will thus specify the 2023 season as system 1 and the comparison season, which ranges from 1871 to 2022, as system 2. Figure 5 displays these estimated paradox probabilities (12) where estimates are obtained using plug-in, sample means and standard deviations are substituted for population parameters. These estimators are $\sqrt{n}$-consistent so that the reliability results of Theorem 4.5

(BABIP) while this analysis focused on batting averages. BABIP increased by 0.0061 from 2022 to 2023.




Figure 5: Comparison of paradox probabilities and empirical exceedance rates relative to the 2023 season, based on data from 1871-2022. The blue line shows estimated paradox probabilities via plug-in using Theorem 4.3. The black dashed line displays a smoothed empirical exceedance rate for Arraez's 2023 batting average.

apply. For comparison, the figure also shows a smoothed empirical curve representing the proportion of seasons in which the maximum batting average exceeded Arraez's 2023 mark. We observe that the estimated paradox probabilities track closely with the empirical exceedance rate. Both quantities increase as we move backward in time, reinforcing the idea that earlier eras, with higher means, larger variances, and smaller populations, were more conducive to producing extreme individual batting averages, even when overall talent levels may have been lower.

Yan et al. [2025a] studied historical batting averages and era-adjusted batting averages through Full House Modeling for Luis Arraez's 2023 seasons and Jeff McNeil's MLB leading 2022 season in which he hit 0.326. They found that when one accounts for park and era effects then Arraez and McNeil had very similar performances, despite Arraez's large lead in raw batting average. These authors also found that assuming normality for batting averages among full-time players might not be appropriate [Yan et al., 2025b]. Even so, we evaluate $\widehat{F}(Y_{(n_i),i})^{n_i}$, the observed percentile of the distribution for the maximum batting average $Y_{(n_i),i}$ in season i. These percentiles are not uniformly distributed as seen in Figure 6.

Interestingly, the percentile corresponding to Luis Arraez's 2023 MLB leading batting average is largely without recent precedent. See Table 1 for specifics. This table contains information from all seasons in which an MLB leading batter's extreme batting average percentile exceeds that of Luis Arraez in 2023. Notice that there is a significant drop off in




Figure 6: Uniform Q-Q plot for extreme batting averages defined as $\widehat{F}(Y_{(n_i),i})^{n_i}$ where i denotes the season, n_i denotes the number of full-time players in season i, $Y_{(n_i),i}$ denotes the highest batting average for full-time players in season i, and $\widehat{F}$ is the distribution function of the normal distribution with sample mean and standard deviation estimators substituted for the population parameters.

the frequency of such seasons after Rogers Hornsby's modern day (post-1900) record 0.424 season from 1924. It is also interesting to note that the four most recent seasons in this table are from international players or players that would not have been allowed to play baseball had segregation continued (same with Rod Carew in 1977). Again, extreme performances were more common when the talent pool was more restricted and batting strategies and training methods were more variable, not when MLB expanded the number of teams [Gould, 1996, Eck, 2020a,b, Yan et al., 2025b]. This table also includes a column for p-values from a Shapiro-Wilk test [Shapiro and Wilk, 1965] on the normality of batting average for full-time players. These results support the findings in Yan et al. [2025b]. Moreover, the values $\widehat{F}(Y_{(n_i),i})^{n_i}$ do not follow a uniform distribution as indicated in Figure 6.

Year	Name	MLB leading BA	n	Shapiro Wilk p-value	Extreme Percentile
1876	Ross Barnes	0.429	40	0.0043	0.9791
1881	Cap Anson	0.399	42	0.3213	0.9665
1884	Fred Dunlap	0.412	110	0.0066	0.9722
1887	Tip O'Neill	0.435	112	0.0352	0.9848
1901	Nap Lajoie	0.426	104	0.3360	0.9760
1904	Nap Lajoie	0.376	104	0.0402	0.9861
1905	Cy Seymour	0.377	112	0.0437	0.9602
1908	Honus Wagner	0.354	115	0.0462	0.9614
1909	Ty Cobb	0.377	111	0.0937	0.9673
1910	Nap Lajoie	0.383	118	0.0082	0.9775
1911	Ty Cobb	0.420	115	0.0416	0.9663
1912	Ty Cobb	0.409	126	0.0001	0.9730
1913	Ty Cobb	0.390	119	0.0132	0.9570
1916	Tris Speaker	0.386	114	0.0003	0.9915
1917	Ty Cobb	0.383	117	0.0890	0.9724
1918	Ty Cobb	0.382	109	0.4605	0.9575
1922	George Sisler	0.420	118	0.0174	0.9671
1924	Rogers Hornsby	0.424	118	0.1749	0.9863
1941	Ted Williams	0.406	110	0.0104	0.9946
1957	Ted Williams	0.388	117	0.0534	0.9885
1977	Rod Carew	0.388	196	0.2965	0.9935
1980	George Brett	0.390	184	0.0400	0.9960
1985	Wade Boggs	0.368	190	0.0417	0.9885
1988	Wade Boggs	0.366	180	0.0573	0.9789
1994	Tony Gwynn	0.394	153	0.0392	0.9631
2002	Barry Bonds	0.370	209	0.0102	0.9647
2004	Ichiro Suzuki	0.372	207	0.0622	0.9634
2023	Luis Arraez	0.354	205	0.0055	0.9564

Table 1: Batting average performances that are at least as extreme as Luis Arraez in 2023. Extreme as defined by the value $\widehat{F}(Y_{(n_i),i})^{n_i}$ where i denotes the season, n_i denotes the number of full-time players in season i, $Y_{(n_i),i}$ denotes the highest batting average for full-time players in season i, and $\widehat{F}$ is the distribution function of the normal distribution with sample mean and standard deviation estimators substituted for the population parameters. This table also includes the year, name, and raw batting average of extreme performers. It also includes the number of full-time players (n), and a Shapiro Wilk p-value corresponding to a test against normality for the batting average distribution for full-time players.

Despite the simplifying assumption of normality, the paradox probability analysis and associated diagnostics (Figures 5 and 6, Table 1) illustrate a key insight that under the empirical structure of 2023, which is characterized by low variance and high competition levels, Luis Arraez's 0.354 batting average represents an extremely rare achievement. Yet

paradoxically, it appears modest when compared to higher averages in historically weaker environments.

This demonstrates a central tenet of Full House Modeling: observed records are shaped as much by the structure of the evaluation system as by individual aptitude. Paradox probabilities offer a formal measure of this distortion. Without such contextualization, modern performances may be systematically undervalued. In particular, the modern structure of MLB makes it increasingly unlikely that even players with exceptional latent ability will produce historically dominant observed outcomes.

6 Discussion

The aptitude-achievement paradox arises when an individual from one context demonstrates higher latent aptitude but simultaneously lower observed achievement than an individual from another context. Although widely encountered in education, labor economics, psychometrics, medicine, and sports analytics, measuring the frequency of this paradox under realistic data-generating regimes is nontrivial. In this article, we formalize the Full House Methodology and investigate the probabilities of observing paradoxes between latent aptitude and observed achievement in comparative analysis of individuals across contexts.

Full House Modeling provides a direct comparison of individuals' latent aptitudes and observed achievements across different contexts, which is a novel contribution to latent trait modeling. While traditional latent trait models [Bartholomew et al., 2011] often focus on estimating underlying traits from observed categorical data , our approach specifically quantifies the probability of paradoxical outcomes in comparative settings. Full House Modeling formalizes the comparison of individuals' performance across distinct contexts based on the large source population size N, thereby offering a more nuanced understanding of individual differences and the impact of contextual factors on observed outcomes. In parallel, the causal bootstrap [Imbens and Menzel, 2018] highlights that valid causal inference similarly requires situating the observed sample within large source population to properly account for sampling uncertainty. By comparison, the Rasch model [Bond and Fox, 2013] ensures that comparisons of individuals' abilities are independent of the specific items administered, and that comparisons of item difficulties are independent of the sample of individuals tested, but it does not explicitly leverage the source population structure. Likewise, the bayesian hierarchical modeling [Berry et al., 1999] with the application in comparison of individuals' latent aptitudes typically captures population-level variation through seasonal or group-level random effects rather than through an explicit embedding into a large source population. Full House Modeling enhances broader statistical modeling practices by providing a framework for robust comparative analysis, particularly in situations where direct, context-independent comparisons are challenging.

We derive closed-form expressions for the paradox probability in two regimes: (i) when observed achievements are i.i.d., and (ii) when context-specific summaries are independent normals with distinct means and variances. The probability rises with population-size asymmetry (large N_1, small N_2) and sampling asymmetry (small n_1, large n_2). Moreover, for a fixed population-size ratio N_2/N_1, it is unchanged by proportional rescaling of the sample sizes (n_1, n_2). We also develop bounds for more general settings and show that, when

observed achievement is normal, a plug-in estimator yields reliable paradox-probability estimates.

Our findings align with a growing body of evidence that real-world evaluative systems routinely substitute *proxies* for the constructs they claim to measure, with consequential distortion. The GRADE algorithm of Waters and Miikkulainen [2014] shows how admissions workflows may privilege machine-learned surrogates over nuanced information (e.g., personal statements), reinforcing pre-existing filters. Relatedly, Alvero et al. [2020] document that seemingly innocuous features (e.g., the word "baseball") can act as strong demographic signals, raising concerns about *indirect inference* in which protected or contextual attributes are inferred through correlated proxies. A broader survey by Baker and Hawn [2022] warns that such systems may conceal "unknown unknowns," particularly for under-examined subpopulations. Complementing these mechanism-level concerns, the meta-analytic evidence in Walton and Spencer [2009] demonstrates that outcome-based measures (grades, test scores) can systematically understate the aptitude of marginalized groups through identity-contingent dynamics.

By elevating probability paradox from an anecdotal concern to a model-based, estimable quantity, our framework clarifies when cross-context comparisons are trustworthy, when they are fragile, and how to report that fragility responsibly. This closes the loop between principled latent measurement, realistic selection processes, and interpretable comparative claims.

References

AJ Alvero, Noah Arthurs, Anthony Lising Antonio, Benjamin W Domingue, Ben Gebre-Medhin, Sonia Giebel, and Mitchell L Stevens. Ai and holistic review: informing human reading in college admissions. In *Proceedings of the AAAI/ACM Conference on AI, Ethics, and Society*, pages 200–206, 2020.

Dimitris Askitis. Logarithmic concavity of the inverse incomplete beta function with respect to the first parameter. *Dimitris Askitis*, 2021.

Ryan S Baker and Aaron Hawn. Algorithmic bias in education. *International Journal of Artificial Intelligence in Education*, pages 1–41, 2022.

David J Bartholomew, Martin Knott, and Irini Moustaki. *Latent variable models and factor analysis: A unified approach*. John Wiley & Sons, 2011.

Scott M Berry, C Shane Reese, and Patrick D Larkey. Bridging different eras in sports. *Journal of the American Statistical Association*, 94(447):661–676, 1999.

Trevor G Bond and Christine M Fox. *Applying the Rasch model: Fundamental measurement in the human sciences*. Psychology Press, 2013.

Adrian Burgos Jr, Shen Yan, and Daniel J Eck. A change in the lineup? a historical foundation for a cross-era statistical model to assess baseball's all-time greats. *NINE: A Journal of Baseball History and Culture*, 2025. Accepted.

Anthony Castrovince. All of your questions about the new baseball rules, answered. *MLB.com*, February 2023. URL https://www.mlb.com/news/mlb-new-rules-for-2023-faq.

Raj Chetty, John N Friedman, and Jonah E Rockoff. Measuring the impacts of teachers i: Evaluating bias in teacher value-added estimates. *American economic review*, 104(9): 2593–2632, 2014.

Elisha Cohen, Adrian Burgos Jr., Shen Yan, and Daniel J Eck. Contextualizing historical impact: An era-adjusted modeling approach. 2025. Submitted.

Daniel J Eck. Challenging nostalgia and performance metrics in baseball. *Chance*, 33(1): 16–25, 2020a.

Daniel J Eck. Challenging WAR and other statistics as era-adjustment tools. *FanGraphs*, 2020b. https://community.fangraphs.com/challenging-war-and-other-statistics-as-era-adjustment-tools/.

Gene H Golub and Charles F Van Loan. *Matrix computations*. Johns Hopkins University press, 2013.

Stephen Jay Gould. *Full House: The Spread of Excellence from Plato to Darwin*. Harvard University Press, 1996.

Paul W Holland and Howard Wainer. *Differential item functioning*. Routledge, 2012.

Lisa I Iezzoni et al. *Risk adjustment for measuring health care outcomes*, volume 2. Health Administration Press Chicago, 1997.

Guido Imbens and Konrad Menzel. A causal bootstrap. Technical report, National Bureau of Economic Research, 2018.

Maurice G Kendall. A new measure of rank correlation. *Biometrika*, 30(1-2):81–93, 1938.

Jouni Kerman. A closed-form approximation for the median of the beta distribution. *arXiv preprint arXiv:1111.0433*, 2011.

Frederic M Lord. A paradox in the interpretation of group comparisons. *Psychological bulletin*, 68(5):304, 1967.

Samuel Sanford Shapiro and Martin B Wilk. An analysis of variance test for normality (complete samples). *Biometrika*, 52(3/4):591–611, 1965.

Edward H Simpson. The interpretation of interaction in contingency tables. *Journal of the Royal Statistical Society: Series B (Methodological)*, 13(2):238–241, 1951.

Robert H Somers. A new asymmetric measure of association for ordinal variables. *American sociological review*, pages 799–811, 1962.

Gregory M Walton and Steven J Spencer. Latent ability: Grades and test scores systematically underestimate the intellectual ability of negatively stereotyped students. *Psychological science*, 20(9):1132–1139, 2009.

Austin Waters and Risto Miikkulainen. Grade: Machine learning support for graduate admissions. *Ai Magazine*, 35(1):64–64, 2014.

Shen Yan, Adrian Burgos, Jr, Christopher Kinson, and Daniel J Eck. Comparing baseball players across eras via novel full house modeling. *The Annals of Applied Statistics*, 19(2): 1778–1799, 2025a.

Shen Yan, Adrian Burgos Jr., Christopher Kinson, and Daniel J Eck. Supplement to "comparing baseball players across eras via novel Full House Modeling". *Annals of Applied Statistics*, 2025b. Forthcoming.