A Formal Investigation of Paradoxes Between Latent Aptitude and Observed Achievement via Full House Methodology

Daniel J. Eck[1], Shen Yan[1*]

1. Department of Statistics, University of Illinois Urbana-Champaign

Abstract

We study a paradox in which an individual with greater latent aptitude exhibits lower observed achievement than an individual with lesser aptitude when comparisons are made across contexts. We formalize this phenomenon as the aptitude-achievement paradox and introduce its probability as an explicit inferential target. To study this quantity, we employ Full House Modeling, a framework that derives latent aptitude from observed achievement by embedding observed samples within larger source populations. We derive closed-form paradox probabilities when achievement distributions are identical across contexts and obtain explicit expressions and bounds when they differ, including tractable results under normality with reliable plug-in estimation. Our results show that paradox risk depends primarily on population and sampling ratios rather than absolute sizes. An application to Major League Baseball contextualizes historically extreme batting performances, showing how modern achievements may appear less exceptional after accounting for population depth and measurement environments. By elevating aptitude-achievement reversals from anecdotal concern to an estimable

*email addresses: Daniel J. Eck (dje13@illinois.edu); Shen Yan (shenyan3@illinois.edu)

quantity, this work provides a formal framework for quantifying structural reversals in cross-context comparisons under explicit models of population depth, selection, and measurement.

Highest-Priority Revision Checklist for JRSS-B Submission

These revisions are aimed at improving both:

1. overall fit with the stated JRSS-B scope, and
2. reviewer perception of contribution density and methodological importance.

Tier 1: Highest Priority Revisions

1. **Compress the application sections.**
    - Retain only the most compelling illustrations of:
        - paradox probabilities,
        - latent-observed reversals,
        - and empirical interpretability.

Tier 2: Important Revisions

1. **Sharpen the inferential framing in Section 4.**
    - Emphasize:
        - computational tractability,
        - operational estimation,
        - and practical implementation.
2. **Tighten the asymptotic rigor in Theorem 4.5.**
    - Reduce heuristic EVT-style arguments where possible.

3. **Reduce defensive discussion of assumptions.**

    - Shorten repeated caveats regarding:
        - rank preservation,
        - idealization,
        - and selection assumptions.
    - Replace apologetic framing with concise scope clarification.

4. **Remove or reduce conjectural discussion.**

    - In particular, consider moving the conjecture following Proposition 3.5 to the Discussion or Supplementary Materials.

Tier 3: Secondary but Valuable Revisions

1. **Shorten the Discussion section considerably.**

    - Avoid re-running the Introduction.
    - Focus on:
        - broader methodological implications,
        - selective observation,
        - and future directions.

2. **Reduce repeated philosophical exposition.**

    - Compress repeated conceptual discussion of:
        - aptitude vs. achievement,
        - population depth,
        - and contextual comparison.

3. **Potentially revise the title toward a more methodological/statistical foundations framing.**

   - Current title emphasizes "paradoxes" and may sound overly conceptual.
   - Consider framing around:
     - selective observation,
     - latent-observed ranking reversals,
     - or comparative inference.

1 Introduction

Comparisons of outcomes across differing contexts, schools, regions, eras, institutions, and tasks are central to education and psychometrics [Holland and Wainer, 2012, Bond and Fox, 2013], labor economics [Nickell and Layard, 1999], medicine, and sports analytics [Berry et al., 1999, Schell, 2013, 2016, Winston et al., 2022, Yan et al., 2025]. These problems have led to the development of a variety of widely used modeling frameworks that estimate relative ability, including empirical-Bayes shrinkage [Efron and Morris, 1975], Bayesian hierarchical models [Raudenbush and Bryk, 1986, Berry et al., 1999], Bradley-Terry-Luce and paired comparisons models [Bradley and Terry, 1952, Bradley, 1976, 1984, David, 1988], Elo-type dynamic rating systems [Elo, 1978, Graepel et al., 2007], Gaussian-process and state-space formulations [Chu and Ghahramani, 2005, Houlsby et al., 2012], item-response and latent class models [McDonald, 1999, Embretson and Reise, 2000, Hagenaars and McCutcheon, 2002, Holland and Wainer, 2012, Bond and Fox, 2013], factor models [McDonald, 1999, Bartholomew et al., 2011], and domain-specific metrics [Baumer and Matthews, 2014, Baumer et al., 2015, Yu et al., 2016].

Despite their differences, these approaches share a common structural feature: ability is inferred relative to a fixed or implicitly comparable set of observed competitors. While such models adjust outcomes within a given context, they do not typically explicitly account for

changes in the size or structure of the eligible population, nor for context-specific distortions in how latent ability is expressed through observed outcomes. Consequently, when competition depth, selection mechanisms, or measurement conditions vary across contexts, observed dominance may not reliably reflect the corresponding latent ordering.

These limitations give rise to a new inferential target: the probability that cross-context comparisons reverse latent aptitude orderings. Specifically, an individual drawn from a deeper or more competitive population may possess higher latent aptitude yet exhibit lower observed achievement than a counterpart from a shallower or more weakly filtered context. We term this phenomenon the aptitude-achievement paradox. Formally, let individual j in context i have observed $\mathbb{R}$-valued achievement $Y_{i,j}$ and latent $\mathbb{R}$-valued aptitude $X_{i,j}$, and let individual n in context m have observed achievement $Y_{m,n}$ and latent aptitude $X_{m,n}$. The probability of an aptitude-achievement paradox is

$$\mathrm{P}\left(X_{i,j} > X_{m,n}, Y_{i,j} < Y_{m,n}\right). \tag{1}$$

This quantity captures the structural risk that comparisons based on observed outcomes reverse the corresponding latent ordering across contexts.

Such reversals arise naturally when measurement systems differ across contexts or when observation occurs through non-random selection mechanisms shaped by institutional, competitive, or environmental constraints. Existing methods provide limited tools for quantifying the frequency of these reversals.

Related work has studied disagreement between observed rankings and target orderings through measures such as Kendall's τ [Kendall, 1938] and Somers' D [Somers, 1962], while latent-trait and comparative ability models—including Rasch models [Bond and Fox, 2013], Bayesian hierarchical formulations [Raudenbush and Bryk, 1986, Berry et al., 1999], and Elo-type rating systems [Elo, 1978, Graepel et al., 2007]—infer underlying ability from observed outcomes. However, these approaches generally operate relative to the observed comparison

pool and do not explicitly model how differences in eligible population depth, selection intensity, or measurement environments distort the relationship between latent aptitude and observed achievement.

To address these limitations, we develop a structural framework for cross-context comparison that explicitly models population depth, selective observation, and context-specific measurement processes. This framework, which we refer to as Full House Modeling (FHM), was initially proposed in an applied setting comparing baseball players across eras [Yan et al., 2025], and is formalized here for comparative analysis across heterogeneous contexts. Under FHM, individuals are drawn from a common latent aptitude distribution, while contexts differ in (i) the size of the eligible population, (ii) the mechanism through which individuals are selected into observation, and (iii) the mapping from latent aptitude to observed achievement. We model measurement through context-specific rank-preserving transformations between latent aptitude and observed achievement.

In this paper, we develop the theory of FHM under rank-based selection mechanisms, particularly top-n selection, where tractable results are available. These assumptions are most natural in highly selective settings where observed individuals represent the upper tail of a larger latent population. Comparisons of athletes across eras provide a canonical example, since entry into professional leagues closely approximates selection of the most talented individuals from large underlying populations.

Yan et al. [2025] introduced Full House Modeling as an operational methodology for cross-era comparison in baseball analytics. The present paper develops the first formal statistical theory for this framework. In particular, our contributions are as follows:

- We formalize the aptitude-achievement paradox as an explicit inferential target for cross-context comparison and study its probability under selective observation.
- We develop the probabilistic foundations of Full House Modeling, including a characterization result showing that the induced latent-observed coupling is the unique rank-preserving monotone transformation consistent with the model assumptions.

- Under identical observed achievement distributions across contexts, we derive explicit and distribution-free paradox probabilities and characterize their dependence on population and sampling structure.
- Under heterogeneous achievement distributions, we derive general bounds and obtain tractable asymptotic approximations in Gaussian settings, including explicit remainder bounds and plug-in consistency results.
- We show through theoretical analysis and sensitivity experiments that paradox probabilities depend primarily on ratios of eligible population sizes and sampling fractions rather than their absolute magnitudes.

Proofs of mathematical results are provided in the Supplementary Materials.

2 Model Setting

We now motivate the structure of the Full House Model. Let N_i denote the size of population i. We distinguish between a *population*, consisting of all individuals N_i in context i, and a *system*, which denotes the subset of individuals whose outcomes are observed or recorded. The system is determined by a selection mechanism that maps the full population into an observed sample of size n_i. We suppose that every individual $k = 1, \ldots, N_i$ has an underlying latent $\mathbb{R}$-valued aptitude $X_{i,1}, \ldots, X_{i,N_i} \overset{iid}{\sim} F_X$. Let $\mathbf{X}_{i,-j}$ be the vector of all latent aptitudes not including the component j. We formalize the selection mechanism through an indicator function $S_{i,j} = 1_i\left(X_{i,j}, \mathbf{X}_{i,-j}\right)$, where $S_{i,j} = 1$ indicates inclusion in the observed system and a value of $S_{i,j} = 0$ indicates exclusion. This function may depend on the individual's aptitude relative to others in the population.

We now suppose that an individual j included in system i has an observed $\mathbb{R}$-valued statistic $Y_{i,j} \overset{iid}{\sim} F_{Y_i}$, $j = 1, \ldots, n_i$, with $n_i \leq N_i$. Under Full House Modeling, the observed values $Y_{i,j}$ are assumed to preserve the ranking induced by the latent aptitudes among the selected individuals.

Let $X_{i,(j)}$ and $Y_{i,(j)}$ denote the jth order statistic for system i. Here, larger values correspond to higher aptitude or achievement, so that $X_{i,(j)} > X_{i,(k)}$ whenever $j > k$. We form (Y, X) pairs of random variables for each individual included in the system. These pairs are of the form $(Y_{i,(j)}, X_{i,(r_{i,j})})$, where $r_{i,j}$ indexes the rank of that individual's latent aptitude within the full population of size N_i.

As an example, consider the selection rule $S_{i,j} = 1 \iff X_{i,j} \geq X_{i,(N_i - n_i + 1)}$, which selects the top n_i individuals in population i. In this case, the pairs take the form $(Y_{i,(j)}, X_{i,(N_i - n_i + j)})$ for $j = 1, \ldots, n_i$.

The key feature of Full House Modeling is the explicit connection between the elements forming the $(Y_{i,(j)}, X_{i,(r_{i,j})})$ pairs. This construction can be understood as a rank-preserving transformation between latent aptitude and observed achievement. Based on the probability integral transformation, Full House Modeling specifies this connection as

$$\begin{aligned} Y_{i,(j)} &= F^{-1}_{Y_{i,(j)}} \circ F_{X_{i,(r_{i,j})}} \left(X_{i,(r_{i,j})} \right), \\ X_{i,(r_{i,j})} &= F^{-1}_{X_{i,(r_{i,j})}} \circ F_{Y_{i,(j)}} \left(Y_{i,(j)} \right), \end{aligned} \tag{2}$$

where $S_{i,j} = 1$, and both F_{Y_i} and F_X are continuous and invertible. We summarize Full House Modeling in the following definition.

Definition 2.1 (Full House Model)**.** *Consider populations indexed by i with sizes N_i and system sizes n_i, where $N_i \geq n_i$. Let latent variables $X_{i,j} \overset{iid}{\sim} F_X$, $j = 1, \ldots, N_i$, and define the selection indicators $S_{i,j} := 1_i(X_{i,j}, \mathbf{X}_{i,-j})$, where $S_{i,j} = 1$ indicates inclusion in the observed system. Let $Y_{i,j} \overset{iid}{\sim} F_{Y_i}$, $j = 1, \ldots, n_i$, denote observed outcomes for the selected individuals. Form pairs $(Y_{i,(j)}, X_{i,(r_{i,j})})$, $j = 1, \ldots, n_i$, corresponding to the individuals for which $S_{i,j} = 1$. Then the Full House Model generates $Y_{i,(j)}$ values according to (2), where F_{Y_i} and F_X are continuous and invertible.*

The Full House Model defines a joint distribution over (X, Y) together with selection indicators $S_{i,j}$. Latent variables X are drawn i.i.d. from F_X, and the selection indicators $S_{i,j}$

determine inclusion in the observed system. Conditional on inclusion, observed outcomes Y are generated via a rank-preserving monotone coupling with X. In particular, the dependence between X and Y is induced through a quantile-matching transformation of their order statistics, ensuring that higher latent aptitude corresponds to higher observed achievement within each system. Conditional on the latent population and the selection mechanism, observed outcomes are generated through a deterministic rank-preserving coupling.

The Full House Model induces structural missingness by construction: for any individual j such that $S_{i,j} = 0$, the corresponding observed statistic $Y_{i,j}$ is undefined. In subsequent sections, we consider probabilities involving both latent and observed quantities. To formalize this setting, we treat $Y_{i,j}$ as a random variable defined only on the event $\{S_{i,j} = 1\}$. For example, $\mathrm{P}\left(X_{i,j} \in A,\ Y_{i,j} \in B,\ S_{i,j} = 1\right)$ is well-defined for Borel sets $A, B \subset \mathbb{R}$, whereas $Y_{i,j}$ is not defined on $\{S_{i,j} = 0\}$. This perspective allows selection mechanisms to be incorporated into probabilistic statements without imputing counterfactual outcomes.

Definition 2.1 assumes a common latent aptitude distribution F_X across contexts, while allowing the observed achievement distributions F_{Y_i} to vary by system. This reflects a modeling choice in which the process generating latent aptitude is taken to be context-neutral, whereas the expression and measurement of aptitude may differ across contexts.

Direct evaluation of the order-statistic distributions in (2) is often cumbersome. However, under continuity, they admit a convenient representation in terms of Beta distributions. In particular,

$$F_{X_{i,(r_{i,j})}}\left(X_{i,(r_{i,j})}\right) = F_{U_{i,(r_{i,j})}} \circ F_X\left(X_{i,(r_{i,j})}\right), \quad U_{i,(r_{i,j})} \sim \mathrm{Beta}(r_{i,j},\, N_i - r_{i,j} + 1), \tag{3}$$

and similarly,

$$F_{Y_{i,(j)}}\left(Y_{i,(j)}\right) = F_{U_{i,(j)}} \circ F_{Y_i}\left(Y_{i,(j)}\right), \quad U_{i,(j)} \sim \mathrm{Beta}(j,\, n_i - j + 1). \tag{4}$$

This representation allows (2) to be computed via F_X, F_{Y_i}, and known Beta distributions,

bypassing explicit order-statistic calculations. It also implies that latent aptitude can be recovered up to a monotone transformation without explicit knowledge of F_X.

Combining (3) and (4) yields

$$X_{i,(r_{i,j})} = F_X^{-1}\Big(F_{U_{i,(r_{i,j})}} \circ F_{U_{i,(j)}} \circ F_{Y_i}\big(Y_{i,(j)}\big)\Big).$$

Define $W_{i,j} = F_{U_{i,(r_{i,j})}} \circ F_{U_{i,(j)}} \circ F_{Y_i}(Y_{i,(j)})$. Since F_X is monotone, so is its inverse, which implies

$$W_{i_1,j_1} \geq W_{i_2,j_2} \implies F_X^{-1}(W_{i_1,j_1}) \geq F_X^{-1}(W_{i_2,j_2}).$$

Thus the ordering of latent aptitudes is invariant to F_X, and F_X acts as a structural nuisance parameter for comparative analysis. In practice, Full House Modeling can therefore be implemented via observed-to-rank mappings without requiring explicit specification of the latent distribution.

The representation in (2) may appear as a modeling choice. The following result shows that, under natural invariance and rank-based assumptions, this construction is in fact essentially unavoidable.

Theorem 2.2 (Characterization of Full House Coupling)**.** *Fix a population i. Let $X_{i,1}, \ldots, X_{i,N_i}$ be i.i.d. from a continuous distribution F_X. Suppose there exists a set $\mathcal{R}_i \subset \{1, \ldots, N_i\}$ such that $S_{i,j} = 1 \iff r_{i,j} \in \mathcal{R}_i$, where $r_{i,j}$ is the rank of $X_{i,j}$ among $(X_{i,1}, \ldots, X_{i,N_i})$.*

For each observed rank j, suppose there exists a measurable strictly increasing function $\psi_{i,j} : \mathbb{R} \to \mathbb{R}$ such that $Y_{i,(j)} = \psi_{i,j}\big(X_{i,(r_{i,j})}\big)$ a.s., and assume $F_{Y_{i,(j)}}$ is continuous. Then, for each observed rank j,

$$Y_{i,(j)} = F_{Y_{i,(j)}}^{-1} \circ F_{X_{i,(r_{i,j})}}\big(X_{i,(r_{i,j})}\big) \quad a.s.$$

Moreover, this representation is unique up to null sets and coincides with that induced by the Full House Model in Definition 2.1.

2.1 Interpretation and scope of the rank-preserving assumption

The coupling in Definition 2.1 is intentionally structural rather than measurement-error based. Under Full House Modeling, observed outcomes are generated through a rank-preserving transformation of latent aptitude among selected individuals, so that stochasticity enters through the latent population and the selection mechanism rather than through additive observational noise. The framework is not intended as a literal noiseless measurement model. Rather, the framework is intended as an idealized abstraction of settings in which observed rankings primarily reflect latent ordering after strong selection into the observed system.

This abstraction is most plausible in highly selective environments, where observed individuals represent the upper tail of a much larger latent population and observed outcomes are designed to discriminate among top performers. Professional sports provide a canonical example: although season-level outcomes contain substantial variability, long-run elite performance is strongly constrained by selection into the league itself. In such settings, rank-preserving approximations provide a tractable framework for studying how population depth and selection interact with observed achievement.

More generally, Full House Modeling should be viewed as a structural model for latent-observed comparison under selective observation, rather than as a literal generative model for noisy repeated measurements. Extending the framework to incorporate stochastic measurement error, imperfect ranking preservation, or probabilistic selection mechanisms is an important direction for future work.

3 Paradoxes Under Identical Achievement Distributions

In this section, we derive explicit probabilities of the aptitude–achievement paradox, as defined in (1), within a Full House Modeling framework involving two populations. We focus on the setting in which the observed achievement distributions are identical across contexts,

that is, $F_{Y_i} = F_Y$ for $i = 1, 2$ in Definition 2.1. The following definition formalizes the notion of paradox used throughout the remainder of the article.

Definition 3.1. *Consider the Full House Model in Definition 2.1 with two populations denoted by $i = 1, 2$. Let $j \in \{1, \ldots, n_1\}$ and $k \in \{1, \ldots, n_2\}$ denote fixed observed ranks in the two populations, where n_i is the number of observed individuals in population i. Thus, $Y_{1,(j)}$ and $Y_{2,(k)}$ are the jth and kth order statistics of the observed achievements in populations 1 and 2, respectively, and $X_{1,(r_{1,j})}$ and $X_{2,(r_{2,k})}$ are the corresponding latent aptitudes of those same individuals. We say that an aptitude-achievement paradox has occurred when both $X_{1,(r_{1,j})} > X_{2,(r_{2,k})}$ and $Y_{1,(j)} < Y_{2,(k)}$ have occurred. The aptitude-achievement paradox probability is defined as*

$$\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right). \tag{5}$$

As a simple illustration, consider the case in which the observed individuals in population i are exactly those with the top n_i latent aptitude values, so that $r_{i,t} = N_i - n_i + t$, for $t = 1, \ldots, n_i$. In particular, if we take $j = n_1$ and $k = n_2$, then $Y_{1,(n_1)}$ and $Y_{2,(n_2)}$ are the largest observed achievements in the two populations, and their corresponding latent aptitudes are $X_{1,(r_{1,n_1})} = X_{1,(N_1)}$ and $X_{2,(r_{2,n_2})} = X_{2,(N_2)}$.

If, moreover, $Y_{1,(n_1)} < Y_{2,(n_2)}$ a.s., then the achievement inequality in (5) holds automatically, so the aptitude–achievement paradox event reduces to $\{X_{1,(N_1)} > X_{2,(N_2)}\}$. Hence in this case,

$$\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})},\, Y_{1,(j)} < Y_{2,(k)}\right) = \mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}\right).$$

Under the conditions described below, this probability tends to 1 as $N_1 \to \infty$. This example shows that the paradox can arise even in a particularly simple setting where the observed individuals are the top latent-aptitude performers in each population.

We now introduce a convenient representation which allows the paradox probability to be expressed in terms of the latent aptitudes alone.

Lemma 3.2. *Consider the Full House Model in Definition 2.1 with two populations, where* $F_{Y_i} = F_Y$ *for* $i = 1, 2$. *Then,*

$$\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right) = \mathrm{P}\left(X_{2,(r_{2,k})} < X_{1,(r_{1,j})} < h_{j,k}\left(X_{2,(r_{2,k})}\right)\right), \tag{6}$$

$$\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right) = \mathrm{P}\left(g_{j,k}\left(Y_{2,(k)}\right) < Y_{1,(j)} < Y_{2,(k)}\right) \tag{7}$$

where

$$h_{j,k}(\cdot) = F^{-1}_{X_{1,(r_{1,j})}} \circ F_{Y_{1,(j)}} \circ F^{-1}_{Y_{2,(k)}} \circ F_{X_{2,(r_{2,k})}}(\cdot), \qquad g_{j,k}(\cdot) = F^{-1}_{Y_{1,(j)}} \circ F_{X_{1,(r_{1,j})}} \circ F^{-1}_{X_{2,(r_{2,k})}} \circ F_{Y_{2,(k)}}(\cdot).$$

Our first formal result shows that, under a monotonicity condition on the Beta-induced distortions, the paradox probability admits a closed-form expression.

Definition 3.3. *Let* $\alpha, \beta > 0$. *Then the regularized incomplete beta function is defined as*

$$I_{\alpha,\beta}(x) = \frac{1}{B(\alpha,\beta)} \int_0^x u^{\alpha-1}(1-u)^{\beta-1} du,$$

where $B(\alpha,\beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$ *is the beta coefficient.*

Theorem 3.4. *Consider the Full House Model in Definition 2.1 with two populations, where* $F_{Y_i} = F_Y$ *for* $i = 1, 2$. *Let* $j \in \{1, \ldots, n_1\}$ *and* $k \in \{1, \ldots, n_2\}$ *denote observed ranks, with corresponding latent ranks* $r_{1,j}$ *and* $r_{2,k}$. *Let* $I_{\alpha,\beta}$ *denote the regularized incomplete beta function as in Definition 3.3. Suppose that*

$$I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1}(t) - I_{r_{1,j},N_1-r_{1,j}+1} \circ I^{-1}_{r_{2,k},N_2-r_{2,k}+1}(t) \;\geq\; 0, \quad \text{for all } t \in (0,1). \tag{8}$$

Then the paradox probability admits the explicit form

$$\mathrm{P}\big(X_{1,(r_{1,j})} > X_{2,(r_{2,k})},\, Y_{1,(j)} < Y_{2,(k)}\big)$$

$$= \sum_{p=j}^{n_1} \binom{n_1}{p} \frac{B(k+p,\, n_1+n_2-p-k+1)}{B(k, n_2-k+1)} - \sum_{p=r_{1,j}}^{N_1} \binom{N_1}{p} \frac{B(r_{2,k}+p,\, N_1+N_2-p-r_{2,k}+1)}{B(r_{2,k}, N_2-r_{2,k}+1)}.$$

Remarks:

1. The paradox probability in Theorem 3.4 is distribution free. Once the Full House Model is specified, it depends only on the population sizes N_i, system sizes n_i, the system inclusion function, and the order-statistic indices. This follows from the Beta representations (3)–(4), which eliminate dependence on F_Y. In particular,

$$\begin{aligned} h_{j,k}(x) &= F^{-1}_{X_{1,(r_{1,j})}} \circ I_{j,n_1-j+1} \circ \left(F_Y \circ F_Y^{-1}\right) \circ I^{-1}_{k,n_2-k+1} \circ F_{X_{2,(r_{2,k})}}(x) \\ &= F^{-1}_{X_{1,(r_{1,j})}} \circ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1} \circ F_{X_{2,(r_{2,k})}}(x), \end{aligned}$$

   so the dependence on F_Y cancels.

2. The latent aptitude distribution F_X likewise plays no role in the final expression. Although Lemma 3.2 represents the event in terms of latent order statistics, the monotone coupling induced by Full House Modeling ensures that all dependence on F_X cancels after transformation. Thus F_X acts as a structural nuisance parameter for paradox probabilities.

3. Combining these observations, the paradox probability admits the representation

$$\int_0^1 \left\{ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1}(t) - I_{r_{1,j},N_1-r_{1,j}+1} \circ I^{-1}_{r_{2,k},N_2-r_{2,k}+1}(t) \right\}^+ dt.$$

   The positive-part operator reflects that the integrand may change sign. The technical condition (8) ensures that the integrand is nonnegative for all $t \in (0,1)$, allowing it to be removed and yielding the closed-form expression in Theorem 3.4. We characterize settings where this condition holds in subsequent results.

A natural case of interest is the comparison of individuals with extreme latent aptitudes

across populations. The following corollary provides a closed-form expression for the paradox probability in this setting under a Full House Model with $F_{Y_i} = F_Y$ for $i = 1, 2$.

Corollary 3.4.1. *Suppose that the conditions of Theorem 3.4 hold and that the selection indicators satisfy $S_{i,(N_i)} = 1$ for $i = 1, 2$, that is, the highest-aptitude individual in each population is included in the observed system. Then,*

$$\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \begin{cases} \frac{n_2}{n_1+n_2} - \frac{N_2}{N_1+N_2}, & \text{if } \frac{n_1}{N_1} \leq \frac{n_2}{N_2}, \\ 0, & \text{otherwise.} \end{cases}$$

Intuitively, this is the difference between two 'lottery' probabilities: the probability that individual 2 wins the observed achievement comparison (first term) minus the probability that individual 2 wins the latent aptitude comparison (second term). The paradox probability is positive precisely when the sampling ratio favors individual 2 more than the population ratio does. An analogous result holds for comparisons of individuals with the lowest latent aptitudes:

Corollary 3.4.2. *Suppose that the conditions of Theorem 3.4 hold and that the selection indicators satisfy $S_{i,(1)} = 1$ for $i = 1, 2$, that is, the lowest-aptitude individual in each population is included in the observed system. Then,*

$$\mathrm{P}\left(X_{1,(1)} > X_{2,(1)}, Y_{1,(1)} < Y_{2,(1)}\right) = \begin{cases} \frac{N_2}{N_1+N_2} - \frac{n_2}{n_1+n_2}, & \text{if } \frac{n_1}{N_1} \geq \frac{n_2}{N_2}, \\ 0, & \text{otherwise.} \end{cases}$$

We now demonstrate a setting in which technical condition (8) holds. This setting is based on a configuration of (n_i, N_i) for $i = 1, 2$ under the assumption that the systems are populated with individuals from the source populations who have the highest aptitudes.

Proposition 3.5. *Consider the Full House Model in Definition 2.1 with two populations, where $F_{Y_i} = F_Y$ for $i = 1, 2$, and suppose that the selection indicators satisfy $S_{i,j} = 1 \iff$*

$X_{i,j} \geq X_{i,(N_i - n_i + 1)}$ *for* $i = 1, 2$. *Denote by* I *the regularized incomplete beta function. Suppose that* $\frac{n_1}{n_2} \leq 1$ *and* $\frac{N_2}{N_1} \leq 1$, *and set* $k = j$. *Under these assumptions, we conclude that technical condition* (8) *holds and the conclusion of Theorem 3.4 applies.*

The conditions in Proposition 3.5 suggest that the technical condition is driven by the relative comparison of sampling and population ratios. In particular, the requirement $\frac{n_1}{n_2} \leq 1$ and $\frac{N_2}{N_1} \leq 1$ indicates that sampling distortion dominates population distortion in a coordinatewise sense.

4 Paradox Under Different Distributions

In this section we will provide aptitude-achievement paradox probability results in a Full House Modeling setting involving two populations when we relax the specification that $F_{Y_1} = F_{Y_2}$ in Definition 2.1. We first provide bounds for aptitude-achievement paradox probability in general settings. We then provide explicit aptitude-achievement paradox probability results when Y_1 and Y_2 are normally distributed with different means and variances.

4.1 General bounds

The general bounds we derive require a relaxed form of first-order stochastic dominance under which the distributions of Y_1 and Y_2 are 'close' to each other. More formally, we introduce first-order ε-weak stochastic dominance, which is necessary for our aptitude-achievement paradox probability results.

Definition 4.1. *We will say that* Y_2 *is first order* ε*-weakly stochastically dominant over* Y_1 *if, for some* $\varepsilon > 0$, $F_{Y_2}(y) \leq F_{Y_1}(y) \leq F_{Y_2}(y) + \varepsilon$ *for all* y, *and there exists some* y *for which* $F_{Y_2}(y) < F_{Y_1}(y)$.

The condition $F_{Y_2}(y) \leq F_{Y_1}(y)$, for all y is exactly the usual first-order stochastic dominance condition [Quirk and Saposnik, 1962], so Y_1 is stochastically larger than Y_2. The

additional upper bound $F_{Y_1}(y) \leq F_{Y_2}(y) + \varepsilon$ does not reverse the dominance direction; rather, it requires that the pointwise gap between the two distribution functions be uniformly bounded by ε. Thus Y_2 dominates Y_1, but only by at most ε in the Kolmogorov distance on the one-sided scale. The strict inequality at some y excludes the degenerate case $F_{Y_1} \equiv F_{Y_2}$.

Under first-order ε-weak stochastic dominance, we derive a general lower and upper bound on the aptitude-achievement paradox probability when $F_{Y_1} \neq F_{Y_2}$.

Proposition 4.2. *Consider the Full House Model in Definition 2.1 with two populations $i = 1, 2$. Suppose that the selection indicators satisfy $S_{i,(N_i)} = 1$. Further suppose that Y_2 is first order ε-weakly stochastically dominant over Y_1 for some $\varepsilon > 0$ and $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$. Then,*

$$\begin{aligned} \frac{n_2}{n_1 + n_2} - \frac{N_2}{N_1 + N_2} &\leq \mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\ &\leq n_2 \sum_{i=0}^{n_1} \frac{\binom{n_1}{i}\varepsilon^i}{n_1 + n_2 - i} - \frac{N_2}{N_1 + N_2} \end{aligned} \tag{9}$$

Proposition 4.2 provides bounds on the paradox probability under general achievement distributions. For example, when $Y_i \sim N(\mu_i, \sigma^2)$, $i = 1, 2$, with $\mu_1 < \mu_2$ and common variance, ε can be bounded explicitly as a function of $(\mu_2 - \mu_1)/\sigma$. Beyond such bounds, we can obtain explicit aptitude-achievement paradox probabilities when $Y_i \sim N(\mu_i, \sigma_i^2)$, $i = 1, 2$, allowing variances to differ across contexts.

4.2 Normal distributions

This section develops explicit aptitude-achievement paradox probabilities (5) when a Full House Model in Definition 2.1 specifies $Y_i \sim N(\mu_i, \sigma_i^2)$, $i = 1, 2$. We focus on systems where $S_{i,(N_i)} = 1$ for $i = 1, 2$. The outline is similar to Section 3 in the sense that a technical condition is needed to prove the main result, and further results provide configurations satisfying this condition. However, with $Y_i \sim N(\mu_i, \sigma_i^2)$, the technical condition yields a remainder term that prevents an exact closed-form expression. We show that this remainder

is negligible under mild conditions on population fractions and distributional parameters.

Similar to the proof of Theorem 3.4, Lemma 3.2 will provide a representation more conducive to formal integration needed to prove our main result. Only this time we will use (7) in Lemma 3.2 to represent the aptitude-achievement paradox probability as

$$\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \mathrm{P}\left(g_{n_1,n_2}\left(Y_{2,(n_2)}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right),$$

where, from the definition of g in Lemma 3.2 applied to our current setting, we have

$$g_{n_1,n_2}(x) = F_{Y_1}^{-1}\left(F_{Y_2}(x)^{\frac{n_2 N_1}{N_2 n_1}}\right) = \Phi^{-1}\left(\Phi\left(\frac{x-\mu_2}{\sigma_2}\right)^{\frac{n_2 N_1}{N_2 n_1}}\right)\sigma_1 + \mu_1, \tag{10}$$

and Φ defines the cumulative distribution function of the standard normal distribution. Integrating over the joint order-statistic distribution yields

$$\mathrm{P}\left(g_{n_1,n_2}(Y_{2,(n_2)}) < Y_{1,(n_1)} < Y_{2,(n_2)}\right) = I + R, \tag{11}$$

where

$$I := \int_{-\infty}^{\infty}\left(F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2 N_1}{N_2}}\right) dF_{Y_2,(n_2)}(x), \tag{12}$$

and

$$R := -\int_{A^c}\left(F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2 N_1}{N_2}}\right) dF_{Y_2,(n_2)}(x), \tag{13}$$

with

$$A = \left\{x : F_{Y_1}^{-1}\left(F_{Y_2}(x)^{\frac{n_2 N_1}{N_2 n_1}}\right) \leq x\right\}. \tag{14}$$

Under the Gaussian specification,

$$A = \left\{x \in \mathbb{R} : \Phi\left(\frac{x-\mu_2}{\sigma_2}\right)^{\frac{n_2}{N_2}} \leq \Phi\left(\frac{x-\mu_1}{\sigma_1}\right)^{\frac{n_1}{N_1}}\right\}.$$

The following lemma shows that the remainder term is negligible under mild parameter

restrictions.

Lemma 4.3. *Assume the setting leading to the decomposition (11), and let R denote the corresponding remainder term (13). If $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$, $\sigma_1 \leq \sigma_2$, and $\mu_1 \leq \mu_2$, then*

$$|R| \leq \left(\frac{1}{2}\right)^{n_2}. \tag{15}$$

Therefore, although the remainder term in (13) arises from the restriction of the integration domain to A, it is exponentially small in n_2 under a familiar ordering of population fractions (see Proposition 4.2) and a simple parametric configuration. Consequently, the aptitude–achievement paradox probability is well approximated by the unrestricted integral given by (12) with R contributing only a negligible correction for moderately large n_2.

We now introduce a multivariate normal representation allowing a closed-form expression of the principal term. Let $\mathbf{W} \sim \mathcal{N}_{n_1+n_2-1}(\mathbf{0}, \mathbf{\Sigma})$, with block covariance matrix

$$\mathbf{\Sigma} = \begin{bmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{12}^\top & \Sigma_{22} \end{bmatrix}, \tag{16}$$

where

$$\Sigma_{11} = I_{n_1} + \frac{\sigma_2^2}{\sigma_1^2} J_{n_1}, \qquad \Sigma_{12} = \frac{\sigma_2}{\sigma_1} \mathbf{1}_{n_1} \mathbf{1}_{n_2-1}^\top, \qquad \Sigma_{22} = I_{n_2-1} + J_{n_2-1}. \tag{17}$$

where I_k denotes the $k \times k$ identity matrix and $J_k = \mathbf{1}_k \mathbf{1}_k^\top$ denotes the $k \times k$ matrix of ones. With this structure we now provide an explicit aptitude-achievement paradox probability, the main result of this section.

This setup yields an aptitude-achievement paradox result when cross-system achievement follows different normal distributions. The result is essentially closed form under parametric specification and a ratio sample size condition that allow the remainder term (13) to be bounded above by a term decaying exponentially in n_2.

Theorem 4.4. *Assume a Full House Model in Definition 2.1 and additionally specify that* $Y_i \sim N(\mu_i, \sigma_i^2)$*, and suppose that the selection indicators satisfy* $S_{i,(N_i)} = 1$ *for* $i = 1, 2$*. Let* $\mathbf{W} \sim \mathcal{N}_{(n_1+n_2-1)}(\mathbf{0}, \boldsymbol{\Sigma})$ *where* $\boldsymbol{\Sigma}$ *is as defined in (16) and (17). Then,*

$$\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = n_2 F_{\mathbf{W}}(\mathbf{w}) - \frac{N_2}{N_1 + N_2} + R, \tag{18}$$

where R *is defined in (13),* $F_{\mathbf{W}}$ *is the distribution function of* $\mathbf{W}$*, and the first* n_1 *elements of* $\mathbf{w}$ *are* $\frac{\mu_2 - \mu_1}{\sigma_1}$ *and the last* $n_2 - 1$ *elements are 0. Moreover, if* $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$*,* $\sigma_1 \leq \sigma_2$*, and* $\mu_1 \leq \mu_2$*, then* $|R| \leq \left(\frac{1}{2}\right)^{n_2}$*.*

We now show that plug-in estimators obtained by replacing μ_i and σ_i, $i = 1, 2$, with consistent estimators yield asymptotically valid estimates of the paradox probability in (18).

Theorem 4.5. *Assume the setting of Theorem 4.4 with* $\sigma_1 < \sigma_2$*, and* $\mu_1 < \mu_2$*, and let* $\widehat{F}_{\mathbf{W}}(\mathbf{w})$ *and* $\widehat{\mathrm{P}}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right)$*, respectively, denote the plug-in estimator of* $F_{\mathbf{W}}(\mathbf{w})$ *and the paradox probability in (18) obtained by replacing* μ_i, σ_i *with consistent estimators* $\hat{\mu}_i, \hat{\sigma}_i$*,* $i = 1, 2$*. Suppose*

$$\frac{\hat{\sigma}_1}{\hat{\sigma}_2} - \frac{\sigma_1}{\sigma_2} = o_P\left(\frac{1}{\log n_1}\right), \qquad \frac{\hat{\mu}_2 - \hat{\mu}_1}{\hat{\sigma}_2} - \frac{\mu_2 - \mu_1}{\sigma_2} = o_P\left(\frac{1}{\sqrt{\log n_1}}\right).$$

Then, $\frac{\widehat{F}_{\mathbf{W}}(\mathbf{w})}{F_{\mathbf{W}}(\mathbf{w})} \xrightarrow{P} 1$ *and*

$$\widehat{\mathrm{P}}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) - \mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) \xrightarrow{P} 0.$$

Remark: The rates in Theorem 4.5 arise from the logarithmic scaling appearing in Gaussian extreme-value approximations (see Embrechts et al. [2013]) and are relatively weak. In particular, they are satisfied whenever

$$\hat{\mu}_i - \mu_i = O_P(n_i^{-1/2}), \qquad \hat{\sigma}_i - \sigma_i = O_P(n_i^{-1/2}),$$

as holds for standard plug-in estimators such as maximum likelihood estimators, provided that $n_2 = O(n_1)$.

5 Empirical Illustrations

We briefly illustrate the interpretability and operational estimation of paradox probabilities under several representative configurations.

5.1 Sensitivity Analysis for Theorem 3.4

Under the setting of Theorem 3.4, we will specify $X_1 \overset{d}{=} X_2 \overset{d}{=} X$, $Y_1 \overset{d}{=} Y_2 \overset{d}{=} Y$, where $X \sim \text{Pareto}(\alpha)$ and $Y \sim \mathcal{N}(0,1)$. Without loss of generality, we set $j = n_1 - 1$ and $k = n_2 - 1$, corresponding to the selection of the second-best performer in the aptitude-achievement paradox probability, as defined in Definition 3.1.

In each simulation, we randomly sample n_1 and n_2 observations from the distribution of Y, and compute their latent aptitudes using Equation (2). Based on these values, we directly calculate paradox probabilities according to Definition 3.1. We repeat this simulation 10000 times and report the average.

Figure 1 presents the results of the sensitivity analysis for the aptitude-achievement paradox probability described in Theorem 3.4. In Figure 1 (A), we fix $n_1 = 40$ and $n_2 = 60$, and vary N_1 and N_2 from 200 to 2000. In Figure 1 (B), we fix $N_1 = 10^3$ and $N_2 = 10^3$, and vary n_1 and n_2 from 20 to 200. In Figure 1 (C), we fix $n_1 = 40$ and $N_2 = 10^3$, and vary n_2 from 20 to 200 and N_1 from 200 to 2000. In Figure 1 (D), we fix $N_1 = 10^3$ and $n_2 = 60$, and vary N_2 from 200 to 2000 and n_1 from 20 to 200.

As shown in Figure 1 (A), a small N_2 and large N_1 increase the paradox probability by making $\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}\right) \to 1$. Similarly, in Figure 1 (B), a small n_1 combined with a large n_2 leads to a higher aptitude-achievement paradox probability since it drives $\mathrm{P}\left(Y_{1,(j)} < Y_{2,(k)}\right) \to 1$.




Figure 1: Sensitivity Analysis for Paradox Probability

5.2 Ratio of N_2 and N_1

Here we investigate the effect of the ratio between N_2 and N_1 on aptitude-achievement paradox probabilities under the specifications of Corollary 3.4.1. Let $c = \frac{N_2}{N_1}$, and vary c from 0.1 to 1 in increments of 0.1. In each simulation, we fix $n_1 = 60$ and $n_2 = 40$, and vary N_1 from 100 to 2000. We repeat this simulation 10000 times and report the average paradox probability.

Figure 2 presents the results. As shown in the figure, the paradox probability decreases as c increases, which is reasonable since $\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}\right)$ decreases as N_2 increases. Importantly, Figure 2 also demonstrates that the paradox probability remains constant when the ratio between N_2 and N_1 is fixed. This implies that it is unnecessary to focus on the exact values of N_1 and N_2 when computing the paradox probability; rather, the ratio N_2/N_1 alone determines its value.




Figure 2: Effect of the ratio between N_2 and N_1 on aptitude-achievement paradox probabilities under the specifications of Corollary 3.4.1.

5.3 Batting Averages in Major League Baseball

We illustrate Theorem 4.4 and the plug-in guarantees of Theorem 4.5 using Major League Baseball (MLB), a setting in which latent batting aptitude is indirectly observed through season-level batting averages and where both population structure and measurement environments vary substantially across eras. Figure 3 displays smoothed estimates of the mean and standard deviation of batting averages for full-time players from 1871–2023.

We treat the 2023 season as system 1 and each season from 1871–2022 as system 2. Seasonal batting averages for full-time players are modeled as Gaussian with means and standard deviations estimated via plug-in; the consistency guarantees of Theorem 4.5 therefore apply. Figure 3 shows that the 2023 offensive environment exhibited both relatively low mean batting average and reduced variability compared with many historical eras. Under




Figure 3: Seasonal means and standard deviations of batting averages for full-time players from 1871–2023. Smoothing is applied to both trends.

our specifications, we additionally have $\hat{\sigma}_1 \leq \hat{\sigma}_2$, $\hat{\mu}_1 \leq \hat{\mu}_2$, and $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$, implying that the remainder term (13) in Theorem 4.4 is negligible. Talent-pool estimates N_i are taken from Yan et al. [2025], which includes sensitivity analyses demonstrating robustness to reasonable perturbations of these estimates.

Figure 4 shows that estimated paradox probabilities closely track empirical exceedance patterns across eras, indicating that the proposed framework captures structural differences in competitive depth and measurement environment rather than merely reproducing descriptive trends. In particular, the framework explains why Luis Arraez's 2023 batting average may appear modest under raw historical comparison while nevertheless corresponding to unusually high latent aptitude after accounting for population depth and contextual variation. Under Full House Modeling, Arraez's 2023 season ranks 68th all-time among full-time batters after adjustment [Yan et al., 2025, Eck et al., 2025], compared with a raw observed




Figure 4: Comparison of estimated paradox probabilities and empirical exceedance rates relative to the 2023 season, based on data from 1871–2022. The blue line shows plug-in paradox probability estimates from Theorem 4.4; the dashed black line shows a smoothed empirical exceedance rate for Luis Arraez's 2023 batting average.

rank of 341. This gap illustrates how cross-context comparisons based solely on observed outcomes can induce substantial latent-observed ranking reversals, precisely the phenomenon quantified by the paradox probability.

6 Discussion

The aptitude-achievement paradox arises when an individual from one context demonstrates higher latent aptitude but simultaneously lower observed achievement than an individual from another context. Although widely encountered in education, labor economics, psychometrics, medicine, and sports analytics, measuring the frequency of this paradox under realistic data-generating regimes is nontrivial. In this article, we formalize Full House Modeling and investigate the probabilities of observing paradoxes between latent aptitude

and observed achievement in comparative analysis of individuals across contexts.

This work is a theoretical clarification of Full House Modeling and provides the definition and solutions for an inferential target, the aptitude-achievement paradox, which extend beyond the baseball motivations in Yan et al. [2025]. In doing so, we note the baseball origins of this work, as articulated by Stephen Jay Gould in his 1996 book *Full House: The Spread of Excellence from Plato to Darwin* (p. 122), from which this framework takes its name: "Baseball is a truly remarkable system for statisticians, manifesting two properties devoutly to be wished, but not often encountered, in actual data: an institution that has worked by the same rules for a century, and has compiled complete data (nothing major missing) on all measurable aspects of its history." In this way, baseball, with its century of consistent rules and comprehensive measurement, served as an ideal proving ground for the Full House framework [Yan et al., 2025]. Gould's discussion emphasized the possibility that population structure and selection may substantially influence observed achievement distributions across eras. The present work formalizes one probabilistic framework through which such effects can be quantified and analyzed.

We derive closed-form expressions for paradox probabilities under Full House data generating models in two regimes: (i) when observed achievements are i.i.d., and (ii) when context-specific summaries are independent normals with distinct means and variances. The probability rises with population-size asymmetry (large N_1, small N_2) and sampling asymmetry (small n_1, large n_2). Moreover, for a fixed population-size ratio N_2/N_1, it is unchanged by proportional rescaling of the sample sizes (n_1, n_2). We also develop bounds for more general settings and show that, when observed achievement is normal, plug-in estimators yield reliable paradox-probability estimates.

The present analysis focuses on rank-preserving structural couplings and highly selective observation regimes, particularly top-n selection mechanisms. Extending the framework to settings involving stochastic measurement error, imperfect ranking preservation, probabilistic selection, or heterogeneous latent aptitude distributions remains an important direction

for future work. Likewise, while the present paper studies pairwise paradox probabilities, broader ranking-level functionals may also be of interest.

Beyond baseball, the aptitude-achievement paradox has potential implications for educational evaluation and admissions. When students are compared across institutions of differing selectivity, observed performance metrics may systematically misrepresent underlying aptitude due to differences in competition depth, selection mechanisms, and measurement environments. The methodology developed here provides a formal way to quantify such latent-observed reversals under explicit assumptions on selection and measurement processes. More broadly, the framework highlights how cross-context comparisons based solely on observed outcomes can become sensitive to differences in population structure, selection intensity, and measurement environment.

By formalizing aptitude-achievement reversals as an estimable probabilistic quantity, this work provides a framework for studying how population depth, selective observation, and measurement structure jointly shape cross-context comparisons. More broadly, the results illustrate how latent-observed ranking reversals can arise systematically under heterogeneous selection and measurement regimes.

Acknowledgements

We are grateful to Gökçe Dayanıklı, Karl Oskar Ekvall, Eren C Kızıldağ, Jingbo Liu, Susu Zhang and many others for helpful comments, great discussions, and inspiration. We acknowledge the use of ChatGPT and Claude to aid with proofreading and writing clarity.

Supplementary Materials

The Supplementary Materials contain proofs of mathematical results, together with auxiliary Monte Carlo simulations provided solely for numerical intuition for Theorem 4.4.

References

David J Bartholomew, Martin Knott, and Irini Moustaki. *Latent variable models and factor analysis: A unified approach*. John Wiley & Sons, 2011.

Benjamin S Baumer and Gregory J Matthews. A statistician reads the sports page: there is no avoiding WAR. *Chance*, 27(3):41–44, 2014.

Benjamin S Baumer, Shane T Jensen, and Gregory J Matthews. openwar: An open source system for evaluating overall player performance in major league baseball. *Journal of Quantitative Analysis in Sports*, 11(2):69–84, 2015.

Scott M Berry, C Shane Reese, and Patrick D Larkey. Bridging different eras in sports. *Journal of the American Statistical Association*, 94(447):661–676, 1999.

Trevor G Bond and Christine M Fox. *Applying the Rasch model: Fundamental measurement in the human sciences*. Psychology Press, 2013.

Ralph A. Bradley. Science, statistics, and paired comparisons. *Biometrics*, 32(2):213–239, 1976. doi: 10.2307/2529494.

Ralph A. Bradley. Paired comparisons: Some basic procedures and examples. In P. R. Krishnaiah and P. K. Sen, editors, *Handbook of Statistics*, volume 4, pages 299–326. Elsevier, Amsterdam, 1984.

Ralph A. Bradley and Milton E. Terry. The rank analysis of incomplete block designs: I. the method of paired comparisons. *Biometrika*, 39:324–345, 1952. doi: 10.1093/biomet/39.3-4.324.

Wei Chu and Zoubin Ghahramani. Preference learning with gaussian processes. In *Proceedings of the 22nd International Conference on Machine Learning*, pages 137–144. ACM, 2005. doi: 10.1145/1102351.1102369.

Herbert A. David. *The Method of Paired Comparisons*. Chapman and Hall, London, 2 edition, 1988.

Daniel Eck, Jun Hee Kim, Shen Yan, Muskan Bhatla, and Zheer Wang. *fullhouse: Package for Estimating Talent and Era-Adjusted Performance Metrics*, 2025. URL https://github.com/DEck13/fullhouse. R package version 1.1.

Bradley Efron and Carl Morris. Data analysis using stein's estimator and its generalizations. *Journal of the American Statistical Association*, 70(350):311–319, 1975.

Arpad E. Elo. *The Rating of Chessplayers, Past and Present*. Arco Publishing, 1978.

Paul Embrechts, Claudia Klüppelberg, and Thomas Mikosch. *Modelling extremal events: for insurance and finance*, volume 33. Springer Science & Business Media, 2013.

Susan E Embretson and Steven P Reise. *Item response theory for psychologists*. Psychology Press, 2000.

Thore Graepel, Tom Minka, and R TrueSkill Herbrich. A bayesian skill rating system. *Advances in Neural Information Processing Systems*, 19(569-576):7, 2007.

Jacques A Hagenaars and Allan L McCutcheon. *Applied latent class analysis*. Cambridge University Press, 2002.

Paul W Holland and Howard Wainer. *Differential item functioning*. Routledge, 2012.

Neil Houlsby, Ferenc Huszar, Zoubin Ghahramani, and Jose Hernández-lobato. Collaborative gaussian processes for preference learning. *Advances in neural information processing systems*, 25, 2012.

Maurice G Kendall. A new measure of rank correlation. *Biometrika*, 30(1-2):81–93, 1938.

Roderick P McDonald. *Test theory: A unified treatment*. psychology press, 1999.

Stephen Nickell and Richard Layard. Labor market institutions and economic performance. *Handbook of labor economics*, 3:3029–3084, 1999.

James P Quirk and Rubin Saposnik. Admissibility and measurable utility functions. *The Review of Economic Studies*, 29(2):140–146, 1962.

Stephen Raudenbush and Anthony S Bryk. A hierarchical model for studying school effects. *Sociology of education*, pages 1–17, 1986.

Michael J Schell. *Baseball's all-time best hitters*. Princeton University Press, 2013.

Michael J Schell. *Baseball's all-time best sluggers*. Princeton University Press, 2016.

Robert H Somers. A new asymmetric measure of association for ordinal variables. *American sociological review*, pages 799–811, 1962.

Wayne L Winston, Scott Nestler, and Konstantinos Pelechrinis. *Mathletics: How Gamblers, Managers, and Fans use Mathematics in Sports*. Princeton University Press, 2022.

Shen Yan, Adrian Burgos, Jr, Christopher Kinson, and Daniel J Eck. Comparing baseball players across eras via novel full house modeling. *The Annals of Applied Statistics*, 19(2): 1778–1799, 2025.

Amy Zhao Yu, Shahar Ronen, Kevin Hu, Tiffany Lu, and César A Hidalgo. Pantheon 1.0, a manually verified dataset of globally famous biographies. *Scientific data*, 3(1):1–16, 2016.