A Formal Investigation of Paradoxes Between Latent Aptitude and Observed Achievement via Full House Methodology

Daniel J. Eck[1], Shen Yan[1*]

1. Department of Statistics, University of Illinois Urbana-Champaign

Abstract

We study a paradox in which an individual with greater latent aptitude exhibits lower observed achievement than an individual with lesser aptitude when comparisons are made across contexts. We formalize this phenomenon as the aptitude-achievement paradox and introduce its probability as an explicit inferential target. To study this quantity, we employ Full House Modeling, a framework that derives latent aptitude from observed achievement by embedding observed samples within larger source populations. We derive closed-form paradox probabilities when achievement distributions are identical across contexts and obtain explicit expressions and bounds when they differ, including tractable results under normality with reliable plug-in estimation. Our results show that paradox risk depends primarily on population and sampling ratios rather than absolute sizes. An application to Major League Baseball contextualizes historically extreme batting performances, showing how modern achievements may appear less exceptional after accounting for population depth and measurement environments. By elevating aptitude-achievement reversals from anecdotal concern to an estimable

*Daniel J. Eck (dje13@illinois.edu); email addresses: Shen Yan (shenyan3@illinois.edu)

quantity, this work provides a principled foundation for fairer and more interpretable cross-context comparisons.

1 Introduction

Comparisons of performances across differing contexts, schools, regions, eras, institutions, and tasks are central to education and psychometrics [Holland and Wainer, 2012, Bond and Fox, 2013], labor economics [Nickell and Layard, 1999], medicine, and sports analytics [Berry et al., 1999, Schell, 2013, 2016, Winston et al., 2022, Yan et al., 2025]. These problems have led to the development of a variety of widely used modeling frameworks that estimate relative ability, including empirical-Bayes shrinkage [Efron and Morris, 1975], Bayesian hierarchical models [Raudenbush and Bryk, 1986, Berry et al., 1999], Bradley-Terry-Luce and paired comparisons models [Bradley and Terry, 1952, Bradley, 1976, 1984, David, 1988], Elo-type dynamic rating systems [Elo, 1978, Graepel et al., 2007], Gaussian-process and state-space formulations [Chu and Ghahramani, 2005, Houlsby et al., 2012], item-response and latent class models [McDonald, 1999, Embretson and Reise, 2000, Hagenaars and McCutcheon, 2002, Holland and Wainer, 2012, Bond and Fox, 2013], factor models [McDonald, 1999, Bartholomew et al., 2011], and domain-specific metrics [Baumer and Matthews, 2014, Baumer et al., 2015, Yu et al., 2016].

Despite their differences, these approaches share a common structural assumption: ability is inferred relative to a fixed or implicitly comparable set of observed competitors. While such models adjust outcomes within a given context, they do not explicitly account for changes in the size or structure of the eligible population, nor for context-specific distortions in how latent ability is transformed into observed outcomes. As a consequence, they may not be able to distinguish observed dominance from underlying aptitude when competition depth, selection mechanisms, or measurement conditions vary across contexts.

These limitations give rise to a previously unquantified inferential target: the probabil-

ity that cross-context comparisons reverse latent orderings. Specifically, an individual drawn from a deeper or more competitive population may possess higher latent aptitude yet exhibit lower observed achievement than a counterpart from a shallower or more weakly filtered context. We term this phenomenon the aptitude-achievement paradox. Formally, let individual j in context i have an observed $\mathbb{R}$-valued statistic $Y_{i,j}$ and an underlying latent $\mathbb{R}$-valued aptitude $X_{i,j}$, and let individual n in context m have an observed $\mathbb{R}$-valued statistic $Y_{m,n}$, and an underlying latent $\mathbb{R}$-valued aptitude $X_{m,n}$. The probability of aptitude-achievement paradox is

$$\mathrm{P}\left(X_{i,j} > X_{m,n}, Y_{i,j} < Y_{m,n}\right) \tag{1}$$

This quantity captures the structural risk that cross-context comparisons based on observed outcomes reverse the true latent ordering.

The paradox cannot arise if a single, strictly increasing, context-invariant measurement map relates aptitude to achievement and if selection into tasks is ignorable. In practice, however, both measurement heterogeneity (non-invariance, differential item functioning) and non-ignorable selection (who attempts which tasks, faces which opponents, receives which treatment) are pervasive. These features can reverse cross-context rankings even when the latent ranking is unambiguous, echoing classic reversal phenomena such as Simpson's [Simpson, 1951] and Lord's paradoxes [Lord, 1967] while implicating modern concerns about fairness. What practitioners lack is a principled way to quantify how often such reversals should occur under an explicit model of aptitude, measurement, and selection.

A large class of methods quantifies how often observed orderings disagree with a target ordering, closely related to the phenomenon studied here. Classic examples include rank-discordance measures such as Kendall's τ [Kendall, 1938] and Somers' D [Somers, 1962], which summarize the frequency of concordant and discordant pairs based on observed rankings. Because these indices operate entirely on observed outcomes, they neither place individuals on a common latent scale nor disentangle the effects of measurement non-invariance or selection, motivating models that make the measurement and selection processes explicit.

Traditional latent trait models [Bartholomew et al., 2011] focus on estimating underlying abilities from observed categorical or discrete outcomes. For example, the Rasch model [Bond and Fox, 2013] ensures that comparisons of individuals' abilities are invariant to the specific items administered, under the assumption of no differential item functioning (DIF) [Holland and Wainer, 2012]. However, such models do not explicitly leverage information about the size or structure of the source population from which individuals are drawn. Likewise, Bayesian hierarchical and state-space models for latent ability—including Elo-type rating systems [Berry et al., 1999, Elo, 1978, Graepel et al., 2007]—typically condition on the observed competitive or testing pool and capture population-level variation through random effects or dynamic evolution, without modeling variation in eligible population depth or selection into observation across contexts. In parallel, the causal bootstrap [Imbens and Menzel, 2018] emphasizes that valid causal inference requires explicitly situating the observed sample within a larger source population, rather than treating the realized sample as the population of interest.

To address these limitations, we develop a framework for cross-context comparison that explicitly accounts for population depth and selection mechanisms. This framework, which we refer to as Full House Modeling (FHM), was initially proposed in an applied setting comparing baseball players across eras [Yan et al., 2025], and is developed here as a general approach for cross-contextual evaluation. This framework separates latent aptitude from observed achievement by modeling how three features jointly shape observed rankings: (i) the size of the eligible population in each context, (ii) the mechanism by which individuals are selected or observed, and (iii) the context-specific mapping from latent aptitude to observable performance. Under FHM, individuals are drawn from a common latent ability distribution, while contexts differ in population depth and in how aptitude is filtered and measured. This formulation makes explicit the structural mechanisms through which observed performance can be inflated or suppressed independently of true underlying aptitude.

In this article, we formalize Full House Modeling and study the probability of aptitude–

achievement paradoxes (1) arising in cross-context comparisons. We derive explicit paradox probabilities in two settings: first, when observed achievement is independent and identically distributed, and second, when observed achievement follows independent but context-specific normal distributions. We also establish bounds on paradox probabilities in more general settings and show that, under normality, these probabilities can be consistently estimated using plug-in methods. A sensitivity analysis reveals that paradox risk depends on ratios of eligible population sizes rather than their absolute magnitudes. Proofs of mathematical results are provided in online Supplementary Materials.

2 Model Setting

We now motivate the structure of the Full House Model. Let N_i denote the size of population i. We suppose that every individual $k = 1, \ldots, N_i$ has an underlying latent $\mathbb{R}$-valued aptitude $X_{i,1}, \ldots, X_{i,N_i} \overset{iid}{\sim} F_X$. Let $\mathbf{X}_{i,-j}$ be the vector of all latent aptitudes not including the component j. We then define an indicator function $1_i\left(X_{i,j}, \mathbf{X}_{i,-j}\right)$ as a determined and known system inclusion function for population i where a 1 indicates inclusion and a 0 indicates exclusion. We now suppose that an individual j included in system i has an observed $\mathbb{R}$-valued statistic $Y_{i,j} \overset{iid}{\sim} F_{Y_i}$, $j = 1, \ldots, n_i$, $n_i \leq N_i$. As we will see later, the observed $Y_{i,j}$ values are rank-preserving realizations of $X_{i,j}$, provided that $1_i\left(X_{i,j}, \mathbf{X}_{i,-j}\right) = 1$. Therefore $Y_{i,1}, \ldots, Y_{i,n_i} \overset{iid}{\sim} F_{Y_i}$.

Let $X_{i,(j)}$ and $Y_{i,(j)}$ denote the jth order statistic for system i. Here, larger numbers will correspond to a notion of "better." Thus $X_{i,(j)} > X_{i,(k)}$ whenever $j > k$, for example. We now form (Y, X) pairs of random variables for each individual included in the system. These pairs are of the form $(Y_{i,(j)}, X_{i,(r_{i,j})})$ where $r_{i,j}$ indexes the rank of that individual's latent aptitude within the full population of size N_i. As an example, let $1_i\left(X_{i,j}, \mathbf{X}_{i,-j}\right) = 1(X_{i,j} \geq X_{i,(N_i - n_i + 1)})$, then the $(Y_{i,(j)}, X_{i,(r_{i,j})})$ pairs take the form $(Y_{i,(j)}, X_{i,(N_i - n_i + j)})$ for $j = 1, \ldots, n_i$.

The key feature of Full House Modeling is the explicit connection between the elements

forming the $(Y_{i,(j)}, X_{i,(r_{i,j})})$ pairs. Full House Modeling states this connection as

$$\begin{aligned} Y_{i,(j)} &= F^{-1}_{Y_{i,(j)}} \circ F_{X_{i,(r_{i,j})}} \left(X_{i,(r_{i,j})} \right), \\ X_{i,(r_{i,j})} &= F^{-1}_{X_{i,(r_{i,j})}} \circ F_{Y_{i,(j)}} \left(Y_{i,(j)} \right), \end{aligned} \tag{2}$$

where $1_i(X_{i,(r_{i,j})}, \mathbf{X}_{i,-(r_{i,j})}) = 1$, and both F_{Y_i} and F_X are continuous and invertible. We summarise Full House Modeling in the definition below.

Definition 2.1 (Full House Model)**.** *Consider populations with sizes N_i and system sizes n_i, $N_i \geq n_i$. Let a system inclusion function $1_i(\cdot, \cdot)$ be specified. Let $X_{i,j} \overset{iid}{\sim} F_X$, $j = 1, \ldots, N_i$. Let $Y_{i,j} \overset{iid}{\sim} F_{Y_i}$, $j = 1, \ldots, n_i$. Form pairs $(Y_{i,(j)}, X_{i,(r_{i,j})})$, $j = 1, \ldots, n_i$ for all individuals such that $1_i(X_{i,(r_{i,j})}, \boldsymbol{X}_{i,-(r_{i,j})}) = 1$. Then the Full House Model generates $Y_{i,(j)}$ values following (2) where F_{Y_i} and F_X are continuous and invertible.*

Remark The Full House Model induces structural missingness by construction: for any individual j such that $1_i(X_{i,j}, \mathbf{X}_{i,-j}) = 0$, the corresponding observed statistic $Y_{i,j}$ is undefined. In subsequent sections, we consider probabilities involving both latent and observed quantities. To formalize this setting, we treat unobserved $Y_{i,j}$ values as not measurable with respect to the observable probability space, reflecting the fact that they lie outside the observable σ-field generated by the system. This perspective allows selection mechanisms to be incorporated into probabilistic statements without imputing counterfactual outcomes.

Definition 2.1 assumes a common latent aptitude distribution F_X across contexts, while allowing the observed achievement distributions F_{Y_i} to vary by system. This reflects a modeling choice in which the process generating latent aptitude is taken to be context-neutral, whereas the expression and measurement of aptitude may differ across contexts.

The probability integral transformations in (2) involve the distribution functions of order statistics, namely $F_{X_{i,(r_{i,j})}}$ and $F_{Y_{i,(j)}}$, which may be cumbersome to evaluate directly. An equivalent and more convenient representation follows from the fact that, under continuity, the probability integral transform of an order statistic can be expressed through a Beta

distribution. In particular,

$$F_{X_{i,(r_{i,j})}}\big(X_{i,(r_{i,j})}\big) = F_{U_{i,(r_{i,j})}} \circ F_X\big(X_{i,(r_{i,j})}\big)\,, \quad U_{i,(r_{i,j})} \sim \mathrm{Beta}(r_{i,j},\, N_i - r_{i,j} + 1)\,, \tag{3}$$

and similarly,

$$F_{Y_{i,(j)}}\big(Y_{i,(j)}\big) = F_{U_{i,(j)}} \circ F_{Y_i}\big(Y_{i,(j)}\big)\,, \quad U_{i,(j)} \sim \mathrm{Beta}(j,\, n_i - j + 1)\,. \tag{4}$$

This allows (2) to be computed via F_X, F_{Y_i}, and known Beta distributions, bypassing explicit order-statistic calculations. The representations (3) and (4) yield an additional advantage: although F_X may be difficult to specify, latent aptitude can be recovered up to monotone transformation without explicit knowledge of F_X.

Combining (3) and (4) yields

$$X_{i,(r_{i,j})} = F_X^{-1}\Big(F_{U_{i,(r_{i,j})}} \circ F_{U_{i,(j)}} \circ F_{Y_i}\big(Y_{i,(j)}\big)\Big)\,.$$

Define $W_{i,j} = F_{U_{i,(r_{i,j})}} \circ F_{U_{i,(j)}} \circ F_{Y_i}(Y_{i,(j)})$. Monotonicity of F_X implies

$$W_{i_1,j_1} \geq W_{i_2,j_2} \implies F_X^{-1}(W_{i_1,j_1}) \geq F_X^{-1}(W_{i_2,j_2}),$$

so the ordering of latent aptitudes is invariant to F_X. Thus F_X is a structural nuisance for comparative analysis, and FHM can be implemented in practice via observed-to-rank mappings without necessitating a true latent distribution.

3 Paradoxes Under Identical Achievement Distributions

In this section, we derive explicit probabilities of the aptitude–achievement paradox, as defined in (1), within a Full House Modeling framework involving two populations. We focus on the setting in which the observed achievement distributions are identical across contexts,

that is, $F_{Y_i} = F_Y$ for $i = 1, 2$ in Definition 2.1. The following definition formalizes the notion of paradox used throughout the remainder of the article.

Definition 3.1. *Consider the Full House Model in Definition 2.1 with two populations denoted by $i = 1, 2$. We say a aptitude-achievement paradox has occurred when both $X_{1,(r_{1,j})} > X_{2,(r_{2,k})}$ and $Y_{1,(j)} < Y_{2,(k)}$ have occurred. The aptitude-achievement paradox probability is defined as*

$$\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right). \tag{5}$$

As a simple illustrative example of the above, let $r_{i,t} = N_i - n_i + t$ and set $j = n_1$ and $k = n_2$, and further suppose that $Y_{1,j} < Y_{2,k}$ a.s. Then the aptitude-achievement paradox probability (5) is equivalent to $\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}\right)$, and we have $\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}\right) \to 1$ when $N_1 \to \infty$. We now introduce a convenient representation which allows the paradox probability to be expressed in terms of the latent aptitudes alone.

Lemma 3.2. *Consider the Full House Model in Definition 2.1 with two populations, where $F_{Y_i} = F_Y$ for $i = 1, 2$. Then,*

$$\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right) = \mathrm{P}\left(X_{2,(r_{2,k})} < X_{1,(r_{1,j})} < h_{j,k}\left(X_{2,(r_{2,k})}\right)\right), \tag{6}$$

$$\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right) = \mathrm{P}\left(g_{j,k}\left(Y_{2,(k)}\right) < Y_{1,(j)} < Y_{2,(k)}\right) \tag{7}$$

where $h_{j,k}(\cdot) = F^{-1}_{X_{1,(r_{1,j})}} \circ F_{Y_{1,(j)}} \circ F^{-1}_{Y_{2,(k)}} \circ F_{X_{2,(r_{2,k})}}(\cdot), g_{j,k}(\cdot) = F^{-1}_{Y_{1,(j)}} \circ F_{X_{1,(r_{1,j})}} \circ F^{-1}_{X_{2,(r_{2,k})}} \circ F_{Y_{2,(k)}}(\cdot)$.

Our first formal result provides conditions that allow for an explicit aptitude-achievement paradox probability result. Those conditions require the following definition of a regularized incomplete beta function, the cumulative distribution function of a Beta(α, β) distribution.

Definition 3.3. *Let $\alpha, \beta > 0$. Then the regularized incomplete beta function is defined as*

$$I_{\alpha,\beta}(x) = \frac{1}{B(\alpha,\beta)} \int_0^x u^{\alpha-1}(1-u)^{\beta-1} du,$$

where $B(\alpha,\beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$ is the beta coefficient.

Theorem 3.4. *Consider the Full House Model in Definition 2.1 with two populations, where $F_{Y_i} = F_Y$ for $i = 1, 2$. Denote by I the regularized incomplete beta function. Further assume that*

$$I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1}(t) - I_{r_{1,j},N_1-r_{1,j}+1} \circ I^{-1}_{r_{2,k},N_2-r_{2,k}+1}(t) \geq 0, \qquad \text{for all } t \in (0,1). \tag{8}$$

Under these assumptions,

$$\begin{aligned}
&\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right) \\
&= \sum_{p=j}^{n_1} \binom{n_1}{p} \frac{B(k+p, n_2-n_1-p+k+1)}{B(k, n_2-k+1)} - \sum_{p=r_{1,j}}^{N_1} \binom{N_1}{p} \frac{B(r_{2,k}+p, N_2-N_1-p+r_{2,k}+1)}{B(r_{2,k}, N_2-r_{2,k}+1)}.
\end{aligned}$$

Remarks:

1. The paradox probability in Theorem 3.4 is distribution free. Once the Full House Model is specified, the result depends only on the population sizes N_i, system sizes n_i, the system inclusion function, and the indices of the order statistics under comparison. As established in Section 2, the Beta representations (3)–(4) imply that the distributions of observed order statistics can be expressed independently of the underlying form of F_Y. Lemma 3.2 leverages this structure to eliminate F_Y entirely from the paradox probability calculation,

$$\begin{aligned}
h_{j,k}(x) &= F^{-1}_{X_{1,(r_{1,j})}} \circ F_{Y_{1,(j)}} \circ F^{-1}_{Y_{2,(k)}} \circ F_{X_{2,(r_{2,k})}}(x) \\
&= F^{-1}_{X_{1,(r_{1,j})}} \circ I_{j,n_1-j+1} \circ \left(F_Y \circ F_Y^{-1}\right) \circ I^{-1}_{k,n_2-k+1} \circ F_{X_{2,(r_{2,k})}}(x)
\end{aligned}$$

$$= F^{-1}_{X_{1,(r_{1,j})}} \circ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1} \circ F_{X_{2,(r_{2,k})}}(x).$$

2. Likewise, the latent aptitude distribution F_X plays no role in the final expression for the paradox probability. Although Lemma 3.2 expresses the event in terms of latent order statistics, $\mathrm{P}\left(X_{2,(r_{2,k})} < X_{1,(r_{1,j})} < h_{j,k}\left(X_{2,(r_{2,k})}\right)\right)$. the monotone coupling induced by Full House Modeling ensures that all integrals involving F_X reduce to functions of Beta distributions and order indices. As shown in the proof of Theorem 3.4, a change of variables removes dependence on F_X, rendering it a structural nuisance parameter for paradox probabilities.

3. Combining these observations, the paradox probability admits a closed-form expression involving only regularized incomplete beta functions. In particular, the probability simplifies to $\int_0^1 \left\{ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1}(t) - I_{r_{1,j},N_1-r_{1,j}+1} \circ I^{-1}_{r_{2,k},N_2-r_{2,k}+1}(t) \right\}^+ dt$. The technical condition (8) ensures that the integrand is nonnegative for all $t \in (0,1)$, guaranteeing that this expression defines a valid probability. In subsequent sections, we characterize when this condition holds and examine its empirical implications.

Often it is of interest to compare the individuals with the highest aptitudes in their respective populations. Below we state the aptitude-achievement paradox probability in such setting using a Full House Model with two populations with the specification that $F_{Y_i} = F_Y$ for $i = 1, 2$.

Corollary 3.4.1. *Suppose that the conditions of Theorem 3.4 hold. Suppose that* $1_j(X_{j,(N_j)}, \boldsymbol{X}_{j,-(N_j)}) = 1$, $j = 1, 2$. *Then,*

$$\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \begin{cases} \frac{n_2}{n_1+n_2} - \frac{N_2}{N_1+N_2}, & \text{if } \frac{n_1}{N_1} \leq \frac{n_2}{N_2} \\ 0, & \text{otherwise} \end{cases}.$$

Intuitively, this is the difference between two 'lottery' probabilities: the probability that individual 2 wins the observed achievement comparison (first term) minus the probability

that individual 2 wins the latent aptitude comparison (second term). The paradox probability is positive precisely when the sampling ratio favors individual 2 more than the population ratio does. A similar result holds for individuals with the lowest latent aptitudes in the population.

Corollary 3.4.2. *Suppose that the conditions of Theorem 3.4 hold. Suppose that* $1_j(X_{j,(N_j)}, \boldsymbol{X}_{j,-(N_j)}) = 1$, $j = 1, 2$. *Then,*

$$\mathrm{P}\left(X_{1,(1)} > X_{2,(1)}, Y_{1,(1)} < Y_{2,(1)}\right) = \begin{cases} \frac{N_2}{N_1+N_2} - \frac{n_2}{n_1+n_2}, & \text{if } \frac{n_1}{N_1} \geq \frac{n_2}{N_2} \\ 0, & \text{otherwise} \end{cases}.$$

We now demonstrate a setting in which technical condition (8) holds. This setting is based on a configuration of n_j, N_j, $j = 1, 2$ under the assumption that the systems are populated with individuals from the source populations who have the highest aptitudes.

Proposition 3.5. *Consider the Full House Model in Definition 2.1 with two populations, where* $F_{Y_i} = F_Y$ *for* $i = 1, 2$, *and let the system inclusion function be* $1_i(X_{i,j} \geq X_{i,(N_i - n_i + 1)})$. *Denote by* I *the regularized incomplete beta function. Suppose that* $\frac{n_1}{n_2} \leq 1$ *and* $\frac{N_2}{N_1} \leq 1$, *and set* $k = j$. *Under these assumptions, we conclude that technical condition* (8) *holds and the conclusion of Theorem 3.4 applies.*

4 Paradox Under Different Distributions

In this section we will provide aptitude-achievement paradox probability results in a Full House Modeling setting involving two populations when we relax the specification that $F_{Y_1} = F_{Y_2}$ in Definition 2.1. We first provide bounds for aptitude-achievement paradox probability in general settings. We then provide explicit aptitude-achievement paradox probability results when Y_1 and Y_2 are normally distributed with different means and variances.

4.1 General bounds

The general bounds we derive require a relaxed form of first-order stochastic dominance under which the distributions of Y_1 and Y_2 are 'close' to each other. More formally, we introduce first-order ε-weak stochastic dominance, which is necessary for our aptitude-achievement paradox probability results.

Definition 4.1. *We will say that Y_2 is first order ε-weakly stochastically dominant over Y_1 if, for some $\varepsilon > 0$, $F_{Y_2}(y) \leq F_{Y_1}(y) \leq F_{Y_2}(y) + \varepsilon$ for all y, and there exists some y for which $F_{Y_2}(y) < F_{Y_1}(y)$.*

Under first-order ε-weak stochastic dominance, we derive a general lower and upper bound on the aptitude-achievement paradox probability when $F_{Y_1} \neq F_{Y_2}$.

Proposition 4.2. *Consider the Full House Model in Definition 2.1 with two populations $i = 1, 2$. Suppose that the system inclusion mechanism has the property that $1_i\left(X_{i,(N_i)}, \boldsymbol{X}_{i,-(N_i)}\right) = 1$. Further suppose that Y_2 is first order ε-weakly stochastically dominant over Y_1 for some $\varepsilon > 0$ and $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$. Then,*

$$\begin{aligned} \frac{n_2}{n_1 + n_2} - \frac{N_2}{N_1 + N_2} &\leq \mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\ &\leq n_2 \sum_{i=0}^{n_1} \frac{\binom{n_1}{i}\varepsilon^i}{n_1 + n_2 - i} - \frac{N_2}{N_1 + N_2} \end{aligned} \tag{9}$$

Proposition 4.2 provides bounds on the paradox probability under general achievement distributions. For example, when $Y_i \sim N(\mu_i, \sigma^2)$, $i = 1, 2$, with $\mu_1 < \mu_2$ and common variance, ε can be bounded explicitly as a function of $(\mu_2 - \mu_1)/\sigma$. Beyond such bounds, we can obtain explicit aptitude-achievement paradox probabilities when $Y_i \sim N(\mu_i, \sigma_i^2)$, $i = 1, 2$, allowing variances to differ across contexts.

4.2 Normal distributions

This section develops explicit aptitude-achievement paradox probabilities (5) when a Full House Model in Definition 2.1 specifies $Y_i \sim N(\mu_i, \sigma_i^2)$, $i = 1, 2$. We focus on systems where $1_i(X_{i,(N_i)}, \mathbf{X}_{i,-(N_i)}) = 1$ for $i = 1, 2$. The outline is similar to Section 3: a technical condition is needed to prove the main result, and further results provide configurations satisfying this condition. However, with $Y_i \sim N(\mu_i, \sigma_i^2)$, the technical condition yields a remainder term that prevents an exact closed-form expression. We show that this remainder is negligible under mild conditions on population fractions and distributional parameters.

Similar to the proof of Theorem 3.4, Lemma 3.2 will provide a representation more conducive to formal integration needed to prove our main result. Only this time we will use (7) in Lemma 3.2 to represent the aptitude-achievement paradox probability as

$$\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \mathrm{P}\left(g_{n_1,n_2}\left(Y_{2,(n_2)}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right),$$

where, from the definition of g in Lemma 3.2 applied to our current setting, we have

$$g_{n_1,n_2}(x) = F_{Y_1}^{-1}\left(F_{Y_2}(x)^{\frac{n_2 N_1}{N_2 n_1}}\right) = \Phi^{-1}\left(\Phi\left(\frac{x-\mu_2}{\sigma_2}\right)^{\frac{n_2 N_1}{N_2 n_1}}\right)\sigma_1 + \mu_1, \tag{10}$$

and Φ defines the cumulative distribution function of the standard normal distribution. From here, we have that

$$\begin{aligned}\mathrm{P}\left(g_{n_1,n_2}\left(Y_{2,(n_2)}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right) &= \mathrm{P}\left(F_{Y_1}^{-1}\left(F_{Y_2}(Y_{2,(n_2)})^{\frac{n_2 N_1}{N_2 n_1}}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\ &= \int_A \int_{F_{Y_1}^{-1}\left(F_{Y_2}(x)^{\frac{n_2 N_1}{N_2 n_1}}\right)}^{x} dF_{Y_1,(n_1)}(y) dF_{Y_2,(n_2)}(x),\end{aligned}$$

where $A := A(\mu_1, \mu_2, \sigma_1, \sigma_2, n_1, n_2, N_1, N_2)$ is the range of integration so that the inner inte-

gral above is properly specified. More formally,

$$
\begin{aligned}
A &= \left\{\omega : F_{Y_1}^{-1}\left(F_{Y_2}(Y_{2,(n_2)}(\omega))^{\frac{n_2N_1}{N_2n_1}}\right) \leq Y_{2,(n_2)}(\omega)\right\} \\
&= \left\{\omega : \Phi\left(\frac{Y_{2,(n_2)}(\omega)-\mu_2}{\sigma_2}\right)^{\frac{n_2}{N_2}} \leq \Phi\left(\frac{Y_{2,(n_2)}(\omega)-\mu_1}{\sigma_1}\right)^{\frac{n_1}{N_1}}\right\},
\end{aligned} \tag{11}
$$

where the second equality in (11) follows from (10). Continuing with the probability calculation above

$$
\begin{aligned}
&\int_A \int_{F_{Y_1}^{-1}\left(F_{Y_2}(x)^{\frac{n_2N_1}{N_2n_1}}\right)}^{x} dF_{Y_1,(n_1)}(y)dF_{Y_2,(n_2)}(x) \\
&= \int_A F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}} dF_{Y_2,(n_2)}(x) \\
&= \int_{-\infty}^{\infty} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}} dF_{Y_2,(n_2)}(x) - \int_{A^c} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}} dF_{Y_2,(n_2)}(x).
\end{aligned} \tag{12}
$$

The left-hand side of (12) can (and will) be further evaluated and leads to the desired explicit aptitude-achievement paradox probability. The right-hand side of (12) represents a remainder term $R := R(\mu_1, \mu_2, \sigma_1, \sigma_2, n_1, n_2, N_1, N_2)$,

$$
R = -\int_{A^c} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}} dF_{Y_2,(n_2)}(x), \tag{13}
$$

where the set A in (13) is defined in (11). The remainder R is negligible when n_2 is moderately large, as we now show. From (11) and (13), we have

$$
\begin{aligned}
|R| &= \left|\int_{A^c} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}} dF_{Y_2,(n_2)}(x)\right| \\
&\leq \int_{A^c} \left|F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}}\right| dF_{Y_2,(n_2)}(x) \\
&\leq \mathrm{P}\left(\Phi\left(\frac{Y_{2,(n_2)}-\mu_2}{\sigma_2}\right)^{\frac{n_2}{N_2}} \geq \Phi\left(\frac{Y_{2,(n_2)}-\mu_1}{\sigma_1}\right)^{\frac{n_1}{N_1}}\right).
\end{aligned}
$$

Now, when we specify that $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$ as in Proposition 4.2, we have

$$\begin{aligned} \mathrm{P}\left(\Phi\left(\frac{Y_{2,(n_2)}-\mu_2}{\sigma_2}\right)^{\frac{n_2}{N_2}} \geq \Phi\left(\frac{Y_{2,(n_2)}-\mu_1}{\sigma_1}\right)^{\frac{n_1}{N_1}}\right) &\leq \mathrm{P}\left(\Phi\left(\frac{Y_{2,(n_2)}-\mu_2}{\sigma_2}\right) \geq \Phi\left(\frac{Y_{2,(n_2)}-\mu_1}{\sigma_1}\right)\right) \\ = \mathrm{P}\left(\frac{Y_{2,(n_2)}-\mu_2}{\sigma_2} \geq \frac{Y_{2,(n_2)}-\mu_1}{\sigma_1}\right). \end{aligned}$$

Under the assumption that $\sigma_1 \leq \sigma_2$, straightforward algebra yields

$$\mathrm{P}\left(\frac{Y_{2,(n_2)}-\mu_2}{\sigma_2} \geq \frac{Y_{2,(n_2)}-\mu_1}{\sigma_1}\right) = \mathrm{P}\left(Y_{2,(n_2)} \leq \left(\frac{\mu_2}{\sigma_2}-\frac{\mu_1}{\sigma_1}\right)\left(\frac{1}{\sigma_2}-\frac{1}{\sigma_1}\right)^{-1}\right).$$

Under the additional assumption that $\mu_1 \leq \mu_2$, the right-hand side is bounded above by $\mathrm{P}\left(Y_{2,(n_2)} \leq \mu_2\right)$. Putting it all together, we have $|R| \leq \mathrm{P}\left(Y_{2,(n_2)} \leq \mu_2\right) \leq \left(\frac{1}{2}\right)^{n_2}$, when $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$, $\sigma_1 \leq \sigma_2$, and $\mu_1 \leq \mu_2$. Therefore, although the remainder term in (13) arises from the restriction of the integration domain to A, it is exponentially small in n_2 under a familiar ordering of population fractions (see Proposition 4.2) and a simple parametric configuration. Consequently, the aptitude–achievement paradox probability is well approximated by the unrestricted integral given by the left-hand side of (12)with R contributing only a negligible correction for moderately large n_2.

We now introduce a multivariate normal representation that allows this principal term to be expressed in closed form. Let $\mathbf{W} \sim \mathcal{N}_{(n_1+n_2-1)}(\mathbf{0}, \mathbf{\Sigma})$ and covariance matrix

$$\mathbf{\Sigma} = \begin{bmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{bmatrix}, \tag{14}$$

defined with the following structure

$$\Sigma_{11} = \begin{bmatrix} \frac{\sigma_2^2}{\sigma_1^2}+1 & \frac{\sigma_2^2}{\sigma_1^2} & \dots & \frac{\sigma_2^2}{\sigma_1^2} \\ \frac{\sigma_2^2}{\sigma_1^2} & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \frac{\sigma_2^2}{\sigma_1^2} \\ \frac{\sigma_2^2}{\sigma_1^2} & \dots & \frac{\sigma_2^2}{\sigma_1^2} & \frac{\sigma_2^2}{\sigma_1^2}+1 \end{bmatrix}_{n_1 \times n_1} \quad \Sigma_{12}^T = \Sigma_{21} = \begin{bmatrix} \frac{\sigma_2}{\sigma_1} & \dots & \dots & \frac{\sigma_2}{\sigma_1} \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ \frac{\sigma_2}{\sigma_1} & \dots & \dots & \frac{\sigma_2}{\sigma_1} \end{bmatrix}_{(n_2-1)\times n_1}$$

$$\Sigma_{22} = \begin{bmatrix} 2 & 1 & \dots & 1 \\ 1 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & 1 \\ 1 & \dots & 1 & 2 \end{bmatrix}_{(n_2-1)\times(n_2-1)} . \tag{15}$$

With this structure we now provide an explicit aptitude-achievement paradox probability, the main result of this section.

Theorem 4.3. *Assume a Full House Model in Definition 2.1 and additionally specify that* $Y_i \sim N(\mu_i, \sigma_i^2)$*, and suppose that the system inclusion mechanism has the property that* $1_i(X_{i,(N_i)}, \boldsymbol{X}_{i,-(N_i)}) = 1$ *for* $i = 1, 2$*. Let* $\mathbf{W} \sim \mathcal{N}_{(n_1+n_2-1)}(\mathbf{0}, \boldsymbol{\Sigma})$ *where* $\boldsymbol{\Sigma}$ *is as defined in (14) and (15). Then,*

$$\begin{aligned} &\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\ &\quad = n_2 F_{\mathbf{W}}(\mathbf{w} = (\frac{\mu_2-\mu_1}{\sigma_1}, \dots, \frac{\mu_2-\mu_1}{\sigma_1}, 0, \dots, 0)) - \frac{N_2}{N_1+N_2} + R, \end{aligned} \tag{16}$$

where R *is defined in (13),* $F_{\mathbf{W}}$ *is the distribution function of* $\mathbf{W}$*, and the first* n_1 *elements of* $\mathbf{w}$ *are* $\frac{\mu_2-\mu_1}{\sigma_1}$ *and the last* $n_2 - 1$ *elements are 0.*

Moreover, when $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$*,* $\sigma_1 \leq \sigma_2$*, and* $\mu_1 \leq \mu_2$*, we have that*

$$|R| \leq \left(\frac{1}{2}\right)^{n_2}. \tag{17}$$

We now show that plug-in estimators of the normal distribution parameters μ_i and σ_i,

$i = 1, 2$, yield reliable estimates of aptitude-achievement paradox probabilities (16). Our approach requires standard properties of equicorrelated covariance matrices. For matrices of the form $\Sigma = \rho \mathbf{1}\mathbf{1}^T + I_n$ (where $\mathbf{1}$ is the vector of ones), we have $\det(\Sigma) = \rho n + 1$ and the inverse follows from the Woodbury identity [Golub and Van Loan, 2013]. The explicit form of Σ^{-1} is given in the proof of Theorem 4.5 (Supplementary Materials).Before we state our result, we introduce the Lemma 4.4

Lemma 4.4. *Let* $\mathbf{W} \sim \mathcal{N}_{n_1+n_2-1}(\mathbf{0}, \mathbf{\Sigma})$ *be a multivariate normal random vector with mean vector* $\mathbf{0}$ *and covariance matrix* $\mathbf{\Sigma}$ *as defined in* (14) *and* (15). *Define* $c := c_{n_1+n_2-1}$ *as the maximal conditional one-dimensional Gaussian probability arising in the sequential factorization of the joint CDF, that is,*

$$
\begin{aligned}
c = \max \Big\{ & \mathrm{P}(W_{n_1+n_2-1} \le w_{n_1+n_2-1} \,|\, W_1 = w_1, \ldots, W_{n_1+n_2-2} = w_{n_1+n_2-2}), \\
& \ldots, \mathrm{P}(W_2 \le w_2 \mid W_1 = w_1), \mathrm{P}(W_1 \le w_1) \Big\}.
\end{aligned} \tag{18}
$$

Then $c \in (0, 1)$ *for all* n_1, n_2, *and* $\mathbf{F}_{\mathbf{W}}(\mathbf{w}) \le c^{n_1+n_2-1}$, *where* $\mathbf{F}_{\mathbf{W}}(\mathbf{w})$ *denotes the cumulative distribution function of a mean-zero multivariate normal random vector with covariance matrix* $\mathbf{\Sigma}$, *evaluated at* $\mathbf{w} \in \mathbb{R}^{n_1+n_2-1}$.

Theorem 4.5. *Assume a Full House Model in Definition 2.1 and additionally specify that* $Y_i \sim N(\mu_i, \sigma_i^2)$, *and suppose that the system inclusion mechanism has the property that* $1_i(X_{i,(N_i)}, \boldsymbol{X}_{i,-(N_i)}) = 1$ *for* $i = 1, 2$. *Let* $\mathbf{W} \sim \mathcal{N}_{(n_1+n_2-1)}(\mathbf{0}, \mathbf{\Sigma})$ *where* $\mathbf{\Sigma}$ *is as defined in* (14) *and* (15). *Suppose that* $\hat{\mu}_1$, $\hat{\mu}_2$, $\hat{\sigma}_1$, *and* $\hat{\sigma}_2$ *are, respectively, root sample-size consistent estimators of* μ_1, μ_2, σ_1, *and* σ_2. *Then, with* c *defined as in the Lemma 4.4, we have*

$$
\begin{aligned}
& \mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)} \mid \hat{\mu_1}, \hat{\mu_2}, \hat{\sigma_1}, \hat{\sigma_2}\right) \\
& \quad - \mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)} \mid \mu_1, \mu_2, \sigma_1, \sigma_2\right) \\
& = O_P(n_1^{\frac{1}{2}} n_2 c^{n_1+n_2-3} + n_1 n_2^{\frac{1}{2}} c^{n_1+n_2-3}).
\end{aligned}
$$




Figure 1: Sensitivity Analysis for Paradox Probability

5 Examples

5.1 Sensitivity Analysis for Theorem 3.4

Under the setting of Theorem 3.4, we will specify $X_1 \overset{d}{=} X_2 \overset{d}{=} X$, $Y_1 \overset{d}{=} Y_2 \overset{d}{=} Y$, where $X \sim \text{Pareto}(\alpha)$ and $Y \sim \mathcal{N}(0,1)$. Without loss of generality, we set $j = n_1 - 1$ and $k = n_2 - 1$, corresponding to the selection of the second-best performer in the aptitude-achievement paradox probability, as defined in Definition 3.1.

In each simulation, we randomly sample n_1 and n_2 observations from the distribution of Y, and compute their latent aptitudes using Equation (2). Based on these values, we directly calculate paradox probabilities according to Definition 3.1. We repeat this simulation 10000 times and report the average.

Figure 1 presents the results of the sensitivity analysis for the aptitude-achievement

paradox probability described in Theorem 3.4. In Figure 1 (A), we fix $n_1 = 40$ and $n_2 = 60$, and vary N_1 and N_2 from 200 to 2000. In Figure 1 (B), we fix $N_1 = 10^3$ and $N_2 = 10^3$, and vary n_1 and n_2 from 20 to 200. In Figure 1 (C), we fix $n_1 = 40$ and $N_2 = 10^3$, and vary n_2 from 20 to 200 and N_1 from 200 to 2000. In Figure 1 (D), we fix $N_1 = 10^3$ and $n_2 = 60$, and vary N_2 from 200 to 2000 and n_1 from 20 to 200.

As shown in Figure 1 (A), a small N_2 and large N_1 increase the paradox probability by making $\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}\right) \to 1$. Similarly, in Figure 1 (B), a small n_1 combined with a large n_2 leads to a higher aptitude-achievement paradox probability since it drives $\mathrm{P}\left(Y_{1,(j)} < Y_{2,(k)}\right) \to 1$.

5.2 Ratio of N_2 and N_1

Here we investigate the effect of the ratio between N_2 and N_1 on aptitude-achievement paradox probabilities under the specifications of Corollary 3.4.1. Let $c = \frac{N_2}{N_1}$, and vary c from 0.1 to 1 in increments of 0.1. In each simulation, we fix $n_1 = 60$ and $n_2 = 40$, and vary N_1 from 100 to 2000. We repeat this simulation 10000 times and report the average paradox probability.

Figure 2 presents the results. As shown in the figure, the paradox probability decreases as c increases, which is reasonable since $\mathbb{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}\right)$ decreases as N_2 increases. Importantly, Figure 2 also demonstrates that the paradox probability remains constant when the ratio between N_2 and N_1 is fixed. This implies that it is unnecessary to focus on the exact values of N_1 and N_2 when computing the paradox probability; rather, the ratio N_2/N_1 alone determines its value.

5.3 Recursive Full House Models

In this section we will investigate a simple extension to Full House Modeling in which Full House Models are repeatedly applied to individuals as they progress through iterations of some generic process. The general setup is that, in the first round, latent aptitude $X_j \overset{iid}{\sim}$




Figure 2: Effect of the ratio between N_2 and N_1 on aptitude-achievement paradox probabilities under the specifications of Corollary 3.4.1.

F_X, $j = 1, \ldots, N$ will be generated and then a label will be randomly assigned to each individual. There will be m labels and each label will designate the outcome distribution. $F_{Y_{i,j}}$, $i = 1, \ldots, n$, $j = 1, \ldots, n_i$. Then outcomes will be realized as

$$Y_{i,(r_{i,(j)})} = F^{-1}_{Y_{i,j}} \circ I^{-1}_{r_{i,(j)}, n_i - r_{i,(j)}+1} \circ I_{j,N-j+1} \circ F_X(X_{(j)}), \tag{19}$$

where $r_{i,(j)}$ is the rank of $X_{(j)}$ among the n_i latent aptitudes also assigned label i.

In the next iteration, all observations will be combined and then sorted in descending order by $Y_{i,(j)}$. The labels will be assigned in descending order, the largest n_m values of $Y_{i,(j)}$ are assigned label m, the next largest n_{m-1} values of $Y_{i,(j)}$ are assigned label $m-1$, and so on. We will setup these simulations so that Y_i stochastically dominates Y_j whenever $i > j$ so that higher outcomes are associated with higher labels. Then new Y values are generated

according to (19). The process will repeat until a pre-assigned stopping rule is satisfied. Note that this simulation setup assumes an immutable latent aptitude.

5.3.1 Stop when highest latent aptitude is among the highest-performing group

Here we consider a recursive Full House Modeling setup described in Section 5.3 in which the recursion stops at the first iteration when $X_{(N)}$ is assigned label m. Here we specify that $N = 100,000$, $m = 200$, $n_i = 500$, $i = 1, \ldots, m$. We will assume that $X_j \overset{iid}{\sim}$ Pareto(1.16), $j = 1, \ldots, N$. And we will assume that $Y_{i,j} \overset{iid}{\sim} N(\mu_i, \sigma_i)$, $j = 1, \ldots, n_i$ and $i = 1, \ldots, m$. We will specify that the μ_is as a set of 200 equally spaced values between 60 and 80 with μ_i increasing in i, ie $\mu_i = 60 - 20/199 + 20i/199$, $i = 1, \ldots, m$. We will fix $\sigma_i = \sigma$ and consider $\sigma \in \{0.5, 1, 2\}$. These simulations will be repeated 500 times.

To illustrate the procedure, consider a simplified scenario in which all $X_{(j)}$ values are realized at their median value. With this simplification, $Y_{i,(r_{i,(j)})}$ in (19) is approximately $F^{-1}_{Y_{i,j}}((r_{i,(j)} - 1/3)/(n_i + 1/3))$ [Kerman, 2011]. And we have that $Y_{i,(n_i)} - Y_{i+1,(n_{i+1}-1)} > \mu_{i+1} - \mu_i$ for all considered values of σ. The individual with latent ability $X_{(N)}$ will consistently attain the value $Y_{i,(n_i)}$, regardless of their assigned group. Consequently, this individual's position among the m groups will improve as the simulation advances.

Results can be seen in Figure 3. These Monte Carlo simulations reveal that it typically takes several rounds before the individual with the top aptitude belongs to the group that allows for the highest performances. On average, with σ set to 0.5, 1, and 2, it takes 2.61, 4.19, and 7.46 rounds, respectively, for the individual with the top aptitude to join the highest-performing group (respective standard errors: 0.05, 0.09, 0.18). Deriving the distribution of this hitting time analytically under explicit FHM assumptions, and characterizing how it depends on σ and grouping structure, remains an open problem.

While the setting is stylized and assumes fixed latent aptitude, it illustrates how structural features of evaluative systems, not just randomness, can delay recognition of true potential. Whereas the paradox probability quantifies cross-sectional reversal risk, recursive Full House




Figure 3: The number of rounds that it takes for $X_{(N)}$ to be assigned label m in a recursive Full House Modeling framework in Section 5.3 with specifications in Section 5.3.1.

Modeling captures temporal persistence: the hitting time measures how many rounds elapse before latent aptitude and observed achievement align. These findings resonate with a growing empirical literature on bias and distortion in algorithmic and educational evaluation systems, which we discuss further in the Discussion.

5.4 Batting Averages in Major League Baseball

We now apply Theorem 4.3 and the reliability guarantees in Theorem 4.5 to evaluate the contextual extremeness of observed outcomes in a real-world system where the underlying distribution of aptitude is not directly observable and evolves over time. Major League Baseball (MLB) offers a canonical example of such a system. Here, player skill (latent batting aptitude) is inferred only indirectly through season-level observed outcomes (batting averages), which are shaped by changing population structures and environmental variance. In particular, MLB introduced a series of rule changes in 2023 intended to increase in-game action, including a restriction on defensive positioning designed to raise batting averages




Figure 4: Seasonal means and standard deviations of batting averages for full-time players from 1871-2023. Smoothing is applied to both trends.

on balls in play (BABIP) [Castrovince, 2023]. Figure 4 presents smoothed estimates of the mean and standard deviation of batting averages for full-time players from 1871 to 2023.

The 2023 season saw an increase in mean batting average for full-time players of 0.0057 (corresponding to an additional hit about every four baseball games) with a reduction in variability. A two-sample test for differences in batting averages among full-time players from the 2023 and 2022 seasons has a p-value of 0.039. Thus, rule changes seemed to be effective in yielding increased batting averages[1] Moreover, the 2023 MLB batting leader Luis Arraez had a 0.354 batting average which was the first batting average to exceed .350 (a mark of excellence) since Josh Hamilton in 2010.

We now apply Theorem 4.3 to quantify paradox probabilities for extreme batting aver-

[1]Technically, Castrovince [2023] referred to MLB's desire to increase batting averages on balls in play (BABIP) while this analysis focused on batting averages. BABIP increased by 0.0061 from 2022 to 2023.




Figure 5: Comparison of paradox probabilities and empirical exceedance rates relative to the 2023 season, based on data from 1871–2022. The blue line shows estimated paradox probabilities via plug-in using Theorem 4.3. The black dashed line displays a smoothed empirical exceedance rate for Arraez's 2023 batting average.

ages, treating the 2023 season as system 1 and each season from 1871 to 2022 as system 2. Seasonal batting averages of full-time players are assumed normally distributed, with means and standard deviations estimated via plug-in; the reliability guarantees of Theorem 4.5 apply. Figure 4 shows that both the average and variability of batting averages in 2023 are among the lowest observed, indicating a historically suppressed environment. Moreover, with our specifications we have that $\hat{\sigma}_1 \leq \hat{\sigma}_2$, $\hat{\mu}_1 \leq \hat{\mu}_2$, and $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$, so the remainder term (13) in Theorem 4.3 is negligible. The talent pool estimates N_i are taken from Yan et al. [2025], which includes sensitivity analyses showing that qualitative conclusions are robust to reasonable perturbations of these estimates.

Figure 5 displays the resulting paradox probabilities alongside a smoothed empirical exceedance rate, the proportion of historical seasons in which the maximum batting average exceeded Arraez's 2023 mark. The estimated paradox probabilities closely track this bench-

mark. Both quantities increase as one moves backward in time, reflecting that earlier eras were structurally more conducive to extreme observed batting averages due to higher means, larger variances, and smaller effective populations.

Under Full House Modeling, Arraez's 2023 performance ranks 68th all-time among full-time batters after accounting for population depth and contextual effects [Yan et al., 2025, Eck et al., 2025], in contrast to his raw observed rank of 341. This discrepancy illustrates how observed dominance can be distorted by changes in population structure and measurement environments. Paradox probabilities provide a formal way to quantify this distortion, clarifying why exceptional modern performances may appear historically unremarkable when evaluated without contextual adjustment.

6 Discussion

The aptitude-achievement paradox arises when an individual from one context demonstrates higher latent aptitude but simultaneously lower observed achievement than an individual from another context. Although widely encountered in education, labor economics, psychometrics, medicine, and sports analytics, measuring the frequency of this paradox under realistic data-generating regimes is nontrivial. In this article, we formalize the Full House Methodology and investigate the probabilities of observing paradoxes between latent aptitude and observed achievement in comparative analysis of individuals across contexts.

This work is a theoretical clarification of Full House Modeling and provides the definition and solutions for an inferential target, the aptitude-achievement paradox, which extend beyond the baseball motivations in Yan et al. [2025]. In doing so, we note the baseball origins of this work, as articulated by Stephen Jay Gould in his 1996 book *Full House: The Spread of Excellence from Plato to Darwin* (p. 122), from which this framework takes its name: "Baseball is a truly remarkable system for statisticians, manifesting two properties devoutly to be wished, but not often encountered, in actual data: an institution that has

worked by the same rules for a century, and has compiled complete data (nothing major missing) on all measurable aspects of its history." In this way, baseball, with its century of consistent rules and comprehensive measurement, served as an ideal proving ground for the Full House framework [Yan et al., 2025]. Gould concluded his reflection on baseball's statistical regularity with the admission that his sense of calculating "something quite general about the nature of systems" was "just a feeling, not a proof." Here, our work offers more than a feeling, providing rigorous proofs that yield explicit and easy-to-calculate aptitude-achievement paradox probabilities.

We derive closed-form expressions for paradox probabilities under Full House data generating models in two regimes: (i) when observed achievements are i.i.d., and (ii) when context-specific summaries are independent normals with distinct means and variances. The probability rises with population-size asymmetry (large N_1, small N_2) and sampling asymmetry (small n_1, large n_2). Moreover, for a fixed population-size ratio N_2/N_1, it is unchanged by proportional rescaling of the sample sizes (n_1, n_2). We also develop bounds for more general settings and show that, when observed achievement is normal, plug-in estimators yield reliable paradox-probability estimates.

Beyond baseball, the recursive Full House Modeling example in Section 5.3.1 speaks to broader concerns in educational evaluation and admissions. When students transition across institutions of varying selectivity, from less selective undergraduate programs to more selective graduate programs, or vice versa, observed performance metrics may systematically misrepresent underlying aptitude due to differences in both competition depth and measurement environments. The framework developed here provides a principled way to quantify such distortions, though substantial work remains to characterize the specific selection mechanisms and measurement properties that govern real educational pipelines. Our hope is that this methodology can eventually inform more equitable evaluation practices across educational and professional contexts where cross-institutional comparisons are routine but poorly understood.

By elevating aptitude-achievement paradoxes from an anecdotal concern to a model-based, estimable quantity, our framework clarifies when cross-context comparisons are trustworthy, when they are fragile, and how to report that fragility responsibly. This closes the loop between principled latent measurement, realistic selection processes, and interpretable comparative claims.

Acknowledgements

We are grateful to ... and many others for helpful comments, great discussions, and inspiration. We acknowledge the use of ChatGPT-5.2 and Claude 4.5 to aid with proofreading and writing clarity.

Supplementary Materials

The Supplementary Materials contain proofs of mathematical results, together with auxiliary Monte Carlo simulations provided solely for numerical intuition for Theorem 4.3.

References

David J Bartholomew, Martin Knott, and Irini Moustaki. *Latent variable models and factor analysis: A unified approach*. John Wiley & Sons, 2011.

Benjamin S Baumer and Gregory J Matthews. A statistician reads the sports page: there is no avoiding WAR. *Chance*, 27(3):41–44, 2014.

Benjamin S Baumer, Shane T Jensen, and Gregory J Matthews. openwar: An open source system for evaluating overall player performance in major league baseball. *Journal of Quantitative Analysis in Sports*, 11(2):69–84, 2015.

Scott M Berry, C Shane Reese, and Patrick D Larkey. Bridging different eras in sports. *Journal of the American Statistical Association*, 94(447):661–676, 1999.

Trevor G Bond and Christine M Fox. *Applying the Rasch model: Fundamental measurement in the human sciences*. Psychology Press, 2013.

Ralph A. Bradley. Science, statistics, and paired comparisons. *Biometrics*, 32(2):213–239, 1976. doi: 10.2307/2529494.

Ralph A. Bradley. Paired comparisons: Some basic procedures and examples. In P. R. Krishnaiah and P. K. Sen, editors, *Handbook of Statistics*, volume 4, pages 299–326. Elsevier, Amsterdam, 1984.

Ralph A. Bradley and Milton E. Terry. The rank analysis of incomplete block designs: I. the method of paired comparisons. *Biometrika*, 39:324–345, 1952. doi: 10.1093/biomet/39.3-4.324.

Anthony Castrovince. All of your questions about the new baseball rules, answered. *MLB.com*, February 2023. URL https://www.mlb.com/news/mlb-new-rules-for-2023-faq.

Wei Chu and Zoubin Ghahramani. Preference learning with gaussian processes. In *Proceedings of the 22nd International Conference on Machine Learning*, pages 137–144. ACM, 2005. doi: 10.1145/1102351.1102369.

Herbert A. David. *The Method of Paired Comparisons*. Chapman and Hall, London, 2 edition, 1988.

Daniel Eck, Jun Hee Kim, Shen Yan, Muskan Bhatla, and Zheer Wang. *fullhouse: Package for Estimating Talent and Era-Adjusted Performance Metrics*, 2025. URL https://github.com/DEck13/fullhouse. R package version 1.1.

Bradley Efron and Carl Morris. Data analysis using stein's estimator and its generalizations. *Journal of the American Statistical Association*, 70(350):311–319, 1975.

Arpad E. Elo. *The Rating of Chessplayers, Past and Present*. Arco Publishing, 1978.

Susan E Embretson and Steven P Reise. *Item response theory for psychologists*. Psychology Press, 2000.

Gene H Golub and Charles F Van Loan. *Matrix computations*. Johns Hopkins University press, 2013.

Thore Graepel, Tom Minka, and R TrueSkill Herbrich. A bayesian skill rating system. *Advances in Neural Information Processing Systems*, 19(569-576):7, 2007.

Jacques A Hagenaars and Allan L McCutcheon. *Applied latent class analysis*. Cambridge University Press, 2002.

Paul W Holland and Howard Wainer. *Differential item functioning*. Routledge, 2012.

Neil Houlsby, Ferenc Huszar, Zoubin Ghahramani, and Jose Hernández-lobato. Collaborative gaussian processes for preference learning. *Advances in neural information processing systems*, 25, 2012.

Guido Imbens and Konrad Menzel. A causal bootstrap. Technical report, National Bureau of Economic Research, 2018.

Maurice G Kendall. A new measure of rank correlation. *Biometrika*, 30(1-2):81–93, 1938.

Jouni Kerman. A closed-form approximation for the median of the beta distribution. *arXiv preprint arXiv:1111.0433*, 2011.

Frederic M Lord. A paradox in the interpretation of group comparisons. *Psychological bulletin*, 68(5):304, 1967.

Roderick P McDonald. *Test theory: A unified treatment*. psychology press, 1999.

Stephen Nickell and Richard Layard. Labor market institutions and economic performance. *Handbook of labor economics*, 3:3029–3084, 1999.

Stephen Raudenbush and Anthony S Bryk. A hierarchical model for studying school effects. *Sociology of education*, pages 1–17, 1986.

Michael J Schell. *Baseball's all-time best hitters*. Princeton University Press, 2013.

Michael J Schell. *Baseball's all-time best sluggers*. Princeton University Press, 2016.

Edward H Simpson. The interpretation of interaction in contingency tables. *Journal of the Royal Statistical Society: Series B (Methodological)*, 13(2):238–241, 1951.

Robert H Somers. A new asymmetric measure of association for ordinal variables. *American sociological review*, pages 799–811, 1962.

Wayne L Winston, Scott Nestler, and Konstantinos Pelechrinis. *Mathletics: How Gamblers, Managers, and Fans use Mathematics in Sports*. Princeton University Press, 2022.

Shen Yan, Adrian Burgos, Jr, Christopher Kinson, and Daniel J Eck. Comparing baseball players across eras via novel full house modeling. *The Annals of Applied Statistics*, 19(2): 1778–1799, 2025.

Amy Zhao Yu, Shahar Ronen, Kevin Hu, Tiffany Lu, and César A Hidalgo. Pantheon 1.0, a manually verified dataset of globally famous biographies. *Scientific data*, 3(1):1–16, 2016.