A Formal Investigation of Paradoxes Between Latent Aptitude and Observed Achievement via Full House Methodology

Daniel J. Eck[1], Shen Yan[1*]

1. Department of Statistics, University of Illinois Urbana-Champaign

Abstract

1 Introduction

Comparisons of performances across differing contexts, schools, regions, eras, institutions, and tasks are central to education and psychometrics [Holland and Wainer, 2012, Bond and Fox, 2013], labor economics [Nickell and Layard, 1999], medicine, and sports analytics [Berry et al., 1999, Schell, 2013, 2016, Winston et al., 2022, Yan et al., 2025]. These problems have led to the development of a variety of widely used modeling frameworks that estimate relative ability, including empirical-Bayes shrinkage [Efron and Morris, 1975], Bayesian hierarchical models [Raudenbush and Bryk, 1986, Berry et al., 1999], Bradley-Terry-Luce and paired comparisons models [Bradley and Terry, 1952, Bradley, 1976, 1984, David, 1988], Elo-type dynamic rating systems [Elo, 1978, Graepel et al., 2007], Gaussian-process and state-space formulations [Chu and Ghahramani, 2005, Houlsby et al., 2012], item-response and latent class models [McDonald, 1999, Embretson and Reise, 2000, Hagenaars and McCutcheon, 2002, Holland and Wainer, 2012, Bond and Fox, 2013], factor models [McDonald, 1999, Bartholomew et al., 2011], and domain-specific metrics [Baumer and Matthews, 2014, Baumer et al., 2015, Yu et al., 2016].

Despite their differences, these approaches share a common structural assumption: ability is inferred relative to a fixed or implicitly comparable set of observed competitors. While such models adjust outcomes within a given context, they do not explicitly account for changes in the size or structure of the eligible population, nor for context-specific distortions in how latent ability is transformed into observed outcomes. As a consequence, they may not be able to distinguish observed dominance from underlying aptitude when competition depth, selection mechanisms, or measurement conditions vary across contexts.

These limitations give rise to a previously unquantified inferential target: the probability that cross-context comparisons reverse latent orderings. Specifically, an individual drawn

*Daniel J. Eck (dje13@illinois.edu); email addresses: Shen Yan (shenyan3@illinois.edu)

from a deeper or more competitive population may possess higher latent aptitude yet exhibit lower observed achievement than a counterpart from a shallower or more weakly filtered context. We term this phenomenon the aptitude-achievement paradox. Formally, let individual j in context i have an observed $\mathbb{R}$-valued statistic $Y_{i,j}$ and an underlying latent $\mathbb{R}$-valued aptitude $X_{i,j}$, and let individual n in context m have an observed $\mathbb{R}$-valued statistic $Y_{m,n}$, and an underlying latent $\mathbb{R}$-valued aptitude $X_{m,n}$. The probability of aptitude-achievement paradox is

$$\mathrm{P}\left(X_{i,j} > X_{m,n}, Y_{i,j} < Y_{m,n}\right) \tag{1}$$

This quantity captures the structural risk that cross-context comparisons based on observed outcomes reverse the true latent ordering.

The paradox cannot arise if a single, strictly increasing, context-invariant measurement map relates aptitude to achievement and if selection into tasks is ignorable. In practice, however, both measurement heterogeneity (non-invariance, differential item functioning) and non-ignorable selection (who attempts which tasks, faces which opponents, receives which treatment) are pervasive. These features can reverse cross-context rankings even when the latent ranking is unambiguous, echoing classic reversal phenomena such as Simpson's [Simpson, 1951] and Lord's paradoxes [Lord, 1967] while implicating modern concerns about fairness. What practitioners lack is a principled way to quantify how often such reversals should occur under an explicit model of aptitude, measurement, and selection.

A large class of methods quantifies how often observed orderings disagree with a target ordering, closely related to the phenomenon studied here. Classic examples include rank-discordance measures such as Kendall's τ [Kendall, 1938] and Somers' D [Somers, 1962], which summarize the frequency of concordant and discordant pairs based on observed rankings. Because these indices operate entirely on observed outcomes, they neither place individuals on a common latent scale nor disentangle the effects of measurement non-invariance or selection, motivating models that make the measurement and selection processes explicit.

Traditional latent trait models [Bartholomew et al., 2011] focus on estimating underlying abilities from observed categorical or discrete outcomes. For example, the Rasch model [Bond and Fox, 2013] ensures that comparisons of individuals' abilities are invariant to the specific items administered, under the assumption of no differential item functioning (DIF) [Holland and Wainer, 2012]. However, such models do not explicitly leverage information about the size or structure of the source population from which individuals are drawn. Likewise, Bayesian hierarchical and state-space models for latent ability—including Elo-type rating systems [Berry et al., 1999, Elo, 1978, Graepel et al., 2007]—typically condition on the observed competitive or testing pool and capture population-level variation through random effects or dynamic evolution, without modeling variation in eligible population depth or selection into observation across contexts. In parallel, the causal bootstrap [Imbens and Menzel, 2018] emphasizes that valid causal inference requires explicitly situating the observed sample within a larger source population, rather than treating the realized sample as the population of interest.

To address these limitations, we develop a framework for cross-context comparison that explicitly accounts for population depth and selection mechanisms. This framework, which we refer to as Full House Modeling (FHM), was initially proposed in an applied setting comparing baseball players across eras [Yan et al., 2025], and is developed here as a general

approach for cross-contextual evaluation. This framework separates latent aptitude from observed achievement by modeling how three features jointly shape observed rankings: (i) the size of the eligible population in each context, (ii) the mechanism by which individuals are selected or observed, and (iii) the context-specific mapping from latent aptitude to observable performance. Under FHM, individuals are drawn from a common latent ability distribution, while contexts differ in population depth and in how aptitude is filtered and measured. This formulation makes explicit the structural mechanisms through which observed performance can be inflated or suppressed independently of true underlying aptitude.

In this article, we formalize Full House Modeling and study the probability of aptitude–achievement paradoxes (1) arising in cross-context comparisons. We derive explicit paradox probabilities in two settings: first, when observed achievement is independent and identically distributed, and second, when observed achievement follows independent but context-specific normal distributions. We also establish bounds on paradox probabilities in more general settings and show that, under normality, these probabilities can be consistently estimated using plug-in methods. A sensitivity analysis reveals that paradox risk depends on ratios of eligible population sizes rather than their absolute magnitudes. Proofs of main results appear in the Mathematical Appendix, with supplementary technical results provided in online Supplementary Materials.

2 Model Setting

We now motivate the structure of the Full House Model. Let N_i denote the size of population i. We suppose that every individual $k = 1, \ldots, N_i$ has an underlying latent $\mathbb{R}$-valued aptitude $X_{i,1}, \ldots, X_{i,N_i} \overset{iid}{\sim} F_X$. Let $\mathbf{X}_{i,-j}$ be the vector of all latent aptitudes not including the component j. We then define an indicator function $1_i\left(X_{i,j}, \mathbf{X}_{i,-j}\right)$ as a determined and known system inclusion function for population i where a 1 indicates inclusion and a 0 indicates exclusion. We now suppose that an individual j included in system i has an observed $\mathbb{R}$-valued statistic $Y_{i,j} \overset{iid}{\sim} F_{Y_i}$, $j = 1, \ldots, n_i$, $n_i \leq N_i$. As we will see later, the observed $Y_{i,j}$ values are rank-preserving realizations of $X_{i,j}$, provided that $1_i\left(X_{i,j}, \mathbf{X}_{i,-j}\right) = 1$. Therefore $Y_{i,1}, \ldots, Y_{i,n_i} \overset{iid}{\sim} F_{Y_i}$.

Let $X_{i,(j)}$ and $Y_{i,(j)}$ denote the jth order statistic for system i. Here, larger numbers will correspond to a notion of "better." Thus $X_{i,(j)} > X_{i,(k)}$ whenever $j > k$, for example. We now form (Y, X) pairs of random variables for each individual included in the system. These pairs are of the form $(Y_{i,(j)}, X_{i,(r_{i,j})})$ where $r_{i,j}$ indexes the rank of that individual's latent aptitude within the full population of size N_i. As an example, let $1_i\left(X_{i,j}, \mathbf{X}_{i,-j}\right) = 1(X_{i,j} \geq X_{i,(N_i - n_i + 1)})$, then the $(Y_{i,(j)}, X_{i,(r_{i,j})})$ pairs take the form $(Y_{i,(j)}, X_{i,(N_i - n_i + j)})$ for $j = 1, \ldots, n_i$.

The key feature of Full House Modeling is the explicit connection between the elements forming the $(Y_{i,(j)}, X_{i,(r_{i,j})})$ pairs. Full House Modeling states this connection as

$$\begin{aligned} Y_{i,(j)} &= F^{-1}_{Y_{i,(j)}} \circ F_{X_{i,(r_{i,j})}}\left(X_{i,(r_{i,j})}\right), \\ X_{i,(r_{i,j})} &= F^{-1}_{X_{i,(r_{i,j})}} \circ F_{Y_{i,(j)}}\left(Y_{i,(j)}\right), \end{aligned} \tag{2}$$

where $1_i(X_{i,(r_{i,j})}, \mathbf{X}_{i,-(r_{i,j})}) = 1$, and both F_{Y_i} and F_X are continuous and invertible. We summarise Full House Modeling in the definition below.

Definition 2.1 (Full House Model)**.** *Consider populations with sizes N_i and system sizes n_i, $N_i \geq n_i$. Let a system inclusion function $1_i(\cdot,\cdot)$ be specified. Let $X_{i,j} \overset{iid}{\sim} F_X$, $j = 1,\ldots,N_i$. Let $Y_{i,j} \overset{iid}{\sim} F_{Y_i}$, $j = 1,\ldots,n_i$. Form pairs $(Y_{i,(j)}, X_{i,(r_{i,j})})$, $j = 1,\ldots,n_i$ for all individuals such that $1_i(X_{i,(r_{i,j})}, \boldsymbol{X}_{i,-(r_{i,j})}) = 1$. Then the Full House Model generates $Y_{i,(j)}$ values following (2) where F_{Y_i} and F_X are continuous and invertible.*

Remark The Full House Model induces structural missingness by construction: for any individual j such that $1_i(X_{i,j}, \mathbf{X}_{i,-j}) = 0$, the corresponding observed statistic $Y_{i,j}$ is undefined. In subsequent sections, we consider probabilities involving both latent and observed quantities. To formalize this setting, we treat unobserved $Y_{i,j}$ values are not measurable with respect to the observable probability space, reflecting the fact that they lie outside the observable σ-field generated by the system. This perspective allows selection mechanisms to be incorporated directly into probabilistic statements without imputing counterfactual outcomes.

Definition 2.1 assumes a common latent aptitude distribution F_X across contexts, while allowing the observed achievement distributions F_{Y_i} to vary by system. This reflects a modeling choice in which the process generating latent aptitude is taken to be context-neutral, whereas the expression and measurement of aptitude may differ across contexts.

The probability integral transformations in (2) involve the distribution functions of order statistics, namely $F_{X_{i,(r_{i,j})}}$ and $F_{Y_{i,(j)}}$, which may be cumbersome to evaluate directly. An equivalent and more convenient representation follows from the fact that, under continuity, the probability integral transform of an order statistic can be expressed through a Beta distribution. In particular,

$$F_{X_{i,(r_{i,j})}}\left(X_{i,(r_{i,j})}\right) = F_{U_{i,(r_{i,j})}} \circ F_X\left(X_{i,(r_{i,j})}\right), \quad U_{i,(r_{i,j})} \sim \text{Beta}(r_{i,j},\, N_i - r_{i,j} + 1), \tag{3}$$

and similarly,

$$F_{Y_{i,(j)}}\left(Y_{i,(j)}\right) = F_{U_{i,(j)}} \circ F_{Y_i}\left(Y_{i,(j)}\right), \quad U_{i,(j)} \sim \text{Beta}(j,\, n_i - j + 1). \tag{4}$$

This representation allows the mapping in (2) to be implemented using the base distributions F_X and F_{Y_i} together with known Beta distributions, avoiding the need to work explicitly with the distributions of order statistics.

The representations (3) and (4) yield both theoretical and practical advantages for implementing the Full House Modeling flow (2). Although FHM is conceptually formulated in terms of a latent aptitude distribution F_X, this distribution may be difficult or impossible to specify in many applications. The Beta representations allow latent aptitude to be recovered up to a monotone transformation without requiring explicit knowledge of F_X.

Specifically, combining (3) and (4) yields

$$X_{i,(r_{i,j})} = F_X^{-1}\left(F_{U_{i,(r_{i,j})}} \circ F_{U_{i,(j)}} \circ F_{Y_i}\left(Y_{i,(j)}\right)\right).$$

Define

$$W_{i,j} = F_{U_{i,(r_{i,j})}} \circ F_{U_{i,(j)}} \circ F_{Y_i}\left(Y_{i,(j)}\right).$$

For any two cross-context observations W_{i_1,j_1} and W_{i_2,j_2} computed from observed achievements, monotonicity of F_X implies

$$W_{i_1,j_1} \geq W_{i_2,j_2} \implies F_X^{-1}(W_{i_1,j_1}) \geq F_X^{-1}(W_{i_2,j_2}).$$

Consequently, the ordering of latent aptitudes is invariant to the specification of F_X. While F_X is required to recover absolute latent scales, it is a structural nuisance for ranking and comparative analysis. In practice, this allows Full House Modeling to be implemented via a mapping from observed achievement to latent aptitude ranks, even though the conceptual data-generating process flows from aptitude to achievement.

3 Paradoxes Under Identical Achievement Distributions

In this section, we derive explicit probabilities of the aptitude–achievement paradox, as defined in (1), within a Full House Modeling framework involving two populations. We focus on the setting in which the observed achievement distributions are identical across contexts, that is, $F_{Y_i} = F_Y$ for $i = 1, 2$ in Definition 2.1. The following definition formalizes the notion of paradox used throughout the remainder of the article.

Definition 3.1. *Consider the Full House Model in Definition 2.1 with two populations denoted by $i = 1, 2$. We say a aptitude-achievement paradox has occurred when both $X_{1,(r_{1,j})} > X_{2,(r_{2,k})}$ and $Y_{1,(j)} < Y_{2,(k)}$ have occurred. The aptitude-achievement paradox probability is defined as*

$$\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right). \tag{5}$$

As a simple illustrative example of the above, let $r_{i,t} = N_i - n_i + t$ and set $j = n_1$ and $k = n_2$, and further suppose that $Y_{1,j} < Y_{2,k}$ a.s. Then the aptitude-achievement paradox probability (5) is equivalent to $\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}\right)$, and we have $\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}\right) \to 1$ when $N_1 \to \infty$. We now introduce a convenient representation which allows the paradox probability to be expressed in terms of the latent aptitudes alone.

Lemma 3.2. *Consider the Full House Model in Definition 2.1 with two populations, where $F_{Y_i} = F_Y$ for $i = 1, 2$. Then,*

$$\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right) = \mathrm{P}\left(X_{2,(r_{2,k})} < X_{1,(r_{1,j})} < h_{j,k}\left(X_{2,(r_{2,k})}\right)\right), \tag{6}$$

$$\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right) = \mathrm{P}\left(g_{j,k}\left(Y_{2,(k)}\right) < Y_{1,(j)} < Y_{2,(k)}\right) \tag{7}$$

where $h_{j,k}(\cdot) = F^{-1}_{X_{1,(r_{1,j})}} \circ F_{Y_{1,(j)}} \circ F^{-1}_{Y_{2,(k)}} \circ F_{X_{2,(r_{2,k})}}(\cdot)$, $g_{j,k}(\cdot) = F^{-1}_{Y_{1,(j)}} \circ F_{X_{1,(r_{1,j})}} \circ F^{-1}_{X_{2,(r_{2,k})}} \circ F_{Y_{2,(k)}}(\cdot)$.

Our first formal result provides conditions that allow for an explicit aptitude-achievement paradox probability result. Those conditions require the following definition.

Definition 3.3. *Let $\alpha, \beta > 0$. Then the regularized incomplete beta function is defined as*

$$I_{\alpha,\beta}(x) = \frac{1}{B(\alpha,\beta)} \int_0^x u^{\alpha-1}(1-u)^{\beta-1} du,$$

where $B(\alpha,\beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$ is the beta coefficient.

Note that the regularized incomplete beta function is the cumulative distribution function of a Beta(α, β) distribution.

Theorem 3.4. *Consider the Full House Model in Definition 2.1 with two populations, where $F_{Y_i} = F_Y$ for $i = 1, 2$. Denote by I the regularized incomplete beta function. Further assume that*

$$I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1}(t) - I_{r_{1,j},N_1-r_{1,j}+1} \circ I^{-1}_{r_{2,k},N_2-r_{2,k}+1}(t) \geq 0, \qquad \text{for all } t \in (0,1). \tag{8}$$

Under these assumptions,

$$\begin{aligned}
&\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right) \\
&= \sum_{p=j}^{n_1} \binom{n_1}{p} \frac{B(k+p, n_2-n_1-p+k+1)}{B(k, n_2-k+1)} - \sum_{p=r_{1,j}}^{N_1} \binom{N_1}{p} \frac{B(r_{2,k}+p, N_2-N_1-p+r_{2,k}+1)}{B(r_{2,k}, N_2-r_{2,k}+1)}.
\end{aligned}$$

Remarks:

1. The paradox probability in Theorem 3.4 is distribution free. Once the Full House Model is specified, the result depends only on the population sizes N_i, system sizes n_i, the system inclusion function, and the indices of the order statistics under comparison. As established in Section 2, the Beta representations (3)–(4) imply that the distributions of observed order statistics can be expressed independently of the underlying form of F_Y. Lemma 3.2 leverages this structure to eliminate F_Y entirely from the paradox probability calculation,

$$\begin{aligned}
h_{j,k}(x) &= F^{-1}_{X_{1,(r_{1,j})}} \circ F_{Y_{1,(j)}} \circ F^{-1}_{Y_{2,(k)}} \circ F_{X_{2,(r_{2,k})}}(x) \\
&= F^{-1}_{X_{1,(r_{1,j})}} \circ I_{j,n_1-j+1} \circ \left(F_Y \circ F_Y^{-1}\right) \circ I^{-1}_{k,n_2-k+1} \circ F_{X_{2,(r_{2,k})}}(x) \\
&= F^{-1}_{X_{1,(r_{1,j})}} \circ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1} \circ F_{X_{2,(r_{2,k})}}(x).
\end{aligned}$$

2. Likewise, the latent aptitude distribution F_X plays no role in the final expression for the paradox probability. Although Lemma 3.2 expresses the event in terms of latent order statistics, $\mathrm{P}\left(X_{2,(r_{2,k})} < X_{1,(r_{1,j})} < h_{j,k}\left(X_{2,(r_{2,k})}\right)\right)$. the monotone coupling induced by Full House Modeling ensures that all integrals involving F_X reduce to functions of Beta distributions and order indices. As shown in the proof of Theorem 3.4, a change of variables removes dependence on F_X, rendering it a structural nuisance parameter for paradox probabilities.

3. Combining these observations, the paradox probability admits a closed-form expression involving only regularized incomplete beta functions. In particular, the probability simplifies to

$$\int_0^1 \left\{ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1}(t) - I_{r_{1,j},N_1-r_{1,j}+1} \circ I^{-1}_{r_{2,k},N_2-r_{2,k}+1}(t) \right\}^+ dt.$$

   The technical condition (8) ensures that the integrand is nonnegative for all $t \in (0,1)$, guaranteeing that this expression defines a valid probability. In subsequent sections, we characterize when this condition holds and examine its empirical implications.

Often it is of interest to compare the individuals with the highest aptitudes in their respective populations. Below we state the aptitude-achievement paradox probability in such setting using a Full House Model with two populations with the specification that $F_{Y_1} = F_Y$ for $i = 1, 2$.

Corollary 3.4.1. *Suppose that the conditions of Theorem 3.4 hold. Suppose that* $1_1(X_{1,(N_1)}, \boldsymbol{X}_{1,-(N_1)}) = 1_2(X_{2,(N_2)}, \boldsymbol{X}_{2,-(N_2)}) = 1$. *Then,*

$$\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \begin{cases} \frac{n_2}{n_1+n_2} - \frac{N_2}{N_1+N_2}, & \text{if } \frac{n_1}{N_1} \leq \frac{n_2}{N_2} \\ 0, & \text{otherwise} \end{cases}.$$

A similar result holds for the individuals with the lowest latent aptitudes in the population.

Corollary 3.4.2. *Suppose that the conditions of Theorem 3.4 hold. Suppose that* $1_1(X_{1,(1)}, \boldsymbol{X}_{1,-(1)}) = 1_2(X_{2,(1)}, \boldsymbol{X}_{2,-(1)}) = 1$. *Then,*

$$\mathrm{P}\left(X_{1,(1)} > X_{2,(1)}, Y_{1,(1)} < Y_{2,(1)}\right) = \begin{cases} \frac{N_2}{N_1+N_2} - \frac{n_2}{n_1+n_2}, & \text{if } \frac{n_1}{N_1} \geq \frac{n_2}{N_2} \\ 0, & \text{otherwise} \end{cases}.$$

We now demonstrate a setting in which technical condition (8) holds. This setting is based on a configuration of n_j, N_j, $j = 1, 2$ under the assumption that the systems are populated with individuals from the source populations who have the highest aptitudes.

Proposition 3.5. *Consider the Full House Model in Definition 2.1 with two populations, where* $F_{Y_i} = F_Y$ *for* $i = 1, 2$, *and let the system inclusion function be* $1_i(X_{i,j} \geq X_{i,(N_i - n_i + 1)})$. *Denote by* I *the regularized incomplete beta function. Suppose that* $\frac{n_1}{n_2} \leq 1$ *and* $\frac{N_2}{N_1} \leq 1$, *and set* $k = j$. *Under these assumptions, we conclude that technical condition* (8) *holds and the conclusion of Theorem 3.4 applies.*

4 Paradox Under Different Distribution

In this section we will provide aptitude-achievement paradox probability results in a Full House Modeling setting involving two populations when we relax the specification that $F_{Y_1} = F_{Y_2}$ in Definition 2.1. We first provide bounds for aptitude-achievement paradox probability in general settings. We then provide explicit aptitude-achievement paradox probability results when Y_1 and Y_2 are normally distributed with different means and variances.

4.1 General bounds

The general bounds we derive require a relaxed form of first-order stochastic dominance that constrains the distributions of Y_1 and Y_2 are 'close' to each other. More formally, we introduce first-order ε-weak stochastic dominance, which is necessary for our aptitude-achievement paradox probability results.

Definition 4.1. *We will say that Y_2 is first order ε-weakly stochastically dominant over Y_1 if, for $\varepsilon > 0$, $\mathrm{P}(Y_2 \leq y) \leq \mathrm{P}(Y_1 \leq y) \leq \mathrm{P}(Y_2 \leq y) + \varepsilon$ for all y and $\mathrm{P}(Y_2 \leq y) < \mathrm{P}(Y_1 \leq y) \leq \mathrm{P}(Y_2 \leq y) + \varepsilon$ for some y.*

Under first-order ε-weak stochastic dominance, we derive a general lower and upper bound on the aptitude-achievement paradox probability when $F_{Y_1} \neq F_{Y_2}$.

Proposition 4.2. *Suppose that the system inclusion mechanism has the property that $1_i\left(X_{i,(N_i)}, \boldsymbol{X}_{i,-(N_i)}\right) = 1$, for $i = 1, 2$, and that the Full House Model is of the form*

$$\begin{aligned} Y_{i,j} &\sim F_{Y_i}(y), \qquad i = 1, 2, \qquad j = 1, \ldots, n_i, \\ X_{i,j} &\sim F_X(x), \qquad i = 1, 2, \qquad j = 1, \ldots, N_i, \end{aligned}$$

where $F_{Y_i}^{-1}(y)$ and $F_X^{-1}(x)$ exist. Further suppose that Y_2 is first order ε-weakly stochastically dominant over Y_1 for some $\varepsilon > 0$ and $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$. Then,

$$\begin{aligned} \frac{n_2}{n_1 + n_2} - \frac{N_2}{N_1 + N_2} &\leq \mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\ &\leq n_2 \sum_{i=0}^{n_1} \frac{\binom{n_1}{i}\varepsilon^i}{n_1 + n_2 - i} - \frac{N_2}{N_1 + N_2} \end{aligned} \tag{9}$$

These bounds are a prescription under general distributions for achievement in Full House Modeling. However, we can still obtain explicit aptitude-achievement paradox probabilities when we specify $Y_i \sim N(\mu_i, \sigma_i^2)$, $i = 1, 2$.

4.2 Normal distributions

This section will develop explicit aptitude-achievement paradox probabilities (5) when a Full House Model in Definition 2.1 additionally specifies that $Y_i \sim N(\mu_i, \sigma_i^2)$, $i = 1, 2$. We will focus on the case where the system inclusion mechanism has the property that $1_i(X_{i,(N_i)}, \mathbf{X}_{i,-(N_i)}) = 1$ for $i = 1, 2$.

The outline for our results is similar to that in Section 3 where a technical condition is needed to prove the main result – like condition (8) in Theorem 3.4 –and further results provide configurations of the sample sizes that satisfy the technical condition. However, in this setting with $Y_i \sim N(\mu_i, \sigma_i^2)$, $i = 1, 2$, the corresponding technical condition yields an additional remainder term R that prevents an exact closed-form expression. We show below that this remainder term is negligible under mild and interpretable conditions on population fractions and distributional parameters.

Similar to the proof of Theorem 3.4, Lemma 3.2 will provide a representation more conducive to formal integration needed to prove our main result. Only this time we will use (7) in Lemma 3.2 to represent the aptitude-achievement paradox probability as

$$\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \mathrm{P}\left(g_{n_1,n_2}\left(Y_{2,(n_2)}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right),$$

where, from the definition of g in Lemma 3.2 applied to our current setting, we have

$$g_{n_1,n_2}(x) = F_{Y_1}^{-1}\left(F_{Y_2}(x)^{\frac{n_2 N_1}{N_2 n_1}}\right) = \Phi^{-1}\left(\Phi\left(\frac{x - \mu_2}{\sigma_2}\right)^{\frac{n_2 N_1}{N_2 n_1}}\right)\sigma_1 + \mu_1, \tag{10}$$

and Φ defines the cumulative distribution function of the standard normal distribution. From here, we have that

$$\begin{aligned}
&\mathrm{P}\left(g_{n_1,n_2}\left(Y_{2,(n_2)}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \mathrm{P}\left(F_{Y_1}^{-1}\left(F_{Y_2}(Y_{2,(n_2)})^{\frac{n_2N_1}{N_2n_1}}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\
&\quad = \int_A \int_{F_{Y_1}^{-1}\left(F_{Y_2}(x)^{\frac{n_2N_1}{N_2n_1}}\right)}^{x} dF_{Y_1,(n_1)}(y) dF_{Y_2,(n_2)}(x),
\end{aligned}$$

where $A := A(\mu_1, \mu_2, \sigma_1, \sigma_2, n_1, n_2, N_1, N_2)$ is the range of integration so that the inner integral above is properly specified. More formally,

$$\begin{aligned}
A &= \left\{\omega : F_{Y_1}^{-1}\left(F_{Y_2}(Y_{2,(n_2)}(\omega))^{\frac{n_2N_1}{N_2n_1}}\right) \le Y_{2,(n_2)}(\omega)\right\} \\
&= \left\{\omega : \Phi\left(\frac{Y_{2,(n_2)}(\omega) - \mu_2}{\sigma_2}\right)^{\frac{n_2}{N_2}} \le \Phi\left(\frac{Y_{2,(n_2)}(\omega) - \mu_1}{\sigma_1}\right)^{\frac{n_1}{N_1}}\right\},
\end{aligned} \tag{11}$$

where the second equality in (11) follows from (10). Continuing with the probability calculation above

$$\begin{aligned}
&\int_A \int_{F_{Y_1}^{-1}\left(F_{Y_2}(x)^{\frac{n_2N_1}{N_2n_1}}\right)}^{x} dF_{Y_1,(n_1)}(y) dF_{Y_2,(n_2)}(x) \\
&= \int_A F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}} dF_{Y_2,(n_2)}(x) \\
&= \int_{-\infty}^{\infty} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}} dF_{Y_2,(n_2)}(x) - \int_{A^c} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}} dF_{Y_2,(n_2)}(x).
\end{aligned} \tag{12}$$

The left-hand side of (12) can (and will) be further evaluated and leads to the desired explicit aptitude-achievement paradox probability. The right-hand side of (12) represents a remainder term $R := R(\mu_1, \mu_2, \sigma_1, \sigma_2, n_1, n_2, N_1, N_2)$,

$$R = -\int_{A^c} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}} dF_{Y_2,(n_2)}(x), \tag{13}$$

where the set A in (13) is defined in (11). The remainder term may seem unintuitive. However, we will now motivate a simple setting that allows for R to be negligible when n_2 is moderately large.

From (11) and (13), we have

$$\begin{aligned}
|R| &= \left|\int_{A^c} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}} dF_{Y_2,(n_2)}(x)\right| \\
&\le \int_{A^c} \left|F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}}\right| dF_{Y_2,(n_2)}(x) \\
&\le \mathrm{P}\left(\Phi\left(\frac{Y_{2,(n_2)} - \mu_2}{\sigma_2}\right)^{\frac{n_2}{N_2}} \ge \Phi\left(\frac{Y_{2,(n_2)} - \mu_1}{\sigma_1}\right)^{\frac{n_1}{N_1}}\right).
\end{aligned}$$

Now, when we specify that $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$ as in Proposition 4.2, we have

$$\begin{aligned}
&\mathrm{P}\left(\Phi\left(\frac{Y_{2,(n_2)}-\mu_2}{\sigma_2}\right)^{\frac{n_2}{N_2}} \geq \Phi\left(\frac{Y_{2,(n_2)}-\mu_1}{\sigma_1}\right)^{\frac{n_1}{N_1}}\right) \\
&\quad \leq \mathrm{P}\left(\Phi\left(\frac{Y_{2,(n_2)}-\mu_2}{\sigma_2}\right) \geq \Phi\left(\frac{Y_{2,(n_2)}-\mu_1}{\sigma_1}\right)\right) \\
&\quad = \mathrm{P}\left(\frac{Y_{2,(n_2)}-\mu_2}{\sigma_2} \geq \frac{Y_{2,(n_2)}-\mu_1}{\sigma_1}\right).
\end{aligned}$$

Under the assumption that $\sigma_1 \leq \sigma_2$, straightforward algebra yields

$$\mathrm{P}\left(\frac{Y_{2,(n_2)}-\mu_2}{\sigma_2} \geq \frac{Y_{2,(n_2)}-\mu_1}{\sigma_1}\right) = \mathrm{P}\left(Y_{2,(n_2)} \leq \left(\frac{\mu_2}{\sigma_2}-\frac{\mu_1}{\sigma_1}\right)\left(\frac{1}{\sigma_2}-\frac{1}{\sigma_1}\right)^{-1}\right).$$

Under the additional assumption that $\mu_1 \leq \mu_2$, the right-hand side is bounded above by

$$\mathrm{P}\left(Y_{2,(n_2)} \leq \mu_2\right).$$

Putting it all together, we have

$$|R| \leq \mathrm{P}\left(Y_{2,(n_2)} \leq \mu_2\right) \leq \left(\frac{1}{2}\right)^{n_2},$$

when $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$, $\sigma_1 \leq \sigma_2$, and $\mu_1 \leq \mu_2$. Therefore, although the remainder term R in (13) arises from the restriction of the integration domain to A, it is exponentially small in n_2 under a familiar ordering of population fractions (see Proposition 4.2) and a simple parametric configuration. Consequently, the aptitude–achievement paradox probability is well approximated by the unrestricted integral given by the left-hand side of (12)with R contributing only a negligible correction for moderately large n_2. We now introduce a multivariate normal representation that allows this principal term to be expressed in closed form.

Let $\mathbf{W} \sim \mathcal{N}_{(n_1+n_2-1)}(\mathbf{0}, \mathbf{\Sigma})$ and covariance matrix

$$\mathbf{\Sigma} = \begin{bmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{bmatrix}, \tag{14}$$

defined with the following structure

$$\begin{aligned}
\Sigma_{11} &= \begin{bmatrix} \frac{\sigma_2^2}{\sigma_1^2}+1 & \frac{\sigma_2^2}{\sigma_1^2} & \cdots & \frac{\sigma_2^2}{\sigma_1^2} \\ \frac{\sigma_2^2}{\sigma_1^2} & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \frac{\sigma_2^2}{\sigma_1^2} \\ \frac{\sigma_2^2}{\sigma_1^2} & \cdots & \frac{\sigma_2^2}{\sigma_1^2} & \frac{\sigma_2^2}{\sigma_1^2}+1 \end{bmatrix}_{n_1 \times n_1} \qquad \Sigma_{12}^T = \Sigma_{21} = \begin{bmatrix} \frac{\sigma_2}{\sigma_1} & \cdots & \cdots & \frac{\sigma_2}{\sigma_1} \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ \frac{\sigma_2}{\sigma_1} & \cdots & \cdots & \frac{\sigma_2}{\sigma_1} \end{bmatrix}_{(n_2-1)\times n_1} \\
&\qquad\qquad \Sigma_{22} = \begin{bmatrix} 2 & 1 & \cdots & 1 \\ 1 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & 1 \\ 1 & \cdots & 1 & 2 \end{bmatrix}_{(n_2-1)\times(n_2-1)}.
\end{aligned} \tag{15}$$

With this structure we now provide an explicit aptitude-achievement paradox probability, the main result of this section.

Theorem 4.3. *Assume a Full House Model in Definition 2.1 and additionally specify that* $Y_i \sim N(\mu_i, \sigma_i^2)$*, and suppose that the system inclusion mechanism has the property that* $1_i(X_{i,(N_i)}, \boldsymbol{X}_{i,-(N_i)}) = 1$ *for* $1 = 1, 2$*. Let* $\mathbf{W} \sim \mathcal{N}_{(n_1+n_2-1)}(\mathbf{0}, \boldsymbol{\Sigma})$ *where* $\boldsymbol{\Sigma}$ *is as defined in (14) and (15). Then,*

$$\begin{aligned} &\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\ &\quad = n_2 F_{\mathbf{W}}(\mathbf{w} = (\frac{\mu_2 - \mu_1}{\sigma_1}, \ldots, \frac{\mu_2 - \mu_1}{\sigma_1}, 0, \ldots, 0)) - \frac{N_2}{N_1 + N_2} + R, \end{aligned} \tag{16}$$

where R *is defined in (13),* $F_{\mathbf{W}}$ *is the distribution function of* $\mathbf{W}$*, and the first* n_1 *elements of* $\mathbf{w}$ *are* $\frac{\mu_2-\mu_1}{\sigma_1}$ *and the last* $n_2 - 1$ *elements are 0.*

Moreover, we have that

$$|R| \leq \left(\frac{1}{2}\right)^{n_2}, \tag{17}$$

when $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$*,* $\sigma_1 \leq \sigma_2$*, and* $\mu_1 \leq \mu_2$*.*

We now show that plug-in estimators of the normal distribution parameters μ_i and σ_i, $i = 1, 2$, yield reliable estimates of the paradox probability (16). Our approach requires properties for matrices of the form

$$\Sigma = \begin{bmatrix} \rho + 1 & \rho & \ldots & \rho \\ \rho & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \rho \\ \rho & \ldots & \rho & \rho + 1 \end{bmatrix}_{n \times n}.$$

These properties are

$$\det(\Sigma) = \rho n + 1, \tag{18}$$

and

$$\Sigma^{-1} = \frac{1}{\rho n + 1} \begin{bmatrix} \rho(n-1) + 1 & -\rho & \ldots & -\rho \\ -\rho & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & -\rho \\ -\rho & \ldots & -\rho & \rho(n-1) + 1 \end{bmatrix}_{n \times n}, \tag{19}$$

where (19) follows from an application of the Woodbury identity [Golub and Van Loan, 2013].

Before we state our result, we introduce the Lemma 4.4

Lemma 4.4. *Let* $\mathbf{W} \sim \mathcal{N}_{n_1+n_2-1}(\mathbf{0}, \boldsymbol{\Sigma})$ *be a multivariate normal random vector with mean vector* $\mathbf{0}$ *and covariance matrix* $\boldsymbol{\Sigma}$ *as defined in (14) and (15). Let* $\Phi_{\boldsymbol{\Sigma}}(\mathbf{w})$ *denote the cumulative distribution function of a mean-zero multivariate normal random vector with covariance*

matrix $\boldsymbol{\Sigma}$*, evaluated at* $\mathbf{w} \in \mathbb{R}^{n_1+n_2-1}$*. Define* $c := c_{n_1+n_2-1}$ *as the maximal conditional one-dimensional Gaussian probability arising in the sequential factorization of the joint CDF, that is,*

$$
\begin{aligned}
c = \max \Big\{ & \mathrm{P}(W_{n_1+n_2-1} \le w_{n_1+n_2-1} \,|\, W_1 = w_1, \ldots, W_{n_1+n_2-2} = w_{n_1+n_2-2}), \\
& \mathrm{P}(W_{n_1+n_2-2} \le w_{n_1+n_2-2} \,|\, W_1 = w_1, \ldots, W_{n_1+n_2-3} = w_{n_1+n_2-3}), \\
& \ldots, \mathrm{P}(W_2 \le w_2 \mid W_1 = w_1), \mathrm{P}(W_1 \le w_1) \Big\}.
\end{aligned} \tag{20}
$$

Then $c \in (0,1)$ *for all* n_1*,* n_2*, and*

$$
\Phi_{\boldsymbol{\Sigma}}(\mathbf{w}) \le c^{n_1+n_2-1} \tag{21}
$$

Proof. The proof can be found in the Supplementary Materials. □

Theorem 4.5. *Assume a Full House Model in Definition 2.1 and additionally specify that* $Y_i \sim N(\mu_i, \sigma_i^2)$*, and suppose that the system inclusion mechanism has the property that* $1_i(X_{i,(N_i)}, \boldsymbol{X}_{i,-(N_i)}) = 1$ *for* $1 = 1, 2$*. Let* $\mathbf{W} \sim \mathcal{N}_{(n_1+n_2-1)}(\mathbf{0}, \boldsymbol{\Sigma})$ *where* $\boldsymbol{\Sigma}$ *is as defined in (14) and (15). Suppose that* $\hat{\mu}_1$*,* $\hat{\mu}_2$*,* $\hat{\sigma}_1$*, and* $\hat{\sigma}_2$ *are, respectively, root sample-size consistent estimators of* μ_1*,* μ_2*,* σ_1*, and* σ_2*. Then*

$$
\begin{aligned}
& \mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)} \mid \hat{\mu}_1, \hat{\mu}_2, \hat{\sigma}_1, \hat{\sigma}_2\right) \\
& \quad - \mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)} \mid \mu_1, \mu_2, \sigma_1, \sigma_2\right) \\
& = O_P(n_1^{\frac{1}{2}} n_2 c^{n_1+n_2-3} + n_1 n_2^{\frac{1}{2}} c^{n_1+n_2-3}),
\end{aligned}
$$

where c *is defined in the Lemma 4.4.*

Proof. The proof can be found in the Supplementary Materials. □

5 Examples

5.1 Sensitivity Analysis for Theorem 3.4

In this section, we conduct a sensitivity analysis to evaluate the influence of the parameters N_1, N_2, n_1, and n_2 in Theorem 3.4.

Under the setting of Theorem 3.4, we assume $X_1 = X_2 = X$, $Y_1 = Y_2 = Y$, where $X \sim \text{Pareto}(\alpha)$ and $Y \sim \mathcal{N}(0,1)$. Without loss of generality, we set $j = n_1 - 1$ and $k = n_2 - 1$, corresponding to the selection of the second-best performer in the paradox probability, as defined in Definition 3.1.

In each simulation, we randomly sample n_1 and n_2 observations from the distribution of Y, and compute their latent aptitudes using Equation (2). Based on these values, we evaluate the paradox probability according to Definition 3.1. We repeat this simulation 10000 times and report the average paradox probability.




Figure 1: Sensitivity Analysis for Paradox Probability

Figure 1 presents the results of the sensitivity analysis for the paradox probability described in Theorem 3.4. In Figure 1 (A), we fix $n_1 = 40$ and $n_2 = 60$, and vary N_1 and N_2 from 200 to 2000. In Figure 1 (B), we fix $N_1 = 10^3$ and $N_2 = 10^3$, and vary n_1 and n_2 from 20 to 200. In Figure 1 (C), we fix $n_1 = 40$ and $N_2 = 10^3$, and vary n_1 from 20 to 200 and N_2 from 200 to 2000. In Figure 1 (D), we fix $N_1 = 10^3$ and $n_2 = 60$, and vary N_1 from 200 to 2000 and n_2 from 20 to 200.

As shown in Figure 1 (A), a small N_2 and large N_1 increase the paradox probability by making $\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}\right)$ approach 1. Similarly, in Figure 1 (B), a small n_1 combined with a large n_2 leads to a higher paradox probability, since it drives $\mathrm{P}\left(Y_{1,(j)} < Y_{2,(k)}\right)$ towards 1.

We also investigate the effect of the ratio between N_2 and N_1 on the paradox probability. Let $c = \frac{N_2}{N_1}$, and vary c from 0.1 to 1 in increments of 0.1. Without loss of generality, we set $j = n_1 - 1$ and $k = n_2 - 1$, corresponding to the selection of the second-best performer in the paradox probability, as defined in Definition 3.1.

In each simulation, we fix $n_1 = 60$ and $n_2 = 40$, and vary N_1 from 100 to 2000. We repeat this simulation 10000 times and report the average paradox probability.




Figure 2: Effect of the ratio between N_2 and N_1 on the paradox probability

Figure 2 presents the results illustrating the effect of the ratio between N_2 and N_1 on the paradox probability under different settings. As shown in the figure, the paradox probability decreases as c increases, which is reasonable since $\mathbb{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}\right)$ decreases as N_2 increases.

Figure 2 also demonstrates that the paradox probability remains constant when the ratio between N_2 and N_1 is fixed. This implies that it is unnecessary to focus on the exact values of N_1 and N_2 when computing the paradox probability; rather, the ratio N_2/N_1 alone determines its value.

5.2 Recursive Full House Models

In this section we will investigate a simple recursive Full House Models in which Full House Models are repeatedly applied to individuals as they progress through iterations of some generic process. The general setup is that, in the first round, latent aptitude $X_j \overset{iid}{\sim} F_X$, $j = 1, \ldots, N$ will be generated and then a label will be randomly assigned to each individual. There will be m labels and each label will designate the outcome distribution. $F_{Y_{i,j}}$, $i = 1, \ldots, n$, $j = 1, \ldots, n_i$. Then outcomes will be realized as

$$Y_{i,(r_{i,(j)})} = F^{-1}_{Y_{i,j}} \circ I^{-1}_{r_{i,(j)}, n_i - r_{i,(j)}+1} \circ I_{j,N-j+1} \circ F_X(X_{(j)}), \tag{22}$$

where $r_{i,(j)}$ is the rank of $X_{(j)}$ among the n_i latent aptitudes also assigned label i.

In the next iteration, all observations will be combined and then sorted in descending order by $Y_{i,(j)}$. The labels will be assigned in descending order, the largest n_m values of $Y_{i,(j)}$ are assigned label m, the next largest n_{m-1} values of $Y_{i,(j)}$ are assigned label $m-1$, and so on. We will setup these simulation so that Y_i stochastically dominates Y_j whenever $i > j$ so that higher outcomes are associated with higher labels. Then new Y values are generated according to (22). The process will repeat until a pre-assigned stopping rule is satisfied. Note that this simulation setup assumes an immutable latent aptitude.

5.2.1 Stop when highest latent aptitude is included in the highest-performing group

Here we consider a recursive Full House Modeling setup described in Section 5.2 in which the recursion stops at the first iteration when $X_{(N)}$ is assigned label m. Here we specify that $N = 100,000$, $m = 200$, $n_i = 500$, $i = 1, \ldots, m$. We will assume that $X_j \overset{iid}{\sim}$ Pareto(1.16), $j = 1, \ldots, N$. And we will assume that $Y_{i,j} \overset{iid}{\sim} N(\mu_i, \sigma_i)$, $j = 1, \ldots, n_i$ and $i = 1, \ldots, m$. We will specify that the μ_is as a set of 200 equally spaced values between 60 and 80 with μ_i increasing in i, ie $\mu_i = 60 - 20/199 + 20i/199$, $i = 1, \ldots, m$. We will fix $\sigma_i = \sigma$ and consider $\sigma \in \{0.5, 1, 2\}$. These simulation will be repeated 500 times.

To illustrate the procedure, consider a simplified scenario in which all $X_{(j)}$ values are realized at their median value. With this simplification, $Y_{i,(r_{i,(j)})}$ in (22) is approximately $F^{-1}_{Y_{i,j}}((r_{i,(j)} - 1/3)/(n_i + 1/3))$ [Kerman, 2011]. And we have that $Y_{i,(n_i)} - Y_{i+1,(n_{i+1}-1)} > \mu_{i+1} - \mu_i$ for all considered values of σ. The individual with latent ability $X_{(N)}$ will consistently attain the value $Y_{i,(n_i)}$, regardless their assigned group. Consequently, this individual's position among the m groups will improve as the simulation advances.

Results can be seen in Figure 3. These Monte Carlo simulations reveal that it typically takes several rounds before the individual with the top aptitude belongs to the group that allows for the highest performances. On average, with σ set to 0.5, 1, and 2, it takes 2.61, 4.19, and 7.46 rounds, respectively, for the individual with the top aptitude to join the highest-performing group (respective standard errors: 0.05, 0.09, 0.18).

These findings resonate with several empirical studies examining bias and distortion in real-world evaluative systems. The GRADE algorithm developed by Waters and Miikkulainen [2014] demonstrates how admission systems can bypass nuanced information (e.g., personal statements) in favor of machine-learned proxies, potentially reinforcing existing filters. Similarly, Alvero et al. [2020] show that seemingly innocuous features (such as the word baseball) can serve as strong demographic signals. This raises concerns about indirect inference. A broader review by Baker and Hawn [2022] warns that algorithmic systems may harbor unknown unknowns, especially in global or under-examined subpopulations. Finally, the meta-analyses in Walton and Spencer [2009] provide direct evidence that outcome-based measures (e.g., grades, test scores) can systematically understate the aptitude of marginalized groups due to identity-based threat.

These phenomena of latent talent being misclassified, sorted, and repeatedly filtered through context-specific outcome layers is precisely what this presentation of recursive Full House Modeling aims to formalize and study. At any stage in the selection process, you cannot be sure that the highest marks correspond to individuals with the highest ability.




Figure 3: The number of rounds that it takes for $X_{(N)}$ to be assigned label m in a recursive Full House Modeling framework in Section 5.2 with specifications in Section 5.2.1.

While the setting we consider is stylized and assumes fixed latent aptitude, it serves to illustrate how structural features of evaluative systems, not just randomness, can delay or obscure the recognition of true potential.

5.3 Batting Averages in Major League Baseball

We now apply Theorem 4.3 and the reliability guarantees in Theorem 4.5 to evaluate the contextual extremeness of observed outcomes in a real-world system where the underlying distribution of aptitude is not directly observable and evolves over time. Major League Baseball (MLB) offers a canonical example of such a system. Here, player skill (latent batting aptitude) is inferred only indirectly through season-level observed outcomes (batting averages), which are shaped by changing population structures and environmental variance. See Figure 4 presents smoothed estimates of the mean and standard deviation of batting averages for full-time players from 1871 to 2023.

In particular, MLB introduced a series of rule changes in 2023 intended to increase in-game action, including a restriction on defensive positioning designed to raise batting averages on balls in play (BABIP) [Castrovince, 2023].

First, the 2023 mean batting average for full-time players increased by 0.0057 (corresponding to an additional hit about every four baseball games) with a reduction in variability. A two-sample tests for differences in batting averages among full-time players from the 2023 and 2022 seasons has a p-value of 0.039. Thus, the rule changes seemed to be effective in




Figure 4: Seasonal means and standard deviations of batting averages for full-time players from 1871-2023. Smoothing is applied to both trends.

yielding increased batting averages[1] Moreover, the 2023 MLB batting leader Luis Arraez had a 0.354 batting average which was the first batting average to exceed .350, a mark of excellence, since Josh Hamilton in 2010.

Now we turn to applying the results of Theorem 4.3 in this example to contextualize the effect of these rule changes. Here we assume that seasonal batting averages of full-time players are normally distributed. The true means and standard deviations of these distributions are unknown. Thus, we estimate them via plug-in sample mean and sample standard-deviation and invoke the reliability guarantees from Theorem 4.5. Notably, we see from Figure 4 that both the average and variability of batting averages in 2023 are among the lowest observed in the dataset, indicating a historically suppressed environment for batting averages of full-time batters.

We now apply Theorem 4.3 to quantify paradox probabilities for extreme batting averages when comparing the 2023 season to all prior seasons. We treat the 2023 season as system 1 and each season from 1871 to 2022 as system 2. Figure 5 displays the resulting paradox probabilities (16), estimated via plug-in using sample means and standard deviations. Because these estimators are $\sqrt{n}$-consistent, the reliability guarantees of Theorem 4.5 apply.

[1]Technically, Castrovince [2023] referred to MLB's desire to increased batting averages on balls in play (BABIP) while this analysis focused on batting averages. BABIP increased by 0.0061 from 2022 to 2023.




Figure 5: Comparison of paradox probabilities and empirical exceedance rates relative to the 2023 season, based on data from 1871-2022. The blue line shows estimated paradox probabilities via plug-in using Theorem 4.3. The black dashed line displays a smoothed empirical exceedance rate for Arraez's 2023 batting average.

For comparison, Figure 5 also reports a smoothed empirical exceedance rate, defined as the proportion of historical seasons in which the maximum batting average exceeded Luis Arraez's 2023 mark. The estimated paradox probabilities closely track this empirical benchmark. Both quantities increase steadily as one moves backward in time, reflecting the fact that earlier eras—characterized by higher means, larger variances, and smaller effective populations—were structurally more conducive to producing extreme observed batting averages, even if underlying talent levels were lower.

Under Full House Modeling, Arraez's 2023 performance ranks 68th all-time among full-time batters after accounting for population depth and contextual effects [Yan et al., 2025, Eck et al., 2025], in contrast to his raw observed rank of 341. This discrepancy illustrates how observed dominance can be distorted by changes in population structure and measurement environments. Paradox probabilities provide a formal way to quantify this distortion, clarifying why exceptional modern performances may appear historically unremarkable when evaluated without contextual adjustment.

6 Discussion

The aptitude-achievement paradox arises when an individual from one context demonstrates higher latent aptitude but simultaneously lower observed achievement than an indi-

vidual from another context. Although widely encountered in education, labor economics, psychometrics, medicine, and sports analytics, measuring the frequency of this paradox under realistic data-generating regimes is nontrivial. In this article, we formalize the Full House Methodology and investigate the probabilities of observing paradoxes between latent aptitude and observed achievement in comparative analysis of individuals across contexts.

Full House Modeling provides a direct comparison of individuals' latent aptitudes and observed achievements across different contexts, which is a novel contribution to latent trait modeling. While traditional latent trait models [Bartholomew et al., 2011] often focus on estimating underlying traits from observed categorical data , our approach specifically quantifies the probability of paradoxical outcomes in comparative settings. Full House Modeling formalizes the comparison of individuals' performance across distinct contexts based on the large source population size N, thereby offering a more nuanced understanding of individual differences and the impact of contextual factors on observed outcomes. In parallel, the causal bootstrap [Imbens and Menzel, 2018] highlights that valid causal inference similarly requires situating the observed sample within large source population to properly account for sampling uncertainty. By comparison, the Rasch model [Bond and Fox, 2013] ensures that comparisons of individuals' abilities are independent of the specific items administered, and that comparisons of item difficulties are independent of the sample of individuals tested, but it does not explicitly leverage the source population structure. Likewise, the bayesian hierarchical modeling [Berry et al., 1999] with the application in comparison of individuals' latent aptitudes typically captures population-level variation through seasonal or group-level random effects rather than through an explicit embedding into a large source population. Full House Modeling enhances broader statistical modeling practices by providing a framework for robust comparative analysis, particularly in situations where direct, context-independent comparisons are challenging.

We derive closed-form expressions for the paradox probability in two regimes: (i) when observed achievements are i.i.d., and (ii) when context-specific summaries are independent normals with distinct means and variances. The probability rises with population-size asymmetry (large N_1, small N_2) and sampling asymmetry (small n_1, large n_2). Moreover, for a fixed population-size ratio N_2/N_1, it is unchanged by proportional rescaling of the sample sizes (n_1, n_2). We also develop bounds for more general settings and show that, when observed achievement is normal, a plug-in estimator yields reliable paradox-probability estimates.

Our findings align with a growing body of evidence that real-world evaluative systems routinely substitute *proxies* for the constructs they claim to measure, with consequential distortion. The GRADE algorithm of Waters and Miikkulainen [2014] shows how admissions workflows may privilege machine-learned surrogates over nuanced information (e.g., personal statements), reinforcing pre-existing filters. Relatedly, Alvero et al. [2020] document that seemingly innocuous features (e.g., the word "baseball") can act as strong demographic signals, raising concerns about *indirect inference* in which protected or contextual attributes are inferred through correlated proxies. A broader survey by Baker and Hawn [2022] warns that such systems may conceal "unknown unknowns," particularly for under-examined subpopulations. Complementing these mechanism-level concerns, the meta-analytic evidence in Walton and Spencer [2009] demonstrates that outcome-based measures (grades, test scores) can systematically understate the aptitude of marginalized groups through identity-contingent

dynamics.

By elevating probability paradox from an anecdotal concern to a model-based, estimable quantity, our framework clarifies when cross-context comparisons are trustworthy, when they are fragile, and how to report that fragility responsibly. This closes the loop between principled latent measurement, realistic selection processes, and interpretable comparative claims.

Supplementary Materials

The Supplementary Materials contain proofs of Lemma 4.4 and Theorem 4.5, together with auxiliary Monte Carlo simulations provided solely for numerical intuition for Theorem 4.3.

Mathematical Appendix

This appendix contains proofs of mathematical results in Section 3 and the general bounds in Section 4.

Proof of Lemma 3.2. Define $f_{1,j}(\cdot) = F^{-1}_{Y_{1,(j)}} \circ F_{X_{1,(r_{1,j})}}(\cdot)$ and $f_{2,k}(\cdot) = F^{-1}_{Y_{2,(k)}} \circ F_{X_{2,(r_{2,k})}}(\cdot)$. Then we have

$$\begin{aligned}
&\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right) \\
&\quad = \mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}), f_{1,j}(X_{1,(r_{1,j})}) < f_{2,k}(X_{2,(r_{2,k})})\right) \\
&\quad = \mathrm{P}\left(X_{2,(r_{2,k})} < X_{1,(r_{1,j})} < h_{j,k}\left(X_{2,(r_{2,k})}\right)\right),
\end{aligned}$$

where $h_{j,k}(\cdot) = (f^{-1}_{1,j} \circ f_{2,k})(\cdot)$. A similar technique can be applied to the second equation to obtain the result. □

Proof of Theorem 3.4. First, Lemma 3.2 allows us to write

$$\mathrm{P}\left(X_{1,(r_{1,j})} > X_{2,(r_{2,k})}, Y_{1,(j)} < Y_{2,(k)}\right) = \mathrm{P}\left(X_{2,(r_{2,k})} < X_{1,(r_{1,j})} < h_{j,k}\left(X_{2,(r_{2,k})}\right)\right),$$

where $h_{j,k}(\cdot) = F^{-1}_{X_{1,(r_{1,j})}} \circ F_{Y_{1,(j)}} \circ F^{-1}_{Y_{2,(k)}} \circ F_{X_{2,(r_{2,k})}}(\cdot)$. We can rewrite $h_{j,k}(\cdot)$ into a form that will allow for specific computation involving Beta distributions. To see this, consider the distribution $F_{Y_{(j)}}$ for some $j = 1, \ldots, n_1$. The random variable $Y_{(j)} = F^{-1}_Y \circ U_{(j)}$ where $U_{(j)} \sim \mathrm{Beta}(j, n_1 - j + 1)$.

$$\begin{aligned}
&\mathrm{P}\left(X_{2,(r_{2,k})} < X_{1,(r_{1,j})} < h_{j,k}\left(X_{2,(r_{2,k})}\right)\right) \\
&= \int_{-\infty}^{+\infty} \left\{ \int_x^{F^{-1}_X \circ I^{-1}_{r_{1,j}, N_1 - r_{1,j}+1} \circ I_{j, n_1 - j+1} \circ I^{-1}_{k, n_2 - k+1} \circ I_{r_{2,k}, N_2 - r_{2,k}+1} \circ F_X(x)} f_{X_{1,(r_{1,j})}}(y)\, dy \right\} f_{X_{2,(r_{2,k})}}(x)\, dx \\
&= \int_{-\infty}^{+\infty} \left\{ \int_x^{F^{-1}_X \circ I^{-1}_{r_{1,j}, N_1 - r_{1,j}+1} \circ I_{j, n_1 - j+1} \circ I^{-1}_{k, n_2 - k+1} \circ I_{r_{2,k}, N_2 - r_{2,k}+1} \circ F_X(x)} d\, I_{r_{1,j}, N_1 - r_{1,j}+1}(F_X(y)) \right\} f_{X_{2,(r_{2,k})}}(x)\, dx
\end{aligned}$$

$$
\begin{aligned}
&= \int_{-\infty}^{+\infty} \left\{ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1} \circ I_{r_{2,k},N_2-r_{2,k}+1} \circ F_X(x) - I_{r_{1,j},N_1-r_{1,j}+1} \circ F_X(x) \right\}^+ \; f_{X_{2,(r_{2,k})}}(x)\; dx \\
&= \int_{-\infty}^{+\infty} \left\{ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1} \circ I_{r_{2,k},N_2-r_{2,k}+1} \circ F_X(x) - I_{r_{1,j},N_1-r_{1,j}+1} \circ F_X(x) \right\}^+ \; d\; I_{r_{2,k},N_2-r_{2,k}+1}(F_X(x)) \\
&= \int_0^1 \left\{ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1}(t) - I_{r_{1,j},N_1-r_{1,j}+1} \circ I^{-1}_{r_{2,k},N_2-r_{2,k}+1}(t) \right\}^+ \; dt.
\end{aligned}
$$

Condition (8) states that

$$
\begin{aligned}
&\int_0^1 \left\{ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1}(t) - I_{r_{1,j},N_1-r_{1,j}+1} \circ I^{-1}_{r_{2,k},N_2-r_{2,k}+1}(t) \right\}^+ \; dt \\
&= \int_0^1 \left\{ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1}(t) - I_{r_{1,j},N_1-r_{1,j}+1} \circ I^{-1}_{r_{2,k},N_2-r_{2,k}+1}(t) \right\} \; dt.
\end{aligned}
$$

Thus,

$$
\begin{aligned}
&\int_0^1 \left\{ I_{j,n_1-j+1} \circ I^{-1}_{k,n_2-k+1}(t) - I_{r_{1,j},N_1-r_{1,j}+1} \circ I^{-1}_{r_{2,k},N_2-r_{2,k}+1}(t) \right\}^+ \; dt \\
&\quad = \int_0^1 I_{j,n_1-j+1}(x)\; dI_{k,n_2-k+1}(x) - \int_0^1 I_{r_{1,j},N_1-r_{1,j}+1}(x)\; dI_{r_{2,k},N_2-r_{2,k}+1}(x) \\
&\quad = \int_0^1 I_x(j, n_1-j+1) \frac{x^{k-1}(1-x)^{n_2-k}}{B(k, n_2-k+1)}\; dx \\
&\qquad - \int_0^1 I_x(r_{1,j}, N_1-r_{1,j}+1) \frac{x^{r_{2,k}-1}(1-x)^{N_2-r_{2,k}}}{B(k, N_2-k+1)}\; dx \\
&\quad = \int_0^1 \sum_{p=j}^{n_1} \left(\begin{array}{c} n_1 \\ p \end{array} \right) x^p (1-x)^{n_1-p} \frac{x^{k-1}(1-x)^{n_2-k}}{B(k, n_2-k+1)}\; dx \\
&\qquad - \int_0^1 \sum_{p=r_{1,j}}^{N_1} \left(\begin{array}{c} N_1 \\ p \end{array} \right) x^p (1-x)^{N_1-p} \frac{x^{r_{2,k}-1}(1-x)^{N_2-r_{2,k}}}{B(r_{2,k}, N_2-r_{2,k}+1)}\; dx \\
&\quad = \sum_{p=j}^{n_1} \left(\begin{array}{c} n_1 \\ p \end{array} \right) \frac{B(k+p, n_2+n_1-p-k+1)}{B(j, n_2-k+1)} \\
&\qquad - \sum_{p=r_{1,j}}^{N_1} \left(\begin{array}{c} N_1 \\ p \end{array} \right) \frac{B(r_{2,k}+p, N_2+N_1-p-r_{2,k}+1)}{B(r_{2,k}, N_2-r_{2,k}+1)}.
\end{aligned}
$$

□

Proof of Proposition 3.5. Specification of $1_i(X_{i,j} \geq X_{i,(N_i-n_i+1)})$ implies that $r_{i,j} = N_i+1-j$. When either $n_1 = N_1$ and $n_2 = N_2$ or $n_1 = n_2$ and $N_1 = N_2$ we have, for all $x \in (0,1)$, that

$$
I_{N_2-j,j+1} \circ I^{-1}_{N_1-j,j+1} \circ I_{n_1-j,j+1} \circ I^{-1}_{n_2-j,j+1}(t) \geq t.
$$

The left hand side of the above equation increases whenever n_1 or N_1 are increased [Askitis, 2021], or n_2 or N_1 are decreased. Therefore, $I_{N_2-j,j+1} \circ I^{-1}_{N_1-j,j+1} \circ I_{n_1-j,j+1} \circ I^{-1}_{n_2-j,j+1}(t) \geq t$

whenever $\frac{n_1}{n_2} \leq 1$ and $\frac{N_2}{N_1} \leq 1$. Observe that

$$\begin{aligned} & I_{N_2-j,j+1} \circ I^{-1}_{N_1-j,j+1} \circ I_{n_1-j,j+1} \circ I^{-1}_{n_2-j,j+1}(t) \geq t, \\ & \quad \implies I_{n_1-j,j+1} \circ I^{-1}_{n_2-j,j+1}(t) \geq I_{N_1-j,j+1} \circ I^{-1}_{N_2-j,j+1}(t), \\ & \quad \implies I_{n_1-j,j+1} \circ I^{-1}_{n_2-j,j+1}(x) - I_{N_1-j,j+1} \circ I^{-1}_{N_2-j,j+1}(t) \geq t. \end{aligned}$$

Technical condition (8) holds and the conclusion of Theorem 3.4 applies. □

Proof of Proposition 4.2. First, in combination with the Definition 4.1 and the assumption $\frac{n_1}{N_1} \leq \frac{n_2}{N_2}$, we have

$$F_{Y_1}(x)^{\frac{n_1}{N_1}} \geq F_{Y_2}(x)^{\frac{n_2}{N_2}} \implies x \geq F^{-1}_{Y_1}\left(F_{Y_2}(x)^{\frac{n_2 N_1}{N_2 n_1}}\right).$$

By applying the Equation 7 from the Lemma 3.2 we have

$$\begin{aligned} & \mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \mathrm{P}\left(g_{n_1,n_2}\left(Y_{2,(n_2)}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\ & = \mathrm{P}\left(F^{-1}_{Y_1}\left(F_{Y_2}(Y_{2,(n_2)})^{\frac{n_2 N_1}{N_2 n_1}}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right) \\ & = \int_{-\infty}^{+\infty} \int_{F^{-1}_{Y_1}\left(F_{Y_2}(x)^{\frac{n_2 N_1}{N_2 n_1}}\right)}^{x} dF_{Y_1,(n_1)}(y) dF_{Y_2,(n_2)}(x) \\ & = \int_{-\infty}^{+\infty} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2 N_1}{N_2}} dF_{Y_2,(n_2)}(x) \\ & = \int_0^1 F_{Y_1}\left(F^{-1}_{Y_2}(t^{\frac{1}{n_2}})\right)^{n_1} - t^{\frac{N_1}{N_2}} dt. \end{aligned}$$

Based on the Definition 4.1, we have

$$\begin{aligned} & \mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \int_0^1 F_{Y_1}\left(F^{-1}_{Y_2}(t^{\frac{1}{n_2}})\right)^{n_1} - t^{\frac{N_1}{N_2}} dt \\ & \geq \int_0^1 F_{Y_2}\left(F^{-1}_{Y_2}(t^{\frac{1}{n_2}})\right)^{n_1} - t^{\frac{N_1}{N_2}} dt \\ & = \frac{n_2}{n_1+n_2} - \frac{N_2}{N_1+N_2}, \end{aligned}$$

and

$$\begin{aligned} & \mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \int_0^1 F_{Y_1}\left(F^{-1}_{Y_2}(t^{\frac{1}{n_2}})\right)^{n_1} - t^{\frac{N_1}{N_2}} dt \\ & \leq \int_0^1 \left(t^{1/n_2} + \varepsilon\right)^{n_1} - t^{\frac{N_1}{N_2}} dt \\ & = n_2 \sum_{i=0}^{n_1} \frac{\binom{n_1}{i}\varepsilon^i}{n_1+n_2-i} - \frac{N_2}{N_1+N_2}. \end{aligned}$$

□

Proof of Theorem 4.3. Equation 7 from Lemma 3.2 yields

$$g_{n_1,n_2}(x) = F_{Y_1}^{-1}\left(F_{Y_2}(x)^{\frac{n_2N_1}{N_2n_1}}\right) = \Phi^{-1}\left(\Phi\left(\frac{x-\mu_2}{\sigma_2}\right)^{\frac{n_2N_1}{N_2n_1}}\right)\sigma_1 + \mu_1.$$

Let A and R be as defined, respectively, in (11) and (13). We have,

$$\begin{aligned}
&\mathrm{P}\left(X_{1,(N_1)} > X_{2,(N_2)}, Y_{1,(n_1)} < Y_{2,(n_2)}\right) = \mathrm{P}\left(g_{n_1,n_2}\left(Y_{2,(n_2)}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right)\\
&\quad= \mathrm{P}\left(F_{Y_1}^{-1}\left(F_{Y_2}(Y_{2,(n_2)})^{\frac{n_2N_1}{N_2n_1}}\right) < Y_{1,(n_1)} < Y_{2,(n_2)}\right)\\
&\quad= \int_A \int_{F_{Y_1}^{-1}\left(F_{Y_2}(x)^{\frac{n_2N_1}{N_2n_1}}\right)}^{x} dF_{Y_1,(n_1)}(y)dF_{Y_2,(n_2)}(x)\\
&\quad= \int_A F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}}\, dF_{Y_2,(n_2)}(x)\\
&\quad= \int_{-\infty}^{\infty} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}}\, dF_{Y_2,(n_2)}(x) - \int_{A^c} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}}\, dF_{Y_2,(n_2)}(x)\\
&\quad= \int_{-\infty}^{\infty} F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}}\, dF_{Y_2,(n_2)}(x) + R.
\end{aligned}$$

We continue with the left-hand side of the last equality,

$$\begin{aligned}
&\int_{-\infty}^{\infty}\left[F_{Y_1}(x)^{n_1} - F_{Y_2}(x)^{\frac{n_2N_1}{N_2}}\right] dF_{Y_2,(n_2)}(x) = \int_0^1 \left[F_{Y_1}\left(F_{Y_2}^{-1}(t^{1/n_2})\right)^{n_1} - t^{N_1/N_2}\right] dt\\
&\quad= \int_0^1 \left(\Phi\left(\frac{\Phi^{-1}(t^{1/n_2})\sigma_2 + \mu_2 - \mu_1}{\sigma_1}\right)\right)^{n_1} dt \;-\; \frac{N_2}{N_1+N_2}\\
&\quad= \int_{-\infty}^{+\infty} \Phi\left(\frac{\sigma_2 x + \mu_2 - \mu_1}{\sigma_1}\right)^{n_1} \left(n_2\Phi(x)^{n_2-1}\phi(x)\right)\, dx \;-\; \frac{N_2}{N_1+N_2}\\
&\quad= n_2\int_{-\infty}^{+\infty} \mathbb{P}\left(\begin{array}{l} Z_{1,1} - \frac{\sigma_2}{\sigma_1}X \le \frac{\mu_2-\mu_1}{\sigma_1}, \ldots, Z_{1,n_1} - \frac{\sigma_2}{\sigma_1}X \le \frac{\mu_2-\mu_1}{\sigma_1},\\ Z_{2,1} - X \le 0, \ldots, Z_{2,n_2-1} - X \le 0 \end{array} \;\middle|\; X = x\right)\phi(x)\, dx \;-\; \frac{N_2}{N_1+N_2}\\
&\quad= n_2\,\mathbb{P}\left(\begin{array}{l} Z_{1,1} - \frac{\sigma_2}{\sigma_1}X \le \frac{\mu_2-\mu_1}{\sigma_1}, \ldots, Z_{1,n_1} - \frac{\sigma_2}{\sigma_1}X \le \frac{\mu_2-\mu_1}{\sigma_1},\\ Z_{2,1} - X \le 0, \ldots, Z_{2,n_2-1} - X \le 0 \end{array}\right) \;-\; \frac{N_2}{N_1+N_2}\\
&\quad= n_2\, F_{\mathbf{W}}\left(\mathbf{w} = (\frac{\mu_2-\mu_1}{\sigma_1}, \ldots, \frac{\mu_2-\mu_1}{\sigma_1}, 0, \ldots, 0)\right) \;-\; \frac{N_2}{N_1+N_2}.
\end{aligned}$$

The distribution function $F_{\mathbf{W}}$ corresponds to random variable $\mathbf{W} \sim \mathcal{N}_{(n_1+n_2-1)}(\mathbf{0}, \mathbf{\Sigma})$ where $\mathbf{\Sigma}$ is as defined in (14) and (15). The first n_1 elements of $\mathbf{w}$ is $\frac{\mu_2-\mu_1}{\sigma_1}$ and the last $n_2 - 1$ elements are 0.

The bound on the remainder (17) follows from the argument in Section 4.2. □

References

AJ Alvero, Noah Arthurs, Anthony Lising Antonio, Benjamin W Domingue, Ben Gebre-Medhin, Sonia Giebel, and Mitchell L Stevens. Ai and holistic review: informing human reading in college admissions. In *Proceedings of the AAAI/ACM Conference on AI, Ethics, and Society*, pages 200–206, 2020.

Dimitris Askitis. Logarithmic concavity of the inverse incomplete beta function with respect to the first parameter. *Dimitris Askitis*, 2021.

Ryan S Baker and Aaron Hawn. Algorithmic bias in education. *International Journal of Artificial Intelligence in Education*, pages 1–41, 2022.

David J Bartholomew, Martin Knott, and Irini Moustaki. *Latent variable models and factor analysis: A unified approach*. John Wiley & Sons, 2011.

Benjamin S Baumer and Gregory J Matthews. A statistician reads the sports page: there is no avoiding WAR. *Chance*, 27(3):41–44, 2014.

Benjamin S Baumer, Shane T Jensen, and Gregory J Matthews. openwar: An open source system for evaluating overall player performance in major league baseball. *Journal of Quantitative Analysis in Sports*, 11(2):69–84, 2015.

Scott M Berry, C Shane Reese, and Patrick D Larkey. Bridging different eras in sports. *Journal of the American Statistical Association*, 94(447):661–676, 1999.

Trevor G Bond and Christine M Fox. *Applying the Rasch model: Fundamental measurement in the human sciences*. Psychology Press, 2013.

Ralph A. Bradley. Science, statistics, and paired comparisons. *Biometrics*, 32(2):213–239, 1976. doi: 10.2307/2529494.

Ralph A. Bradley. Paired comparisons: Some basic procedures and examples. In P. R. Krishnaiah and P. K. Sen, editors, *Handbook of Statistics*, volume 4, pages 299–326. Elsevier, Amsterdam, 1984.

Ralph A. Bradley and Milton E. Terry. The rank analysis of incomplete block designs: I. the method of paired comparisons. *Biometrika*, 39:324–345, 1952. doi: 10.1093/biomet/39.3-4.324.

Anthony Castrovince. All of your questions about the new baseball rules, answered. *MLB.com*, February 2023. URL https://www.mlb.com/news/mlb-new-rules-for-2023-faq.

Wei Chu and Zoubin Ghahramani. Preference learning with gaussian processes. In *Proceedings of the 22nd International Conference on Machine Learning*, pages 137–144. ACM, 2005. doi: 10.1145/1102351.1102369.

Herbert A. David. *The Method of Paired Comparisons*. Chapman and Hall, London, 2 edition, 1988.

Daniel Eck, Jun Hee Kim, Shen Yan, Muskan Bhatla, and Zheer Wang. *fullhouse: Package for Estimating Talent and Era-Adjusted Performance Metrics*, 2025. URL https://github.com/DEck13/fullhouse. R package version 1.1.

Bradley Efron and Carl Morris. Data analysis using stein's estimator and its generalizations. *Journal of the American Statistical Association*, 70(350):311–319, 1975.

Arpad E. Elo. *The Rating of Chessplayers, Past and Present*. Arco Publishing, 1978.

Susan E Embretson and Steven P Reise. *Item response theory for psychologists*. Psychology Press, 2000.

Gene H Golub and Charles F Van Loan. *Matrix computations*. Johns Hopkins University press, 2013.

Thore Graepel, Tom Minka, and R TrueSkill Herbrich. A bayesian skill rating system. *Advances in Neural Information Processing Systems*, 19(569-576):7, 2007.

Jacques A Hagenaars and Allan L McCutcheon. *Applied latent class analysis*. Cambridge University Press, 2002.

Paul W Holland and Howard Wainer. *Differential item functioning*. Routledge, 2012.

Neil Houlsby, Ferenc Huszar, Zoubin Ghahramani, and Jose Hernández-lobato. Collaborative gaussian processes for preference learning. *Advances in neural information processing systems*, 25, 2012.

Guido Imbens and Konrad Menzel. A causal bootstrap. Technical report, National Bureau of Economic Research, 2018.

Maurice G Kendall. A new measure of rank correlation. *Biometrika*, 30(1-2):81–93, 1938.

Jouni Kerman. A closed-form approximation for the median of the beta distribution. *arXiv preprint arXiv:1111.0433*, 2011.

Frederic M Lord. A paradox in the interpretation of group comparisons. *Psychological bulletin*, 68(5):304, 1967.

Roderick P McDonald. *Test theory: A unified treatment*. psychology press, 1999.

Stephen Nickell and Richard Layard. Labor market institutions and economic performance. *Handbook of labor economics*, 3:3029–3084, 1999.

Stephen Raudenbush and Anthony S Bryk. A hierarchical model for studying school effects. *Sociology of education*, pages 1–17, 1986.

Michael J Schell. *Baseball's all-time best hitters*. Princeton University Press, 2013.

Michael J Schell. *Baseball's all-time best sluggers*. Princeton University Press, 2016.

Edward H Simpson. The interpretation of interaction in contingency tables. *Journal of the Royal Statistical Society: Series B (Methodological)*, 13(2):238–241, 1951.

Robert H Somers. A new asymmetric measure of association for ordinal variables. *American sociological review*, pages 799–811, 1962.

Gregory M Walton and Steven J Spencer. Latent ability: Grades and test scores systematically underestimate the intellectual ability of negatively stereotyped students. *Psychological science*, 20(9):1132–1139, 2009.

Austin Waters and Risto Miikkulainen. Grade: Machine learning support for graduate admissions. *Ai Magazine*, 35(1):64–64, 2014.

Wayne L Winston, Scott Nestler, and Konstantinos Pelechrinis. *Mathletics: How Gamblers, Managers, and Fans use Mathematics in Sports*. Princeton University Press, 2022.

Shen Yan, Adrian Burgos, Jr, Christopher Kinson, and Daniel J Eck. Comparing baseball players across eras via novel full house modeling. *The Annals of Applied Statistics*, 19(2): 1778–1799, 2025.

Amy Zhao Yu, Shahar Ronen, Kevin Hu, Tiffany Lu, and César A Hidalgo. Pantheon 1.0, a manually verified dataset of globally famous biographies. *Scientific data*, 3(1):1–16, 2016.