

Testing for the prevalence or transience of a shock effect in the post-shock setting

Abstract

We provide a hypothesis test for the existence or transience of a shock in the post-shock setting of [Lin and Eck \[2021\]](#).

1 Setting

1.1 Model setup

The assumed autoregressive models considered are that in [Lin and Eck \[2021\]](#), we describe details here. Let $I(\cdot)$ be an indicator function, T_i be the time length of the time series i for $i = 1, \dots, n+1$, and T_i^* be the time point just before the one when the shock is known to occur, with $T_i^* < T_i$. For $t = 1, \dots, T_i$ and $i = 1, \dots, n+1$, the model \mathcal{M}_1 is defined as

$$\mathcal{M}_1: y_{i,t} = \eta_i + \alpha_i D_{i,t} + \phi_i y_{i,t-1} + \theta_i' \mathbf{x}_{i,t} + \varepsilon_{i,t} \quad (1)$$

where $D_{i,t} = I(t = T_i^* + 1)$ and $\mathbf{x}_{i,t} \in \mathbb{R}^p$ with $p \geq 1$. We assume that the $\mathbf{x}_{i,t}$ s are fixed. Let $|x|$ denote the absolute value of x for $x \in \mathbb{R}$. For $i = 1, \dots, n+1$ and $t = 1, \dots, T_i$, the random effects structure for \mathcal{M}_1 is:

$$\begin{aligned} \eta_i &\stackrel{iid}{\sim} \mathcal{F}_\eta \text{ with } E_{\mathcal{F}_\eta}(\eta_i) = 0, \text{Var}_{\mathcal{F}_\eta}(\eta_i) = \sigma_\eta^2 \\ \phi_i &\stackrel{iid}{\sim} \mathcal{F}_\phi \text{ where } |\mathcal{F}_\phi| < 1, \\ \theta_i &\stackrel{iid}{\sim} \mathcal{F}_\theta \text{ with } E_{\mathcal{F}_\theta}(\theta_i) = \mu_\theta, \text{Var}_{\mathcal{F}_\theta}(\theta_i) = \Sigma_\theta^2 \\ \alpha_i &\stackrel{iid}{\sim} \mathcal{F}_\alpha \text{ with } E_{\mathcal{F}_\alpha}(\alpha_i) = \mu_\alpha, \text{Var}_{\mathcal{F}_\alpha}(\alpha_i) = \sigma_\alpha^2 \\ \varepsilon_{i,t} &\stackrel{iid}{\sim} \mathcal{F}_{\varepsilon_i} \text{ with } E_{\mathcal{F}_{\varepsilon_i}}(\varepsilon_{i,t}) = 0, \text{Var}_{\mathcal{F}_{\varepsilon_i}}(\varepsilon_{i,t}) = \sigma_i^2, \\ \eta_i &\perp\!\!\!\perp \alpha_i \perp\!\!\!\perp \phi_i \perp\!\!\!\perp \theta_i \perp\!\!\!\perp \varepsilon_{i,t}, \end{aligned}$$

where $\perp\!\!\!\perp$ denotes the independence operator. We also consider a modeling framework where the shock effects are linear functions of covariates with an additional additive mean-zero error. For $i = 1, \dots, n+1$, the random effects structure for this model (model \mathcal{M}_2) is:

$$\begin{aligned} \mathcal{M}_2: \quad y_{i,t} &= \eta_i + \alpha_i D_{i,t} + \phi_i y_{i,t-1} + \theta_i' \mathbf{x}_{i,t} + \varepsilon_{i,t} \\ \alpha_i &= \mu_\alpha + \delta_i' \mathbf{x}_{i,T_i^*+1} + \tilde{\varepsilon}_i, \end{aligned} \quad (2)$$

where the added random effects are

$$\begin{aligned} \tilde{\varepsilon}_i &\stackrel{iid}{\sim} \mathcal{F}_{\tilde{\varepsilon}} \text{ with } E_{\mathcal{F}_{\tilde{\varepsilon}}}(\tilde{\varepsilon}_i) = 0, \text{Var}_{\mathcal{F}_{\tilde{\varepsilon}}}(\tilde{\varepsilon}_i) = \sigma_{\tilde{\varepsilon}}^2 \\ \eta_i &\perp\!\!\!\perp \alpha_i \perp\!\!\!\perp \phi_i \perp\!\!\!\perp \theta_i \perp\!\!\!\perp \varepsilon_{i,t} \perp\!\!\!\perp \tilde{\varepsilon}_i. \end{aligned}$$

We further define $\tilde{\alpha}_i = \mu_\alpha + \delta_i' \mathbf{x}_{i,T_i^*+1}$. We will investigate the post-shock aggregated estimators in \mathcal{M}_2 in settings where δ_i is either fixed or random. We let \mathcal{M}_{21} denote model \mathcal{M}_2 with $\delta_i = \delta$ for $i = 1, \dots, n+1$, where δ is a fixed unknown parameter. We let \mathcal{M}_{22} denote model \mathcal{M}_2 with the following random effects :

$$\begin{aligned} \delta_i &\stackrel{iid}{\sim} \mathcal{F}_\delta \text{ with } E_{\mathcal{F}_\delta}(\delta_i) = \mu_\delta, \text{Var}_{\mathcal{F}_\delta}(\delta_i) = \Sigma_\delta \\ \delta_i &\perp\!\!\!\perp \tilde{\varepsilon}_i. \end{aligned}$$

We further define the parameter sets

$$\begin{aligned} \Theta &= \{(\eta_i, \phi_i, \theta_i, \alpha_i, \mathbf{x}_{i,t}, y_{i,t-1}, \delta_i) : t = 1, \dots, T_i, i = 2, \dots, n+1\} \\ \Theta_1 &= \{(\eta_i, \phi_i, \theta_i, \alpha_i, \mathbf{x}_{i,t}, y_{i,t-1}, \delta_i) : t = 1, \dots, T_i, i = 1\} \end{aligned} \quad (3)$$

where Θ and Θ_1 can adapt to \mathcal{M}_1 by dropping δ_i . We assume this for notational simplicity.

1.2 Forecast

In our post-shock setting we consider the following candidate forecasts:

$$\begin{aligned} \text{Forecast 1 : } \hat{y}_{1,T_1^*+1}^1 &= \hat{\eta}_1 + \hat{\phi}_1 y_{1,T_1^*} + \hat{\theta}_1' \mathbf{x}_{1,T_1^*+1}, \\ \text{Forecast 2 : } \hat{y}_{1,T_1^*+1}^2 &= \hat{\eta}_1 + \hat{\phi}_1 y_{1,T_1^*} + \hat{\theta}_1' \mathbf{x}_{1,T_1^*+1} + \hat{\alpha}, \end{aligned}$$

where $\hat{\eta}_1$, $\hat{\phi}_1$, and $\hat{\theta}_1$ are all OLS estimators of η_1 , ϕ_1 , and θ_1 , respectively, and $\hat{\alpha}$ is some form of estimator for the shock effect of time series of interest, i.e., α_1 .

Throughout the rest of this article we highlight when the information from the time series donor pool, indexed by $\{y_{i,t} : t = 1, \dots, T_i, i = 2, \dots, n+1\}$, can be used to construct a shock effect estimator $\hat{\alpha}$ in which Forecast 2 beats Forecast 1 and vice-versa. We will consider the different dynamic panel models \mathcal{M}_1 , \mathcal{M}_{21} , and \mathcal{M}_{22} . We want to determine which forecast is appropriate over a horizon while the methods in [Lin and Eck \[2021\]](#) were only appropriate in the nowcasting setting in which prediction was only focused on the response immediately following the shock.

1.3 Multi-horizon forecast evaluation

We now consider an approach for comparing the forecasts over a specified time horizon after a shock has occurred by borrowing information across similar forecasts.

[Quaedvlieg \[2021\]](#) provided a methodology for comparing forecasts jointly across all horizons of a forecast path, $h = 1, \dots, H$. In our post-shock setting, we want to compare the forecasts

$$\hat{y}_{1,t}^{1,h} \text{ and } \hat{y}_{1,t}^{2,h}$$

where $\hat{y}_{1,t}^{1,h}$ is the forecast for y_t that accounts for the yet-to-be observed structural shock and is based on the information set \mathcal{F}_{t-h} , and $\hat{y}_{1,t}^{2,h}$ is defined similarly for the forecast that does not include any shock effect information. We will compare these forecasts in terms of their loss differential

$$d_t = L_{1,t} - L_{2,t},$$

where $L_{j,t} = L(y_t, \hat{y}_{j,t})$, $j = 1, 2$, and L is a loss function. Hypothesis tests in [Quaedvlieg \[2021\]](#) are with respect to $E(d_t) = \mu_t$. Conditions for these tests require conditions of [Giacomini and White \[2006\]](#).

Note: We need more formality for constructing $\hat{y}_{1,t}^{1,h}$. We could use the forecasts in [Lin and Eck \[2021\]](#) and then consider h -ahead methods after adjusting for the shock. Or we could consider aggregation approaches which average all post-shock responses of the series in the donor pool.

We will consider the average superior predictive ability (aSPA) to assess whether or not a shock is permanent or transitory. The aSPA investigates forecast comparisons based on their weighted average loss difference

$$\mu^{(AVG)} = \mathbf{w}^T \mu = \sum_{h=1}^H w_h \mu^h$$

with weights \mathbf{w} that sum to one. Note that aSPA requires the user to take a stand on the relative importance of under-performance at one horizon against out-performance at another, and note that it is likely that $\mu^h > 0$ for h closer to 1 since the user expects that a structural shock will occur and the structural shock is taken into account by forecast 1.

References

- Raffaella Giacomini and Halbert White. Tests of conditional predictive ability. *Econometrica*, 74(6): 1545–1578, 2006.
- Jilei Lin and Daniel J Eck. Minimizing post-shock forecasting error through aggregation of outside information. *International Journal of Forecasting*, 2021. Accepted.
- Rogier Quaadvlieg. Multi-horizon forecast comparison. *Journal of Business & Economic Statistics*, 39 (1):40–53, 2021.