Narcissus and Covenant

One Half, Two Ways



A chapter for *HexOct: The Hexagonality Law*. Theorem: the HexOct h-tile's piece at the end of the walk, Narcissus, is its own mirror image and fills space alone; two of them joined on their hexagons make the twisted cube when put on straight and Covenant, a parallelogram dodecahedron that fills space by the fcc lattice, when turned 60°; and the Dehn invariants of all HexOct pieces are exact linear functions of the stretch.

DICTO (artist, Osaka, Japan) · contact@dictospheres.com · https://dictospheres.com
4 November 2026 · View in app: Polyhedraverse, DICTO's pieces (Narcissus; Narcissus and cubes; RD/TRD half-cell with ⟳ Turn) · Licence CC BY-SA 4.0 · Scripts and data: research studies 2026-11-04-dehn-screen, 2026-11-04-halves, 2026-11-04-lqg-bridge.
The ideas are DICTO's: that the h-tile's piece is one piece, not a pair (and its name, Narcissus); that it might be "halved" and turned; the name Covenant; the RD/TRD half that can be put on straight or twisted, in a two-colour livery. Computation, proofs and drafting with Claude (Anthropic).

DRAFT for DICTO's reading. Not published. Chapter for the next version of *HexOct: The Hexagonality Law*. Delete this box at publication. Open: the constructive search whether the h-piece tiles alone for $5/24 \le \lambda < 3/8$ (both necessary tests pass); the Narcissus–cube common-dissection search (running); prior-art search done for Covenant (type classical: Fedorov, Grünbaum) and Narcissus (type very likely in Goldberg 1978, to be read); the banded twisted RD still to search. Figures 1–4 are working renders, to be redrawn for print.

Abstract. At the end of the HexOct walk (#75, $\lambda \ge 3/8$) the octahedral pieces vanish and each layer tile is two copies of one piece: a cube in a c layer, and in an h layer a ten-cornered piece we call Narcissus. Narcissus is its own mirror image (three mirror planes), so copies of one shape fill space. Two copies joined on their hexagonal faces give two solids. Put on straight they make the twisted cube of #86; turned by 60° they make Covenant, a convex dodecahedron of twelve parallelograms, which is a parallelohedron and fills space by translation along the fcc lattice. The same law of two ways, straight or twisted, one free choice per layer, holds for the classical half of the rhombic dodecahedron (whose two ways are the RD and the TRD), as it holds for the octet pyramids in #92. We also give the Dehn invariants of every HexOct piece exactly: $36(5/24 - \lambda) \otimes \beta$ for the h pieces (zero from $\lambda = 5/24$ on) and $24(3/8 - \lambda) \otimes \beta$ for the c pieces, with $\beta = \arccos(1/\sqrt{3})$.

Cut a solid in half, and you can put the halves back two ways: as they were, or with one turned. For most solids the turned version has a dent. For a few it does not, and for fewer still it fills space. This chapter is about the few.




Figure 1. Narcissus: a regular hexagon on top (gold), three quadrilaterals (purple) and six triangles (teal) running down to one corner on a ball one layer below. Ten corners; volume √2/4, half the volume per ball.

1. Main theorem

Theorem. Let N (Narcissus) be the tetrahedral piece of the HexOct h-tile for λ ≥ 3/8, and C the cube of edge 1/√2 (ball spacing 1). (a) N has the symmetry C_{3v}; in particular it is its own mirror image. The h-tile is two copies of N, so copies of N alone fill space. (b) Two copies of N joined on their hexagonal faces, the upper one turned over: put on straight they form the twisted cube of #86 (11 corners, 6 squares and 6 triangles); turned by 60° about the hexagon's centre they form **Covenant**, a convex dodecahedron with 12 parallelogram faces and 14 corners. Covenant is the zonohedron of three orthogonal segments of length 1/√2 and one along their body diagonal of a third of its length; it fills space by translation along the fcc lattice, one Covenant per ball. (c) Every stacking of c and h layers is filled by copies of N (h layers) and C (c layers). (d) The Dehn invariant of every piece of the HexOct tiles is exactly (s ⊗ β) with β = arccos(1/√3), where s = 36(5/24 − λ) for the h pieces with λ < 5/24, s = 0 for the h pieces with λ ≥ 5/24, and s = 24(3/8 − λ) for the c pieces (tetrahedral pieces with the opposite sign of the octahedral ones).

> *One self-mirroring piece, joined to itself two ways, gives two different space-fillers; and the stretch at which the HexOct pieces could start to fill space on their own is fixed by an exact straight line.*

2. Narcissus

At λ = 3/8 the octahedral pieces of #75 shrink to nothing; for every larger λ the h-tile is the same two pieces (the walk saturates). Their corners are: a regular hexagon of side 1/√3 at the height of the next layer (three ball centres of that layer and three holes between them), three corners half a layer lower under the three holes, and one corner on the ball below the hexagon's centre. The faces are one hexagon, three quadrilaterals and six triangles; the dihedral angles are 90°, 120° and 135° only; the volume is √2/4 and the Mahler product exactly 34/3 (rational affine image). Three mirror planes contain its axis; the two copies in an

h-tile are the same piece, one turned over. Copies of one piece therefore fill space (Theorem (a)). Within HexOct, Narcissus fills only the h layers: in a c layer the piece is the cube (Theorem (c), checked over all 32 five-layer contexts at λ = 3/8 and 2/5). The every-stacking statement is #86's, read with its tiles halved: two Narcissus make the twisted cube and two cubes the brick of #86's pair A.




Figure 2. Two Narcissus (gold, purple) on their hexagons. Straight: the twisted cube. Turned 60°: Covenant, whose faces line up across the seam into twelve parallelograms.

3. Covenant


Turned by 60°, the union of two Narcissus is convex and centrally symmetric, and all twelve faces are parallelograms. Its edges run in four directions: three mutually orthogonal of length 1/√2 (the small fcc cube's edges) and one along the cube's body diagonal of a third of the diagonal's length. Every convex zonohedron of the five parallelohedron types fills space by translation (Fedorov; see Grünbaum [G]); Covenant's six face-to-face translations all have length 1 and generate the fcc lattice (twelve nearest neighbours at 1, six at √2). In the family of zonohedra of a cube's three edges and c times its body diagonal, c = 0 is the cube, c = 1 the rhombic dodecahedron; Covenant is c = 1/3. Its faces are not rhombi (edges 1/√2 and 1/√6), so it is none of the isozonohedra Grünbaum lists. Caution: in Covenant's tiling only the lower Narcissus of each pair has its corner on a ball; it is an fcc fundamental domain, not a ball-by-ball c-layer cell.

4. The same law for the rhombic dodecahedron

Cut across its three-fold axis through the centre, the rhombic dodecahedron falls into two congruent halves with a regular hexagon as cut, and the trapezo-rhombic dodecahedron (TRD) is made of the very same half: put back straight, the halves make the RD; one turned by 60°, the TRD (classical: Coxeter; the TRD is the RD with one half turned). Read as a standing option, the twist is a free choice per layer and never per cell: twisting one cell's half in a packing leaves gaps and overlaps, and a whole layer twists only if the layers above re-seat over the other holes; with each ball's halves turned to face its actual neighbours, every stacking tested (fcc, hcp, ABCB, ABCACB and the Fibonacci approximant ABACBCAC) is filled with no overlap. This is the law of #92 (one straight-or-twisted choice per layer, for the octet pyramids) for a second set of pieces. In Polyhedraverse, the half wears a two-colour seam: alternating colours across the seam mean the RD, matching colours the TRD (Figure 3).

Which solids can be halved and turned. A band (a prism) put between the two halves keeps every turn convex exactly when the central cut equals the solid's shadow along the

axis. The RD satisfies this on its 2-, 3- and 4-fold axes; the truncated octahedron and the cube only on their 2- and 4-fold axes (cut/shadow 0.84 and 0.75 on the 3-fold axis, which is why the truncated octahedron's turned halves are dented). Only a hexagonal cut gives a genuine choice in a tiling, so the banded RD (half, hexagonal prism, half; straight or turned) is a family of space-fillers for every band height and every stacking. A leaning band dents at every lean; leaning a whole layer (a shear) keeps the tiling. Among the classical solids only the cuboctahedron (to J27) and the RD (to the TRD) turn convexly. The Dehn invariant is additive, so every rejoining of two Dehn-zero halves is Dehn-zero; what decides whether a turned solid can tile is the local count of dihedral angles round an edge. The turned B3 permutohedron, for instance, is Dehn-zero but fails that count (a 135° edge with no 90° partner). Halves of different solids with the same cut also rejoin: Narcissus and the RD half share their hexagon, and both of their joins (straight and turned) have only 90°, 120° and 135° dihedrals, so they pass both tests — candidates still to be decided.



◇ DICTO ◇

Figure 3. The RD/TRD half with a two-colour seam. Put on straight: gold meets gold (TRD); turned 60°: gold meets purple (RD).

5. Exact Dehn invariants of the HexOct pieces

Every dihedral angle of every HexOct piece is a rational multiple of π plus a multiple of β = arccos(1/√3): 70.53° = π − 2β, 109.47° = 2β, 125.26° = π − β, 144.74° = π/2 + β, and 90°, 120°, 135°. Squared edge lengths are rational in lattice coordinates, so each length is r√d with r rational and d squarefree. The Dehn invariant is therefore $(\Sigma\, c_e\, r_e\sqrt{d_e}) \otimes \beta$ with integer c_e, and is zero exactly when each rational sum $\Sigma\, c_e\, r_e$ over one d vanishes. Computed in exact fractions over the whole walk, only d = 1 occurs, and the sums are the straight lines of Theorem (d): 36(5/24 − λ) for the h pieces below 5/24 and 0 from 5/24 on; 24(3/8 − λ) for the c pieces. The octahedral and tetrahedral pieces of each tile carry opposite values, so every whole tile has Dehn invariant zero, as it must. The two special stretches of HexOct are the two zeros: 5/24, where the h pieces could begin to fill space alone, and 3/8, where the c pieces become cubes and the h pieces Narcissus.



Figure 4. Eight layers of a Fibonacci stacking (types c h c c h c h c) filled by Narcissus (gold, h layers) and cubes (purple, c layers); 112 pieces, no overlaps (checked).

6. What is known and what is new

Known: the RD/TRD relation and the Voronoi cells of the close packings (Coxeter; standard), the ortho/gyro operation (Johnson 1966), the space-filling of every parallelohedron-type zonohedron (Fedorov; Grünbaum), the stackings filled by #86's pair A, and the per-layer law for the octet pyramids (#92). Known in type (prior-art search, 5 November 2026): every convex zonohedron with four zones is combinatorially Kepler's rhombic dodecahedron and a parallelohedron (Fedorov 1885; Grünbaum 2010), so Covenant belongs to a classical type; in its family (the cube's three edge directions plus a fourth along a body diagonal, at a fraction c of the diagonal) the fraction c is an affine invariant, c = 1 being Kepler's dodecahedron and c = 1/3 Covenant. Narcissus, half of Covenant cut through its hexagonal waist, has seven faces and ten corners; its combinatorial type is that of half the rhombic dodecahedron and is very likely among Goldberg's space-filling heptahedra (1978; the full list not yet read). New within the sources checked: Narcissus as this particular metric heptahedron, self-mirroring and filling space alone; its two joins, the twisted cube and Covenant as the c = 1/3 member of the four-zone family; the "cut equals shadow" criterion for banded turns; and the exact Dehn lines of the HexOct pieces. Open: whether the h-piece fills space alone for $5/24 \leq \lambda < 3/8$, where both necessary conditions (Dehn zero, the edge-angle count) hold.

References

[G] B. Grünbaum, The Bilinski dodecahedron and assorted parallelohedra, zonohedra, monohedra, isozonohedra and otherhedra (2010). — E. S. Fedorov, Начала учения о фигурах (Elements of the Study of Figures), St Petersburg (1885). — M. Goldberg, On the space-filling heptahedra, Geometriae Dedicata 7 (1978) 175–184. — N. W. Johnson, Convex polyhedra with regular faces (1966). — H. S. M. Coxeter, Regular Polytopes. — DICTO, HexOct (#75), The HexOct Family (#86, draft), Aristotle's Pyramid in Layers (#92 v2).